

IMMUNOREGULATION

Volume 2. Supplement 1. April 2018

ISSN: 2588-5200

ABSTRACT BOOK



www.icia.ir





Contents

Allergy and Immunotherapy of Allergic Disease	17
Cancer Immunology	27
Cancer Immunotherapy	39
Computational Immunology and Systems Biology	47
Diagnostic Methods in Immunology.....	56
Immune Cell Therapy	64
Immunodeficiency.....	73
Immunogenetics	77
Immunohematology.....	86
Immunology of Chemical Victims and Environmental Pollution.....	93
Immunology of Exercise, Aging, and Nutrition	100
Immunology of Infectious Diseases Bacteria and Fungi	106
Immunology of Infectious Diseases Viruses	112
Immunology of Mycobacterium Tuberculosis.....	120
Immunology of Rheumatic Diseases.....	126
Immunomodulation and Immunoregulation	134
Immunoparasitology.....	142
Immunopharmacology and Medicinal Plants	153
Innate Immunity and Inflammation	164
Monoclonal Antibody Diagnostic and Therapeutic	172
Mucosal and Gastrointestinal Immunology.....	181
Nanoimmunology	193
Oral Immunology	201
Psychoneuroimmunology and Immunoendocrinology.....	210
Reproductive Immunology	216
Research and Development	228
Stem Cells Based Immunotherapy.....	239
Tolerance and Autoimmunity.....	248
Transplantation	258
Vaccine	263
Veterinary Immunology.....	272
Allergy and Immunotherapy of Allergic Disease	280
Cancer Immunology	290
Cancer Immunotherapy	304
Computational Immunology and Systems Biology	315
Diagnostic Methods in Immunology.....	321
Immune Cell Therapy.....	333
Immunodeficiency.....	339
Immunogenetics	348
Immunohematology.....	358
Immunology of Chemical Victims and Environmental Pollution.....	361
Immunology of Exercise, Aging, and Nutrition	365

Immunology of Infectious Diseases Bacteria and Fungi	369
Immunology of Infectious Diseases Viruses	380
Immunology of Mycobacterium Tuberculosis.....	391
Immunology of Rheumatic Diseases.....	400
Immunomodulation and Immunoregulation	411
Immunoparasitology.....	418
Immunopharmacology and Medicinal Plants	429
Innate Immunity and Inflammation	440
Monoclonal Antibody Diagnostic and Therapeutic	451
Mucosal and Gastrointestinal Immunology.....	462
Nanoimmunology	474
Oral Immunology	482
Psychoneuroimmunology and Immunoendocrinology.....	492
Reproductive Immunology	496
Stem Cells Based Immunotherapy	511
Tolerance and Autoimmunity.....	522
Transplantation	532
Vaccine	543
Veterinary Immunology.....	554
Allergy and Immunotherapy of Allergic Disease	565
Cancer Immunology	607
Cancer Immunotherapy	672
Computational Immunology and Systems Biology	701
Diagnostic Methods in Immunology.....	716
Immune Cell Therapy.....	719
Immunogenetics	724
Immunohematology.....	742
Immunology of Chemical Victims and Environmental Pollution.....	748
Immunology of Exercise, Aging, and Nutrition	755
Immunology of Infectious Diseases Bacteria & Fungi.....	765
Immunology of Infectious Diseases Viruses	784
Immunology of Mycobacterium Tuberculosis.....	818
Immunology of Rheumatic Diseases.....	820
Immunomodulation and Immunoregulation	837
Immunoparasitology.....	841
Immunopharmacology and Medicinal Plants	845
Innate Immunity and Inflammation	890
Monoclonal Antibody Diagnostic and Therapeutic	917
Mucosal and Gastrointestinal Immunology.....	919
Nanoimmunology	928
Psychoneuroimmunology and Immunoendocrinology.....	938
Reproductive Immunology	945
Index.....	1072






14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018 Organizers



Chairman Welcome Message



On behalf of the congress organizers, the ***Iranian Society for Immunology and Allergy (ISIA)*** and ***Shahid Beheshti University of Medical Sciences***, it is my great pleasure to invite you to the “14th International Congress of Immunology and Allergy” (ICIA2018). As the largest meeting in the field of immunology and allergy in Iran, **ICIA2018** is an event which is held biannually.

The theme of the congress is “major breakthroughs and advances in immunology” covering all fields of immunology, ranging from innate and adaptive immunity to emerging diagnostics and therapeutics.

With the participation of invited speakers and chairs from all over the world, the opportunity for discussing science and forming new collaborations is provided. Front-runners in the fields of basic, clinical and diagnostic immunology with keynote lectures will present discoveries in fundamental immunology.

We very much look forward to seeing you in Tehran on 26-28 April 2018, in which we will do our best to make **ICIA2018** the most memorable scientific event for you.

Sincerely,

Saeid Namaki, Ph.D.
Chairman of the Congress

President of ISIA

Welcome Message



As president of the “*Iranian Society for Immunology and Allergy*” (ISIA), it is my pleasure to extend this invitation to you to join us in Tehran in April 2018 at the “**14th international Congress of Immunology and Allergy**” (ICIA2018). Being held every two years and regularly attracting over 1500 delegates, **ICIA2018** will begin to work on 26th April. ISIA together with ICIA’s patrons will work to create a memorable scientific event for participants. The opportunity of exploring the most recent and innovative advances in the field of immunology and allergy from bench to bedside will motivate all attendings with different interests in almost all the fields of modern immunology. Offering a fine balance between basic and applied/translational researchers, we ensure you will totally be satisfied with this scientific session. Like all previous ICIAs, ISIA will follow its missions of fostering young and not so young researchers as scientists. I do hope that you will be able to join us at **ICIA2018** in Tehran and I look forward to seeing you there.

Sincerely,

Mohammad Vodjgani, Ph.D.

President of the Iranian Society for Immunology and Allergy (ISIA)

Head of Scientific Committee Welcome Message



We are delighted to invite you to join the ***“14th International Congress of Immunology and Allergy” (ICIA 2018)*** at Tehran, Iran, from 26 to 28 April 2018. Holding such an invaluable academic and scientific meeting provides a good opportunity for sharing ideas and experiences with the most outstanding domestic and foreign scholars to motivate students and graduates of immunology and all other related fields on their scientific path. Moreover, with the participation of scientists and experts of clinical immunology and allergy covering dimensions of diagnosis, prevention and treatment in the field of immunology along with scholars of both cellular and molecular aspects of basic immunology, practical integration of basic and clinical sciences will be provided helping share information in order to strengthen the relationship between basic and clinical sciences.

Keynote speeches of different panels, presentation of abstracts, advanced courses and workshops will form the core of **ICIA 2018**, which will be the center of attention of all scholars, experts, and students attracted by the field of immunology. Thus, **ICIA2018** strives to take advantage of valuable experiences and ideas to make this meeting more effective.

I hope that with the scientific cooperation of nationwide medical universities and research centers, companies related to immunological and laboratory products, pharmaceutical companies and other scientific organizations, **ICIA2018** becomes a memorable meeting.

Sincerely,

Seyed Mahmoud Masiha Hashemi, Ph.D.
Head of Scientific Committee of the Congress

Head of Executive Committee Welcome Message



I am delighted to warmly welcome you to attend 14th ICIA, held on April 26-28, 2018, in Tehran-Iran .The congress is organized jointly by ***Iranian Society of Immunology and Allergy (ISIA)*** and ***Shahid Beheshti University of Medical Sciences (SBMU)*** accompanied with other national and international outstanding Medical Universities and Scientific Institutes. As the Executive manager, I would like to express my sincere welcome to all of the distinguished guests and participants to **ICIA2018**.

With the purpose of bringing together leading academic scientists, researchers and scholars to exchange and share their experiences and research results on all aspects of Immunology and promoting the international communication and cooperation, it is our best pleasure to hold the **ICIA2018**, which covers various fields of immunology.

We again invite you to join us at **ICIA2018** to have memorable scientific and cultural experiences. All members of **ICIA2018** organizing committee look forward to meeting you in Tehran-Iran.

Mahdi Shabani, Ph.D.

Head of Executive Committee of the Congress

Congress Board



The poster features a blue background with a white floral pattern. At the top left is the logo of the Iranian Society for Immunology and Allergy (ISIA), which consists of a stylized column with two bird-like figures at the top. To the right of this is a large yellow circle containing the text "14th International Congress of Immunology and Allergy" and "ICIA2018" in bold, along with the dates and location. Further right is the logo of Shahid Beheshti University of Medical Sciences, a stylized geometric flower. Below these are six circular portraits of the board members, each with their name and title in yellow text. The members are arranged in two rows of three. The top row includes Saeid Namaki (Congress Chairman), Mohammad Vodjgani (President of The ISIA), and Seyed Mahmoud Masiha Hashemi (Head of Scientific Committee). The bottom row includes Mahdi Shabani (Head of Executive Committee), Mehrnaz Mesdaghi (Head of International Committee), and Tooba Ghazanfari (Head of Third Festival).

Iranian Society for Immunology and Allergy

14th International Congress of Immunology and Allergy
26-28 April 2018, Tehran, I.R.Iran
ICIA2018
چهاردهمین کنفرانس بین‌المللی ایمنولوژی و آلرژی
۲۶ تا ۲۸ بهمن ۱۳۹۶

Shahid Beheshti University of Medical Sciences

Saeid Namaki, Ph.D.
Congress Chairman

Mohammad Vodjgani, Ph.D.
President of The ISIA

Seyed Mahmoud Masiha Hashemi, Ph.D.
Head of Scientific Committee

Mahdi Shabani, Ph.D.
Head of Executive Committee

Mehrnaz Mesdaghi, MD, Ph.D.
Head of International Committee

Tooba Ghazanfari, Ph.D.
Head of Third Festival



Strategic Committee of ICIA2018

Dr. Saeed Namaki

Chairman of the Congress

Dr. Mohammad Vodjgani

President of ISIA

Dr. Seyed Mahmoud Masiha Hashemi,

Head of Scientific Committee of the Congress

Dr. Mahdi Shabani

Head of Executive Committee of the Congress

Dr. Tooba Ghazanfari

Head of Third Festival of Immunology and Allergy

Dr. Mehrnaz Mesdaghi

Head of International Committee of the Congress

Dr. Sara Soudi

Secretary of Junior Members of ISIA

Coordinator of ICIA2018 Accommodation

Dr. Reza Falak

Coordinator of ICIA2018 Workshops

Dr. Farhad Riazi Rad

Coordinator of ICIA2018 Tours

Dr. Samira Rajaei

Coordinator of ICIA2018 Accommodation

Dr. Mohammad Hosein Nicknam

Vice President of ISIA

Dr. Jamshid Hadjati

ISIA Board Member

Dr. Abbas Rezaei

ISIA Board Member

Dr. Mehdi Shekarabi

ISIA Board Member

Dr. Seyed Mohammad Moazzeni

ISIA Board Member

Dr. Reza Mansouri

ISIA Board Member

Dr. Mohammad Mehdi Mohammadi

ISIA Board Member

Scientific Committee of ICIA2018

(National)

Abbaszadegan, Mohammad Reza	Andalib, Ali Reza	Bazargan, Nasrin	Eslami, Mohammad Bagher
Abd Nikfarjam, Bahareh	Anisian, Ali	Behmanesh, Fatemeh	Eslamian, Mohammad-Hossein
Abdolmaleki, Mohsen	Arababadi, Mohammad Kazemi	Behniafard, Nasrin	Esmail, Nafiseh
Abediankenari, Saeid	Arandi, Narges	Behzad Poopak	Esmailnejad, Atefe
Abolhassani, Mohsen	Araسته, Javad	Bemarian, Mohammad Hassan	Esmailzadeh, Abdolreza
Abolnejadian, Farhad	Ardestani, Soosan	Bidad, Katayoon	Esmailzadeh, Hossein
Aghaie, Afsaneh	Arefian, Ehsan	Bozorgmehr, Mahmoud	Falak, Reza
Aghamohamadi, Asghar	Armin, Shahrnaz	Chavoshzadeh, Zahra	Fallahpour, Morteza
Aghdami, Nasser	Arshi, Saba	Cheraghi, Taher	Faridhosseini, Reza
Aghebatimaleki, Leili	Asadi Asl, Sara	Dabaghian, Mehran	Farjadian, Shirin
Aghili, Babak	Asgarin-Omran, Hossein	Dabbaghzadeh, Abbas	Farrokhi, Shokrollah
Ahanchian, Hamid	Ashoori, Elham	Darabi, Behzad	Farsad, Faraneh
Ahmadi, Kazem	Asnaashari, Ehsan	Darvishalipor, Shamila	Farzaneh, Parvaneh
Ahmadiashar, Akefeh	Assarehzadegan, Mohammad Ali	Davoodi, Homa	Fatemeh, Beyzaei
Ajami, Aboulghasem	Ataollahi, Mohammadreza	Delbandi, Ali Akbar	Fathi, Seyed Mohammad
Ajdary, Soheila	Atarod, Lida	Delirezh, Norouz	Fattahi, Javad
Alijani, Ebrahim	Athari, Seyyed Shamsadin	Doroudchi, Mehrnoosh	Fayezi, Abbas
Alimohammadian, Mohammad Hossein	Azadmehr, Abbas	Douraghi, Masoomeh	Fazeli, Bahare
Alsahebfosoul, Fereshteh	Babaie, Delara	Ebrahimi, Esmail	Fazlollahi, Mohammad Reza
Alyasin, Soheyla	Babaloo, Zohreh	Ebrahimi, Marzeih	Fereidouni, Mohammad
Amani, Davar	Badiee, Ali	Ebrahimi, Mohsen	Feyzollahzade, Sadegh
Amin, Reza	Bagheri, Nader	Ebtekar, Massoumeh	Foroughi Mehrnoush
Amini, Abbas Ali	Baharvand, Hosein	Eftekharian, Mohammad Mahdi	Foroughi, Farshad
Amirchaghmaghi, Elham	Bahrami, Ahmad	Ehsani Arkani, Mohammad Javad	Forouzan, Karimi
Amirghofran, Zahra	Bani Hashemi, Seyd Reza	Emam, Mahdi	Fotohi
Amiri, Mohammad Mehdi	Baradaran, Behzad	Entezami, Kobra	Ganjalkhani Hakemi, Mazdak
Amirzargar, Ali Akbar	Barzegari, Saeed	Erfani, Nasrollah	Ganji, Ali
Anani Sarab, Gholamreza	Bayanolhagh, Saeed	Eskandari, Nahid	Ghaderi, Abbas
			Ghadiri, Ata Allah

Ghaemi, Amir	Hosseinzadeh, Morteza	Khodadadi, Ali	Mohammadi, Mohammad Mehdi
Ghafari, Javad	Houshmand, Hamidreza	Khorrami, Samaneh	Mohammadi, Mojgan
Ghafourian Borujerdnia, Mehri	Idali, Farah	Khosravi, Afra	Mohammadi-Shahrokhi, Vahid
Ghaleh Farsa, Ghasem	Isanezhad, Amin	Khosravi, Alireza	Mohammadnia, Mousa
Ghaneie, Mostafa	Jabbari Azad, Farahzad	Kiani, Jafar	Mohammadreza, Rahmani
Gharagozloo, Marjan	Jadidiniaragh, Farhad	Kokhaei, Parviz	Mohammadzadeh, Adel
Gharagozlou, Mohammed	Jafarzadeh, Abdollah	Langroudi, Ladan	Mohammadzadeh, Iraj
Gharebaghian, Ahmad	Jalali, Amir	Mahboudi, Fereidoun	Moin, Mostafa
Gharesifard, Behrouz	Jalali, Seyed Amir	Mahdavian, Seyed Alireza	Mojtabavi, Nazanin
Ghatrehsamani, Mahdi	Jalili, Ali	Mahmoodzadeh-Niknam, Hamid	Mojtahedi, Zahra
Ghazanfari, Tooba	Jamshidian, Azam	Mahmoudi, Mahdi	Mokhtari Azad, Talaat
Ghazavi, Ali	Javad Ahmadian, Haris	Mahmoudi, Mahmoud	Mollanouri Shamsi, Mahdieh
Gholamin, Mehran	Jazayeri, Mirhadi	Majidi, Jafar	Momen, Tooba
Gholamzad, Mehrdad	Jeddi Tehrani, Mahmoud	Mansouri, Davoud	Moniri, Afshin
Gholijani, Nasser	Kafami, Laya	Mansouri, Mahboubeh	Moravej, Ali
Ghorban Khodayar	Kafshdozan, Khatere	Mansouri, Reza	Mortaz, Esameil
Givtaj, Nozar	Kalantar, Kurosh	Marashi, Mahdi	Mortazavi, Seyed Hamidreza
Golsazshirazi, Forough	Kalantari, Arash	Marjani, Majid	Mosaffa, Nariman
Gorgin, Ali	Kalantari, Tahereh	Masjedi, Mohsen	Mosayebi, Ghasem
Hadinedoushan, Hossein	Kamali-Sarvestani, Eskandar	Masoudinejad, Ali	Mostafaie, Ali
Hadjati, Jamshid	Kardor, Gholamali	Massoud, Amir Hossein	Mostafazadeh, Amrollah
Haghmorad, Dariush	Karimi, Mohammad Hossein	Mehdipour, Fereshteh	Motedayen, Mohammad Hassan
Haji Molla Hoseini, Mostafa	Kavosi, Kaveh	Mehrzad, Jalil	Mousavi Shabestari, Tahereh
Hajibehzad, Mehdi	Kawosi, Hoda	Memarian, Ali	Mousavi, Seyed Fazlullah
Hajighasemi, Fatemeh	Kazemi, Bahram	Mesdaghi, Mehrnaz	Movahedi, Masoud
Hajilooi, Mehrdad	Kazemi, Tohid	Mirahmadian, Mahroo	Nabavi, Mohammad
Hamedifar, Haleh	Keramatipour, Mohammad	Mirshafiey, Abbass	Naderi, Nadereh
Hamidieh, Amir Ali	Keshavarz, Maryam	Moaidi Esfahani, Behjat Sadat	Nadji, Seyed Alireza
Harandi, Ali	Keyhani, Abdolhossein	Moazzeni, Seyed Mohammad	Namaki, Saeed
Hashemi, Seyed Mahmoud	Khaghanzadeh, Narges	Modabber, Farrokh	Namdar Ahmadabad, Hassan
Hassan, Zuhair Mohammad	Khaki, Pejvak	Modarressi, Mohammad Hossein	Nasiri Kalmarzi, Rasoul
Heidarzadeh Arani, Marzieh	Khalili, Abbas	Mofazzal Jahromi, Mirza Ali	Namaki, Saeid
Hekmatdooft, Azita	Khansari, Nematollah	Moghbeli, Meysam	Negar, Seyed
Hojjat-Farsangi, Mohammad	Khazaei, Hossein Ali	Mohammadi, Marjan	Nejatollahi, Foroogh
Hosseinali Izad, Maryam	Kheirandish, Maryam		

Nicknam, Mohammad Hosein	Riazi Rad, Farhad	Shabani, Mahdi	Tahoori, Mohammad Taher
Nikbakht Brujeni, Gholamreza	Rostamian, Abdolrahman	Shafi Mojadadi, Mohammad	Tajbakhsh, Hassan
Nikbin, Behrouz	Rostamian, Abdolrahman	Shahabi, Shahram	Tajik, Nader
Nikoo, Shohreh	Rostaminejad, Mohammad	Shahbazi, Mehdi	Taleghani, Feryal
Nikoueinejad, Hassan	Saadat, Farshid	Shahidi, Minoo	Tavakol Afshari, Jalil
Noorbakhsh, Farshid	Saboor Yaraghi, Ali Akbar	ShahMohammadi, Shohreh	Tavakol, Marziyeh
Noorbala, Ahmad-Ali	Sabzehei, Mohammad Kazem	Shahrokhi, Somayeh	Tavakoli, Maryam
Nosratabadi, Reza	Sadeghi, Rokhsareh	Shahsavari, Farhad	Tebyanian, Majid
Nouri, Hamid Reza	Sadeghi-Shabestari, Mahnaz	Shaiegan, Mojgon	Tehrani, Mohsen
Nourizadeh, Maryam	Sadighimoghaddam, Bizhan	Shakib Jafari, Reza	Vaezadeh, Farhad
Ostadali, Mohammadreza Reza	Sadri, Homa	Shakiba, Yadollah	Varahram, Mohammad
Pakravan, Nafiseh	Saeidi, Mohsen	Shams, Ali	Varasteh, Abdolreza Reza
Pakzad, Parviz	Saeidi, Reza	Shamsdin, Seyedeh Azra	Varzi, Ali Mohammad
Parvaneh, Nima	Safa, Omid	Shamsi, Ali	Vaziri, Behrouz
Pezeshki, Mohammad	Safari, Elahe	Shariat, Mansoureh	Vodjgani, Mohammad
Pishdadian, Abbas	Safari, Mojgan	Sharifi, Nourieh	Yadegarynia, Davood
Pourfathollah, Ali Akbar	Saghafi, Shiva	Sharifpaghaleh, Ehsan	Yaraee, Roya
Pourpak, Zahra	Sahebi, Leyla	Sharifzadeh, Sedigheh	Yari, Fatemeh
Poyan Mehr, Mehrdad	Sahraeian, Mohammad Ali	Sheikhi, Abdolkarim	Yazdani, Reza
Rabbani, Mohammad	Salari, Farhad	Sheikhian, Ali	Yazdani, Yaghoob
Rafati, Sima	Salehi, Eisa	Shekarabi, Mehdi	Yeganeh, Farshid
Rafatpanah, Hoshang	Salekmoghadam, Alireza	Shirzad, Hedayatollah	Yosefi, Bahman
Rafiei Tehrani, Shahanaz	Salimi, Vahid	Shokohi, Shervin	Younesi, Vahid
Rafiei, Alireza Reza	Salimian, Jafar	Shokrgozar, Mohammad Ali	Yousefi, Mehdi
Rahbarizadeh, Fatemeh	Samadi, Morteza	Shokri, Fazel	Zahed, Ghazal
Rahimi, Roghayeh	Samiei, Afshin	Mohammad Ali Shokrgozar	Zahra Faghih
Rajaei, Samira	Samira Mohammadi-Yeganeh	Soleiman jahi, Horie	Zamani, Alireza
Rasaie, Mohammad Javad	Sandoghchian Shotorbani, Siamak	Soleimanifar, Narges	Zamani, Mohammad Ali
Rashno, Mohammad	Sankian, Mojtaba	Solgi, Ghasem	Zamani, Zahra
Razavi, Seyed Alireza	Sardari, Soroush	Soroush, Mohammadreza	Zandieh, Fariborz
Razmkhah, Mahboobeh	Sarrafnejad, Abdolfattah	Soudi, Sara	Zarnani, Amir-Hassan
Rekabi, Mahsa	Sattari, Mandana	Tabaresi, Payam	Zavaranhosseini, Ahmad
Rezaei, Abbas	Sedghy, Farnaz	Taghadosi, Mahdi	Zavvar, Mahdi
Rezaei, Nima	Seyedi, Shahram	Taghvaei, Behrang	
		Taheri, Tahereh	



Scientific Committee of ICIA2018 (International)

Ciacchi, Carolina

Harandi, Ali M

Jalili, Ahmad

Kabelitz, Dietrich

Kheradmand, Farrah

Kropf, Pascale

Manjili, Masoud H

Modabber, Farrokh

Nademi, Zohreh

Rajalingam, Raja

Sadegh-Nasseri, Scheherazade

Schmidt, Reinhold E.

Simon, Hans-Uwe

Stear, Michael J

Wahn, Ulrich

Scientific Committee of ICIA2018

Workshops

Dr. Ahad Zare
Dr. Ali Mostafaie
Dr. Ali Norozi Aghideh
Dr. Ali Pormohammad
Dr. Amir Jalali
Dr. Babak Arjmand
Dr. Ebrahim Faghihloo
Dr. Ehsan Jan Zamin
Dr. Elham Ashouri
Dr. Farhad Jadidi-Niaragh
Dr. Fatemeh Faraji
Dr. Flora Forouzesh
Dr. Forouzan Karimi
Dr. Hossein Asgarian-Omran
Dr. Jafar Salimian
Dr. Kamran Mansouri
Dr. Kaveh Baghaei
Dr. Koroush Kalantar
Dr. Mahmood Bozorgmehr
Dr. Majid Khoshmirsafa
Dr. Maryam Khairandish
Dr. Marzieh Ebrahimi
Dr. Mirhadi Jazayeri

Dr. Maryam Kheirandish
Dr. Mitra Nourbakhsh
Dr. Mohamad Keramatipour
Dr. Mohamad Reza Haghsheenas
Dr. Mohammad Davoodzadeh Gholami
Dr. Mohammad Javad Nasiri
Dr. Mohammad Mehdi Amiri
Dr. Mohammad Roostami Nejad
Dr. Nariman Mosaffa
Dr. Parvane Farzane
Dr. Raheleh Shokouhi Shoormasti
Dr. Reza Falak
Dr. Reza Zolfaghari Emameh
Dr. Rozita Khanafari
Dr. Saeid Abediankenari
Dr. Sattar Gorgani-Firuzjaee
Dr. Siavash Ahmadi Nourbakhsh
Dr. Vahid Jajarmi
Dr. Vahid Salimi
Dr. Zahra Alizadeh
Dr. Zohreh Mohamadi
Dr. Mohammad Davoodzadeh Gholami

Executive Committee

Abassi, Ardashir	Eskandarian, Maryam	Jafarpour, Roghaye
Abbasi-kenarsari, Hajar	Esmaeili, Seyed Alireza	Jafarzadeh, Fatemeh
Abdolmohammadi, Kamal	Ezzatifar, Fatemeh	Javidan, moslem
Abedini, Fateme	Faghih, Manizhe	Johari nad, Fatemeh
Afzali, Shima	Fakhari, Yaser	Kakadezfuli, Neda
Aghamajidi, Azin	Fallah Mehrjardi, Keyvan	Karimi, Mozhd
Ahmadi Fesharaki, Raoufe	Fathi, Farshid	Katebi, Asal
Ahmadifard, Reza	Fathollahi, Anwar	Kazemi Sefat, Nazanin Atieh
Alimohammadi, Masoumeh	Fattahi, Pooya	Kazemi, Mohammad Hossein
Alimohammadi, Reza	Gaeini, Amirhossein	Keshavarzian, Nafise
Alinezhad, Maede	Ghahremani, Maryam	Khabareh, Maryam
Alirezaee, Atefe	Ghasemi, Nazanin	Khakpoor Koosheh, Mohammad
Asadirad, ALI	Ghasiyari, Haniye	Khan Nazer, Nikoo
Asgari, Marzieh	Ghazanfari, Hadi	Khani chamani, Fateme
Asghari, Mohadeseh	Gholami, Parastou	Kheiri, Farzaneh
Askarzadeh, Monireh	Golmohammad pour, Ladan	Khoshmirsafa, Majid
Atashzar, Mohammad reza	Habibzadeh, Sima	Khosroopour, Zahra
Azempour, Elham	Haghdooft, Sepideh	Khubraftar, Saeedeh
Azizi, Mahdieh	Hajatbeigi, Baran	Kiaa, Vahid
Babaei, Erfan	Haji Khodaverdi Khani, Hedieh	Lotfinejad, Sedighe
Bagheralmoosavi, Sarvin	Hajivalili, Mahsa	Mahmoudi, Mohammad
Bagheri, Salman	Hajmahmoodi, Somayeh	Mami, Fatemeh
Baghernezhad, Zainab	Hasanzadeh, Mohammad Ali	Mami, Somaye
Baradaran, Behnoosh	Hashemi, Zahra	Mansourabadi, Amirhossein
Bazargan, Mahsa	Hassan Pour, Zahra	Mardani, Fatemeh
Beheshti, Elahe	Heidari, Maryam	Mashhadi, Fatemeh
Changaei, Mostafa	Heidari, Neda	Masoudnia, Mina
Dalil Roofchayee, Neda	Heidari, Sahel	Masoumi, Farima
Davoudzade Gholami, Mohammad	Hesseini, Vahedeh	Masoumikarimi, Masoumeh
Delsuz, Rezaee	Hosseini, Maryam	Mazarzaei, Abdulbasat
Doroudian, Fateme	Iranshahi, Nasrin	Mehdinejadani, Kobra
Eshghi, Fateme	Jafari, Davood	Mehdinejadani, Shayesteh



Mohammadi kordkheyli, Marziyeh	Rahimi, Arezou	Seyyednajib, Mirali
Mohammadi, Maryam	Rahmani, Hoda	Shahidi, Maryam
Mojtahedi, Haniye	Ramezani, Mahya	Shahir, Mehri
Moloudi Zargari, Milad	Rammezani, Khadijeh	Shams, Forough
Momeni, Zahra	Rashidi, Nesa	Sheikhi, Shima
Moradi, Maryam	Rasouli, Shima	Shirazi
Mortezagholi, Sahar	Rastegari-Pouyani, Mohsen	Shojaei, Samaneh
Moulazade, Alireza	Rezaei Lotfi, Alaleh	Shojaei, Zeinab
Mousavi, Seyed Mohammad Javad	Rezaei, Ramazan	Shokri, Mohammad Reza
Mousavinasab, Fatemehsadat	Rostami, Elham	Shokrollahi, Mahdiah
Namazi, Shideh	Rostamian, Hosein	Taene, Abdulghader
Nazari, Farzad	Rostamzadeh, Zainab	Taghavi, Mahsa
Nejatbakhsh Samimi, Leila	Sadeghi, Atefe	Tavakkoli, Sajjad
Niknam, Bahare	Sadeghi, Somaye	Touzandehjani, Dania
Nikoonezhad, Maryam	Sadegholvad, Mona	Vafa, Ashraf
Norouzbeigi, Mina	Sadri, Maryam	Vaziri Nezam Doust, Fereshte
Omid, Zahra	Saeidi Brojeni, Ali	Yafthian, Atefeh
Paydarfar, Shadi	Safaei, Elmira	Yousefi, Roya
Pormohammad, Ali	Safarzadeh, Mehrnoush	Zadian, Seyed Sajjad
Pouya, Sedigheh	Salehi, Saeedeh	Zand, Nafiseh

Invited Speakers



The poster features a blue background with a subtle pattern of white molecular structures. At the top, there are three logos: the Society for Immunology and Allergy, the ICIA2018 logo, and the Shariati School of Medical Sciences. Below these is the IUIS logo. The title "Invited Speakers" is prominently displayed in the center. The names and affiliations of the invited speakers are listed in two columns, each accompanied by a circular portrait photo.

Speaker Name	Affiliation
Prof. Carolina Ciacchi	University of Salerno, Italy
Dr. Ali M. Harandi	University of Gothenburg, Sweden
Dr. Ahmad Jalili	Baergersteck Medical Center, Switzerland
Prof. Dieter Kabelitz	University of Kiel, Germany
Prof. Farrah Kheradmand	Baylor College of Medicine, US
Dr. Pascale Kroft	Imperial College London, UK
Prof. Masoud H. Manjili	Virginia Commonwealth University, US
Dr. Zohreh Naderi	Newcastle upon Tyne University, UK
Dr. Raja Rajalingam	University of California, US
Prof. Hans-Uwe Simon	University of Bern, Switzerland Editor in chief of Allergy
Prof. Scheherazade Sadegh-Nasseri	Johns Hopkins University, US
Prof. Michael James Stear	La Trobe University, Australia
Prof. Reinhold E. Schmidt	Hannover Medical School, Germany
Prof. Ulrich Wahn	Charité-Berlin, Germany Editor in Chief of Pediatric Allergy and Immunology



Allergy and Immunotherapy of Allergic Disease

Oral Presentation

14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Allergy and immunotherapy\Oral\5314

5314

Status of IL-10 gene expression in patients with allergic rhinitis who received immunotherapy and symbiotic

Sajad Dehnavi¹, Mojgan Mohammadi^{2*}, Farahzad Jabbari Azad², Reza Farid Hosseini², Jalil Tavakol Afshari³, Maryam Khoshkhooi², Hamid Ahanchian², Nasrin Moazzen¹, Maryam Davarpanah¹, Amirabbas Salmani¹

1. Department of Immunology, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

2. Allergy Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

3. Immunology Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

Background: Allergic Rhinitis (AR) as the most common allergic disease has an increasing incidence in all over the world. Annually, billion dollars are paid for various types of allergic diseases. There are several lines of evidence showing the effects of symbiotic in the treatment of allergic diseases. We aimed in this study to evaluate the effects of symbiotic on the gene expression of IL-10 as one of regulatory T cell (Treg) cytokines in patients with AR who received immunotherapy.

Methods: Twenty patients with AR who underwent cluster immunotherapy were randomized to receive symbiotic and placebo for two months. Before and after the procedure, clinical symptoms were evaluated and whole blood samples were collected to extract RNA from peripheral blood mononuclear cells (PBMCs), then cDNA was synthesized and finally gene expression was evaluated by SYBR Green real-time PCR technique.

Results: Along with improvement of clinical symptoms and quality of life, gene expression of IL-10 increased in both symbiotic and placebo groups, this increase was more evident in symbiotic group but the difference was not significant ($p=0.72$).

Conclusion: Our study showed that administration of symbiotic may have a positive effect on Treg cells leading to improved clinical symptoms and quality of life in AR patients who underwent immunotherapy. However, increasing the sample size might help us get a better conclusion.

Keywords: Allergic Rhinitis, Immunotherapy, Symbiotic, IL-10

ICIA2018\Allergy and immunotherapy\Oral\7586

7586

PD-1 and Tim-3 immune checkpoint receptors are more expressed on T-CD4⁺ lymphocytes of patients with asthma

Ali Mosayebian¹, Zohreh Koohini¹, Hadi Hossein-Nataj¹, Saeid Abediankenari^{1,2}, Siavash Abedi^{3*}, Hossein Asgarian-Omran^{1,2*}

1. Department of Immunology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

2. Immunogenetics Research Center, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

3. Department of Internal Medicine, Pulmonary and Critical Care Division, Mazandaran University of Medical Sciences, Sari, Iran

Background: Asthma is a chronic disorder of the airways characterized by reversible airway obstruction, inflammation and bronchial hyperresponsiveness. Different immune cells and molecules have been shown to be involved in the pathogenesis mechanisms of asthma. T cell immunoglobulin and mucin domain-containing molecule-3 (Tim-3) and programmed death-1 (PD-1) are two surface immune checkpoint receptors expressed mainly on Th1 and Th2 cells, respectively which play an important role in regulating the activity of these cells. Due to the limited studies on the roles of Tim-3 and PD-1 in the pathogenesis and development of asthma, in the present study, we studied the expression of Tim-3 and PD-1 on CD4⁺ T cells of asthmatic patients.

Methods: A total of 37 asthmatic patients and 32 healthy controls were enrolled in this study. Fresh peripheral blood mononuclear cells (PBMC) were collected from all subjects and were used for subsequent experiments. The frequency of Tim-3⁺/ PD-1⁺/ CD4⁺ T cells was determined by three-color flowcytometry method. To evaluate the Th1/Th2 ratio, PBMCs were stimulated with PMA/ionomycin for 18 h. IFN- γ and IL-4 were measured in culture supernatants by ELISA. Serum total IgE was also measured in all samples.

Results: Significant increase in the percentage and absolute count of Tim-3⁺/ PD-1⁺/ CD4⁺, Tim-3⁺/CD4⁺ and PD-1⁺/CD4⁺ T cells was found in asthmatic patients compared to healthy controls. The IFN- γ /IL-4 ratio (Th1/Th2 ratio) was significantly higher in healthy controls than that of asthmatic patients.

Conclusion: Our data showing the increased expression of PD-1 and Tim-3 on CD4⁺ T cells of patients with asthma suggest the potential roles of these immune checkpoint receptors on the pathogenesis of asthma and promotion of Th2-mediated immune responses.

Keywords: Asthma, Tim-3, PD-1, Th1/Th2 ratio

ICIA2018\Allergy and immunotherapy\Oral\7636

7636

Identification and Characterization of major allergens from elm (*Ulmus glabra*) pollen grains

Elham Mahlooji¹, Mohammad Reza Mozdianfard¹, Marzieh Heidarzadeh Arani², Mostafa Moin³, Maryam Sharifshoushtari^{3,4}, Morteza Djamali⁵, Reza Mansouri⁶, Zahra pourpak³, Hossein Batooli⁷, Gholam Ali Kardar³

1. Department of Chemical Engineering, Faculty of Engineering, University of Kashan, IR Iran

2. Pediatric Department, Faculty of Medicine, Kashan University of Medical Sciences, Kashan

3. Immunology, Asthma and Allergy Research Institute, Children's Medical Center, Tehran University of Medical Sciences, Tehran, IR Iran

4. Department of Biology, Science and Research Branch, Islamic Azad University, Tehran, Iran

5. Institut Méditerranéen d'Ecologie et de Paléoécologie (IMEP UMR CNRS 6116), Europôle Méditerranéen de l'Arbois, Bâtiment Villemin BP80, 13545 Aix-en-Provence, Cedex 04, France

6. Department of Physics, Sharif University of Technology, Tehran, 14588-89694, Iran and Saeedi Institute for Advanced Studies, University of Kashan, Kashan, 87317-51167, Iran

7. Isfahan Research Center of Natural Sources and Agriculture, Kashan Station, Kashan, Iran

Background: Elm (*Ulmus*) tree pollen is a known aeroallergen, found commonly in the semi-arid climate of Kashan-Iran, planted primarily for its shade and aesthetic appeal. It pollinates from early February to mid March and is distributed widely throughout the world's temperate and tropical regions. Four species have been identified in Iran, namely: *U. glabra*, *U. minor*, *U. boissieri* and *U. umbraculifera* with the former being an important species in the northern forests of Iran. In this research, we identified and characterized major allergens in elm pollen.

Methods: Physical appearance specifications of *U. glabra* species were examined using SEM. Pollen protein was analyzed using SDS PAGE while, total protein content was measured by Bradford assay. Total IgE was examined by taking venous blood from 18 patients (8 male and 10 female), sensitive to elm with positive skin prick test (SPT). Major allergens were identified by immunoblot technique on pooled patients' sera (using 5% pollen extract taken from locally collected pollen). Allergenic proteins were later purified using gel filtration chromatography.

Results: Pollen grains were oblate-spheroidal to angular and square in form and contained 4-6 elliptical pores with thick and rugulate exine, sized 16-50 X 18-50 μm . Four components showed IgE-binding activity and were therefore identified as the elm pollen allergens, having molecular weight ranged between 39-70 kDa. Average total IgE was 162.98 IU/mL which reacted with a distinct major band at nearly 55 kDa. Immunoblot analysis identified a novel allergen protein, named *Ulm g1*.

Conclusion: The 55 kDa allergen was the major allergen of *U. glabra*. Moreover, combination of total IgE and immunoblot techniques, could be effectively used in the diagnosis of elm pollen allergy.

Key words: Pollen grains, Elm tree, allergen, seasonal allergy, *Ulm g1*

ICIA2018\Allergy and immunotherapy\Oral\7879
7879

Loratadine counter-regulates YKL-40 release from human monocytic cell line (THP-1) but not human epithelial cells (A549)

Zeynab Baghernejadan ¹, Mostafa Haji Molla Hoseini ²

1. Department of Medical Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

2. Department of Medical Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Background: Recent research has found the evidence of chitinase and chitinase like proteins contributions to the mechanisms of allergic response and their association with atopy. The chitinase like protein, YKL-40, plays an important role in the pathogenesis of allergic diseases. Allergens induce its production from macrophages and epithelial cells.

Some investigations indicate that loratadine has anti-inflammatory effects apart from its H1 receptors anatagonistic activity. We sought to know whether human epithelial cells and macrophages stimulated for YKL-40 synthesis by host dust mite (HDM) extract, might be influenced by loratadin. We used the human pulmonary type II epithelial cell line A549 and the human monocytic cell line (THP-1) for our in vitro study.

Method: After dose finding studies, using neutral red cytotoxicity assay, the cells were pre-incubated with loratadin (0.5 µg/mL) and then activated by HDM (Greer Lab). Also in post-treatment conditions; A549 and THP-1 cells were stimulated with HDM extract followed by incubation with loratadine. Then, the amounts of IL-8 and YKL-40 released into the cell supernatants were determined by a specific ELISA assay. The classic chemotaxis assay was performed, using modified Boyden chambers as well.

Results: Loratadine has no effect on IL-8 production and neutrophil chemotaxis under these conditions. Unlike A549, THP-1 cells stimulated with HDM significantly increased the release of YKL-40. Pre-incubation with loratadin diminished the YKL-40 release from THP-1 cells in a significant manner.

Conclusions: Loratadin might exert anti-inflammatory effects beyond its H1-receptor antagonistic activity in the context of inflammatory disorders of the respiratory tract such as bronchial asthma and allergic rhinitis.

Keywords: Loratadine, YKL-40, A549, THP-1

ICIA2018\Allergy and immunotherapy\Oral\7996
7996

T cell-specific transcription factors gene are differently expressed in Allergic Rhinitis

Soraya Bilvaieh¹, Hamid reza Mortazavi¹, Farhad Salari¹, Ali Grgin karaji¹.

1. Department of Immunology, School of Medicine, Kermanshah University of Medical Sciences

Background: Allergic rhinitis (AR) is the most common form of IgE-mediated disease. T cells subsets have a main role in induction and control of AR. The aim of this study was to evaluate the level of GATA-3, ROR γ t and FoxP3 (the hallmark of Th2, Th17, and T regulatory cells, respectively) gene expression in peripheral blood cells of AR patients.

Methods: RNA was extracted from blood cells of 32 AR patients and 20 healthy controls and then transcribed to cDNA. The cDNA was used to measure the level of GATA-3, ROR γ t and FoxP3 gene expression by means of Real Time-PCR and specific primers. In addition, AR patients and healthy controls Total serum IgE was measured by ELISA.

Results: Our results showed that while the level of FoxP3 expression in AR patients was higher than that of control group ($P<0/05$), the level of GATA-3 expression in AR patient was lower than controls ($P<0/05$). On the other hand, the level of ROR γ t expression in patient group was higher than that of control group, although it was not statistically significant. Total serum IgE in AR patients was higher than control group ($P<0/05$).

Conclusion: Taken together these results showed that the level of FoxP3 expression in AR patients was higher but the level of GATA-3 expression was lower than that of healthy controls. These unexpected results may be related to chronic state of the disease and consumption of anti-inflammatory drugs such as Glucocorticoids by AR patients.

Key words: Allergic Rhinitis, GATA-3, ROR γ t, FoxP3, Th2, Th17, T regulatory

ICIA2018\Allergy and immunotherapy\Oral\8089
8089

Epidemiology of Chronic Rhinosinusitis in Southwestern Part of Iran: A GA²LEN Study

Shokrollah Farrokhi¹, Afshin Ostovar², Wytske J. Fokkens³

1. Department of Immunology, Asthma and Allergy, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Research Institute, Bushehr University of Medical Sciences, Bushehr, Iran

2. Department of Otorhinolaryngology, Academic Medical Center, Amsterdam, the Netherlands

3. Department of Epidemiology, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Research Institute, Bushehr University of Medical Sciences, Bushehr, Iran

Background: Population based studies by the European Position Paper on Rhinosinusitis (EPOS) criteria for assessment of the prevalence of chronic rhinosinusitis (CRS) play important roles in the development and promotion of public health policies.

Methods: A multistage, stratified cluster random sampling method was used to select participants randomly from people living in Bushehr, southwestern part of Iran. Standardized questionnaire of GA²LEN were completed by 5201 participants after personal interviewing. Logistic regression analyses were used to examine the association between variables and risk factors of CRS.

Results: The overall prevalence of CRS was 28.4%. It was slightly higher among females (28.3%) than males (28.2%) ($P = 0.4$). CRS was more prevalent in smokers, 25-34 age range group, individuals with a history of asthma, allergic rhinitis or eczema, non-educated persons and health care workers ($P < 0.05$).

Conclusions: Bushehr had the highest prevalence of CRS in the world. Therefore, CRS can pose high economic burden on the population and health system. Our study also provides important information for the development and promotion of public health policies associated with CRS particularly in developing countries.

Key words: chronic rhinosinusitis, EPOS criteria, GA²LEN study, prevalence

ICIA2018\Allergy and immunotherapy\Oral\8244

8244

Evaluation of the efficacy of nicotine in the treatment of allergic asthma in BALB/c mice

Ebrahim Mazloomi¹, Behrooz Ilkhanizadeh², Ahad Zare³, Adel Mohammadzadeh⁴, Nowruz Delirez⁵
Shahram Shahabi^{1*}

1. Cellular and Molecular Research Center, Urmia University of Medical Sciences, Urmia, Iran

2. Department of Pathology, School of Medicine, Urmia University of Medical Sciences, Urmia, Iran

3. Immunology, Asthma and Allergy Research Institute, Tehran University of Medical sciences, Tehran, Iran

4. Departments of Genetics and Immunology, School of Medicine, Urmia University of Medical Sciences, Urmia, Iran

5. Department of Cellular and Molecular Biotechnology, Institute of Biotechnology, Urmia University, Urmia, Iran

Background: Nicotine, an nAChR agonist, shows prominent anti-inflammatory properties, and some studies have shown its suppressive effects on inflammation. Here, we examined whether nicotine as a medicine has beneficial effects on asthma treatment in a mouse model of allergic asthma

Methods: BALB/c mice were sensitized with OVA and alum. Two weeks later, the mice received nicotine in concentrations of 1 and 10 mg/kg three times. After 10 days the mice were challenged with OVA (5 percent) using an ultrasonic nebulizer and sacrificed the next day. To evaluate Treg cells proliferation against the allergen, splenocytes were cultured with OVA; the cells were stained with anti-mouse CD4 and FOXP3 antibodies 72 hours later.

Results: Our results showed that the administration of nicotine reduced lung-tissue inflammation, number of eosinophils in bronchoalveolar fluid, allergen-specific IgE and IL-4 production, while it increased the TGF- β /IL-4 ratio and Treg cells proliferation against the allergen. Our results showed that nicotine exerts its suppressive effects in a dose-dependent manner: administration of 10 mg/kg of nicotine showed more suppressive effects than 1 mg/kg

Conclusion: These data suggest that nicotine, by decreasing allergic inflammation severity and potentiating Treg cells proliferation against the allergen, could be used as a medicine in the treatment of allergic asthma by.

Keywords: Allergic asthma, anti-inflammatory cholinergic pathway, nicotine, nAChRs

ICIA2018\Allergy and immunotherapy\Oral\8594
 8594

Ash (*Fraxinus excelsior*) pollen grains Proteomic approach of most common allergens sensitization pattern in Iranian allergic patients

Maryam Sharifshoushtari^{1,2}, Ahmad Majd³, Mostafa Moin^{2,4}, Zahra pourpak^{2,4}, Taher Nejdassattari¹, Shaghayegh Tajik², Mohsen Badalzadeh², Mohamad Reza Fazlollahi², Raheleh Shokohi Shoormasti², Behnoosh Tayebi⁵, Zakiyeh Azadi², Ahad Zare², Gholam Ali Kardar³

1. Department of Biology, Science and Research Branch, Islamic Azad University, Tehran, Iran
2. Immunology, Asthma and Allergy Research Institute, Children's Medical Center, Tehran University of Medical Sciences, Tehran, Iran
3. Department of Plant Biology, faculty of biology science, Islamic Azad University, North Tehran Branch, Tehran, Iran
4. Department of Immunology and Allergy, Children Medical Center, Tehran University of Medical Sciences, Tehran, Iran
5. Royan institute_ institute for Stem Cell Biology and Technology (RI-SCBT), Tehran, IRAN

Background: Ash (*Fraxinus excelsior*) is a popular tree, climate change patterns and as a cause of winter-spring pollinosis in some regions worldwide. The aim of this research we investigated to determine the profile of main allergen components diagnosis for patients with sensitivity to ash pollen implicated in the common ash pollen allergy.

Methods: Ash pollen count analysis and data were documented from 2010 through 2017 in Tehran. Pollen extracts were resolved by one and two-dimensional electrophoresis and assayed with sera of allergic subjects. Sera from three groups, group one: (n=19), patients polysensitized with ash positive specific IgE, group two: (n=10), patients were ash negative, but positive to other pollens and group three: (n=9), sera control. All serum samples did immunoblot analysis with 5% pollen extract.

Results: Total pollen counts varied extensively between 87 to 2037 grains/m³/24 h. The allergens with apparent molecular masses ranging from nine to 70 kDa were detected. Generally, twelve spot allergens have been identified in ash pollen while just one allergen has been novel allergen in hsp70 KDa. The interesting result is; all patients of group two sera have shown positive reaction with ash extract in western blot.

Conclusion: According to results, it seems the local *F.excelsior* extract is difference with source extract used for developed RIDA strip and also prick test solution and/or this is cause of different result of commercial RIDA and this research western blot on group two samples. Maybe we should be overview in our diagnosis methods for avoiding to false negative and also false positive results.

Keywords: Proteomic approach, Ash pollen, Mass Spectrophotometry, Allergen

ICIA2018\Allergy and immunotherapy\Oral\9822
9822

An asthma-associated IL4R variant exacerbates airway inflammation by promoting conversion of regulatory T cells to TH17-like cells

Massoud AH^{1,2}, Charbonnier LM^{1,2}, Phipatanakul W^{1,2}, Chatila TA^{1,2}.

1. Division of Immunology, Boston Children's Hospital, Harvard Medical School, Boston, Massachusetts, USA.
2. Department of Pediatrics, Harvard Medical School, Boston, Massachusetts, USA.

Background: Mechanisms by which regulatory T (T_{reg}) cells fail to control inflammation in severe asthma remain poorly understood. A glutamine- (Q) to arginine (R) substitution at position 576 of IL-4Rα (IL-4Rα-Q576R) has been linked to asthma exacerbation and severity.

Methods: By utilizing a mouse model of airway inflammation in IL-4Rα-Q576R mice we demonstrated an aberrant increase of IL-17 and IL-6 expression in mutated Treg cells (by flow cytometry), associated with skewing of Treg cell to the T helper (T_H) 17 cell fate. T_{reg} cell lineage-specific deletion of RAR related orphan receptor C (Rorc) reverses the aggravated airway inflammation in Il4ra^{R576} mice.

Results: We here show the IL-4Rα-R576 mutation enables recruitment of the adaptor GRB2 to promote airway inflammation by a novel mechanism involving IL-4-directed T_{reg} cell differentiation towards the T helper (T_H)17 cell lineage.

Conclusion: This study elucidates a novel mechanism for the development of mixed T_H2 and T_H17 cell responses relevant to asthma and other allergic inflammatory diseases. By activating an additional branch of IL-4R signaling initiated by the adaptor protein GRB2, an IL-4Rα polymorphism associated with asthma exacerbations and severity potentiates T_H17 and to a lesser extent T_H2 responses.

Keywords: Asthma, Regulatory T cells, T_H17



Cancer Immunology

Oral Presentation

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Cancer Immunology\Oral\7634
7634

The significance of CD45RO+ tumor-infiltrating lymphocytes in breast cancer

Simin Ahmadvand^{1, 2}, Zahra Faghih², Akbar Safaei³, Mehdi Montazer³, Abbas Ghaderi^{1, 2*}

1. Department of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

2. Shiraz Institute for Cancer Research, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

3. Department of Pathology, School of Medicine, Shiraz University of Medical Sciences, Iran

Background: Breast cancer (BC) is the most common and the second leading cause of cancer related deaths among women worldwide. The presence of lymphocytic infiltrates in breast tumors have long been proved and seems to have prognostic impact on patients' outcome.

Methods: The study population included 94 females with invasive ductal carcinoma of the breast who received surgical resection between 2009 and 2011 in Shahid Faghihi Hospital of Shiraz University of Medical Sciences, Iran. Four μ m sections were prepared from their FFPE tissue blocks and immunohistochemically stained for CD3, CD8, CD45RO and Foxp3 molecules. Digital photos were taken from both, center (CT) and invasive margins (IM) of tumors by an expert pathologist and number of the positive cells were counted by FIJI image analyzer. Differences in cells' frequencies among different clinicopathological features of the disease as well as recurrence were analyzed by Mann-Whitney U test.

Results: We observed that higher frequencies of CD45RO+ TILs as well as other subpopulations were associated with lymph node involvement, high grade, advanced stages, ER negativity and HER2 positivity. The frequency of CD45RO+ TILs was significantly higher in relapse free patients ($P < 0.05$).

Conclusion: more TILs infiltration was associated with more aggressive features of BC, additionally our results implied that increase in frequency of CD45RO+ TILs which probably enriched for effector/memory cells could reduce the risk of post-operative risk of relapse.

Keywords: Breast cancer, CD45RO+tumor-infiltrating lymphocytes

ICIA2018\Cancer Immunology\Oral\7746
7746

Functional study of CD25⁺FoxP3⁺CD127⁻CD4⁺ cells in tumor draining lymph nodes of breast cancer patients

Zahra Faghih¹, Andisheh Niakan^{1,2}, Abbas Ghaderi^{1,2}

1. Shiraz Institute for Cancer Research, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

2. Department of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Tumor-derived CD4⁺CD25⁺FoxP3⁺ Tregs have been extensively studied in different types of cancer including breast cancer (BC) and in most cases, they are associated with tumor progression mostly through suppressing immune responses at tumor sites. Recently, a FoxP3⁺ subtype with CD4⁺CD25⁻FoxP3⁺ phenotype has been observed in some diseases especially autoimmune diseases. As little is known about their function in cancer patients, in this study, their frequency as well as their ability to produce cytokines were investigated in TDLNs of BC patients.

Methods: Mononuclear cells were isolated from auxiliary lymph nodes of 20 BC patients. Following activation with PMA/Ionomycin in the presence of Golgi inhibitors, the cells were stained for CD25, CD4, CD127, FoxP3, IFN- γ , IL-2, IL-22, IL-17 and IL-10 and acquired on four-color flow cytometer. Data were analyzed by CellQuest Pro software.

Results: Our analysis showed that less than 1% of CD4⁺CD25⁻FoxP3⁺ cells produced cytokines which were significantly lower than CD4⁺CD25⁺FoxP3⁺ regulatory and CD4⁺CD25⁺FoxP3⁻ effector subpopulations. Nevertheless, the MFI of IFN- γ and IL-2 in CD4⁺CD25⁻FoxP3⁺ subset was significantly higher than the MFI of these cytokines in Tregs; but lower than effector subset. On the other hand, the MFI of IL-22 in Tregs was significantly higher than in CD4⁺CD25⁻FoxP3⁺ population. The MFI of IL-10 in CD4⁺CD25⁻FoxP3⁺ subset was significantly higher than in effector cells.

Conclusion: Our results showed that CD4⁺CD25⁻FoxP3⁺ cells in TDLNs of BC patients have an intermediate phenotype between effector and regulatory cells, since they produce more IL-2 and IFN- γ than Tregs and more IL-10 than effector cells. These cells are probably a heterogeneous population of effector and regulatory T cells and/or a subset of dysfunctional Tregs which have lost their CD25 expression because of chronic stimulation, high suppressive function and repeatedly exposure to tumor antigens in tumor microenvironment.

Key words: Breast cancer, TDLNs, Treg, CD4⁺CD25⁻FoxP3⁺ T cells.

ICIA2018\Cancer Immunology\Poster\7457

7457

Invitro increasing cytotoxicity of CMV-specific CD8⁺ T cell in CMV-seropositive colorectal cancer patients

Shaker Ardakani M¹, Fazeli MS², Kokhaei P¹, Shakiba Y³, Tabatabaei Yazdi SM⁴, Abbasian A⁴, Emami AH⁵, Pak F¹, Nourizadeh M⁶

1. Department of Immunology, Semnan University of Medical Sciences, Semnan, Iran.
2. Department of Surgery, Imam Khomeini Complex Hospital, Tehran University of Medical Sciences, Tehran, Iran
3. Regenerative medicine Research center, Kermanshah University of Medical Sciences, Kermanshah, Iran
4. Tehran Blood Transfusion Center, Tehran, Iran.
5. Department of Hematology/Oncology, Imam Khomeini Hospital, Tehran University of Medical Sciences, Tehran, Iran
6. Immunology, Asthma and Allergy Research Institute, Tehran University of Medical Sciences, Tehran, Iran

Background: Colorectal cancer (CRC) is the third most common cancers with a high incidence of mortality and morbidity. A potential oncogenic role of HCMV in different types of cancers including CRC has been recently shown by detecting the CMV peptides in tumor tissues compared to the adjacent normal tissues. Targeting the CMV peptide-expressing tumor cells has been considered as an effective immunotherapy approach. Therefore, we tried to induce CMV-specific CD8⁺ cytotoxic T cells (CTLs) in vitro in CMV-seropositive CRC patients and compare it to healthy controls.

Methods: Sixteen CMV⁺ colorectal cancer patients and eight CMV⁺ healthy controls were enrolled the study. Patients' blood samples were obtained during tumor surgery (before chemotherapy and radiotherapy treatments). Cytotoxicity was measured by detecting the CD107a on the surface of CMV-specific CD8⁺ T cells using flow cytometry before and after 14-day culture of PBMCs in the presence of CMV peptide epitopes and IL-2 cytokine. Data were analyzed using FLOWJO software.

Results: CD107a expression on CD8⁺T cells in both patient and control groups increased significantly after 14-day culture in the presence of CMV peptide epitopes and IL-2 which was an indicator of the potential cytotoxicity function of CTLs.

Conclusion: The results of this study specifies that in vitro stimulation of PBMC in the presence of CMV peptide epitopes and IL-2 can be an applicable method to generate potent CD8⁺ CTLs for future T cell therapy in CRC patients.

Keywords: Colorectal Cancer, CMV, CD107a

ICIA2018\Cancer Immunology\Oral\7845

7845

MicroRNA-330 inhibit colorectal cancer proliferation and migrationBehzad mansoori^{1,2}, Ali Mohammadi¹, Solmaz Shirjang¹, Behzad Baradaran¹*1. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran**2. Student Research Committee, Tabriz University of Medical Sciences, Tabriz, Iran*

Background: miR-330 has been reported in several cancers, the cellular and molecular mechanism of miR-330 in colorectal cancer remains unclear. A recent study suggested that miR-330 could be one of the key regulators of CRC. The aim of this study is to investigate the role of miR-152 in CRC and its mechanisms.

Methods: The pCMV-GFP-miR-330 construct was transfected into SW-480 cells. The miR-330 positive cells were selected by Geneticin antibiotic (G418) for two-weeks. To evaluate the effect of miR-330 on the colorectal cancer cell proliferation and migration, MTT and scratch wound healing assays were performed, respectively. Then, the expression level of miR-330 and the inhibitory effect of miR330 on caspase3 expression was evaluated by the qRT-PCR.

Results: The induction of miR-330 in SW-480 cells was confirmed by GFP channel imaging system and the 10 fold increase in miR-330 expression in stable cells. MTT and wound healing assays showed that miR-330 have anti-proliferation and cell migration effects in stable miR-330 colorectal cancer cell line compared to control group. The result of qRT-PCR on caspase3 expression showed miR-330 could decrease the expression level of miR-330 in stable miR-330 cells compared to control cells.

Conclusions: As the results displayed miRNA-330 could regulate the colorectal cancer proliferation and migration by targeting the caspase3 mRNA. Also the result suggested that miR-330 can be a therapeutic molecule in target therapy of colorectal cancer.

Keywords: miR-330, Colorectal cancer, Proliferation, Migration, Caspase3

ICIA2018\Cancer Immunology\Oral\7911

7911

Increased frequency of CD4⁺ and CD8⁺ T-cell expressing CD11c and PD-1 in lymphoid organs of mice with colorectal cancer

Davood Rostamzadeh^{1,2}, Ahmad Hosseini², Zohreh Babaloo³, Alireza Rezvani⁴, Zahra Mojtahedi²

1. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

2. Shiraz Institute for Cancer Research, School of Medicine, Shiraz University of Medical Science, Shiraz, Iran

3. Tabriz, Iran. Department of Immunology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

4. Department of Hematology, Medical Oncology and Stem Cell Transplantation, Hematology Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

Background: CD11c is an alpha integrin classically employed to define myeloid dendritic cells. Recent reports identified CD11c-expressing CD8⁺ T-cells as a new subset of CD4⁺ regulatory T-cells. There is some evidence that CD11c⁺CD8⁺ T-cells may exert their effector functions or regulatory functions in different conditions. Until now, there have no studies published in which the frequency of CD11c⁺ T-cells in cancer. There is also limit evidence on expression of immune-checkpoint receptor programmed cell death-1 (PD-1) in mouse lymphoid organs, especially in bone marrow. Therefore, we analyzed the frequency of CD11c⁺CD8⁺ and CD11c⁺CD4⁺ T-cells as well as PD-1 expressing CD4⁺ and CD8⁺ T-cells in different tissues of young, mature and colorectal cancer-bearing (CRC) mice.

Methods: C57BL/6 mice divided into three groups, including young mice (3-4 weeks), mature mice (13-14 weeks old) and CRC mice. Mice were sacrificed and frequency of immune cells in bone marrow, lymph nodes and spleen were analyzed by flow cytometry.

Results: In contrast to young and mature mice, CRC mice displayed higher percentage of CD11c⁺CD8⁺ and CD11c⁺CD4⁺ T-cells in bone marrow than those compared to lymph nodes and spleen. Interestingly, higher percentage of CD11c⁺CD8⁺ T-cells, but not CD11c⁺CD4⁺ T-cells was observed in bone marrow of CRC mice in compared to lymph nodes and spleen. We also higher frequency of CD8⁺PD-1⁺ T-cells in BM, spleen and LN of CRC mice compared with young and mature mice.

Conclusion: As T-cell exhaustion is associated with inhibitory receptor PD-1 and according to regulatory roles of CD11c expression CD8⁺ T-cells, elevated percentage of CD11c and PD-1 expressing T-cells are related with dampens antitumor immunity.

Keywords: CD11c⁺CD8⁺ T-cells, CD11c⁺CD4⁺ T-cells, PD-1, Colorectal cancer

ICIA2018\Cancer Immunology\Oral\8329

8329

CD4⁺Treg cells in tumor draining lymph nodes of bladder cancer patientsZahra Faghih¹, Yasmin Vahidi¹, Maryam Fakhimi¹, Ardalan Asadollah pour², Shahriar Zeighami², Ali Ariaifar^{2,3}*1. Shiraz Institute for Cancer Research, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran**2. Department of Urology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran**3. Urology-Oncology Research Center, Shiraz University of Medical Sciences, Shiraz, Iran*

Background: Regulatory T cells (Tregs) are a suppressive subset of CD4⁺ T cells which play an important role in the regulation of immune responses. It has been also demonstrated that Tregs at tumor sites could significantly suppress immune responses, leading to immune tolerance of tumor cells. In the present study, we investigated the frequency of CD4⁺ Tregs in tumor draining lymph nodes of patients with bladder cancer (BLC).

Methods: Mononuclear cells were isolated from lymph nodes of 35 untreated BLC patients subjected to radical cystectomy and stained for CD4, FoxP3, CD25, and CD127. The cells were then acquired on four-color flow cytometer and data were analyzed by CellQuest Pro software. The frequency of different cell subtypes i.e. CD25⁺FoxP3⁺CD127^{low/-} and CD25⁻FoxP3⁺CD127^{low/-} were assessed in CD4⁺ cells.

Results: The frequency of Tregs with CD4⁺CD25⁺CD127^{low/-}Foxp3⁺ phenotype is 57.40±2.92 in draining lymph nodes of BLC patients though no significant difference was found in their prevalence in patients with different clinicopathological characteristics. However, statistical analysis showed that the frequency of CD4⁺CD25⁻CD127^{Low/-}Foxp3⁺ cells significantly increased in involved lymph nodes (p=0.027). Additionally, CD4⁺Foxp3⁺ cells (p=0.021) and CD4⁺CD25⁻CD127^{Low/-}Foxp3⁺ cells (p=0.017) increased in patients with positive perivesical fat invasion.

Conclusion: Collectively, our results indicated that regulatory cells in draining lymph nodes of bladder cancers are probably a heterogeneous population with various phenotypes. Previous studies suggested that CD4⁺CD25⁻CD127^{Low/-}Foxp3⁺ cells can be a subset of dysfunctional Tregs which have lost their CD25 expression because of chronic stimulation, high suppressive function and repeatedly exposure to tumor antigens in tumor microenvironment. Increasing in the frequency of this subsets in patients with involved nodes suggested an inhibitory role for these cells in BLC.

Key words: Bladder cancer, lymph node, regulatory T cells

ICIA2018\Cancer Immunology\Oral \8414

8414

Comparison of the immunomodulatory effects of isolated stage 2 and stage 4 carcinoma associated fibroblasts in mouse breast cancer model

Ladan Langroudi¹, Zuhair MohammadHassan², MasoudSoleimani³, Seyed Mahmoud Hashemi ⁴

1. Department of Immunology, School of Medicine, Kerman University of Medical Sciences, kerman, Iran

2. Department of Immunology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

3. Department of Hematology, Faculty of Medical Sciences Tarbiat Modares University, Tehran, Iran

4. Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: studies have shown that the microenvironment of solid tumors affect the development and progression of cancer. Additionally they are the cells responsible for immune evasion and drug resistance of cancer cells. However the crosstalk between CAFs and the immune system is still unknown.

Methods: in this study, we characterized and compared the phenotype and the immunomodulatory properties of stage 2 and 4 CAFs isolated from mammary tumor of Balb/C mouse.

Results: Flow cytometry of the surface marker panel showed that stage 4 CAFs express significantly higher levels of HLA-DR. Also Higher levels of IL10, COX-2 and MMP9 enzyme were also observed in stage 4 CAFs. Whereas in stage 2 CAFs higher amounts of TFG- β and IDO expression was observed.

Conclusion: CAFs represent the supporting stroma of cancer. They are known responsible for immune evading and growth of cancer. Functional differences showed by their surface markers, cytokine and enzyme production indicates to induction of different microenvironments in their presence. The discrepancy observed in cancer therapies may be attributed to the. Therefore, further studies are required to fully elucidate the role of CAFs in various stages of cancer.

Keywords: Carcinoma Associated Fibroblasts, Breast Cancer, Immunomodulatory Properties

ICIA2018\Cancer Immunology\Oral\8176

8176

Silencing of Tim-3 Expression by miR-326 Affects Apoptosis and Proliferation of Human HL-60 Leukemia Cell Line

Maryam Mohammad-Ganji¹, Mazdak Ganjali khani-Hakemi^{1*}, Vida Homayouni², Abbas Rezaei¹, Hossein Khan-Ahmad³

1. Department of Immunology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

2. Acquired Immunodeficiency Research Center, Isfahan University of Medical Sciences, Isfahan, Iran

3. Department of Genetics and Molecular Biology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

Background: Leukemia Stem Cells (LSCs) are the main reason for drug-resistance and disease relapse in Acute Myeloid Leukemia (AML). Current drugs destroy normal Hematopoietic Stem Cells (HSCs) rather than LSCs. T cell immunoglobulin mucin-3 (TIM-3), an immune regulatory molecule, is a CD34+CD38- LSCs specific surface marker with high expression on these cells compared to HSCs. The interaction between TIM-3 and its ligand, Galectin-9 (Gal-9), mediates signaling pathways involved in apoptosis and proliferation. We hypothesized that miR-326 could have a suppressive activity on TIM-3 expression and hence, affects the proliferation and apoptosis processes in AML cells.

Methods: Bioinformatics predictions were done using Mirwalk and Target Scan softwares. TIM-3 expression was induced on HL-60 cells by PMA. After miR-326 transfection, MTT, q-RT-PCR, flowcytometry were performed to evaluate the cells survival and TIM-3 expression level. Then, after adding recombinant Galectin-9, apoptosis and proliferation rates were measured with Annexin-V and CFSE assays, respectively.

Results: Flow cytometry assay confirmed our bioinformatics prediction of suppressive effect of miR-326 on TIM-3 expression (66.4% silencing) in HL-60 cell line ($p=0.002$). The qRT-PCR results were also confirmatory. Annexin-V and MTT assays showed increased cell apoptosis and decreased cell survival, while data from CFSE assay indicated a severe reduction in HL-60 cells proliferation.

Conclusion: Our results demonstrated that, miR-326 can silence TIM-3 expression in AML HL-60 cells. Moreover, it is shown that miR-326 can enhance AML cells apoptosis and reduce their proliferation and survival and hence, might be considered as a novel target for therapeutic approaches against AML.

Keywords: Acute myeloid leukemia; T cell immunoglobulin mucin-3; microRNA-326; proliferation; apoptosis; .Galectin-9

ICIA2018\Cancer Immunology\Oral\8320

8320

Immunomodulation Effect of Gastro-spheres Through Disturbance of Th17/Treg Balance

Alaleh RezaLotfi^{1,2}, Ghasem Solgi^{*1}, Marzieh Ebrahimi^{*2}

1. Immunology Department, School of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

2. Dept. Stem Cells and developmental Biology, Cell Sciences Research Center, Royan Institute for Stem Cell Biology and Technology, Tehran, Iran

Background: Gastro-spheres with capacity to form tumor, believed to be responsible forescape from immune-mediated destruction. Adaptive immune system plays a major role in gastric cancer including Th17/Treg cells. Th17 producing IL-17 cells considered as crucial cells in pathogenesis of autoimmune diseases. In opposite, Tregs are immune-suppressive cells, prominent with tumor progression. Although it is indicated the prevalence of Treg/Th17 in the most cancers, their interaction with gastro-spheres have been elusive. The object of this study is to find the effect of gastro-spheres on Th17/Treg balance.

Methods: Isolated PBMCs from two healthy donors were cultured with MKN-45 cells and its derived gastro-spheres, as the model for enriched cancer stem cells conditioned medium (CM) through mixed lymphocyte reaction for 5 days at 37°C with 3 times stimulation. Responder groups expansion were evaluated by CFSE and the percentages of CD4⁺CD25⁺FOXP3⁺Treg cells and CD4⁺IL-17⁺ Th17 cells were analyzed by flowcytometry. ELISA measurement also performed for cytokines IL-17, TGF-β and IL-10.

Results: Flowcytometry analysis revealed an increase in percentages of Tregs and Th17 cells in PBMCs treated with gastro-spheres compared to parental cells CM. It was also demonstrated that T cell expansion was insignificantly decreased in gastro-spheres CM compared to parental and control group. ELISA measurement also revealed an increase in concentration of IL-10 and IL-17 but a decrease in TGF-β in gastro-spheres CM.

Conclusion: We concluded that gastro-spheres CM may increase the percentage of Tregs and Th17 cells more than their parental counterpart. This fact exhibited the possible existence of differentiation factors among gastro-spheres secretion, which contribute to T cells differentiation. Furthermore, it has been shown that gastro-spheres secretion can be possibly more immunomodulator than its parental and leads to limited T cell expansion.

Key words: Regulatory T cells, Th17 cells, Gastro-Spheres

ICIA2018\Cancer Immunology\Oral\8485

8485

Snail-1 silencing inhibits mir-34a and Let-7 expression in esophageal squamous cell carcinoma cell lineMaryam Hemmat zadeh^{1,2}, Hamed Mohammadi^{1,2}, Behzad Baradaran^{1,2}

1. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

2. Department of Immunology, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

Background: In cancer cells, the transcription factor Snail-1 promotes epithelial-to-mesenchymal transition (EMT), a process related to the emergence of invasion, and cancer progression. The purpose of this study was to evaluate the effects of the Snail-1 specific small interference RNA (siRNA) on the expression of miR-34a and Let-7 in TE-8 human esophageal squamous cell carcinoma cell line, in vitro.

Methods: TE-8 cells were transfected with the Snail-1 specific siRNA. The relative expression levels of Snail-1 mRNA, miR-34a, and let-7a were investigated by Quantitative Reverse Transcription PCR. Western blotting was carried out to evaluate the Snail-1 protein level. In vitro migration assay of TE-8 was also performed following the treatment with or without the Snail-1 specific siRNA.

Results: It was found that treatment of cancer cells with the Snail-specific siRNA effectively downregulated the expression of Snail-1 in mRNA and protein level. However, it had elevated the expression of miR-34a and let-7a expressions. Furthermore, suppression of Snail-1 led to diminished cell migration.

Conclusion: Our findings imply that increased level of miR-34a and let-7a might be associated with downregulation of Snail-1 and, thereby, subsided cell migration.

Keywords: Snail-1, miR-34a, Let-7a, Esophageal squamous cell carcinoma

ICIA2018\Cancer Immunology\Oral\9772
9772

Safety of intravenously applied mistletoe extract: results from a phase I dose escalation study in patients with advanced cancer

Seyed Ehsan Asadi^{1*}, Ahmad Rahimi², Matin Aghalar², Akram Jamali²

1. *Phd OF Nursing, Sina Hospital, Isfahan, Iran*

2. *Nursing Student OF Dehaghan University, Isfahan, Iran*

Background: Mistletoe extracts have anti-tumor properties and are approved for subcutaneous use in cancer patients. Data on Intravenous application are limited.

Methods: An aqueous extract from pine-mistletoe was used to investigate maximum tolerable dose (MTD) and safety of intravenous application. It was infused once weekly for 3 weeks in patients with advanced cancer. Any type of cancer was included; relevant exclusion criteria were concurrent chemo- or radiation therapy. The classical phase I 3 + 3 dose escalation scheme was followed. Predefined dose groups were 200, 400, 700, 1200 and 2000 mg. Maximum planned dose was 2000 mg. With the MTD three more patients should be treated for 9 weeks in order to evaluate intermediate term tolerability. Weekly during the treatment and 1 week later tolerability, clinical status, safety laboratory parameters and adverse events were documented.

Results: Twenty-four patients (3 in the dose groups 200, 400, 700 and 1200 mg, respectively, 10 in the dose group 2000 mg) were included. MTD was not reached. Because one dose-limiting toxicity (DLT), an allergic reaction, occurred during infusion of 2000 mg, three more patients had to be included in this dose group and tolerated it, as well as the three patients who received 2000 mg for 9 weeks. Occasionally in the dose group 2000 mg mild to moderate fever occurred.

Conclusion: Weekly infusions of 2000 mg of the pine-mistletoe extract were tolerated and can be used in further studies but had a risk for allergic reactions and fever.

Keywords: Cancer, Clinical trial, Intravenous, MTD, Side effects, Viscum album



Cancer Immunotherapy

Oral Presentation

14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Cancer Immunotherapy\Oral\5244

5244

Polyclonal antibody against different extracellular subdomains of HER2 induces tumor growth inhibition in vitro

Reza Hosseini-Ghatar¹, Tahereh Soltantoyeh¹, Motahareh Bahadori², Jalal Khoshnoodi¹, Forough Golsaz-Shirazi¹, Mahmood Jeddi-Tehrani², Mohammad Mehdi Amiri^{1*}, Fazel Shokri^{1,2*}

1. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

2. Monoclonal Antibody Research Center, Avicenna Research Institute, ACECR, Tehran, Iran

Background: Human epidermal growth factor receptor 2 (HER2) has a crucial role in several malignancies and is overexpressed in 20-30% of patients with breast cancer. The extracellular domain of HER2 (HER2-ECD) has been extensively employed as an important target in passive and active immunotherapy. Isolated recombinant prokaryotic HER2-ECD subdomains were previously found to be ineffective to induce anti-tumor antibody response. In this study, we employed recombinant eukaryotic HER2-ECD subdomains to raise anti-HER2 antibodies and to determine their anti-tumor activity in vitro.

Methods: Two paired subdomains of HER2-ECD (DI+II and DIII+IV), which represent Pertuzumab and Trastuzumab binding domains, respectively, along with the full extracellular domain of HER2 were generated in eukaryotic CHO cells. Polyclonal antibodies were raised against these subdomains and characterized using ELISA, flow cytometry, immunoblot and their anti-tumor activity was assessed by XTT assay. The cross-reactivity of these antibodies with other members of the human HER family was also determined.

Results: Anti-DI+II and DIII+IV polyclonal antibodies react with recombinant HER2-ECD and native HER2 expressed on tumor cells similar to Trastuzumab and anti-HER2-ECD antibody. These two polyclonal antibodies were able to inhibit binding of Pertuzumab and Trastuzumab to HER2, respectively, and did not cross-react with other members of HER family. Assessment of the functional activity of these antibodies shows that they inhibit tumor cells growth in vitro, similar to Trastuzumab.

Conclusion: High immunogenicity of human HER2 DI+II and DIII+IV subdomains in rabbits and the tumor inhibitory activity of the purified specific antibodies imply that they might be suitable for active immunotherapy in formulation with appropriate adjuvants and in combination with other HER2 specific therapeutics.

Keywords: Breast cancer, subdomains of HER2, polyclonal antibody, immunotherapy

ICIA2018\Cancer Immunotherapy\Oral\7402

7402

Vaccination with Human Amniotic Epithelial Cells Confer Effective Protection in a Murine Model of Colon Adenocarcinoma

Meraj Tabatabaei¹, Nariman Mosaffa¹, Roya Ghods^{2,3, 4}, Somaieh Kazemnejad⁵, Manijeh Khanmohammadi⁵, Ebrahim Mirzadeghan⁵, Ahmad-Reza Mahmoudi⁴, Mohammad-Reza Bolouri⁴, Reza Falak⁴, Bahareh Keshavarzi¹, Mehrnoosh Ramezani⁶, Amir-Hassan Zarnani^{7,4}

1. Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

2. Oncopathology Research Center, Iran University of Medical Sciences, Tehran, Iran.

3. Department of Molecular Medicine, Faculty of Advanced Technologies in Medicine, Iran University of Medicine Sciences, IUMS, Tehran, Iran.

4. Immunology Research Center (IRC), Iran University of Medical Sciences, Tehran, Iran.

5. Reproductive Biotechnology Research Center, Avicenna Research Institute, ACECR, Tehran, Iran.

6. Department of Biochemistry, school of medicine, Ardabil University of Medical Science, Ardabil, Iran.

7. Reproductive Immunology Research Center, Avicenna Research Institute, ACECR, Tehran, Iran.

Background: The immunologic similarities between cancer development and placentation have helped researchers to unravel molecular mechanisms responsible for carcinogenesis and to take advantage of stem cells from reproductive organs to elicit robust anti-cancer immune responses.

Methods: Human amniotic epithelial cells (hAECs) were isolated and characterized by assessment of their multi-lineage differentiation potential and expression of mesenchymal, hematopoietic and embryonic markers. Their effectiveness as a preventive and therapeutic vaccine in a mouse model of colon cancer was then evaluated by profiling of tumor growth parameters, cross-protective specific CTL and antibody responses, frequency of T cell subsets and cytokine profile.

Results: hAECs expressed markers of mesenchymal and embryonic origin, namely cytokeratin, CD9, CD10, CD29, CD73, CD105, HLA-I, HLA-G, STRO-1, SSEA-4 and OCT-4 with considerable capacity to differentiate toward chondrocytes and osteocytes. Vaccination with both CT26 and hAECs resulted in complete inhibition of tumor development in all vaccinated mice. Similarly, mice vaccinated with hAECs were significantly protected against melanoma, B16F10, challenge. Vaccination with hAECs led to expansion of systemic and splenic cytotoxic T cell population and induced tumor specific cross-protective cellular immunity against CT26 cells with increased splenic IFN- γ production in response to CT26 challenge. Interestingly, cross-reactive antibodies were generated against CT26 cells after vaccination with hAECs. As therapeutic vaccine, hAECs immunization reduced tumor burden leading to expansion of peripheral blood and splenic CD8⁺ T cells and induction of Th1 cytokine profile.

Conclusion: The results of our study clearly demonstrated that hAECs possess the potential for being used as an effective prophylactic vaccine for immune prevention of colon cancer most probably due to the shared antigenic determinants.

Key words: Human amniotic epithelial cells, Cancer vaccine, Colon cancer, Immunotherapy, Anti-tumor immunity

ICIA2018\Cancer Immunotherapy\Oral\7820

7820

Metabolic reprogramming in T cells reduces Treg cells and improves the PD-1 blockade cancer immunotherapy

Maryam Akrami Somea Bozorg, Kenji Chamoto, Tasuku Honjo

Department of Immunology and Genomic Medicine graduate School of Medicine

Background: Tumor microenvironment is profoundly immunosuppressive. FoxP3⁺ regulatory (Treg) T cells represent a major barrier to effective antitumor immunity and immunotherapy. Consequently, there has been considerable interest to target FoxP3⁺Treg cells in tumor. There are various signaling and metabolic pathways to induce instability of FoxP3 expression and to convert FoxP3⁺Treg cells into FoxP3-lost T cells (ex-FoxP3 T cell). Unlike FoxP3⁺Treg cells, the ex-FoxP3 cells express an activated-memory T cell phenotype, and produced inflammatory cytokines. Therefore targeting FoxP3⁺ instability has been potential therapeutic strategy for antitumor immunotherapy. Some reports showed that instability of FoxP3 expression might be associated with PD-1 signaling and metabolic pathways. However, from these aspects, exact connecting pathways between FoxP3 stability and PD-1 blockade is unknown.

Methods: Here we investigated the effect of metabolic modulators in FoxP3 stability during the PD-1 blockade therapy using mice carrying the Foxp3-GFPcre knock-in (KI) locus and Rosa26-tdRFP transgene, which can present ex-Foxp3 cells as a GFP⁺RFP⁺ cells.

Result: We followed the history of Foxp3 expression during the combination therapy of PD-1 blockade and mTOR inhibitor, AMPK inhibitor, HIF-1a inhibitor, and Foxo1 inhibitors. Among them, only Rapamycin reduced the Foxp3⁺ T cells, but not others tested. Interestingly, when Foxp3 was reduced, ex-Foxp3 increased, indicating that Foxp3 cells were converted to ex-Foxp3 by the inhibition of mTOR signaling.

Conclusion: Therefore combination of PD-1 blockade and metabolic modulator targeting Treg is a novel strategy, and our results shed light on improving PD-1 blockade cancer immunotherapy.

Keywords: ex-Foxp3, PD-1, T regulatory

ICIA2018\Cancer Immunotherapy\Oral\7988

7988

Myeloid-Derived Suppressor Cells suppress antitumor immune responses through STAT3 expression in patient with breast cancer

Elham Safarzadeh^{1, 2}, Shahryar Hashemzadeh³, Behzad Mansoori¹, Ali Mohammadi¹, VahidKhazeshahgoli¹, Milad Asadi¹, Elham Bagbani¹, Behzad Baradaran^{1,2*}

1. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

2. Faculty of Medicine, Department of Immunology, Tabriz University of Medical Sciences, Tabriz, Iran

3. General and Vascular Surgery Department of Tabriz University of Medical Sciences, Tabriz, Iran

Background: The capacity of tumors to evade immune-mediated destruction is recently included in the list of hallmarks of tumor development and progression. Given the role of expansion and proliferation of immunosuppressive cells within the tumor microenvironment, these cells remain under investigation in cancers. Myeloid-derived suppressor cells (MDSCs), a heterogeneous population of myeloid cells, are thought to be a major subset of cells that induce an immunosuppressive tumor microenvironment. The purpose of this study was to characterize the different subsets of MDSCs and determine their level and function in the circulation of breast cancer patients.

Methods: We analyzed the frequency of MDSCs in the freshly-isolated peripheral blood mononuclear cell (PBMC) of patients with breast cancer or healthy donor by flow cytometry. These MDSCs were further characterized for phenotype using anti-human monoclonal fluorescently-labeled antibodies. Then effects of MDSCs on the CD3+T cell response were evaluated. STAT3 is considered as an important transcription factor for MDSC expansion and function, so we assessed the effect of inhibition of STAT3 signaling on MDSC suppressive activity.

Results: Our results showed significant increases in circulating HLA-DR – CD33+ MDCSs in breast cancer patients when compared with healthy donors. Moreover, co-culture experiments of magnetically purified HLA-DR – CD33+ MDCSs and CD3+ T cells indicated that accumulation of MDSCs was associated with defective T cell function. Of note, STAT3-targeted siRNA abolished MDSC's suppressive activity.

Conclusion: This finding suggest that MDSCs play an important role in breast cancer and future extensive validation studies and progress in this field may help identify new immunotherapeutic strategies to inhibit MDSC function.

Keywords: Breast cancer, Myeloid-derived suppressor cells, Immunosuppressive cells, tumor microenvironment

ICIA2018\Cancer Immunotherapy\Oral\8076
8076

Inhibition of tumor growth by mouse ROR1 specific antibody in a syngeneic mouse tumor model

HadiHassannia¹, Mohammad Mehdi Amiri^{1*}, FarhadJadidi-Niaragh², Reza Hosseini-Ghatar¹, Jalal Khoshnoodi¹, Ramezan-Ali Sharifian³, Forough Golsaz-Shirazi¹, Mahmood Jeddi-Tehrani⁴, Fazel Shokri^{1,4*}

1. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran.

2. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran.

3. Imam Khomeini Hospital, Tehran University of Medical Sciences, Tehran, Iran.

4. Monoclonal Antibody Research Center, Avicenna Research Institute, ACECR, Tehran, Iran.

Background: Immunotherapy with tumor-associated antigens (TAAs) is a potentially powerful approach to eradicate tumor cells. The receptor tyrosine kinase-like orphan receptor 1 (ROR1) plays a crucial role for survival of tumor cells and is overexpressed in various malignancies. In the present study, we developed a syngeneic mouse tumor model to assess anti-tumor effect of mouse ROR1 specific polyclonal antibody (pAb) in vivo.

Methods: Mouse ROR1 specific antibody was produced in rabbit using recombinant ROR1 protein. Two mouse tumor cell lines, (4T1 and CT26), were transfected with full length mouse ROR1 construct and stable clones were selected and characterized by immunocytochemistry, Western blot and flow cytometry. In vitro and in vivo anti-tumor activities of anti-ROR1 antibody were assessed by XTT and syngeneic BALB/c mouse model, respectively.

Results: We successfully established two mouse ROR1-overexpressing tumor cell lines. The in vitro results indicate that the ROR1p Ab did not significantly inhibit growth of ROR1+ cell lines. One of these cell lines (CT26-ROR1) was implanted in syngeneic BALB/c mice to assess anti-ROR1 tumor inhibitory activity in vivo. The tumor size was significantly reduced in mice treated with ROR1p specific Ab.

Conclusion: Our results demonstrated for the first time tumor inhibitory effect of mouse ROR1 specific antibody in a syngeneic mouse tumor model. This model is a promising tool for preclinical assessment of ROR1 therapeutics and investigation of the underlying molecular mechanisms.

Keywords: ROR1, syngeneic tumor model, anti-ROR1 antibody, tumor-associated antigen

ICIA2018\Cancer Immunotherapy\Oral\8385

8385

Evaluation of anti-tumor responses by combination therapy with oxaliplatin and anti-PD-L1 antibody in a mouse model of colorectal carcinomaSoheila Golchin¹, Seyed Amir Jalali²*1. Department of Immunology, School of Medicine Shahid Beheshti University of Medical Sciences**2. Department of Immunology, School of Medicine Shahid Beheshti University of Medical Sciences*

Background: Chemotherapy has been widely used in cancer treatment. Chemotherapy drugs remain the backbone of current treatment, but are limited by a narrow therapeutic index, significant toxicities, and frequently-acquired resistance. Oxaliplatin (OXP), a platinum-based chemotherapeutic agent, can promote a favourable immune microenvironment and stimulate anti-cancer immune responses. Exposure to OXP markedly increased expression of the T-cell inhibitory molecule programmed death receptor-ligand 1 (PD-L1). Overexpression of immune checkpoint molecules affects tumor-specific T-cell immunity in the tumor microenvironment, and can reshape tumor progression and metastasis. Many tumors can upregulate PD-L1 expression, thus avoiding immune surveillance and elimination. Many molecularly-targeted agents have not shown curative properties when administered as monotherapies. With that in mind, the current opinion is that combining molecularly-targeted anticancer therapy with conventional cytotoxic chemotherapy will improve treatment outcomes. Important considerations regarding optimizing dosage, sequence, and timing of targeted therapies will be required when designing future clinical treatments.

Methods: To determine whether combined treatment affects the local immune cell populations we evaluated the anti-tumor activity of anti-PD-L1 combined with OXP against a murine colon carcinoma in vivo and examined the tumor immune microenvironment by flowcytometric analysis of immune cells from the tumor site, lymph nodes, and spleen.

Results: Our study showed that the in-vivo administration of OXP combined with anti-PD-L1 markedly inhibited the growth of murine colorectal carcinoma tumors in mice. The anti-tumor effect of anti-PD-L1 plus OXP correlated with a marked increase in the number of tumor-infiltrating activated CD8⁺ T cells and a marked decrease in the number of regulatory T cells in drainage lymph nodes and spleen.

Conclusion: Mice treated with anti-PD-L1 combined with OXP had smaller tumors and greater survival rates than mice receiving either anti-PD-L1 or OXP alone. We also found that the timing of drug administrations affected tumor responses.

Keywords: Anti PD-L1, combination therapy, cancer, immunotherapy, chemotherapy, timing

ICIA2018\Cancer Immunotherapy\Oral\8533

8533

Combined blockade of PD-1 and Tim-3 immune checkpoint receptors to restore the function of exhausted CD8⁺ T cells in chronic lymphocytic leukemia

Hadiseh Rezazadeh¹, Hadi Hossein-Nataj¹, Ramin Shekarriz², Mohsen Tehrani¹, Hossein Asgarian-Omran^{1,3*}

1. Department of Immunology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

2. Gastrointestinal Cancer Research Center, Mazandaran University of Medical Sciences, Sari, Iran

3. Immunogenetics Research Center, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

Background: Although, common treatment strategies have improved overall survival in chronic lymphocytic leukemia (CLL), majority of patients continue to relapse and therefore finding a new strategy seems to be necessary. Blocking antibodies against programmed death-1 (PD-1) immune checkpoint receptor have been approved by FDA for treatment of many malignancies. But, in recent studies it was demonstrated that PD-1 blockade alone cannot completely restore the function of exhausted T cells. In the current in vitro survey, combinatorial blockade of PD-1 and T cell immunoglobulin- and mucin-domain-containing molecule-3 (Tim-3) regulatory pathways were done to restore the function of exhausted CD8⁺ T cells in CLL patients.

Methods: Peripheral blood mononuclear cells were separated from fifteen patients with CLL and CD8⁺ T cells were positively isolated using magnetic beads separation method. Purified CD8⁺ T cells were treated with either blocking antibodies against PD-1 and Tim-3 and isotype matched control antibodies. Treated CD8⁺ T cells were stimulated with anti-CD3/CD28 antibodies and PMA/Ionomycin to assess their proliferation capacity and cytokines production by MTT assay and ELISA, respectively. Concentrations of IL-2, IL-10, IFN- γ , and TNF- α were measured in culture supernatants. Degranulation properties of CD8⁺ T cells were also assessed by CD107a degranulation assay.

Results: Our results showed that combined blockade of PD-1 and Tim-3 by inhibitory antibodies reversed the effector functions of exhausted CD8⁺ T cells and improved their proliferation, cytotoxicity and production of pro-inflammatory cytokines.

Conclusion: Considering potential roles of PD-1 and Tim-3 molecules in T cells exhaustion, obtained results from this study in restoring the function of exhausted CD8⁺ T cells could be a potential support in formation of a new treatment option for patients with CLL.

Keywords: Chronic lymphocytic leukemia, exhausted CD8⁺ T cells, Tim-3, PD-1, immunotherapy



Computational Immunology and Systems Biology

Oral Presentation

14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Computational Immunology and Systems Biology\Oral\5311

5311

Modeling the tumor growth and production of anti-tumor cytotoxic T lymphocyte after deletion of Myeloid-Derived Suppressor Cells using Differential equation of system

Zahra Mirsanei¹, Nassim Kheshtchin¹, Samaneh Arab¹, Nazanin mojtavavi², Jamshid Hadjati^{1,*}

1. Department of Immunology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

2. Department of Immunology, School of Medicine, Iran University of Medical Sciences, Tehran, Iran

Background: The tumor microenvironment is a complex tissue that is participated in anti-tumor activity or suppress antitumor immune responses. The ability of MDSCs (Myeloid-Derived Suppressor Cells) to inhibit the cytotoxic immune responses mediated by NK cells and CTL led to the elimination of these cells as a target for immunotherapy that increases the efficiency of these methods. For eliminate MDSC's inhibitory mechanisms, the use of drugs is effective.

Method: 4T1 cells were implanted subcutaneously in right flank of BALB/c mice in two groups: The control group contains 15 mice without any treat and the test group contain 15 mice, which treated by 5-fluorouracil (5-FU) from 5th day (when the tumor was tangible for the first time) by 3 days interval. Tumor growth in the course of treatment and survival of mice after that was followed by measuring tumor size and Immunohistochemistry (IHC) assay was used for counting tumor infiltrating CD8⁺ cells as a result of MDSCs elimination.

Result and conclusion: A class of mathematical models called Ordinary Differential Equation (ODE) model was developed using our aquired empirical data to analyze and simulation the behavior of the immune system in the tumor microenvironment. We show that our model is capable of capturing the observed experimental results, and hence can be potentially used in designing future experiments involving this approach to immunotherapy.

KeyWord: Tumor Modeling, MDSC, ODE

ICIA2018\Immunology of Chemical Victims and Environmental Pollution\Oral\8418
8418

The study of particulate matter effects on Alveolar macrophages inflammatory function in C57BL/6 mice

Hoda Rahmani¹, Somayeh Sadeghi ², Mohsen Roshani³, Davar Amani⁴, Tooba Ghazanfari⁵, Nariman Mosaffa⁶

1. Department of immunology, School of medicine, Shahid Beheshti medical university

2. Department of immunology, School of medicine, Shahid Beheshti medical university

3. Tehran air quality standard agency

4. Department of immunology, School of medicine, Shahid Beheshti medical university

5. Department of immunology, School of medicine, Shahed medical university

6. Department of immunology, School of medicine, Shahid Beheshti medical university

Background: The presence of ambient particulate matter poses more dangers to human health than that of other common air pollutants such as CO₂ and ozone. Epidemiologic studies show the direct correlation between PM and risk of respiratory and cardiovascular disease. Immune system seems to play a critical role in the process of these diseases. Alveolar macrophages represent major population of immune system cells keeping the lung alveoli clean from pathogens and environmental agents. We studied the effect of PMs in two aerodynamic diameters (PM_{2.5} and PM₁₀) on Alveolar macrophages and evaluate the stimulatory effects on cytokine release in culture medium.

Methods: PM collector filters of different station were gifted from Tehran air quality standard agency. Harvested particles (2.5&10µm) were suspended in RPMI at a concentration of 5 µg/ml and 10 µg/ml, and then added to alveolar macrophage culture of C57BL/6 inbred mice. After 24h incubation, vitality of cells was evaluated with MTT test. IL10 and TNF-alpha production were measured in supernatants of cultured cells using enzyme-linked immunosorbent assay (ELISA) kits, after 6, 12 and 24 hour incubation.

Results: viability of PM treated alveolar macrophage significantly increased in comparison with controls. PM_{2.5} activated macrophages vitality significantly increases in comparison with PM₁₀. Also PM caused TNF-alpha production increased after 6, 12 and 24 hours incubation. TNF-alpha reached peaked concentration after 6 hour incubation with pm_{2.5} treatment Pv<0.05. IL10 production decreased after 6 and 12 hours incubation with pm_{2.5} in comparison with control. Pv<0.05. After 24 hours treatment the level of IL10 began increasing.

Conclusion: In this study, PM_{2.5} induces higher inflammatory response in comparison to PM₁₀. In immunophysiological status of lung environment, the inflammatory cytokines such as TNF-a and down regulation of regulatory cytokine (IL10) may harmful for respiratory health and leads to many acute and chronic disorders. Another size of PM our study suggests that PM₁₀ may cause toxic effects on alveolar macrophages as we detected in vitality assay.

Keywords: Particulate matter, Alveolar macrophages, MTT, TNF-alpha, IL10

ICIA2018\Computational Immunology and Systems Biology\Oral\7950

7950

Game Theory Modeling as a Novel Methodology to Describing the Host-Microbe Interactions: Mycobacterium Tuberculosis (Mtb) and Host Immune Defense as an Example

Hiva Sharebiani¹, Shadi Abbasnia¹, Sara Hajimiri¹, Rezvan Pilpa¹, Arezo Firoozeh¹, SamanSoleimanpour², Kiarash Ghazvini², S.A Rahim Rezaee¹

1. *Inflammation and Inflammatory Diseases Research Center, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran.*

2. *Microbiology & Virology Research Center, Bu-Ali Research Institute, Mashhad University of Medical Sciences, Mashhad, Iran.*

Background: The interactions between host and microbe determine the outcome of infections. To understand the underlying pathogenesis of infectious diseases, epigenetic study and then the game theory models which describes the outcome (pay-off) from specific interactions (game) between two individuals (players) were considered. In this study, Mtb and host interactions in tuberculosis manifestation were evaluated as a model.

Methods: The virulence factors (Ag85A, CFP-10 and ESAT-6) and main host defense strategies (CCR1, CCR2, T-bet, iNOS, IDO, TGF- β , MMP3 and 9) were evaluated by real-time-PCR, TaqMan method. Then using these assessing interactions modeled as a dynamic game with complete and perfect information. The pay-off matrix is designed based on gene expression of host and microbe using the Naive Bayes classifier (NBC) by Rapid minerV5.3 software. Extensive form of game modeled via backward induction and detection of SPE strategies by Min-Max method to determination of Nash equilibrium.

Results: Host and Mtb strategies evaluated by Pareto and PSPE efficiency indices as appropriate response markers and classified to dominant, dominated and optimal strategies. Cooperative and grim trigger strategies was defined. Therefore, the mechanisms of leaving the latency or remaining in the same situation, were determined. The optimum levels of gene expression as the best strategies of each player were: Mtb high expression of Ag85 and high level of CCR1 and medium level of CCR2, T-bet, iNOS, IDO and TGF- β that confirmed by MMPs level and patient's outcome as the host protective immune response. The role of each gene in this interaction was investigated. Gene expression changes of TGF- β /IDO, T-bet/iNOS and CCR1/CCR2 were responsible for reactivation or remaining in latency, response to CFP-10: ESAT-6 or Ag85A and patient's outcome, respectively.

Conclusion: A rational way for assessing interaction outcomes for vaccine design, therapeutic strategy or prognosis of an infection is epigenetics studies of the players in a time course and then evaluation of different outcomes by game theory model.

Keywords: Game theory, Nash equilibrium, Interaction, Epigenetic.

ICIA2018\Computational Immunology and Systems Biology\Oral\7971
7971

Bioinformatics approach to identifying molecular biomarkers in multiple sclerosis

Mina Rezaei¹, Shirin Farivar^{2*}

1. Department of cellular and molecular biology, shahid Beheshti University, Tehran, Iran

2. Department of cellular and molecular biology, shahid Beheshti University, Tehran, Iran

Background: Multiple sclerosis (MS) is a chronic autoimmune, inflammatory neurological disease of the central nervous system (CNS). Considering the complexity of its etiopathogenesis, early diagnosis of MS and individualized management are challenging in clinical practice. It is also difficult to obtain CNS samples for diagnostic purposes. Therefore, studies on biomarkers in MS are useful for early prediction and diagnosis, monitoring of disease activity and predicting treatment response.

The present study aimed to analyze the molecular mechanism of MS using microarray analysis combined with bioinformatics techniques.

Methods: We downloaded the gene expression profile of MS from NCBI Geo datasets, Gene Expression Omnibus (GEO) and analyzed the microarray data. Genes are compared with logfc for their expression in normal controls and MS patients and sorted by their p-value.

Result: Finally gene with most modification in multiple sclerosis is selected for further analysis. Also its most related pathways and gene function are identified by DAVID database and Gene Card database and its protein structure such as secondary and domains are identified through ExpASy database.

Conclusion: The result suggested some genes may play central regulatory roles in controlling gene expression in the pathogenesis of MS. Our findings confirm the presence of multiple molecular alterations in MS and indicate the possibility for identifying prognostic factors associated with MS pathogenesis.

Keywords: multiple sclerosis, microarray analysis, RNA and protein structure, Gene expression

ICIA2018\Computational Immunology and Systems Biology\Oral\7980
7980

Dissection of neutrophils responses to sepsis via network analysis: From weighted co-expression networks to protein-protein and gene regulatory networks

Rasoul Godini¹, FatemehHadi¹, Hossein Fallahi¹

1. Department of Biology, Faculty of Science, Razi University

Background: Infection of blood leads to a clinical condition known as sepsis. Sepsis comes with systemic dysregulation of inflammation. Neutrophils are important immune cells in blood that move to the infection site, catch and digest infectious agents. The importance of neutrophils in sepsis has been shown. Herein, we analysed neutrophil responses to sepsis condition through network based bioinformatics.

Methods: Free available dataset GSE49758 was obtained from NCBI. After preprocessing of the genes for adequate variable ones, the matrix analyzed through Weighted Gene Co-expression Network Analysis (WGCNA) method using R program. Septic samples were compared against normal ones to determine differentially expressed genes (DEGs). Using DEGs, the Protein-Protein Interaction (PPI) and Gene Regulatory Network (GRN) were constructed by STRING and ChEA databases, respectively. Visualizing and analyzing of the networks were performed by Cytoscape and Gephi programs software.

Results: WGCNA showed that there are 3 module eigengenes which are related to sepsis with a correlation of 50 percent (p-value less than 0.01). Module membership and gene significance correlation in the detected modules was at least 0.45 (p-value = 3.3e-10). The members of modules were related to inflammatory responses, protein transport and signal transduction. For other types of networks, we obtained a list containing 633 DEGs that resulted in a PPI network constructed of 260 nodes and 524 edges. We detected 9 sub-networks by overlap neighborhood expansion methods in PPI. Different hubs were detected in this method such as SMAD3 and IL1B. GRN was constructed by two differentially expressed transcription factors including BCOR and SMAD3 which regulates 79 and 95 target genes, respectively.

Conclusion: In this study, we comprehensively analyzed responses of neutrophils to sepsis by various network analysis. The results determined genes and molecular processes which could be focused on in further studies.

Keywords: Differentially expressed genes, Protein-protein interaction, Sepsis, WGCNA

ICIA2018\Computational Immunology and Systems Biology\Oral\8463

8463

The power of whole exome sequencing in detection of causative mutation in a lost primary immunodeficient baby based on the whole exome data of its healthy parents

Ali Aghebati Maleki¹, Majid Changi-Ashtiani², Taravat Talebi¹, Neda Mohsenpour¹, Mohammad Shahrooei^{3,4}, Alireza Biglari¹, Tina Shahani¹, Hassan Rokni-Zadeh⁵

1. Department of Genetics and Molecular Medicine, School of Medicine, Zanzan University of Medical Sciences (ZUMS), Zanzan, Iran

2. School of Mathematics, Institute for Research in Fundamental Sciences (IPM), Tehran, Iran.

3. Experimental Laboratory Immunology, Department of Microbiology and Immunology, KU Leuven, Leuven, Belgium

4. Specialised Immunology Laboratory of Dr. Shahrooei, Ahvaz, Iran

5. Department of Medical Biotechnology and Nanotechnology, School of Medicine, Zanzan University of Medical Sciences (ZUMS), Zanzan, Iran

Background: Whole Exome sequencing (WES) is used to efficiently detect common single-nucleotide variants (SNVs) across a broad spectrum of applications such as identification of rare genetic variants that are presumably responsible for complex genetic diseases. Primary immunodeficiency diseases (PIDD) are a group of more than 300 rare, genetic and chronic disorders that the immune system functions improperly. In this study we aimed to find the causative mutation(s) by WES, performed on healthy parent samples of a lost PIDD baby; among the dozens of Iranian families whose children had died due to PIDD and routine diagnostic procedure have not been able to determine the cause of the illness.

Methods: After selecting the appropriate pedigree, extracting gDNA was performed from the parents' blood. Samples were sent to WES and then data were analyzed to find common heterozygous mutations associated with patient phenotype. In order to confirm the results, the candidate mutation segregation in the family was examined by Sanger sequencing method.

Results: Variant calling presented about sixty thousand variants in parent data which differed from the human reference genome. Of these variants, we reached two hundred heterozygous common variants in parent. Eventually, only one strong candidate variant associated with the disease phenotype was considered according to WES pipeline as the only probable cause of the disease that confirmed by sanger sequencing too. Finally, homozygous transversion of G>T at exon 33 of Dock8 gene was identified as causative mutation at lost baby.

Conclusion: Here, we present reliability of WES to infer genomic variants in disorders with genetic heterogeneity, particularly, for detection of causative mutation from healthy parent genomic data. This approach will improve management of PID families when the proband sample is not available. Due to life instability of such patients, this approach is of high value.

Keywords: Primary immunodeficiency, Whole Exome Sequencing, Carrier detection

ICIA2018\Computational Immunology and Systems Biology\Oral\9920
9920

Determination and Ranking of Involved Signaling Pathways in Systemic Lupus Erythematosus Using PBMC Transcriptome Profile by Bayesian Network Approach

Samaneh Maleknia¹, Dr.Ali Shirifi Zarchi², Dr.Kaveh Kavousi¹

1. *Department of Bioinformatics Institute of Biochemistry and Biophysics (IBB); Tehran University (UT)*
2. *Department of computer engineering, Sharif University of Technology*

Background: In recent years, studying immune system related diseases has a great importance in medicinal research. According to these studies, several specific biological pathways are influenced by different types of disordered immune systems. Many of these studies focus on autoimmune diseases. Computational methods have pivotal roles in generating results and making new discoveries. The aim of this study is to nominate signaling pathways that are specially related to “systemic lupus erythematosus” (SLE) with Bayesian network approach.

Methods: The first phase of this study involves performing enrichment analysis on 5 datasets of “PBMC” cells gene expression, with both case and control groups for each dataset. All datasets were downloaded from NCBI. Enrichment analysis was performed based on best-curated signaling pathway databases like KEGG, Reactome, and wiki pathway. More than 20 effective signaling pathways were identified with a significance level of less than 0.05. In the next step, to identify the most affected genes and network modules, Bayesian networks were inferred from involved pathways. These networks were trained using gene expression data.

Results: According to the results, most of these pathways are related directly to the immune system. Sufficient data must be provided to permit evaluation by the reviewers and public reading of the abstracts. Statements such as “additional information to be presented at the meeting” are not acceptable.

Conclusion: Finally, in the current state, the most influenced subnetworks of pathways are nominated, the most influenced genes (by this approach) were ranked and a validation process was used to select the most related ones.

Keywords: Systems biology, Autoimmunity, Bayesian network, Systemic lupus erythematosus

ICIA2018\Computational Immunology and Systems Biology\Oral\8471
8471

Immunology of Glioblastoma

FatemeShaabanpour Aghamaleki¹, Shirin Farivar²

1. Department of molecular and cellular Biology School of Biotechnology and Life Science University of Shahid Beheshti

2. Department of molecular and cellular Biology School of Biotechnology and Life Science University of Shahid Beheshti

Background: glioblastoma is a malignant, high grade and a progressive brain tumor that affects both children and adults. It has a multifactorial pathogenesis such as genetics, epigenetics and environmental factor. But it has genetics originally in 10 percent of pediatric patients. Among this genetics factor, immune system genes may play an important role in glioblastoma pathogenesis, progression, and metastasis. Therefore, identification of these genes can help select the best therapy for glioblastoma.

Methods: Gene expression profiling by array of 12 glioblastoma patients and 3 healthy individuals were extracted from Geo datasets. Genes are compared with logFC for their expression between two groups and sorted by their p-value. Also, the most related pathway, location, function, and protein networks of this gene were identified with STRING, GeneCard and DAVID databases.

Results: several important immune genes associated with glioblastoma tumors are identified such as CXCL2, NFATC1, TGFB2, PTPN9, FCGR1, XCL1, CCL4 and etc. These genes have hyperexpression in glioblastoma patients. According to DAVID database, all of the immune genes that have hyperexpression in glioblastoma play important roles in the inflammatory response, cytokine and growth factor. Also, their related pathways are chemokine, TNF signaling pathways, and G-protein coupled receptor (GPCR) signaling pathway.

Conclusion: Although, there are a wide variety of immune pathways and genes that play important roles in many diseases, some of these pathways and genes are important and specific in some diseases such as glioblastoma. Therefore, identification of these critical and specific immune genes in glioblastoma can lead to appropriate targeting in immunotherapy and chemotherapy.

Keywords: glioblastoma, immune system, glioblastoma immunology, glioblastoma therapy



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Diagnostic Methods in Immunology

Oral Presentation

ICIA2018\Diagnostic Methods in Immunology\Oral\7399
7399

IL-10 as a potential predictor of grave prognosis in premature infant

Farshid Saadat¹, Seyedeh Zohreh Jalali²

1. Department of Immunology, School of Medicine, Guilan University of medical sciences, Rasht, Iran

2. Department of Neonatology, School of Medicine, Guilan University of medical sciences, Rasht, Iran

Background: Prematurity is the direct cause of neonatal death worldwide. To increase survival rates, mechanical ventilation with highly inspired oxygen, corticosteroids and exogenous surfactant therapy are used. The present study was designed to explore the outcome in premature neonate based on the anti inflammatory cytokine levels.

Methods: This was a prospective, randomized investigation of infants who were less than 37 weeks of gestational age. Blood samples were collected on day 1 and plasma cytokine was assayed for interleukin (IL)-10.

Results: This study showed that the plasma IL-10 levels stand as a promising biomarker for predicting prognosis. Our findings show that the higher IL-10 levels statistically associated to poor outcome after excluded other confounders. ($P<0.001$).

Conclusion: Although, further studies are required to elucidate the mechanistic role of IL-10, it might be considered as a valuable marker for outcome assessment.

Keywords: Prematurity, interleukin-10, prognosis

ICIA2018\Diagnostic Methods in Immunology\Oral\7689

7689

Developing a specific anti-TAZ (WWTR1) Polyclonal Antibody

Mahsa Mohammadzadeh¹, Fatemeh Ghaemimanesh², Mehrdad Hashemi¹, Babak Negahdari^{2,3}, Ali Ahmad Bayat², Reza Hadavi², Saeideh Milani², Hodjatallah Rabbani²

1. Department of Genetics, Islamic Azad University, Tehran Medical Sciences Branch, Tehran, Iran

2. Antibody-Antigen Engineering Department, Monoclonal Antibody Research Center, Avicenna Research Institute, ACECR, Tehran, Iran

3. Department of Medical Biotechnology, School of Advanced Science in Medicine, Tehran University of Medical Science, Tehran, Iran

Background: TAZ (WWTR1) is a transcriptional co-activator protein in Hippo signaling pathway which plays a pivotal role in organ size control during embryogenesis. TAZ has been shown to be overexpressed in various human carcinomas such as breast, hepatocellular and lung cancers. The aim of this study was to produce a specific polyclonal antibody (pAb) against TAZ to be used as a research tool for further investigation of TAZ characteristics in human carcinomas.

Methods: A white New Zealand rabbit was immunized using a Keyhole limpet hemocyanin conjugated synthetic peptide of TAZ. Following the raise of antibody titration, the developed polyclonal anti-TAZ antibody (pAb anti-TAZ) was purified using affinity chromatography. After determining the specificity of pAb in ELISA, its reactivity with TAZ molecule was assessed in different cancer cell lines by various immunoassays.

Results: The produced antibody could recognize the immunizing peptide in ELISA. PAb anti-TAZ was able to specifically recognize TAZ in human carcinoma cell lines containing A431, MCF7 and Raji in Western blot, flow cytometry and ICC. Immunohistochemistry results showed that the developed pAb detected TAZ in prostate and bladder carcinoma tissues.

Conclusion: The developed anti-TAZ pAb might be used as a research tool in expression profiling of many cancer cells with application of Western blot, flow cytometry, ICC and IHC.

Keywords: Polyclonal Antibody, TAZ, cancer, detection

ICIA2018\Diagnostic Methods in Immunology\Oral\8069
8069

Evaluation of Phagocytosis in Human Neutrophils using Enhanced Green Fluorescent Protein (EGFP) expressing *E. coli*

Hosain AqaHosaini, Sayyed Hamid Zarkesh Esfahani

1. Department of Biology, Faculty of Sciences, University of Isfahan, Iran

Backgrounds: Phagocytosis plays a very important role in innate immunity and helps body against bacterial infections. Patients who have defect in phagocytosis suffer from recurrent bacterial infections that may be life threatening. It is important to detect the defect in phagocytosis as early as possible in life. Patients who receive immunosuppressive drug may also have suppressed phagocytosis. There are different tests for evaluation of phagocytosis such as NBT and DHR which use chemical compounds not real bacteria to test the phagocytosis. Bacteria such as *E. coli* can be transformed to express Enhanced Green fluorescent protein (EGFP).

To introduce and evaluate a method for testing Neutrophil's phagocytosis using real bacteria.

Methods: EGFP sequence was cloned in an expression vector called pColdI using the enzymes NdeI and XhoI. BL21 strain of *E. coli* was transformed by pColdI containing EGFP. Presence of the EGFP sequence was confirmed by PCR and sequencing. The expression of EGFP in bacteria was detected by fluorescent microscope and Flow cytometry. EGFP expressing bacteria were added to heparinized whole blood of either healthy volunteers or patients receiving immunosuppressive medicines. The engulfment and digestion of fluorescent bacteria was detected by flow cytometry at different time points.

Results: The *E. coli* transformed with vector containing EGFP expressed high levels of green fluorescence as detected by both fluorescence microscope and flow cytometry. Neutrophils that engulfed fluorescent bacteria became highly fluorescent as could be detected by flow cytometry. Digestion of bacteria during the time resulted in decrease in Neutrophil's fluorescence.

Conclusion: EGFP expressing bacteria and flow cytometry technique can be used to evaluate the phagocytosis capability of neutrophils.

Key words: phagocytosis, pColdI, EGFP, *E. coli*, fluorescent, flow cytometry.

ICIA2018\Diagnostic Methods in Immunology\Oral\8088
8088

Severe Decline in Mir-20a and Mir-92a in the Context of the Mir-17-92 Cluster: Ideal Biomarkers of Various COPD Subtypes

Akbar Ghorbani, Alvanegh Mahmood Tavallaei, Houri Edalat^{1*}

1. Human Genetics Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran

Background: chronic obstructive pulmonary disease (COPD) is going to be the third leading cause of death by 2020. Spirometry is laborious that results in a great under-diagnosis of COPD. On the other hand, circulating miRNAs are among the most beneficial feasible non-aggressive biomarkers for the diagnosis and treatment of many diseases. Among members of the significant miR-17-92 cluster, miR-20a and miR-92a are greatly involved in both inflammation and hypoxia (that are known as the main reasons for COPD comorbidities).

Methods: Thus, the expression of these miRNAs was evaluated in 26 COPD patients and 19 normal individuals by a highly sensitive approach called stem-loop RT-qPCR. An explanation of the study design and experimental method.

Results: The expressions of the studied miRNAs were significantly reduced in the serum of the COPD patients relative to that of the controls ($p < 0.001$).

Conclusion: This finding may be attributed to the susceptible genetic background of the COPD group and/or the epigenetic changes caused by smoking. Altogether, our findings indicated that the clusteral expression pattern of miR-17-92 reflects the cluster's ideal position as a biomarker of the pathogenesis of COPD subtypes. Decreased expression of miR-20a in a patient might reflect a progressive stage of the disease. MiR-92a might be used as an early-detection biomarker of COPD. A higher reduction level for miR-92a compared with miR-20a might reveal the chronic nature of the disease. The miRNAs can be used as therapeutic targets specifically in the context of the miR-17-92 cluster to address the various clinic pathological aspects of the disease.

Keywords: COPD, biomarker, treatment, serum, inflammation, stem-loop real-time quantitative PCR (stem-loop RT-qPCR)

ICIA2018\Diagnostic Methods in Immunology\Oral\8233
8233

Total PSA tumor marker detection based on nano-bio-optical sensor

Zahra akbari jonoush¹, Reza Mansouri ¹, Fatemeh Yazdiyan ²

1. Department of Immunology, Shahid Sadoughi University of Medical Sciences & Health Services, Yazd,
2. Faculty of New Science and Technology, University of Tehran, Iran

Background: Nowadays, early cancer detection is crucial for improved prognosis and cancer management. Therefore, identification of sensitive and specific methods for early cancer detection is required. One of the methods for early cancer detection are biosensor and gold nanoparticles.

Methods: Herein, antibody-coated gold nanoparticles were employed in optical immunoassay for detecting total PSA biomarker. To achieve this goal, antibody was covalently conjugated to gold Nano spheres using 11-Mercaptoundecanoic acid (MUA). Finally, serum sample was investigated by localized surface Plasmon resonances. The results obtained by LSPR and Chemiluminescence assay were compared with each other.

Results: The results suggested that Nano bio-optical sensor had good potential and good sensitivity in detecting of total PSA. The proposed Nano bio-optical sensor delivered a good sensitivity with average sensitivity 1.1 nm/ng ml⁻¹, and a low detection limit of 0.9 ng/ml

Conclusion: These results indicated that the Nano bio-optical sensor has a great potential in diagnosis of cancer biomarkers.

Keywords: Nano bio-optical sensor, total PSA biomarker, total PSA monoclonal antibody

ICIA2018\Diagnostic Methods in Immunology\Oral\8656

8656

Determining laboratory reference values of TREC and KREC in different age groups of Iranian healthy individuals

Leila Shakerian¹, Zahra Pourpak¹, Somayeh Shamlou¹, Erna Domsgen², Anoshirvan Kazemnejad³, Hossein Dalili⁴, Lennart Hammarstrom², Maryam Nourizadeh¹

1. Immunology, Asthma and Allergy Research Institute, Tehran University of Medical Sciences, Tehran, Iran

2. Department of Laboratory Medicine, Karolinska Institutet, Karolinska University Hospital Huddinge, Stockholm, Sweden

3. Department of Biostatistics, Tarbiat Modares University, Tehran, Iran

4. Breastfeeding Research Center, Tehran University of Medical Sciences, Tehran, Iran

Background: Assessment of T-cell receptor excision circles (TREC) and kappa-deleting recombination excision circles (KREC) copies has been recently described as biomarkers of thymus and bone marrow's functions, respectively. In this study, first we aimed to explore the effects of demographic variables including age, sex, race as well as weight and height at birth on these two episomal molecules. Secondly, for the first time in our country, we determined the reference values of TREC and KREC copy numbers in different age groups of Iranian healthy individuals as a thresholds for identifying the T cell and B cell lymphopenia.

Methods: TREC and KREC copy numbers were evaluated in 251 dried blood spot (DBS) samples from healthy volunteers (age range: 0-60 years), 11 premature newborns (age range: 30-35 weeks) and also six primary immunodeficiency (PID) patients as disease controls using multiplex quantitative real time PCR.

Results: TREC and KREC copies were notably reduced with increasing the age. We could not find any significant differences between male and female as well as among various ethnicities in Iran.

Conclusion: These finding suggest that demographic variables including age as important interfering factor, should be considered for interpretation of TREC/KREC results.

Keywords: T-cell receptor excision circles, kappa-deleting recombination excision circles, primary immunodeficiency disorders.

ICIA2018\Diagnostic Methods in Immunology\Oral\9813
9813

Development and evaluation of specific scFv antibody against human dentin sialophosphoprotein

Seyed Nooreddin Faraji^{1,3}, Mozafar Mohammadi^{3,4}, Foroogh Nejatollahi^{2,3}

1. School of Advanced Medical Sciences and Technologies, Shiraz University of Medical Sciences, Shiraz, Iran

2. Shiraz HIV/AIDS Research Center, Institute of Health, Shiraz University of Medical Sciences, Shiraz, Iran.

3. Recombinant Antibody Laboratory, Department of Immunology, Shiraz University of Medical Sciences, Shiraz, Iran

4. Applied Biotechnology Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran

Background: Prostate cancer (PCa) is the second leading cause of cancer related death in men. Diagnosis of PCa faces many challenges especially its biomarker-based detection methods. Dentin sialophosphoprotein (DSPP) has been shown with higher specificity and sensitivity for PCa diagnosis compared to current biomarkers.

Methods: A single chain antibody (scFv) against DSPP was isolated using the panning procedure. PCR and DNA fingerprint were done following 4th round of panning. The reactivity of the isolated scFv to DSPP was evaluated by phage ELISA on PCa urines (number: 10) and compared to healthy individuals (number: 10). Following plasmid extraction and DNA sequencing, the 3D structure of isolated scFv was simulated base on homology modeling. Docking of the isolated scFv on DSPP was also simulated to investigate the antigen-antibody interaction.

Results: The frequency of isolated scFv against DSPP was 40%. Phage-ELISA results showed significant difference between PCa patients (OD: 0.92) and healthy individuals (OD: 0.34). Sequence analysis of the isolated scFv showed several amino acid substitutions in CDRs and FRs of both VH and VL. Docking results showed that the scFv can tightly bind to the DSPP by several interaction especially by hydrogen bonds between antigen and antibody interface.

Conclusion: The results of the new isolated anti DSPP-scFv antibody evaluated by experimental and in silico techniques, introduce a specific and sensitive antibody for designing DSPP-based PCa detection methods.

Keywords: Dentin sialophosphoprotein (DSPP), single chain antibody (scFv), prostate cancer (PCa), panning, docking



Immune Cell Therapy

Oral Presentation

14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Immune Cell Therapy\Oral\8000
8000

Interlukine-27, a Key Player in Rejecting 4T1 Breast Carcinoma, an in vitro and in vivo Experiment

Abdolreza Esmailzadeh^{1,2}, Masoumeh Ebtekar¹, Alireza Biglari³, Zuhair M. Hassan Saraf¹, Takayoki Yoshimoto⁴

1. Department of Immunology, Faculty of Medical Sciences, Tarbiat Modares University Tehran, Iran.

2. Cancer Gene Therapy Research Center, Faculty of Medicine, Zanjan University of Medical Sciences, Zanjan, Iran

3. Department of Genetic and Molecular Medicine, Faculty of Medicine, Zanjan University of Medical Sciences, Zanjan, Iran

4. Department of Immunoregulation, Institute of Medical Science, Tokyo Medical University, Tokyo, Japan

Background: Cytokines are the chief participant in regulating immune responses, and cytokine-based cancer therapy have been proved in treating malignancies. Interleukin (IL)-27 is a cytokine from IL-12 family and promotes antitumor immunity by enhancing NK cells, CTLs and through exerting antiangiogenic potential. IL-27 has played an anti-tumoral role in numerous animal tumor models and suppressed tumor development. Here we have assessed whether IL-27 might be a feasible treatment for breast cancer.

Methods: In this study, murine IL-27 gene was cloned into expression vector P3XFLAG-CMV-9, and then recombinant plasmid was cloned into 4T1 cell line. We injected 4T1 cells to BALB/c mice to establish the antitumor activity of IL-27 in a syngeneic mouse model of breast cancer. The immunologic effects, anti proliferative function of IL-27 and tumor size have been evaluated. The injected tumoral mice and 4T1 treated cells with P3XFLAG-CMV-9 + IL-27 exhibited a significant reduction in tumor size and 4T1 cell numbers respectively.

Result and conclusion: Our results for the first time, suggest that IL-27 efficiently displays capability to suppress breast tumor, through upregulating IFN- γ and granzyme B as well as down regulating IL-4. We interestingly demonstrated that IL-27 can hold notable promise for breast cancer immunotherapy.

Keywords: Interleukin (IL)-27, Breast Cancer, immune gene therapy, Anti-tumor

ICIA2018\Immune Cell Therapy\Oral\9758
9758

A new way for miRNA delivery and priming dendritic cells: electroporated exosome with miRNA

Ali asadirad, Seyed Mahmoud Hashemi, Kaveh Baghai , Hosein Ghambarian , Esmail Mortaz, Davar Amani

Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Science

Background: Exosomes, membranous Nanovesicles, naturally carry bio-macromolecules and play pivotal roles in both physiological intercellular crosstalk and disease pathogenesis. Tumor-derived exosomes are proposed as a new type of cancer vaccine.

Methods: In this study we purpose to show that tumor cell-derived exosomes can function as vehicles to deliver exogenous miRNA-155 mimic in to dendritic cells (DC) for simultaneous miRNA delivery and Antigen priming of DC.

For this purpose, miRNA -155 was electroporated into tumor cell-derived exosomes then DCs were pulsed with electroporated exosomes.

Results: DCs pulsed miRNA-155 loaded exosomes can significantly increase the level of maturation and uniformity of DCs. In addition, these DCs in comparison with DCs in control group can proliferate lymphocytes insignificantly.

Conclusion: Furthermore, simultaneous miRNA delivery and antigen priming of DC can improve maturity and uniformity compared to control groups. Similar approaches could be useful in modification of target biomolecules in vitro and in vivo.

ICIA2018\Immune Cell Therapy\Oral\8470

8470

Exosomes from Neuroblastoma Cell-Exposed Natural Killer Cells can educate Naïve NKs to eradicate neuroblastoma tumors in vivo

AlirezaShoae-Hassani, Maryam Behfar, Seyed Abdolreza Mortazavi-Tabatabaei, Rashin Mohseni, Amir Ali Hamidieh

1. Applied Cell Sciences & Tissue Engineering Department, School of Advanced Technologies in Medicine, Tehran University of Medical Sciences, Tehran, Iran

Background: Immune cell-derived exosomes can increase immunity against tumors. On the other hand, tumor-derived exosomes can reduce the immunity and change the tumor microenvironment to further develop and initiate metastasis. These effects take place via alterations in the functions of innate and adoptive immune cells. In this experiment, we have studied the NK cell (NKs) effectiveness on tumor cells after expansion and incubation with exosomes.

Methods: The exosomes were derived from two populations of NKs: 1) Naïve NKs and, 2) neuroblastoma cell-exposed NKs. Also, we studied the effects of neuroblastoma derived exosomes (NB-Ex) on NK function. The molecular load of the characterized exosomes (by means of nanoparticle tracking analysis, flow cytometry, scanning electron microscopy and western blot) from NKs exposed to the neuroblastoma cell showed the expression of NCRs in addition to CD56, NKG2D, and KIR2DL2 receptors. These exosomes were used to treat NKs and then administered to NB tumor cells both in vitro and in vivo.

Results: Our results showed that some educations from exosomes derived from neuroblastoma cell-exposed NK cells activate NK cells to exert more efficient cytotoxicity against NB tumors, while NB-Ex act as tumor promoters by providing a tumor supporting niche.

Conclusion: This method of preparing the exosomes dramatically increases NK-mediated anti tumor effects against neuroblastoma cells.

Keywords: Cancer Therapy, Exosome, Immune Cell Therapy, Natural Killer Cell, Neuroblastoma

ICIA2018\Immune Cell Therapy\Oral\8294
8294

Construction of Chimeric Antigen Receptor (CAR) bearing Anti-Prostate Specific Membrane Antigen (PSMA) nanobody (VHH)

Mahmoud Hassani^{1,2}, Fatemeh Hajari Taheri², Mohsen Abolhassani², Mohammad Hossein Modaresi¹, Zahra Sharifzadeh², Arash Arashkia³

1. Department of Molecular Medicine, School of Advanced Technologies in Medicine, Tehran University of Medical Sciences, Tehran, Iran

2. Hybridoma Lab, Department of Immunology, Pasteur Institute of Iran, Tehran, Iran

3. Department of Molecular Virology, Pasteur Institute of Iran, Tehran, Iran

Background: Adoptive transfer of T cells expressing chimeric antigen receptors (CARs) is a promising anti-cancer therapy. T cells isolated from the body and equipped with CARs; when given back to the patient, these “CAR T cells” recognize and attack cancer cells. In 2017, two CAR T-cells were approved by the FDA, one for the treatment of acute lymphoblastic leukemia and the other for the advanced lymphomas. This technique is being investigated for use in other solid tumors and blood cancers.

Methods: For the first time, we used camelid nanobody (VHH) against prostate specific membrane antigen (PSMA) to construct the CAR T cells. T cells were transfected with construct composed of co-stimulatory molecules (CD28 and CD3 ζ) joined with spacer (CH2-CH3 fragment of IgG1) to nanobody. After confirmation of surface expression, the functions were evaluated when co-cultured with prostate cancer cells.

Results: Our data show that co-culture of CAR-T cells with PSMA⁺ prostate cancer cell (LNCap) not only increased the expression of IL-2 cytokine (about 400 ng/ml), but also expressed T cell activation marker (CD-69) almost 29%. In addition, CAR-T cells proliferated nearly 60% as compared with PSMA negative prostate cancer cell (DU145). This result indicates the specificity of CAR-T cells against PSMA.

Conclusion: The engineered VHH-CAR T cells may be developed to target virtually any tumor associated antigen in the future for the adoptive T-cell immunotherapy of solid tumors.

Keywords: CAR, PSMA, Prostate Cancer, Nanobody, Immunotherapy

ICIA2018\Immune Cell Therapy\Oral\8186

8186

Immunoregulatory function of Treg cells in experimental animal model of inflammatory bowel disease

Kaveh Baghaei¹, Seyed Mahmoud Hashemi², Sedigheh Pouya², Maryam Heidari², Hamid Asadzadeh Aghdaei¹, Mohammad reza Zali³

1. Basic and Molecular Epidemiology of Gastrointestinal Disorder Research Center, Research Institute for Gastroenterology and Liver Disease, Shahid Beheshti University of Medical Science, Tehran, Iran

2. Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences Tehran, Iran

3. Gastroenterology and Liver Disease Research Center, Research Institute for Gastroenterology and Liver Disease, Shahid Beheshti University of Medical Science, Tehran, Iran

Background: T regulatory (Treg) cells have a basic role in maintain stability of immune homeostasis and reduce autoimmune responses by modulating cells of innate and adaptive immune system. The immunoregulatory functions of Treg cells are mediated through different mechanisms including create tolerance against external factors such as bacteria, antigens and allergens. Therefore, the immunological function of these cells in the intestine is very important as a place that is constantly associated with foreign antigens and commensal bacteria. As Treg therapy has the potential to be a clinical treatment in the field of transplantation, its set up in the treatment of the IBD diseases can be much more effective than broad-spectrum immunosuppressive therapies. In current study, beneficial of Treg therapy in acute crohn's disease model in mice is considered.

Methods: Acute colitis was induced in female C57BL/6 mice using DSS in drinking water. Naïve T cell were isolated from spleen by Macs cell separation columns. By using CD3, CD28 and TGF-B, naïve T cells were differentiated to Treg cells. Tregs, 10⁶ cells per mouse, was injected intraperitoneally, on the last day of DSS. Clinical, serological and pathological tests were applied to analyze clinical outcomes.

Results: After receiving T regulatory cells in treated group, the clinical sign of colitis were moderated recovered. Clinical signs include body weight changes, bleeding and stool consistency were significantly changed and histopathology results also showed restriction of inflammation.

Conclusion: Regarding to immunoregulatory effects of Treg in inflammatory reactions, using of Treg could be presented as a new potential for therapeutic approach in IBD diseases. In following, clinical trial and comparative studies could be carrying out to identify effective strategy in order to help patients treatment.

Keywords: Treg, IBD, Acute colitis, Animal model

ICIA2018\Immune Cell Therapy\Oral\7964
7964

Bone marrow dendritic cells derived from CD40 and p19 knockout mice have in-vitro tolerogenic effect

Tahereh Kalantari^{1,2}, Bogoljub Ciric⁴, Eskandar Kamali-Sarvestani³, And Abdol mohamad Rostami⁴

1. Diagnostic Laboratory Sciences and Technology Research Center, School of Paramedical Sciences, Shiraz University of Medical Sciences, Shiraz, Iran.

2. Department of Laboratory Sciences, School of Paramedical Sciences, Shiraz University of Medical Sciences, Shiraz, Iran.

3. Immunology Department, Shiraz University of Medical Sciences, Shiraz 7144155, Iran

4. Department of Neurology, Thomas Jefferson University, Philadelphia, PA 19107, USA

Background: Multiple sclerosis (MS) as an auto immune disease are characterized by perivascular inflammatory lesions, demyelination and axonal damage. The role of dendritic cells (DCs) as antigen presenting cells (APCs) in MS or its animal model EAE (experimental autoimmune encephalomyelitis) development has been shown in different studies. In this study, Potency of Knockout mice (KO) derived BMDCs (Bone Marrow Dendritic Cells) (CD40KO-DCs and p19KO-DCs) were compared to C57BL/6 mice (Cont-DCs) for their tolerogenic/immunogenic efficacy.

Methods: Flow cytometry was used to analyze the expression of maturation markers MHCII and co-stimulatory molecules CD40, CD80 and CD86. Expression of CD40 and p19 was measured by RT-PCR. The ability of DCs to stimulate CD4⁺ T cells of 2D2 mice was evaluated by ³H-thymidine uptake and the endocytic capacity was evaluated in KO derived DCs compared to Cont-DCs using fluorescent OVA.

Results: CD40 expression of DCs obtained from CD40KO was significantly different ($p < 0.05$) to Cont-DCs. IL-6 and IL-12 expressions of KO derived BMDCs were significantly lower when compared with Cont-DCs ($p < 0.05$). The gene expression of CD40 and/or IL-23p19 was measured by quantative RT-PCR in KO derived DCs compared to control group. As compared to Cont-DCs groups, CD40KO-DCs and P19KO-DCs showed a significant difference in mRNA expression of CD40 and p19 respectively ($p < 0.05$). CD40KO-DCs in stimulation of MOG-specific CD4⁺ T cell proliferation were decreased significantly compared to Cont-DCs group ($p < 0.05$). The percentage of endocytosed FITC-OVA was not statistically different in KO derived DCs when compared to Cont-DCs group ($p > 0.05$).

Conclusion: This study shows that BMDCs derived from CD40KO and p19KO mice has a tolerogenic potency in-vitro. We suggest a further in-vivo study using these KO derived DCs to check their tolerogenic effect in therapy of auto immune diseases such as MS.

Keywords: Knockout mice, CD40KO, p19KO, tolerogenic DC.

ICIA2018\Immune Cell Therapy\Oral\7821

7821

Mesenchymal Stem Cells pulsed with 17- β estradiol lead to a more favorable outcome in collagen induced model of Rheumatoid Arthritis

Monireh Jahantigh, Seyyed Meysam Abtahi Froushani, Nahideh Afzale Ahangaran

Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Background: Insufficient of Mesenchymal stem cells (MSCs) reaching to inflamed tissues has limited their potential therapeutic benefits. Some documents indicated that 17- β estradiol augments homing of stem cells. Here, we evaluate the effect of 17- β estradiol treatment of Mesenchymal stem cells (MSC) on their potential for modulation of the animal model of Rheumatoid arthritis (RA).

Methods: Bone marrow-derived MSCs were pulsed with 100 μ M of 17- β estradiol for 24 h. RA was induced by collagen peptide and complete Freund's adjuvant in Wistar rats. Animals were allocated in 3 groups, one week after induction: treated with MSCs alone, treated with MSCs pulsed with estradiol and untreated group. The change in the diameter of wrists and ankles of each rat were recorded every 5 days until 33 days after induction. Then, splenocytes were tested to assess proliferation rate and cytokines production.

Results: The swelling and edema of ankles of RA rats received 17- β estradiol pulsed MSCs were significantly regressed compared to RA rats received MSCs alone. Aside from reducing lymphocyte proliferation, 17- β estradiol pulsed MSCs significantly reduced the production of proinflammatory IL-17 as well as TNF- α through antigen-specific re-stimulation more profound than RA rats treated with MSCs alone. Moreover, the level of anti-inflammatory IL-10 was significantly increased in RA rats treated with 17- β estradiol pulsed MSCs compared to RA rats received MSCs alone. Nevertheless, the levels of IFN- γ didn't show any significant changes between MSCs treated group.

Conclusion: MSCs pulsed with 17- β estradiol lead to a more favorable outcome in RA model compared to un-treated MSCs.

Keywords: 17- β estradiol, Mesenchymal stem cell, Rheumatoid arthritis.

ICIA2018\Immune Cell Therapy\Oral\11082
11082

Mesenchymal Stem Cell-Derived Exosomes Ameliorate Inflammation in Mouse Model of Acute Colitis

Neda Heidari¹, HajarAbbasi¹, Seyed Mahmoud Hashemi¹, Saeed Namaki¹, Mohammad Reza Zali¹

1. Department of Immunology, School of medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: inflammatory bowel disease (IBD) is called a group of diseases which causes chronic and recurrent inflammation of the large intestine and narrow intestine that the imbalanced immune responses have an important role in development and progression of the disease. Studies on mesenchymal stem cells (MSCs) have shown that these cells have immunomodulatory properties that can exert their effect through exosomes secretion. Therefore, this study was designed to investigate the effects of the mesenchymal stem cell-derived Exosome injection in mouse model of acute colitis.

Methods: Acute colitis was induced by 3 % dextran sulfate sodium (DSS) in C57Bl/6. The disease was induced in 1 cycle (4 days use of water containing DSS, followed by 6 days of water). After induction of disease in mice, injection of Exosome (100μl) for 4 times was done intraperitoneally. During the study, changes in body weight, bleeding, stool consistency, disease activity index (DAI), and mortality rate were recorded. After euthanizing the mice, weight, and length of the colon, the percentage of Treg cells were measured. The pathology examination of the colon was also performed.

Results: After Exosomes injection body weight decrease, bleeding, DAI was improved and mortality rates decreased. The stool consistency was improved too. The percentage of Treg cells and the TGF-β production was increased while the level of IL-17 decreased. Pathologic findings showed after injection of Exosomes, infiltration of inflammatory cells and epithelial destruction decreased.

Conclusion: Given that exosome has immunomodulatory effects, Exosomes can be used as a cell-free therapy in the treatment of inflammatory bowel disease.

Keywords: Mesenchymal stem cell, Inflammatory bowel diseases, Dextran sulfate sodium, Exosome



Immunodeficiency

Oral Presentation

14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Immunodeficiency\Oral\8548
8548

Clinical course, laboratory data and cell-surface expression of IL12Rβ1 in MSMD suspected patients with disseminated or atypical local BCGosis

Leila Moradi¹, Leila Parvaneh^{1,4}, Hamid Eshaghi³, Rezayazdani², Nima Parvaneh^{1,2}, Asghar Agha Mohammadi^{1,2}

1. Department of Immunology, Asthma and Allergy, Children's Medical Center, Tehran University of Medical Sciences, Tehran, Iran

2. Research center for immunodeficiency Diseases (RCID), Tehran University of Medical Sciences, Tehran, Iran

3. Department of Infectious Diseases, Children's Medical Center, Tehran University of Medical Sciences, Tehran, Iran

4. Department of Biology, Central Tehran Branch, Islamic Azad University, Tehran, Iran

Background: Autosomal recessive IL12Rβ1 deficiency is known as the most common cause of Mendelian Susceptibility to Mycobacterial Diseases (MSMD). This rare congenital disorder caused by mutations in genes encoding cytokines or receptors of IL12/23-INF-γ immunity axis, characterized by increased susceptibility of these patients to infections due to weakly virulent mycobacteria such as BCG vaccine or environmental mycobacteria (EM) and salmonella.

Methods: Clinical course and immunological laboratory data of Patients with atypical-local or disseminated BCGosis referred to Children's medical center in Tehran were studied to investigate the suspected MSMD patients, after ruling out more common PIDs presented with BCGosis, such as SCID or CGD. Blood samples of healthy controls and 19 patients were analyzed. Lymphocytes from both group were activated and stimulated in vitro with PHA and IL2 and evaluated for flowcytometric IL12Rβ1 expression.

Results: Reduced IL12Rβ1 expression was noticed in 4 patients' blood sample. All of them had consanguineous parents and history of BCG vaccination at birth. Whereas all of them had recurrent episodes of diarrhea, just one had documented history of disseminated salmonellosis. Two patients had recurrent oral candidiasis. There was no evidence of viral infection or malignancies.

Conclusion: Regarding to high frequency of consanguinity in Iran and BCG vaccination program at birth, it seems advisable to investigate IL12Rβ1 deficiency as the most prevalent cause in patients with suspicious MSMD. It may be beneficial postponing BCG vaccination in neonates with family history of documented or suspected diagnosis of a PID or an unusual consequence of BCG vaccination.

Keywords: BCGosis, IL12Rβ1 deficiency, MSMD, primary immunodeficiency, Salmonellosis

ICIA2018\Immunodeficiency\Oral\9878
9878

Evaluation of early and late apoptosis in patients with CVID

Reza Yazdani^{1,2*}, Mazdak Ganjalikhani-Hakemi², Nima Rezaei¹, Fateme Babaha³, Asghar Aghamohammadi¹

1. Research Center for Immunodeficiencies, Pediatrics Center of Excellence, Children's Medical Center, Tehran University of Medical Science, Tehran, Iran

2. Department of Immunology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

3. Department of Immunology, School of Medicine, Tarbiat Modares University, Tehran, Iran

Background: Common variable immunodeficiency (CVID) is a primary immunodeficiency with heterogeneous complications. One of the most important issues with regard to pathogenesis of CVID is defect in survival and differentiation of B-cells. Accordingly, we evaluated apoptosis (early and late apoptosis) with and/or without stimulation in B-cells, as well as measurement of B-cell subsets in CVID patients and controls.

Methods: B-cells were purified from PBMCs by negative selection and were stimulated by anti-IgM and anti-CD40 antibody for 24 hours. We measured spontaneous and induced apoptosis of B cells by annexin V/PI apoptosis detection kit. Four-color flow cytometric immunophenotyping determination of B-cell subsets (transitional, CD21^{low}, naive, marginal zone and memory B cells as well as plasmablasts) were performed using FACS. Moreover, BCL-2 gene expression was measured using real-time PCR.

Results: The expression of annexin V and PI in both unstimulated and stimulated B cells from patients were significantly higher than controls. Moreover, early apoptosis ($P = 0.04$), late apoptosis ($P = 0.04$) and necrosis ($P = 0.03$) in both unstimulated and stimulated B-cells were strongly further in the patients than controls. Furthermore, transcript levels of Bcl-2 were lower in patients than controls. On the other hand, our patients presented a significant reduction in absolute counts and percentage of B-cell subsets in patients than controls.

Conclusions: Our data suggest that increased apoptosis leads to abnormality in B-cell subset numbers, as B-cells could be unable to complete their maturation and differentiation. Thus, apoptosis could be one of the mechanisms involved in the significant reduction of B-cell subsets in these patients.

Keywords: apoptosis, immunophenotyping, common variable immunodeficiency

ICIA2018\Immunodeficiency\Oral\9923
9923

Finding Two Autosomal Dominant Gain-Of-Function Mutations in STAT1 Gene of Three Chronic Mucocutaneous Candidiasis Patients

Vajiheh Ostadi^{1,2}, Roya Sherkat¹, MazdakGhanjalikhan Hakemi², Seyed Mehran Modaressadegi⁴, Anne Puel³

1. *Acquired Immune Deficiency Research Center, Isfahan University of Medical Sciences, Isfahan, Iran*

2. *Department of Immunology, School of Medical University of Isfahan, Iran*

3. *Laboratory of Human Genetics of Infectious Diseases, Necker Branch, Paris, France*

4. *Isfahan University of Medical Sciences, Department of Genetics, School of Medical University of Isfahan, Iran*

Background: Chronic mucocutaneous candidiasis disease (CMCD) is characterized by susceptibility to recurrent or resistant candida infections, typically *Candida albicans*, on cutaneous and mucosal surfaces. In this survey, function and genetics of Th17 cells were studied in susceptible patients to mucocutaneous candidiasis.

Methods: for suspected patients who suffered from chronic candidiasis, CFSE proliferation test with candida antigen, detection of Th17 cells and pSTAT1 were done by flow cytometry method. Also, expression of IL-17A, IL-17F and IL-22 genes were measured with real-time PCR on all suspected patients. At the same time, whole exome sequencing (WES) was done for all patients.

Results: we detected three patients with gain-of-function (GOF) mutation in STAT1 molecule. The frequency of Th17 cells and their proliferation against candida antigen were reduced significantly in these patients compared with healthy controls. IFN- γ stimulation resulted in higher generation of pSTAT1 in the T lymphocyte cultures from patients compared to healthy controls. Gene expression of IL-17A, IL-17F and IL-22 from PBMCs was significantly reduced in these patients.

Conclusion: In summary, our study showed importance of STAT1 molecule in IL-17A T cells differentiation against *Candida*. Since, previous studies are reported STAT1 GOF mutations to be causative in more than half of CMC patients, for finding an effective treatment, more studies are required for displaying of relevance of this molecule with autoimmunity and other clinical symptoms.



Immunogenetics

Oral Presentation

14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Immunogenetics\Oral\7355

7355

KIR2DS1, 2DS5, 3DS1 and KIR2DL5 are associated with the risk of head and neck squamous cell carcinoma in Iranian patients

Shaghik Barani^{1,3}, Bijan Khademi², Abbas Ghaderi^{1,3 *}

1. Department of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz Iran

2. Otolaryngology Research Center, Department of Otolaryngology, Shiraz University of Medical Sciences, Iran

3. Shiraz Institute for Cancer Research, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Activating and inhibitory KIR receptors (aKIR, iKIR) control the development and function of NK cells whose function alterations adjust the tumor microenvironment immunity. This research was conducted to determine the KIRs gene impact on genetic predisposition to Head and Neck Squamous Cell Carcinoma (HNSCC) as the sixth most prevalent cancer worldwide and its clinicopathological features in Iranians.

Methods: KIR genotyping using sequence-specific primers-polymerase chain reaction (SSP-PCR) method was performed to identify the presence of all 16 KIR genes in 285 Head and Neck Squamous Cell Carcinoma (HNSCC) patients, including laryngeal, oral cavity and pharyngeal SCC (L.SCC, O.SCC, P.SCC) and 273 controls (CNs).

Results: Comparison of KIRs gene frequency between HNSCC and CN groups revealed a highly significant increase in KIR2DL5, 2DS1, 2DS5, 3DS1 and CxT4 genotype and a decrease in KIR2DS4 deleted variant and AA genotype carriers. A significant increase was noted in individuals with more inhibitory KIRs than activating KIRs in HNSCC compared with CNs. Individuals with ≥ 4 inhibitory KIR and those with ≥ 5 activating KIRs were significantly more common in HNSCC than CNs. The carrier frequency of KIR2DS1 and C4T4 genotype is lower in young-onset HNSCC compared with late-onset ones. 68 distinct KIR genotypes were identified in 558 individuals. A higher frequency of lymph node metastasis (LNM) was observed in patients with KIR2DL2 gene than those who lacked it, as well as in patients with 5 inhibitory KIR than those with less than it. The frequency of KIR2DL2 and KIR2DS4 deleted variant carriers was increased in patients with advanced and early clinical stages respectively.

Conclusion: Our findings discovered the detrimental impact of KIR2DS1, 2DS5, 3DS1, 2DL5 and CxT4 genotype as well as the positive impact of KIR2DS4del and AA genotype on genetic predisposition to HNSCC and KIRs associations with clinicopathological characteristics.

Keywords: killer immunoglobulin like receptors (KIRs), SSP-PCR, Head and neck squamous cell carcinoma (HNSCC), NK cells

ICIA2018\Immunogenetics\Oral\7394

7394

Association between TRA3IP2 (rs33980500) and left main coronary artery stenosis in Acute Myocardial InfarctionPordel, S.¹, SajediKhanian, M.², Karimi, M.H.³, Nikoo H.⁴, Doroudchi M.¹*1. Department of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran**2. Department of Cardiology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran**3. Fars Blood Transfusion Center, Shiraz, Iran**4. Cardiovascular Research Center, Shiraz University of Medical Sciences, Shiraz, Iran*

Background: There is compelling evidence showing connections between inflammation, immune system and cardiac pathology. IL-17A and its adaptor Act1 protein, which is encoded by TRAF3IP2 gene, may have a role in this process. Due to the association of TRAF3IP2- rs33980500 with inflammation, we assessed this single nucleotide polymorphism (SNP) in acute myocardial infarction (AMI).

Methods: AMI patients (n=201) referring to the affiliated hospitals of Shiraz University of Medical Sciences in a one year period were included in this study. DNA was extracted from venous blood and the genetic variation in the TRAF3IP2-rs33980500 was identified using PCR-RFLP method. Control individuals (n=201) were recruited from among healthy blood donors of the same age range and gender.

Results: No significant difference was observed in allelic and genotype distributions of TRAF3IP2-rs33980500 between patients and controls. However, there was an association between this SNP and left main coronary artery stenosis. MI patients with left main coronary artery stenosis had a significantly higher frequency of T allele ($p=0.03$) and TT genotype ($p=0.01$) compared to patients without this complication.

Conclusion: The association of TT genotype and T allele of TRAF3IP2-rs33980500 SNP with left main coronary artery stenosis is in accordance with its link with inflammation. T allele of TRAF3IP2-rs33980500 SNP is reported to be associated with increased inflammation which is a key factor in atherosclerotic plaque formation, atherosclerosis progression and narrowing of the arteries.

Keywords: Myocardial infarction, Single nucleotide polymorphism, TRAF3IP2-rs33980500

ICIA2018\Immunogenetics\Oral\7876
7876

Assessment of protein prenylation pathway in multiple sclerosis patients

Mohammad Taheri^{1,2}, Mir Davood Omrani^{1,2*}

1. Department of Medical Genetics, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Urogenital Stem Cell Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Multiple sclerosis (MS) is a chronic inflammatory disorder with several genetic and environmental factors being implicated in its pathogenesis. Protein prenylation as one of the important post-translational modifications of proteins has crucial role in immune system regulation.

Methods: In the current case-control study we compared expression of five genes coding for the different subunits of proteins implicated in protein prenylation in 50 Iranian MS patients with those of healthy subjects.

Result: No significant difference has been found in FNTA and PGGT1B expressions between cases and controls. Spearman Correlation analysis between FNTA relative expression and disease duration showed significant correlation in male patients ($r=-0.671$, $P=0.024$) but not female patients ($r=0.253$, $P=0.12$). FNTB expression was significantly higher in MS patients compared with healthy subjects. Spearman Correlation analysis between FNTB relative expression and disease duration showed significant correlation in male patients ($r=-0.876$, $P=0.004$) but not female patients ($r=0.296$, $P=0.06$). RABGGTA was significantly up-regulated in total MS patients, total male patients, female patients aged between 30 and 40 and male patients aged >40 compared with corresponding control groups. RABGGTB was significantly down-regulated in total MS patients, total female patients and female patients aged >40 compared with corresponding control groups.

Conclusion: Totally, we demonstrated dysregulation of protein prenylation pathway in MS patients compared with healthy subjects. Future studies are needed to find the clinical implication of this pathway in MS patients.

ICIA2018\Immunogenetics\Oral\7894

7894

Negative regulation of Semaphorin 3A expression in peripheral blood mononuclear cells using microRNA-497-5p

Shima Shapoori¹, Mazdak Ganjalikhani-hakemi¹, Mahsa Rezaeepoor¹, Fereshteh Alsahebhosoul¹, Sharifeh Khosravi², Masoud Etemadifar³, Marjan Mansourian⁴

1. Department of Immunology, Faculty of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

2. Department of Genetics, Faculty of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

3. Multiple Sclerosis and Neuroimmunology Research Center, Isfahan University of Medical Sciences, Isfahan, Iran

4. Department of Biostatistics and Epidemiology, Faculty of Health, Isfahan University of Medical Sciences, Isfahan, Iran

Background: Semaphorin3A (Sema3A) as a secreted semaphorin, is generated by a wide spectrum of cells in the human body including most of the immune cells. Thus, this protein as an immune modulator could participate in the pathogenesis of autoimmune diseases like Multiple sclerosis (MS). MicroRNAs (miRNAs) which are non-coding and single-stranded RNAs with ~21-25 nucleotides length, modulate target gene expression at the post-transcriptional level. MiRNAs play significant roles in many physiologic processes as well as semaphorins. Accordingly, it proposes that miRNAs may be crucial to modulate the semaphorins functions. Previous findings have proved that miR-497-5p is up-regulated and Sema3A is down-regulated in some autoimmune disorders. Thus, this paper aimed to increase our understanding of correlation between the Sema3A and Mir-497-5p in Peripheral Blood Mononuclear Cells (PBMCs).

Methods: In this study, we cultured PBMCs in optimum condition and transfected miR-497-5p mimic into them using X-treme gene Reagent. There were 4 groups: 1- Positive control, 2- Mock-transfection control, 3- Scrambled control, 4- Mimic transfection which in all groups PBMCs were treated with PHA. After 48 hours, we assessed supernatant and cells by ELISA assay and qRT-PCR, respectively. Also, we evaluated cell viability using MTT assay. All experiments were done in triplicates and P values equal to or less than 0.05 were considered significant. All the statistical analysis were performed through the SPSS 20 and P values equal to or less than 0.05 were considered significant.

Results: We observed down-regulation of Sema3A gene (p value = 0.0001) and its secretion (p value = 0.021) in PBMCs through miR-497-5p transfection. Moreover, transfection with miR-497-5p mimic and down-regulation of Sema3A did not affect viability of PBMCs (p value = 0.156).

Conclusion: Based on the obtained results, we suggest that miR-497-5p has a high suppressive effect on Sema3A expression and both Sema3A and miR-497-5p could be considered as critical targets for further studies on future therapeutic attempts for autoimmune diseases such as MS and rheumatoid arthritis (RA).

Keywords: Autoimmune disease, miRNA, miR-497-5p, Sema3A

ICIA2018\Immunogenetics\Oral\8116

8116

IRF7 gene expression profile and methylation of its promoter region in patients with systemic sclerosis

Ramazan Rezaei^{1,3}, Mahdi Mahmoudi¹, Farhad Gharibdoost¹, Mohammad Vodjgani²

1. Rheumatology Research Center, Tehran University of Medical Sciences,

2. Department of Immunology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

3. Department of Immunology, Medical School, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: The aim of the current study was to evaluate if methylation status of CpG sites of interferon regulatory factor 7 (IRF7) promoter in peripheral blood mononuclear cells (PBMCs) of systemic sclerosis (SSc) patients is involved in pathogenesis of the disease.

Methods: PBMCs were isolated from whole blood of 50 SSc patients and 30 controls. After the extraction of total RNA and DNA contents from PBMCs, complementary DNA (cDNA) was synthesized. Afterwards, quantitative analysis of IRF7 messenger RNA (mRNA) was conducted by real-time polymerase chain reaction (PCR). To evaluate the methylation status of the promoter region of IRF7 gene, PCR products of bisulfite-treated DNA from SSc patients and controls were sequenced.

Results: The mRNA expression of IRF7 in PBMCs from patients compared with controls was significantly upregulated. While limited cutaneous SSc patients expressed the mRNA of IRF7 higher than controls, the diffuse cutaneous SSc group did not demonstrate significantly increased expression in comparison to controls. Insignificant promoter hypomethylation of IRF7 was observed in SSc patients compared with the control group. However, CpG2 hypomethylation was significantly associated with increased SSc risk. Furthermore, overall promoter methylation and mRNA level of IRF7 were significantly correlated with each other. Nonetheless, none of them correlated with Rodnan score of SSc patients. There was significant difference in IRF7 mRNA expression between CpG8 methylated and unmethylated SSc patients. Moreover, the difference of methylation and expression was not significant between anti-nuclear antibody (ANA)-positive and ANA-negative SSc patients.

Conclusions: It is suggested that hypomethylation of the IRF7 promoter might play a role in SSc pathogenesis, probably through promoting the IRF7 expression in PBMCs of patients with SSc.

Keyword: IRF7, systemic sclerosis

ICIA2018\Immunogenetics\Oral\8297

8297

Association between variants of FOXP3 gene and the occurrence of pre-eclampsia in Iranian women

Milad Gholami^{1,2}, Sahar Bayat³, RezaMirfakhraie⁴, Reihaneh Pirjani⁵, Robabeh Taheripannah², Sayyed Mohamad Hossein Ghaderian⁶

1. Department of Biochemistry and Genetics, faculty of Medicine, Arak University of Medical Sciences, Arak, Iran

2. Infertility & Reproductive Health Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

3. Department of Medical Genetics, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

4. Department of Medical Genetics, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

5. Associate professor, Obstetrics and Gynecology Department, Arash hospital, Tehran University of Medical Sciences, Tehran, Iran

6. Urogenital Stem Cell Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Pre-eclampsia (PE), is a pregnancy-specific disorder, with serious consequences for both the mother and the fetus that affecting 4–8% of pregnant women around the world. Despite intense studies, the pathophysiology of PE remains enigmatic. Previous studies suggested that regulatory T cells (Treg) dysfunction is involved in the pathogenesis of PE. We hypothesized that functional variants of the Forkhead Box Protein 3 (FOXP3) gene might be associated with PE via dysregulation of Treg cells.

Methods: We genotyped three variants of FOXP3 by PCR-RFLP and Tetra ARMS-PCR methods in 133 PE patients and 143 healthy age-matched controls.

Results: The genotypic frequencies of rs2232365 were found to be protective from the development of PE under codominant [odds ratio (OR) 0.49, 95 percent confidence interval (CI) 0.28 – 0.87, P-value = 0.043], dominant [odds ratio (OR) 0.54, 95 percent confidence interval (CI) 0.32 – 0.94, P-value = 0.027] and over dominant [odds ratio (OR) 0.57, 95 percent confidence interval (CI) 0.35 – 0.92, P-value = 0.02] models. Moreover, the rs3761548 conferred a risk of PE in recessive model [odds ratio (OR) 2.05, 95 percent confidence interval (CI) 1.08 - 3.88, P-value = 0.025]. However, no mutation was detected in FOXP3 exon2 in any of the studied samples.

Conclusion: Based on our results, thought that FOXP3 variants may be an important contributor for development of PE in Iranian women, probably by altering FOXP3 function and/or its expression.

Keywords: Association, Variants, Pre-eclampsia, FOXP3

ICIA2018\Immunogenetics\Oral\8719
8719

The Relationship PDCD1 polymorphisms with Systemic Lupus Erythematosus in an Iranian Population

Farshid Fathi¹, Hossein Motedayyen², Noushin Lotfi¹, Erfan Sadeghi³, Mehdi Raie⁴, Hossein Hafezi⁵

1. Department of Immunology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

2. Department of Laboratory Medicine, Kashan University of Medical Sciences, Kashan, Iran

3. Department of Biostatistics and Epidemiology, School of Public Health, Isfahan University of Medical Sciences, Isfahan, Iran

4. Health Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran

5. Department of Dermatology, Isfahan University of Medical Sciences, Isfahan, Iran

Background: Systemic lupus erythematosus (SLE) is a heterogeneous autoimmune disease that characterized by the breakdown of tolerance and the production of autoantibodies against self-antigens. PD-1 is one of the most important inhibitory receptors for regulating the immune system and preventing the development of autoimmune disorders. Therefore, we evaluated whether PDCD1 (PD-1) gene polymorphisms are associated with the incidence of SLE disease in the Iranian population.

Methods: Blood samples (5ml) were collected from 120 SLE patients and 120 healthy volunteers'. Genomic DNA extracted by salting out method and genotype determination of PD1.1, PD1.3, PD1.5 and PD1.9 was done by polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP).

Results: Our results showed a significant difference between PD1.5 genotypes in SLE patients and control group ($P = 0.02$). The frequency of genotypes CC, CT and TT were, respectively, 60%, 31.7% and, 8.3% in SLE patients, while those of control group were 42.5%, 46.7% and 10.8% for CC, CT and TT, respectively. In addition, the allelic analysis indicated that there was a significant difference in PD1.5C allele between SLE patients and control group. The frequency of C allele of PD1.5 in SLE patients was significantly higher than the control group ($p = 0.01$). However, there was no difference between PD1.1, PD1.3 and PD1.9 genotypes and alleles in SLE and control group.

Conclusion: Our study suggested that PD-1.5 polymorphisms in PDCD1 gene may increase SLE susceptibility in our Iranian population. However, the PD1.1, PD1.3 and PD1.9 were not correlated to genetic predisposition of SLE in our Iranian population.

Keywords: PDCD1, Polymorphism, Systemic lupus erythematosus, autoimmune disease.

ICIA2018\Immunogenetics\Oral\9950

9950

Variation in the Interleukin 4–Receptor α Gene (I50V) Confers Susceptibility to AsthmaMasouma Mowahedi¹, Morteza Samadi²

1. Department of Immunology, International Campus, Shahid Sadoughi University of Medical Sciences and Health Services, Yazd, Iran

2. Department of Immunology, Faculty of Medicine, Shahid Sadoughi University of Medical Sciences and Health Services, Yazd, Iran

Background: Th17 cells apart from Th2 cells have been found to participate in the development of asthma. Recent studies have implicated polymorphisms in the signaling pathway of interleukin 4 (IL-4) receptor in the progress of inflammation. The aim of the present study was to investigate how to associate the single nucleotide polymorphism (SNP) in the interleukin 4 receptor α chain (IL-4R α) gene rs1805010 (I50V) to total immunoglobulin E (IgE) and IL-17A in the etiology of asthma in an Iranian population.

Methods: To evaluate how associate SNP I50V and asthma, in a case-control study, I50V variant were screened using PCR-RFLP followed by Fisher exact test. To investigate how associate SNP I50V and serum levels of IL-17A and total IgE, first ELISA was performed. To determine how to communicate SNP I50V, IL-17A and total IgE with the severity of the disease, patients were classified into three groups according to GINA criteria: mild, moderate and severe. Next student's t-test or one-way ANOVA were performed.

Results: We found a significant association between SNP I50V with asthma ($p = 0.001$). Despite IL-17A and total IgE in asthmatic patients were significantly higher than the control group ($p = 0.026$ and $p < 0.01$ respectively), neither total IgE nor IL-17A associated to SNP I50V.

Conclusion: I allele is more common in patients and more likely to reflect the importance of Th2 cells in asthma. Despite higher levels of total IgE and IL-17A were in accordance with the disease subgroups, increase of total IgE in tune with the severity of disease while enhancement of IL-17A in serum against the total IgE and according to decrease of disease severity.

Keywords: Asthma, I50V, IL-17A, IgE



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Immunohematology

Oral Presentation

ICIA2018\Immunohematology\Oral\7712
7712

Preliminary investigation of unexpected antibodies against blood group antigens in major β -thalassemia patients in Lorestan province in 2017-2018

Alimohammad Varzi¹, Mona Hemmati Chegeni², Mohammad Ziaee³, Babak Abdolkarrimi⁴

1. Department of Immunology, Lorestan University of Medical Sciences, Khorramabad, Iran

2. Department of Immunology, Lorestan University of Medical Sciences, Khorramabad, Iran

3. Medical School, Lorestan University of Medical Sciences, Khorramabad, Iran

4. Assistant professor of children hematologist and oncologist, Lorestan University of Medical Sciences

Background: Major β -thalassemia is a congenital hemolytic disorder that is classified as a hemoglobinopathy which is caused by defect in β -globin chain synthesis and considered as the most common genetic disorder world-wide. Regular blood transfusion plays a key role in treatment of the patients with this disorder. However, the development of hemolytic alloantibodies and autoantibodies against red blood cells (RBCs) can be culminated to development of different complications in these patients. Therefore, our study is focused on determination of the frequency of RBC alloimmunization and the most common alloantibodies involved among Lorestan patients who have received transfusion regularly.

Methods: We will use and test the information of 91 major β -thalassemia patients of Lorestan who have transfused RBC components. Our clinical and laboratory information including age at the start of transfusions, total number of transfusions, splenectomy status and other variants have been obtained through questionnaires. In this preliminary study alloantibody screening and identification of alloantibodies have been carried out on 15 cases out of 91 thalassemia patients using three cell and 11 cell panel, provided by Iranian blood transfusion organization, respectively.

Results: Out of 15 analyzed samples, two (13.34%) were positive for alloantibodies against C^w and Lu^a. Moreover anti-lewis, anti-kell and anti-Jk^a were detected in one of these two samples.

Conclusion: In this preliminary work, alloantibodies against C^w, kell, lutheran, lewis and kidd were detected in 13.34% of patients. Alloimmunization in transfused major β -thalassemia patients is very common which is an important issue for the blood bank department to find a fully matched blood components easily. However, working on new technology with high-throughput blood group typing that are developing in different research centers could be a good answer for reducing these problems in next future.

Keywords: alloimmunization, β -thalassemia major, regular blood transfusion, antibody identification

ICIA2018\Immunohematology\Oral\8340

8340

Relationship between HPA-1 and HPA5 Gene Polymorphisms and Refractory to Platelet Therapy and Recombinant Factor VII in Glanzmann Thrombasthenia Patients in Southeast of Iran

Shaban Alizadeh¹, Manizheh Habibpour¹, Zahra Kashani Khatib², Akbar Dorgalaleh³, Majid Naderi⁴

1. Department of Hematology, School of Allied Medical Sciences, Tehran University of Medical Sciences, Tehran, Iran

2 Hematology Department, High Institute for Research and Education in Transfusion Medicine, Iranian Blood Transfusion Organization, Tehran, Iran

3. Department of Hematology, School of Allied Medical Sciences, Iran University of Medical Sciences, Tehran, Iran

4. Genetic Researcher Center in Non-Communicable Disease, Zahedan University of Medical Sciences, Zahedan, Iran

Background: Glanzmann Thrombasthenia (GT) is a rare autosomal disease. HPA (Human Platelet Alloantigen) is a surface polymorphic alloantigen of platelets. This study was intended to investigate and compare the polymorphism of HPA-1 and HPA-5 genes in two groups of GT patients, with and without resistance to platelet and recombinant factor VII therapy.

Methods: This case control study was performed on GT patients (n=16) with resistance to platelet therapy and recombinant factor VII and control group of GT patients (n=16) without resistance to platelet therapy and recombinant factor VII. The consent form was completed by each patient. Gene polymorphisms of HPA-1 and HPA-5 were investigated using SSP-PCR, and the obtained data were analyzed using statistical software SPSS16.0.

Results: The results indicated no significant relationship between the studied genes and their resistance to platelet therapy and recombinant factor VII. The frequencies of HPA-1 genotype a/a were 98% and 94% in patient and control groups, respectively. The frequency of allele b was found to be less than allele a. The value of this allele was 4% in patient group and 1% in control group. In addition, the HPA-5a/a (98%) was the most frequent alloantigen in both groups. Seven percent (7%) of the patients had the HPA-5a/b genotype, and the HPA-5b/b was found to be absent in these individuals.

Conclusion: According to the results obtained, it could be concluded that these genes play no role in resistance to platelet therapy.

Keywords: Human platelet Ag-1, Human platelet Ag-5, Platelet therapy, Glanzmann thrombasthenia

ICIA2018\Immunohematology\Oral\8405

8405

Alloimmunization in transfusion dependent thalassemia major patients in South KhorasanLeila Mobasher¹, Masoomeh Asadifard², Gholamreza Anani Sarab¹*1. Department of immunology, Birjand University of Medical Science, Birjand-Iran**2. Blood transfusion center, Birjand-Iran*

Background: Patients with major beta thalassemia require regular blood transfusion in order to overcome deficient hemoglobin production. Multiple transfusion associated with some complications especially erythrocyte alloimmunization. In this study we aimed to perform antibody identification tests for transfusion dependent patients with major thalassemia in South Khorasan during 1395 to 1396.

Methods: This cross-sectional study was carried out on transfusion dependent beta thalassemia major patients registered at the thalassemia clinic in Birjand University of Medical Sciences. The demographic and clinical information of the patients for age, sex and history of transfusion were recorded. Alloantibody screening was done and then positive cases were examined to identify antibodies, using panel cells provided by Iranian Blood Transfusion Organization.

Results: Out of 68 patients recruited in this study, 36(53%) were male and 32 (47%) were female with the mean age of 15 years. Alloimmunization was detected in 3 (4.4%) of patients with male gender. Coincidence of Anti-C and Anti-E was detected in one case and Anti-K developed in the second patient. A cold agglutinin was found in the third case through antibody screening.

Conclusion: Based on our findings, Anti-Rh and Anti-Kell were the most prevalent alloantibodies. Despite multiple transfusions, favorable condition in terms of alloimmunization prevention was achieved. However, the results re-emphasized the importance of red cell antigens typing from the beginning of transfusion and antigen matching, especially for Rh and Kell antigens.

Keywords: Major beta thalassemia, Alloimmunization, Alloantibody screening

ICIA2018\Immunohematology\Oral\8412
8412

Global Plasma Industry: A Narrative on Outlines with Glance at the Endeavors in EMR Members

Afsaneh Aghaie¹, Seyed Mehrdad Jalali², Ali Akbar Pourfathollah^{1,3}

1. Blood Transfusion Research Center, High Institute for Research and Education in Transfusion Medicine, Tehran, Iran

2. Iranian Blood Research and Fractionation Company

3. Department of Immunology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

Background: Well solidified in developed countries, plasma industry is going to penetrate into several emerging economies in any aspects. Plasma supply could be described as the promotor of such industry and nowadays, there are plenty of intentions in order to manufacture products from locally produced plasma in such countries. On the other hand, traditional fractionation method has been modified in order to shortcut the process, enhance the yield or lessen the risk of Transfusion Transmitted Infections (TTI) and it has a great impact on the final product price. In addition, plasma derivatives have mostly had a remarkable share in the global biopharmaceuticals' market. Eastern Mediterranean Region (EMR) consists of various range of developing countries, some of which has magnificent foundation in national plasma supply and years of worthwhile experience in local fractionation and toll manufacturing programs. Great efforts have been made across the region in order to progress the plasma industry in terms of the quality of plasma domestically produced as well the programs align with self-sufficiency in plasma derivatives.

It is to overally review but in brief on outlines and trends in global plasma industry such as plasma supply, industrial process and a short market study followed by a narrative on the endeavors made in EMR members with this regard.

Keywords: EMR, plasma industry, plasma supply, plasma derivatives

ICIA2018\Immunohematology\Oral\8596
8596

Developing hemovigilance in University Teaching hospitals of Birjand – South khorasan

Gholamreza Anani Sarab¹, Hediye Gholampour Sardashti²

1. Department of Immunology. Birjand University of Medical Sciences

2. Department of Laboratory Sciences. Birjand University of Medical Sciences

Background: Blood transfusion monitoring and supervision (hemovigilance) has been established as one of the relatively new branches of transfusion medicine in many countries of the world. After conducting initial studies in the blood transfusion organization and emphasizing the importance of the blood and blood products and surveillance system, the implementation of this important system in health centers was recommended and approved by the Supreme Council of Iranian Blood Transfusion Organization in the winter of 2007. The implementation of this system in Iran since 2009 was started in 50 medical centers all over the country (19 hospitals in Tehran and 31 hospitals in 12 provinces of the country). The aim of this study was to evaluate the monitoring status and the level of completion of hemovigilance forms in teaching hospitals in Brigand.

Methods: The information regarding incomplete/inappropriate blood products requisition forms and blood transfusion monitoring forms were recorded in different wards of the teaching hospitals between the years. The results of the survey in 2014 were compared with the same research results in 2017 and the Summarized data were analyzed.

Results: The most common deficiency in vigilant forms was the neglect in record of vital signs of the patient over time after starting the transfusion. The well documented parameters were pulse pressure, Blood pressure and respiratory rate. Maximum numbers of incomplete forms were received from Intensive Care Units followed by Cardiac Care Unit. Overall, 17% improvement for completing blood request forms, administering blood, monitoring transfusions and being vigilant for the signs and symptoms of adverse reactions was detected in 3 years period from 2015 to 2017.

Conclusion: According to the results of the study, there is a significant improvement in the completion of blood transfusion monitoring form in year 2017 compared to 2015. This is a significant contribution to reducing the error in the process, and allows for safe and effective blood transfusion, system evaluation and collecting data on all types of transfusion-related adverse reactions. However, the study found that different wards do not have the same focus in completing the form, which can be due to the size of the work force or the lack of supervision in some sectors. It is suggested that, in order to improve the quality of supervision, nurses should be offered further courses to increase their awareness of the human and legal responsibilities in blood transfusion safety.

Keywords: hemovigilance, blood transfusion, Birjand

ICIA2018\Immunohematology\Oral\8659

8659

An association study between Lewis Blood group phenotype and IL-6 and IL-8 production by Staphylococcus aureus stimulated PBMC in blood group A humans

Sedigheh Younessi Amirabad¹, Kambiz Bagheri²

1. Department Of Microbiology, Kazerun Branch, Islamic Azad University, Kazerun, Iran

2. Department Of Immunology, Faculty Of Medicine, Kazerun Branch, Islamic Azad University, Kazerun, Iran

Background: Blood antigens are a major category of human antigens which are formed at the level of red blood cells and different epithelial cells. Adults with a Le gene have Le^a or Le^b. If the people are secretor ABHs, their red blood cells are Le^b and if they are non-secretor they will be Le^a.

Methods: In this study, volunteers with a blood type A were classified in to Le^a and Le^b using blood group test. Mononuclear cells of each individual were separated and subdivided into two subtypes: the control (cultured without bacterial stimulation) and test group (cultured in the presence of Staphylococcus aureus extract). The suspension containing bacterial extract was prepared, and added to the mononuclear cell culture. Following the optimization and finding the best dose, duration, and appropriate exciting pattern, the levels of IL-8 and IL-6 in the cultured groups were measured by the ELISA assay.

Findings: The results show that after examining dose and time intervals for incubation and stimulation of PBMCs by bacterial extract, the best dose and time was 50 microliters which fractinated into 2 dose and 24 hours, respectively. In fact, by increasing the dose, the cytotoxic effect of the bacterial extract on the cells in the culture was possible.

Although in some samples, IL-6 and IL-8 were secreted through 24-hour incubation, the growth of Lewis a and Lewis b in both groups was significantly increased by stimulating the culture cells with Staphylococcus aureus.

Conclusion: Repeated tests and cytokine assays identified that there was no significant difference between PBMCs in individuals with Lewis a and Lewis b regarding IL-6 and IL-8 production capacity.

Keywords: Interleukin-6 and Interleukin-8, Blood group Lewis a and Lewis b, PBMCs, Staphylococcus aureus bacteria , blood group A



Immunology of Chemical Victims and Environmental Pollution

Oral Presentation

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Immunology of Chemical Victims and Environmental Pollution\Oral\7858
7858

TCDD effects on induction of regulatory T cells in peripheral blood mononuclear cells in endometriosis patients

Mahsa Tanha¹, Shohreh Nikoo², Tahereh Naji³, Amir-hassan Zarnani⁴

1. Department of Molecular and Cellular Sciences, Faculty of Advanced Sciences & Technology Pharmaceutical Sciences Branch, Islamic Azad University, Tehran - Iran (IAUPS)

2. Immunology Research Center, Iran University of Medical Sciences, Tehran, Iran.

3. Department of Molecular and Cellular Sciences, Faculty of Advanced Sciences & Technology Pharmaceutical Sciences Branch, Islamic Azad University, Tehran - Iran (IAUPS)

4. Reproductive Immunology Research Center, Avicenna Research Institute, ACECR, Tehran, Iran

Background: Endometriosis is a benign and chronic inflammatory disease defined by the presence and development of endometrial tissues outside the uterine cavity. 2,3,7,8-Tetrachlorodibenzo-p-dioxin(TCDD), commonly referred to as dioxin, is considered as one of the main risk factors involved in development of endometriosis. TCDD exerts its impacts through binding the aryl hydrocarbon receptor (AHR). On the other hand, Regulatory T cells (Tregs), have also been suggested to play a part in the immunopathogenesis of endometriosis. Here, we investigated TCDD impacts on Treg induction in peripheral blood mononuclear cells (PBMCs) of endometriosis patients.

Methods: The effect of TCDD on induction of Tregs was assessed by flow cytometry. In addition, the expression of FoxP3 and AhR was evaluated using real-time PCR.

Results: TCDD treatment significantly increased FoxP3 expression in CD4⁺ T cells of endometriosis as compared with the non-endometriosis group. Furthermore, an increase in the expression of FoxP3 and AhR in the presence of dioxin in the endometriosis group was significant compared to the non-endometriosis group. Interestingly, we found a significantly decreased AhR expression in the endometriosis group, however; we observed no marked differences between the endometriosis and non-endometriosis groups with respect to FoxP3 expression.

Conclusion: Our findings demonstrated, for the first time, that TCDD is able to increase the level of FoxP3, a key Treg transcription factor, in peripheral blood T cells of endometriosis patients compared to the non-endometriosis samples. This suggests the existence of an intense feedback to inflammatory responses in endometriosis patients. Collectively, PBMCs of the endometriosis patients respond differently to dioxin in comparison with that of the non-patient individuals.

Keywords: Endometriosis, Dioxin, AhR, Regulatory T cells

ICIA2018\Immunology of Chemical Victims and Environmental Pollution\Oral\7982
7982

Investigation on the effects of aflatoxin B₁ on activation of caspases and ATP depletion in astrocytes

Jalil Mehrzad¹, ParisaVahidi², Amir Mohammad Malvandi³, Saman Hosseinkhani²

1. Department of Microbiology and Immunology, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran

2. Department of Biochemistry, Faculty of Biological Sciences, Tarbiat Modares University, Tehran, Iran

3. International Center for T1D, Pediatric Clinical Research Center Fondazione Romeo edEnricaInvernizzi, Department of Biomedical and Clinical Sciences L. Sacco, University of Milan, Milan, Italy

Background: Aflatoxin B₁ (AFB₁) is a poisonous substance which is classified as group 1 carcinogenic agents by International Agency for Research on Cancer (IARC). Although AFB₁ is implicated as a carcinogen in hepatocellular carcinoma, brain autopsies in affected areas shows its presence in 81% of cases. Considering the importance of AFB₁ impact on the astrocytes, there still exists a scarcity of research on this issue within the literature.

Methods: This research investigates the apoptosis effect of AFB₁ on primary mouse astrocytes in vitro. To this aim, after detection of LC₅₀ of AFB₁ for astrocytes, the astrocytes were exposed to various concentration of AFB₁ for 24, 48 and 72 hours. Concentration of intracellular ATP and caspase-3/7 activity was determined by the phenomenon of bioluminescence and luciferase reactions. Furthermore, cytochrome c release from mitochondria was carried out by western blot and percentage of apoptotic astrocytes was obtained using flow cytometry as well.

Results: The results indicated that, environmentally relevant level of AFB₁ (32 nM) induces apoptosis in astrocytes through ATP depletion and caspases activation. AFB₁-mediated apoptosis in astrocytes occurs through activation of caspase-3/7 with typical cytochrome c release.

Conclusion: Such in vitro effects of AFB₁ on astrocytes at the organelle and protein levels opens new doors to understanding the biological behavior of AFB₁ as well as molecular mechanisms that might translate to in vivo CNS inflammation, infection, and cancer in mammals caused by chronic exposure to AFB₁. Indeed, AFB₁-astrocytes interactions could trigger neurotoxicity in mammals. Further basic studies on the effect of biologically relevant level of AFB₁ on brain immune cells and molecules are in progress.

Keywords: Aflatoxin B₁, Bioluminescence, Astrocytes, Caspases, Neuroinflammation.

ICIA2018\Immunology of Chemical Victims and Environmental Pollution\Oral\8009
8009

Evaluating the effect of Dioxin mediated aryl hydrocarbon receptor activation on T helper cell subsets in an inflammatory condition

Narjes Soleimanifar^{1,2,3}, Behrouz Nikbin^{1,2,3}, Katayoon Bidad⁴, Mahdi Mahmoudi⁵,
Ahmad Reza Jamshidi⁵, Zahra Hosseinihah², Shayan Mostafaei⁵, Mohammad Hossein Nicknam^{1,3}

1. Molecular immunology research center, Tehran University of Medical Sciences, Tehran, Iran
2. Department of Molecular Medicine, School of Advanced Technologies in Medicine, Tehran University of Medical Sciences, Tehran, Iran
3. Department of Immunology, Tehran University of Medical Sciences, Tehran, Iran
4. Immunology, Asthma and Allergy Research Institute, Tehran University of Medical Sciences, Tehran, Iran
5. Rheumatology Research Center, Tehran University of Medical Sciences, Tehran, Iran

Background: Dioxin as an omnipresence environmental toxin is a main activator of aryl hydrocarbon receptor (AhR). AhR is a cytoplasmic transcription factor that regulates toxin-metabolizing enzymes such as cytochrome P450. The widespread expression of this factor in the immune system has shifted the point of AhR studies toward the immunologic role of it. The aim of our study is surveying the effect of Dioxin mediated aryl hydrocarbon receptor activation on T helper cell subset.

Methods: Peripheral blood mononuclear cells were obtained from 12 healthy people, and cultured before and after treatment with dioxin and antagonist for 72 hours. The percentage of Th17, Th22, and Tfoxp3 + cells was measured by flow cytometry. IL17 and IL10 cytokine levels were evaluated in the supernatant of culture media by ELISA method. The rate of expression of genes were evaluated using Real-Time PCR.

Results: Th22 cells were significantly increased by dioxin treatment, whereas Dioxin decreased the percentage of Th17 cells and expression of IL17 in the healthy subjects. The expression level of Aryl hydrocarbon receptor in dioxin treated cells was increased.

Conclusion: AhR can effect on fine regulation of the immune responses by altering the Th17 and enhancing of the Th22 cells.

Keywords: Dioxin, AhR, Th17,Th22.

ICIA2018\Immunology of Chemical Victims and Environmental Pollution\Oral\8618
8618

Seborrheic dermatitis and its relationship with Serum immunoglobulins levels in Sardasht chemical victims (20 years after sulfur mustard exposure)

Sara Parsapour¹

1. Department of imonology, Faculty of Medicine, Shahed University, Tehran, Iran

Background: Seborrheic dermatitis is a very common skin complication that affects seborrheic dermatitis in general, 4% of the world's population, and half of adults also have dandruff, which is actually a mild form of seborrheic dermatitis in the palms of the head. Dermatitis means skin lesions that are red and inflamed, causing itchy skins; seborrheic means that these skin lesions appear in the areas of the skin's oily skin (the location of the sebaceous stem cells), such as the face, the palatal, and the middle chest. The cause of seborrheic dermatitis is a fungus called malassezia. Sulfur mustard is a cytotoxic and vesicant chemical warfare agent it long-term effects include skin, respiratory and eye complications. The most important skin long-term complications include itching, eczema, xerosis, hyperpigmantasion. Serum IgE and IgG4 levels has a direct relationship with skin problems in allergic diseases, and hypersensitivity such as eczema, itching and etc. In this study, the relationship between serum levels of immunoglobulins with seborrheic dermatitis in Sardasht chemical victims was Studied 20 years after exposure to sulfur mustard.

Methods: This research is part of a cohort study of Sardasht chemical victims. A total of 372 male chemical victims exposed to sulfur mustard and 128 male in the control group in age range of 20-60 years old were. Both groups were clinically evaluated by a specialist physician. The sandwich ELISA was used for measure of serum immunoglobulins (ELISA kit : Bethyl, USA). The results were reported Mann-Whitney test and all statistical computing were performed using SPSS version 20

Results: Serum IgM level in chemical victims without seborrheic dermatitis was significantly lower than the control group with seborrheic dermatitis ($P=0.004$).

Conclusion: The results of this study showed that no significant relationship between Serum immunoglobulins levels with seborrheic dermatitis caused by sulfur mustard.

Keywords: immunoglobulins, seborrheic dermatitis, sulfur mustard

ICIA2018\Immunology of Chemical Victims and Environmental Pollution\Oral\9833
9833

Evaluation of MiR-21b and MiR-15b gene expression in Formalin Fixed Paraffin Embedded (FFPE) tissues of sulfur mustard exposed individuals with long-term pulmonary complications

Sahar Salimi¹, Farshid Noorbakhsh², Tooba Ghazanfari^{1*}

1. Department of Immunology, Faculty of Medicine, Shahed University, Tehran, Iran.

2. Department of Immunology, Faculty of Medicine, Tehran University of Medical Sciences. Tehran, Iran

3. Department of research center of regulation the immunology response. Tehran, Iran

Background: Sulfur mustard (SM) is powerful alkylating agent with cytotoxic, carcinogenic and mutagenic features which the toxicity of this substance has been recognized last decade ago. Sulfur Mustard Complications appear in two forms of acute and chronic that affects multiple organs such as lung, eye, skin and etc. Along with, inflammatory cytokines have major roles in acute and chronic inflammatory complications by sulfur mustard. Furthermore, miRNAs has important function in chronic inflammatory diseases. As this regard, miR-21 and miR-15 could potentiate TGF- β signaling pathway and be able to promote fibrosis process in lung tissue. So that, gene expression of miR-21 and miR-15 was evaluated in exposed individuals with sulfur mustard and unexposed people as control group.

Methods: The number of exposed individuals with SM is twenty as case group and the number of unexposed control group twenty.

Total RNA was extracted from FFPE tissue samples of exposed individuals to sulfur mustard and unexposed individuals. Extracted total RNA quantity was measured by Nano drop and its quality evaluated by bioanalyzer. Gene expression of miR-21b and miR-15b was evaluated by Real-time PCR and U6 snRNA was utilized as control gene.

Results: gene expression of miR-21b was decreased significantly in exposed individuals group as compared to unexposed control group. There was no significant differences in gene expression of miR15-b in exposed individuals in compared to unexposed control group.

Conclusion: In regard to decrease expression of miR-21b, it seems that miR-21 have a role to down-regulate TGF- β signaling pathway.

ICIA2018\Immunology of Chemical Victims and Environmental Pollution\Oral\9995
9995

Circulating MSCs in Sulfur Mustard-exposed Patients with Long-term Pulmonary Complications

Sara Ghaffarpour¹, Tooba Ghazanfari^{1, 2}, Eisa Salehi³, Seyed Mahmoud Hashemi⁴, Sussan K. Ardestani⁵, Keivan Gohari Moghadam⁶, Ensieh Sadat Mirsharif¹, Razieh Dilmaghanian¹, S.Faghihzadeh⁷

1. Immunoregulation Research Center, Shahed University, Tehran, Islamic Republic of Iran

2. Department of Immunology, School of Medicine, Shahed University, Tehran, Islamic Republic of Iran

3. Department of Immunology, School of Medicine, Tehran University of Medical Sciences, Tehran, Islamic Republic of Iran

4. Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Islamic Republic of Iran

5. Institute of Biochemistry and Biophysics, Tehran University, Tehran, Islamic Republic of Iran

6. Internal Medicine Department, Faculty of Medicine, Tehran University of Medical Sciences, Tehran, Iran

7. Department of Biostatistics and Social Medicine, Zanjan University of Medical Sciences, Zanjan, Islamic Republic of Iran

Background: Sulfur mustard (SM) is a toxic agent that causes acute and long-term pulmonary complications. Recent evidence has shown the impact of SM on mesenchymal stem cells (MSCs). These cells have a critical role in repairing damaged tissues. In this study, we evaluated the mobilization of MSCs in SM-exposed patients with long-term pulmonary complications.

Methods: Fifty-nine SM-injured patients with prolonged pulmonary complications and 20 healthy individuals were included. Patients were classified based on taking drugs, having comorbidities, and respiratory consequence severity. MSCs with phenotype of CD45-CD44+CD29+CD105+ were measured in peripheral blood by using flow cytometry.

Results: Circulating MSCs were lesser in SM-exposed patients rather than the control group (median = 0.93 and 2.72 respectively, $p = 0.005$). No significant difference was observed in the MSC population between patients taking systemic corticosteroids or antibiotics and did not taking them, but comorbidities like liver and kidney diseases, and asthma had changed the peripheral level of MSCs in SM-exposed subjects. In addition, circulating MSCs did not show association with long-term pulmonary complication severity.

Conclusion: SM exposure causes decline in the circulating MSCs in survivors. Lower level of peripheral MSC population in SM-exposed patients is not affected by taking corticosteroids or antibiotics, but comorbidities are probably involved in MSC trafficking. Decreased circulating MSCs do not associate with pulmonary complications' severity; however, further studies in mustard lung models are required to demonstrate the therapeutic or pathologic role of MSCs in SM injuries.

Keywords: Mesenchymal Stem Cells, Mustard Gas, Lung, Flow Cytometry.



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Immunology of Exercise, Aging, and Nutrition

Oral Presentation

ICIA2018\Immunology of Exercise, Aging, and Nutrition\Oral\7500
7500

Immuno-potential Effects of Donkey's Milk on Human Mononuclear Cells

Seyedeh Forough Sadeghian-Kiadehi¹, Saeid Taghiloo^{1,2}, Esmail Allahmoradi^{1,2}

1. Department of Immunology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

2. Student Research Committee, Mazandaran University of Medical Sciences, Sari, Iran

Background: Nowadays, donkey's milk consumption has been reevaluated for their potential benefits to human health. In infants with intolerance to cow's milk, donkey's milk represents a good alternative due to its chemical characteristics similar to those of human milk. In the present in vitro study, the pro- and anti-inflammatory effects of donkey's milk were evaluated on human peripheral blood mononuclear cells (PBMC).

Methods: PBMCs were isolated from 12 young and 12 aged normal subjects using Lymphosep media. 2×10^5 cells were cultured in 0.2 ml of RPMI medium supplemented with 10 μ l of each the optimal and non-cytotoxic dose of pasteurized donkey's milk, 100 IU/ml penicillin, 100 μ g/ml streptomycin and 10% fetal calf serum in flat-bottom 96-microwell culture plates. PolymyxinB (PMB-50 μ g/mL) was added to all wells to inhibit the possible endotoxin contamination. Lipopolysaccharide (LPS-100 μ g/mL) was also applied as positive control. All cultures were performed in triplicate and plates were incubated at 37°C in a humidified atmosphere with 5% CO₂. After 18 hours, culture supernatants were harvested to measure TNF- α , IL-6, IL-8 and IL-10 by ELISA.

Results: Donkey's milk significantly increased TNF- α (p: 0.01), IL-8 (p< 0.0001), IL-6 (p< 0.0001) and IL-10 (p: 0.01) levels in PBMCs. In addition, the levels of IL-6 (p: 0.002), IL-8 (p: 0.002) and TNF- α (p: 0.002) secreted from aged subjects were significantly higher than young subjects. Contrast with these data, the level of IL-10 was markedly reduced in aged subjects (p: 0.02).

Conclusion: Our findings highlight the immune-potential effects of donkey's milk, and suggest that donkey's milk might be investigated as a beneficial dietary component to up-regulate the immune response in aged people.

Keywords: Donkey's milk, Immuno-potential, aged subject, young subjects, LPS, Cytokine

ICIA2018\Immunology of Exercise, Aging, and Nutrition\Oral\7596

7596

The effect of vitamin D supplements on the frequency of CD4⁺T cell subsets in women with Hashimoto Thyroiditis, a Double-Blind Placebo-Controlled Study

Nodehi M¹, Ajami A¹, Izad M², Asgarian-Omran H¹, chardouli R³, Amouzegar A³, Azizi F³, Hematabadi M⁴, Esfahanian F⁴, Yekaninejad S⁵, Saboor-Yaraghi AA⁶

1. Department of Immunology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran and Molecular and Cell Biology Research Center, Mazandaran University of Medical Sciences, Sari, Iran

2. Department of Immunology, School of Medicine, International Campus, Tehran University of Medical Sciences, Tehran, Iran

3. Endocrine Research Center, Research Institute for Endocrine Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran

4. Department of Endocrinology, Vali-Asr Hospital, Imam Khomeini Complex Hospital, Tehran University of Medical Sciences, Tehran, Iran

5. Department of Epidemiology and Biostatistics, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

6. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

Background: In addition to its known effects on bone metabolism, vitamin D may regulate immune function. Investigations on the pathogenesis of Hashimoto Thyroiditis and the mechanism of vitamin D action have been revealed that vitamin D may be a good choice to reduce damage to the thyroid cells caused by autoreactive immune cells. Considering the important role of CD4⁺ T cells in the pathogenesis of Hashimoto's thyroiditis, the effect of this vitamin on the frequency of Th1, Th2, Th17 and Tr1 cells in patients with Hashimoto Thyroiditis was explored.

Methods: 34 patients aged 20-45 years were introduced by endocrinologists and entered the study. Patients with history of other autoimmune disease, allergy, malignancy, chronic kidney or liver disease were excluded from the study. They were recruited into two groups including 17 patients in each group. Group I and II were treated weekly with 50,000 IU of cholecalciferol and placebo respectively, for 3 months. Before and after supplementation, the frequency of Th1, Th2, Th17 and Tr1 cell was evaluated using flowcytometry method.

Results: The results showed that cholecalciferol supplementation caused a significant decrease in the Th17/Tr1 ratio, however, had no significant effect on percentages of Th1, Th2, Tr1 and Th17 cells in Hashimoto's thyroiditis patients.

Conclusion: In this study that conducted on women with Hashimoto Thyroiditis, supplementation by cholecalciferol for 3 months could change the phenotype of CD4⁺ T cell lymphocytes. Therefore, this vitamin can be considered as a supplement to prevent and treat the disease.

Keywords: Cholecalciferol, Hashimoto Thyroiditis, Th17 Cells, Tr1 Cells, Th1 cells, Th2 cells

ICIA2018\Immunology of Exercise, Aging, and Nutrition\Oral\8228
8228

Investigating the ratio of Th1/Th2 cells after three sessions of endurance exercise as an adjuvant in HSV-gd1 DNA vaccine

Mohammad Taher Afshounpour¹, Mahdieh Molanouri Shamsi¹, Afsaneh Jamali¹, Sara Soudi², Hoorieh Soleimanjahi³, Reza Gharakhanlou¹

1. Department of Physical Education and Sports Science, University of Tarbiat Modares, Tehran, Iran

2. Department of Immunology, University of Tarbiat Modares, Tehran, Iran

3. Department of Virology, University of Tarbiat Modares, Tehran, Iran

Background: The relations between the vaccination, exercise and adjuvant effects of exercise are a new perspective in the exercise immunology, but there are many questions about it and needs extensive research. Therefore, the aim of this study was to investigate the ratio of Th1/Th2 cells after three sessions of endurance exercise as an adjuvant in HSV-gd1 DNA vaccine.

Methods: Balb/c mice (6 to 8 weeks) were randomly assigned into six groups, 1- Endurance exercise, 2- Vaccine, 3-PCDNA, 4- Control, 5- Endurance exercise + Vaccine and 6- Endurance exercise + PCDNA. Endurance exercise was performed sequentially for three days on the animal treadmill. Immediately after the third session, HSV-gD1 DNA vaccine was injected into the tibialis muscle of mice. 12 days after the second vaccine, in spleen INF- γ and IL-4 cytokines were evaluated by ELISA method. One-way ANOVA was used to analyze the data.

Results: The results showed that three sessions of endurance exercise could significantly increase the ratio of Th1/Th2 cells in the HSV-gD1 vaccine model. Also, there is a significant difference between this group and other groups, especially in the vaccinated group.

Conclusion: According to the findings, three sessions of endurance exercise act as an adjuvant in HSV-gd1 DNA vaccine and enhance the immune response.

Keywords: Th1/Th2 cells, endurance exercise, adjuvant, DNA vaccine.

ICIA2018\Immunology of Exercise, Aging, and Nutrition\Oral\8506

8506

Effects of vitamin D on the expression of Th17 cell-related cytokines in experimental autoimmune encephalomyelitis

Sayyed Vahab Azizi¹, Rayhaneh Ahangar-Parvin¹, Marziyeh Mohammadi-Kordkhai²,
Mohammad-Taghi Rezayati¹, Maryam Nemati³, Fatemeh Ayoobi⁴, Zahra Taghipour⁵,
Hossain Khoramdelazad¹, Abdollah Jafarzadeh^{*2}

1. Department of Immunology, Medical School, Rafsanjan University of Medical Sciences, Rafsanjan, Iran.
2. Department of Immunology, Medical School, Kerman University of Medical Sciences, Kerman, Iran.
3. Department of Laboratory Sciences, Para-Medicine School, Kerman University of Medical Sciences, Kerman, Iran.
4. Department of Physiology, Medical School, Rafsanjan University of Medical Sciences, Rafsanjan, Iran.
5. Department of Histology, Medical School, Rafsanjan University of Medical Sciences, Rafsanjan, Iran.

Background: A broad of immunomodulatory properties is attributed to vitamin D. The aim was to evaluate the vitamin D effects on the expression of some Th17 cell-related cytokines in an experimental autoimmune encephalomyelitis (EAE) model.

Methods: The C57Bl/6 mice used for EAE induction by immunization with myelin oligodendroglial glycoprotein mixed with Freund's adjuvant. From day +3 to +30, the mice were administrated by Intraperitoneal injection of PBS or olive oil in control groups and vitamin D in treatment group. At day 31, mice were scarified and the expression of IL-17A and IL-23 in the spinal cord and serum were measured by real time-PCR and ELISA, respectively.

Results: The expression of IL-17, IL-23 P19 and IL-23 P40 in spinal cord and serum IL-17 and IL-23 levels in PBS-administrated EAE mice was significantly higher than healthy group ($P<0.02$, $P<0.001$, $P<0.001$, $P<0.04$ and $P<0.05$ respectively). In EAE mice treated with vitamin D, the expression of aforementioned parameters were significantly decreased as compared with control group ($P<0.05$, $P<0.001$, $P<0.001$, $P<0.04$ and $P<0.02$ respectively).

Conclusion: Vitamin D downregulates the Th17 cell-related cytokines in EAE mice. The possible therapeutic potential of vitamin D for treatment of MS can be consider in future investigation.

Keywords: Experimental autoimmune encephalomyelitis, Vitamin D, IL-17, IL-23

ICIA2018\Immunology of Exercise, Aging, and Nutrition\Oral\9771
9771

Comparative study of the combinatory effects of Docosahexaenoic acid (DHA) and Linoleic acid (LA) with Taxol on the expression of metastasis related genes and microRNAs in triple-negative metastatic breast cancer cell line

M. Javadian^{1,2}, N. Shekari¹, M. Soltani¹, A. Mohammadi¹, B. Baradaran¹, D. Shانهbandi¹, T. Kazemi¹

1. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

2. Student Research Committee, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Involvement of some genes and microRNAs (miRNAs) in metastatic breast cancer has been extensively investigated. However, there are limited numbers of study regarding the effects of standard anti-cancer therapeutic agents and also common dietary supplements on the expression of such genes and miRNAs. In this study, we aimed to investigate the effects of DHA and LA fatty acids on the expression of matrix metalloproteinase (MMP)-9, 2, Talin2, HIF-1 α and vimentin genes, and tumor-suppressor miR-194 and 106b oncomiR, and also their possible synergistic effects with Taxol.

Methods: MDA-MB-231 Cells were cultured under normoxic and hypoxic conditions, and treated for 24 hours with taxol in IC50 concentration, 100 μ M DHA and 50 μ M LA, alone or in different combinations. Cells were harvested, and RNA/miRNA extraction and cDNA synthesis were performed using standard methods. Expression levels of studied genes and miRNAs were analysed using quantitative real-time PCR. Western blotting and scratch test were also performed to confirm findings.

Results: Taxol and DHA significantly down-regulated MMP-9, MMP-2, HIF-1 α and vimentin expression and up-regulated Talin2 in normoxic and hypoxic condition ($p < 0.05$). But our results showed that LA induced up-regulation of MMP-2 and MMP-9, vimentin expression in these conditions ($p < 0.05$). miR-194 and miR-106b, showed up-regulated and down-regulated expression in all treated respectively. The result of western blotting for Talin2 didn't confirm the effects of different treatments, but scratch test showed inhibitory effect of DHA alone and in combination with Taxol.

Conclusion: Antimetastatic effects of DHA and Taxol, alone or in combination, were confirmed in this study. MMP-2, MMP-9, HIF-1 α and vimentin showed down-regulation after treatment. Consistent result were found about down-regulation of miR-106b oncomir and up-regulation of tumor-suppressor miR-194. But their role in altered expression of Talin-2 needs for more investigation.

Keywords: Docosahexaenoic acid, Breast cancer, MicroRNA, Metastasis.



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Immunology of Infectious Diseases

Bacteria and Fungi

Oral Presentation

ICIA2018\Immunology of Infectious Diseases Bacteria and Fungi\Oral\7815
7815

Comparison the percentages of Th1 and Th17 cells in untreated and antimicrobial treated patients with brucellosis

Mina Rahmanpour¹, Sanaz Jourghasemi¹, Ghasem Solgi¹, Fariba Keramat², Mehrdad Hajilooi^{1*}

1. Department of Immunology, School of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

2. Department of Infectious Diseases, School of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran; Brucellosis Research Center, Hamadan University of Medical Sciences, Hamadan, Iran

Background: Brucellosis is a worldwidelly zoonotic infection caused by brucella, facultative intracellular Gram-negative coccobacilli bacteria. Host protection against brucella relies on cell-mediated immunity. T helper type 1 (Th1) cells play a critical role in immunity to brucellosis, while role of T helper type 17 (Th17) cells against the infection remained unknown. In this study aim to compare the percentages of Th1 and Th17 cells in untreated and antimicrobial treated patients with brucellosis.

Methods: Peripheral blood mononuclear cells (PBMCs) isolated from 15 untreated brucellosis patients (new cases) and 15 treated brucellosis patients with antimicrobial treatment (30 study subjects) using ficoll density gradient centrifugation. PBMCs cultured with phorbolmyristate acetate (PMA, 50 ng/ml), ionomycin (1 µg/ml) and brefeldin A (5 µg/ml) for 5 h at 37°C in 5% CO₂. Subsequently, flowcytometry analysis performed for CD4⁺IFN-γ⁺ Th1 and CD4⁺IL-17⁺ Th17 cells.

Results: The percentage of Th1 cells were increased in new cases in comparison to patients under treatment (Th1: 3.57±1.86% vs 2.79±2.25%, P=0.34). Also, there was an increase in the count of Th17 cells among untreated and treated patients (Th17: 0.35±0.14% vs 0.25±0.14%, P=0.07).

Conclusion: In this study, we demonstrated the increase percentages of Th1 and Th17 cells in untreated patients. This results indicate that immune responses related to this cells are important in patients with brucellosis.

Keywords: Brucellosis, Th1 cells, Th17 cells

ICIA2018\Immunology of Infectious Diseases Bacteria and Fungi\Oral\8085
8085

PBMCs of patients with abdominal aortic aneurysm in co-culture with endothelial cells produce lower levels of IL-9 in comparison to control group

Hamid Aria^{1,2}, Mehdi Kalani³, Hossein Hodjati⁴, Mehrnoosh Doroudchi¹

1. Department of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran.

2. Bioinformatics and Computational Biology Research Center, Shiraz University of Medical Sciences, Shiraz, Iran.

3. Prof. Alborzi Clinical microbiology research center, Nemazee hospital, Shiraz University of Medical Sciences, Shiraz, Iran.

4. Department of Vascular Surgery, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran.

Background: Cytokines are major players in inflammatory condition leading to diverse diseases. Abdominal Aortic Aneurysm (AAA) is the dilatation of abdominal aorta, which happens in some of the patients with atherosclerosis. In an attempt to mimic the interaction of blood cells with wounded endothelium, we investigated the effect of CagA⁺ and CagA⁻ *Helicobacter pylori* extracts in co-culture of PBMCs with endothelial cells (EC).

Methods: PBMCs were isolated from 5 men with diagnosed AAA and 5 men with normal/insignificant angiography, CT-scan and ultrasonography results. EC cells were extracted from umbilical cords by collagenase method (HUVEC). PBMCs were cultured in plates with bacterial extract of CagA⁺ and CagA⁻ *Helicobacter pylori* in co-culture with HUVEC cells for 48 hrs. Then, the supernatant were removed to measure IL-2, IL-4, IL-5, IL-6, IL-9, IL-10, IL-13, IL-17A, IL-17F, IL-21, IL-22, IFN- γ and TNF- α using a commercial fluorescent-labeled bead assay.

Results: IL-9 production by patients' PBMCs without antigenic stimulation in co-culture with HUVECs (1403.07 ± 2856.07 Pg/ml) was lower than controls (12412.00 ± 7399.47 Pg/ml) ($P=0.05$). Production of IL-22 by PBMCs in response to CagA⁻ *H. pylori* extract in co-culture with HUVECs (22.82 ± 11.37 Pg/ml) was lower compared to controls (37.21 ± 10.76 Pg/ml) ($P=0.05$).

Conclusion: The difference in the cytokine production pattern in patients with AAA and controls suggest the importance of these factors in the initiation/progression of AAA.

Keywords: Abdominal aortic aneurysm, cytokines, *Helicobacter pylori*, endothelial cells

ICIA2018\Immunology of Infectious Diseases Bacteria and Fungi\Oral\8415
8415

The capability of recombinant Hcc fragments in the activation of T cells was assessed by CD69 expression

Saeedeh Torabi Goudarzi¹, Mehdi Yusefi. PhD^{1,2}, Jafar Majidi. PhD^{1,2}

1. Department of Immunology, Faculty of Medicine Tabriz University of Medical Sciences, Tabriz, Iran

2. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

Background: A protective response against tetanus toxin and toxoid demands efficient specific T cell and B cell responses. Tetanus neurotoxin (TeNT), a 150 kDa polypeptide, is the main cause of tetanus disease. TeNT consists of two structurally distinct chains, a 50 kDa N-terminal light (L) and a 100 kDa C-terminal heavy (H) chain. C-terminal heavy (H) chain (fragment C) has two sub-domains named as proximal HCN and carboxyl sub-domain or HCC. Beside neural binding property, HCC has been recently found as an immunodominant part of TeNT. In the present study, we investigated the effects of recombinant HCC (rHCC) on the capability of recombinant Hcc fragments in the activation of T cells was assessed by CD69 expression.

Methods: In the present study, we isolated human T cells by magnetic-activated cell sorting method (MACS), Purified T cells were co-incubated with recombinant Hcc. The concentration of different cytokines were determined by ELISA and then we were analyzed the expression of markers of lymphocyte T activity by flowcytometry.

Results: The results showed that rHcc enhanced expression of CD69 on the surface of T-cells and promoted differentiation of CD4⁺ T-lymphocytes toward a T-helper1 (T_H1) phenotype and upregulation of interferon (IFN)- γ secretion.

Conclusion: These results indicated that HCC is critical for immunogenic and immunoprotective activity of TeNT and is able to induce T cells and has a stimulatory effect on IFN- γ secretion. This subdomain can propose as an adjuvant and a candidate to stimulate the immune system.

Keywords: Tetanus, Tetanus neurotoxin, Immune Responses, Hcc subdomain

ICIA2018\Immunology of Infectious Diseases Bacteria and Fungi\Oral\9779
9779

Aspergillus fumigatus conidia stimulate cytokines release from epithelial cells (TC-1 JHU-1): a study of modulatory effect of propolis extract

Ali Reza Khosravi¹, Hadis Moradjodaki²

1. Department of Microbiology and Immunology, University of Tehran, Tehran, Iran

2. Department of Microbiology and Immunology, University of Tehran, Tehran, Iran

Background: Aspergillus fumigatus conidia are the most prevalent indoors fungal allergens. The interaction between Aspergillus antigens and lung epithelial cells result in innate immune functions. The association between Aspergillus conidia and allergic reactions, like allergic bronchopulmonary aspergillosis (ABPA) and asthma have been repeatedly reported. The interaction between Aspergillus antigens and lung epithelial cells result in innate immune functions. Since conventional therapies for allergy and asthma are limited, finding new promising treatments are inevitable. This study was designed to evaluate the effect of Aspergillus fumigates conidia on some cytokine releases from mouse epithelial cells and to investigate the effect of propolis on cytokines modulation.

Methods: Cells were divided to two groups, one was exposed to 3×10^4 conidia of Aspergillus fumigatus and another group was treated by propolis (25 µg/ml) as well as exposure to A. fumigatus conidia. Cytokines IL-13, IL-12, IFN γ and IL-17 were measured at times 0, 6 and 12 hours after exposure using ELISA assay.

Results: The results indicated that A. fumigatus could increase the release of understudied cytokines with IL-13 and IL-17 being the most affected one, whilst treatment with propolis decreased the effects of A. fumigatus on IL-13 and IL-17 production. The results showed that propolis has down regulatory effects on Th2 cytokine, IL13, and IL17 production, whereas it caused a significant induction of IL-12, as an important Th1 cytokines by lung epithelial cells (LECs).

Conclusion: With respect to the obtained results, propolis extract might be contributed to decrease Th2 responses in allergic asthma phenomenon. However more investigations must be done in future to fully understand its efficacy.

Keywords: Aspergillus fumigatus, cytokines, epithelial cells, propolis

ICIA2018\Immunology of Infectious Diseases Bacteria and Fungi\Oral\9892
9892

NRWCFAGRR: A Novel Antimicrobial Peptide with Nitric Oxide Induction Activity

Mehdi Ghahary¹, Kazem Ahmadi¹, Mahdi Fasihi-Ramandi¹

1. Molecular Biology Research Center, system biology and poisonings institute, Baqiyatallah University of Medical Sciences, Tehran, Iran

Background: Bacterial resistance against antibiotic has caused many problems in the procedure of treating humans and animals all around the world. Inappropriate utilization of antibiotics can result to resistance in the target microorganism population. Antimicrobial peptides have been introduced as new effective strategies that kill bacteria quickly and cause less antibiotic resistance.

Methods: In the present research we have been studied the cytotoxic effect of NRWC peptide on eukaryotic and prokaryotic cells. Twelve bacterial strains were selected to study the antimicrobial effects of NRWC peptide. MIC and MBC assay were used to study the inhibitory and killing activity of these peptides respectively. Peptide cytotoxic effects were assayed on HeLa cell line and human RBC using MTT assay and Hb release measurement respectively. J774 macrophage cell line was used to measure the amount of nitric oxide production in response to the peptide.

Results: The results showed that NRWCFAGRR peptide in different concentrations had bactericidal and inhibitory effect on the growth of all 12 bacterial strains in a dose dependent manner. It has also been proven that the toxic effect of peptide on human cells is inevitable at the MIC and MBC concentration. The highest amount of nitric oxide produced under the influence of this peptide was observed in a period after 48 hours treatment.

Conclusion: Considering the researches have been conducted in the field of microbial peptides, the peptide introduced in this research has anti-microbial properties that kills some microbes directly and theoretically has the ability to kill some organisms indirectly via induction of nitric oxide in the macrophages.

Keywords: Antibiotics resistance, Nitric oxide, MBC, MIC, Antimicrobial peptide



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Immunology of Infectious Diseases

Viruses

Oral Presentation

ICIA2018\Immunology of Infectious Diseases Viruses\Oral\6358
6358

Designing a DNA Vaccine for Therapeutic Goals in Cancer Mouse Model of Human Papilloma Virus Type-16

Ali Ganji^{1,2}, Hamid Abtahi^{1,2}, Ghasem Mosayebi^{1,2}, Iman Farahani^{1,2}, Ali Ghazavi^{1,2}, Maryam Moazamigoodarzi¹, Mostafa Ejtehadifar¹, Saeed Babaie³

1. Molecular and Medicine Research Center, Arak University of Medical Sciences, Arak, Iran.

2. Department of Microbiology and Immunology, School of Medicine, Arak University of Medical Sciences, Arak, Iran.

3. Department of Anatomy, Arak University of Medical Sciences, Arak, Iran

Background: Infection with human papillomavirus type 16 (HPV 16) causes cervical cancers. Although, many attempts have been made to treat of these infections, but further studies are necessary to find new therapies. So, the aim of this study was setting light on therapeutic potencies of HPV 16-derived DNA vaccines to introduce a treatment based on improving the immune system functions.

Methods: DNA vaccine for optimum expression in bacteria was designed and the recombinant protein was expressed. The designed DNA vaccine was injected via intramuscular route in mouse model of HPV in compare with intra-peritoneal injection of recombinant protein with adjuvant. The induced mice were monitored based on animal survival, tumor size, and immunologic responses.

Results: Comparisons illustrated meaningful decrease in tumor size ($P < 0.05$) of treated with a mixture of DNA vaccine and protein in contrast to the other groups. Leukocyte infiltration and proliferation were significantly showed increased number in the treated group with DNA in comparison with control group ($P < 0.05$). The level of IgG2a showed an increasing in DNA vaccine and mixture of DNA vaccine and protein in comparison with control group ($P < 0.05$). Stimulated splenocytes of the treated mice with mixture of protein and DNA vaccine expressed significantly higher interferon gamma ($\text{INF-}\gamma$) than the untreated group ($P < 0.05$). The meaning percentage of cytotoxic T lymphocytes in treated group with DNA vaccine was more than other groups ($P < 0.05$).

Conclusion: The designed DNA vaccine is suitable candidate for therapeutic goals and results of present work proved that DNA prime/Protein boost approaches or injection of primary doses of DNA vaccine and booster doses of protein can be an impressive way to treatment of those cancers which may be a result of papilloma infections.

Keywords: Recombinant proteins, DNA vaccine, Vaccine therapy, Human papilloma virus

ICIA2018\Immunology of Infectious Diseases Viruses\Oral\7444
7444

Increase of CD4⁺ CD25⁺ CD127^{low} FoxP3⁺ regulatory T cells, but not CD4⁺ CD25⁺ FoxP3⁺ cells in active hepatitis B

Mousa Mohammadnia-Afrouzi¹, Mehdi Shahbazi¹, Hiva Mohammadizad¹, Mojgan Bagherzadeh¹

1. Department of Immunology, School of Medicine, Babol University of Medical Sciences, Babol, Iran

Background: To investigate immune mechanisms underlying the difference between active and inactive hepatitis B, we analyzed regulatory T cell frequency in chronic hepatitis B patients.

Methods: 66 chronic hepatitis B patients were included in the study. Patients were divided in active (32 patients) and inactive (34 patients) groups based on laboratory parameters. Isolation of peripheral blood mononuclear cells (PBMCs) was carried out by using Ficoll-Paque density centrifugation. The frequency of Treg cells and FoxP3 expression level were evaluated by multicolor flowcytometry, after staining PBMCs with fluorochrome-conjugated antibody against CD4, CD25, CD127 and FoxP3. Mean Florescence intensity (MFI) was used as index of FoxP3 expression in Treg Cells.

Results: The frequency of CD4⁺ CD25⁺ CD127^{low} FoxP3⁺ Treg cells was higher in active hepatitis compared to inactive patients. Additionally, there was a positive correlation between HBV viral load and frequency of CD4⁺ CD25⁺ CD127^{low} FoxP3⁺ Treg cells in active patients. There was not any significant differences in other subtypes of Treg cells between groups. However, MFI of FoxP3 was directly correlated with HBV viral load.

Conclusion: Results of this study suggest that dynamic changes in CD4⁺ CD25⁺ CD127^{low} FoxP3⁺ Treg cell frequency and FoxP3 expression show different trends in active and inactive hepatitis B patients.

Keywords: Treg, Hepatitis B, Active, Inactive, FoxP3

ICIA2018\Immunology of Infectious Diseases Viruses\Oral\7488
7488

TIM-3 as a marker of exhaustion in CD8⁺ T cells of active chronic hepatitis B Patients

Hiva Mohammadizad¹, Parizad Bavandpour-Baghshahi¹, Mousa Mohammadnia-Afrouzi¹, Mehdi Shahbazi¹,
Mojgan Bagherzadeh¹

1. Department of Immunology, School of Medicine, Babol University of Medical Sciences, Babol, Iran

Background: Chronic hepatitis B is the most important chronic viral infection affecting 5 percent of the world population. Distinguish of the inactive patients from active, play an important role in treatment of patients. Virus-specific CD8⁺ T cells lose effector function during chronic HBV infection and can express inhibitory receptors, TIM3, PD-1, and CD39, that are key feature of CD8⁺ T Cells exhaustion in humans. In this study, we analyzed the level of PD-1, Tim-3 and CD39 expression in CD8⁺ T cells isolated from peripheral blood mononuclear cells of patients with active and inactive chronic hepatitis B.

Method: In the study, 34 inactive and 32 active chronic hepatitis B patients were included. Isolation of peripheral blood mononuclear cells (PBMCs) was done by Ficoll-Paque density centrifugation. The level of PD-1, Tim-3 and CD39 expression in CD8⁺ T cells of Patients determined by multicolor flo cytometry after staining PBMCs using fluorochrome-conjugated antibody against CD39, Tim3, PD1, CD8.

Results: CD8⁺ T cells frequency was significantly lower in patients with active chronic B hepatitis than inactive hepatitis B ($p = 0.0001$). CD8⁺ TIM3⁺ T cells and also CD8⁺TIM3⁺PD1⁺ T cells were significantly higher in patients with active chronic hepatitis B than inactive ones. However, PD1 and CD39 markers did not show significant differences. In active chronic phase, there was a positive correlation between AFP, ALT, and HBV viral load with CD8⁺ TIM3⁺ T cells frequency.

Conclusion: In active chronic hepatitis B, the number of exhausted T cells, due to the virus involvement, are more than the inactive and TIM3 seems to be a marker of exhaustion in CD8 T cells of these patients.

Keywords: exhausted, active, hepatitis B, CD8, TIM3

ICIA2018\Immunology of Infectious Diseases Viruses\Oral\7757

7755

Blocking of opioid receptors in experimental formalin-inactivated respiratory syncytial virus (FI-RSV) immunopathogenesis: from beneficial to harmful impacts

Vahid Salimi¹, Habibollah Mirzaei², Ali Ramezani², Alireza Tahamtan¹, Fatemeh Adjaminejad¹, Ghazal Fatemi-Nasab¹, Fazel Shokri³, Talat MokhtariAzad¹.

1. Department of Virology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

2. Department of Virology, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz

3. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

Background: Opioid system plays a significant role in pathophysiological processes, such as immune response and impacts on disease severity. Here, we investigated the effect of opioid system on the immunopathogenesis of respiratory syncytial virus (RSV) vaccine (FI-RSV)-mediated illness in a widely used mouse model.

Methods: Female Balb/c mice were immunized at days 0 and 21 with FI-RSV (2×10^6 pfu, i.m.) and challenged with RSV-A2 (3×10^6 pfu, i.n.) at day 42. Nalmefene (NL) as a universal opioid receptors blocker administered at a dose of 1 mg/kg in combination with FI-RSV (FI-RSV+NL), and daily after live virus challenge (RSV+NL). Mice were sacrificed at day 5 after challenge and bronchoalveolar lavage (BAL) fluid and lungs were harvested to measure airway immune cells influx, T lymphocyte subtypes, cytokines/chemokines secretion, lung histopathology, and viral load.

Results: Administration of nalmefene in combination with FI-RSV (FI-RSV+NL-RSV) resulted in the reduction of the immune cells infiltration to the BAL fluid, the ratio of CD4/CD8 T lymphocyte, the level of IL-5, IL-10, MIP-1 α , lung pathology, and restored weight loss after RSV infection. Blocking of opioid receptors during RSV infection in vaccinated mice (FI-RSV-RSV+NL) had no significant effects on RSV immunopathogenesis. Moreover, administration of nalmefene in combination with FI-RSV and blocking opioid receptors during RSV infection (FI-RSV+NL-RSV+NL) resulted in an increased influx of the immune cells to the BAL fluid, increases the level of IFN- γ , lung pathology, and weight loss in compared to control condition.

Conclusion: Although nalmefene administration within FI-RSV vaccine decreases vaccine-enhanced infection during subsequent exposure to the virus, opioid receptor blocking during RSV infection aggravates the host inflammatory response to RSV infection. Thus, caution is required due to beneficial/harmful functions of opioid systems while targeting as potentially therapies.

Keywords: Respiratory syncytial virus, Formalin-inactivated RSV, Immunopathogenesis, Nalmefene.

ICIA2018\Immunology of Infectious Diseases Viruses\Oral\8181
8181

Measles virus infected dendritic cell migration in a 3D in vitro human airway test system

Shaghayegh Derakhshani¹, Andreas Kurz², Sibylle Schneider-Schaulies¹

1. Department of biophysics and biotechnology, University of Würzburg, Germany

2. Department of Virology and Immunobiology, University of Würzburg, Germany

Background: Parameters important in Measles Virus (MV) transmission from dendritic cells (DCs) to epithelial cells such as cytoskeletal or receptor dynamics were determined in 2D systems. It remains unclear whether and to what extent these apply in a more complex environment such as the respiratory tract epithelium. Therefore we analyze these parameters in 3D models of the airway mucosa with high in vitro/in vivo correlation.

Methods: 3D models Fibroblasts are seeded apically onto a decellularized porcine intestine derived scaffold, and H358 epithelial cells are subsequently added; models are cultivated under airlift conditions. Cellular composition of the 3D models was evaluated by immunohistochemistry (IHC) and immunofluorescence (IF). DCs infected with MV (IC323-eGFP) for 24 h were applied basolaterally to the collagen scaffold. DC trafficking towards epithelial cells was recorded by time lapse microscopy. Spread of MV infection to epithelial cells was microscopically monitored and quantified by flowcytometry.

Results: IHC/IF stainings of paraffin embedded models performed 24 h following basolateral DC application allowed for distinction of layers of fibroblasts and epithelial cells with DCs interspersed in the epithelial layer. To visualize DC migration through the models, cells were tracked 30 min following their basolateral application in a 3h 3D time series. Remarkably, MV infected DCs migrated more efficiently towards the apical surface of the models than uninfected cells indicating that MV-infection enhances DC tissue motility. In support of this hypothesis, pAkt levels were found elevated in MV-infected DCs. In addition to that of proteins, lipid sorting is important in membrane dynamics and thereby, in viral transmission. Accordingly, pharmacological inhibition of the acid sphingomyelinase (ASM) in DCs efficiently modulated viral transmission to epithelial cells in 2D and 3D conditions.

Conclusion: In the current study we showed that MV infection enhanced DCs motility in a 3D environment. Moreover, sphingolipid turnover is of obvious importance in viral transmission to epithelial cells. Current analyses address subcellular redistribution of ceramides in infected DCs to further unravel their role in this process.

Keywords: Measles virus, dendritic cells, cell migration, 3D tissue models

ICIA2018\Immunology of Infectious Diseases Viruses\Oral\8255

8255

Induction of inflammatory mediators by Influenza A virus protein PB1-F2 in J774.1 macrophage

Zahra Farzaneh¹, Nasser Gholijani², Kurosh Kalantar¹

1. Department of Immunology, Shiraz University of Medical Sciences, Shiraz, Iran

2. Autoimmune Disease Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Influenza A virus (IAV) has potential to cause mortal pandemics with considerable health and socio-economic burdens. Some of IAV express virulence factor PB1-F2. The PB1-F2 protein has pro-apoptotic activity and contribute to viral pathogenesis by delaying in viral clearance and inducing inflammation. Macrophages are susceptible to IAV infection and produce high levels of inflammatory cytokines and chemokines.

Methods: The cytotoxic effect of different concentrations of PB1-F2 protein on J774.1 macrophage was determined by MTT assay. Then, the pro-inflammatory effect of PB1-F2 on the expression of iNOS, COX-2, IL-6 and TNF- α genes were evaluate by the real-time PCR. Production of key inflammatory mediators including IL-6 and TNF- α were assessed using ELISA assay. In addition, PB1-F2 treated macrophages examined for nitric oxide (NO) production by means of griess reagent.

Results: Treatment of cells with 0.22 and 0.45 $\mu\text{mol/ml}$ of peptide increased NO production to $105.8 \pm 1.24\%$ and $119.5 \pm 12.8\%$ of the LPS-only treated cells, respectively. Peptide at 0.9 $\mu\text{mol/ml}$ concentration significantly increased gene expression of iNOS and COX-2. Gene expression increased to 2.02 ± 0.25 RFC ($p < 0.001$) (iNOS) and 3.81 ± 0.74 RFC ($p < 0.0001$) (COX-2) of the LPS-only treated cells (positive control). Treatment of cells with concentrations 0.45 and 0.9 $\mu\text{mol/ml}$ of peptide significantly induced gene expression of pro-inflammatory cytokines (IL-6, TNF- α). Peptide at these concentrations increased IL-6 gene expression to 2.19 ± 0.73 RFC and 3.65 ± 0.3 RFC. At the same concentrations of peptide, TNF- α gene expression showed an increasing level to 4.15 ± 0.86 RFC and 5.55 ± 0.44 RFC of positive control, respectively. Both of cytokines in protein level also increased.

Conclusion: These finding suggest that PB1-F2 protein is a potent inducer of inflammatory mediators in macrophages that may contribute to diseases severity.

Keywords: Influenza A virus, PB1-F2, Inflammation, Macrophage

ICIA2018\Immunology of Infectious Diseases Viruses\Oral\8539
8539

Investigating the Effects of different vaccination schedules on Hepatitis vaccine induced antibody titers in ICU Personnel of Kowsar Hospital of Sanandaj, 2018

Pejman Allahveisey¹, Jalil Adabi², Asrin Safari², Elahe Amani²

1. *School of nursing, University of medical sciences Iran*

2. *School of nursing, University of medical sciences Kurdistan*

Background: Hepatitis type B has been recognized as one of the infections, which can be easily transformed through infected blood. The rate of this kind of infection transmission was reported to be under 1% in the 1970s. The risk of transmission of Hepatitis infection through infected blood has never been completely eliminated, due to the fact that factors such as Window Period and Low Titer of HBV can cause infection transmission. Vaccination is the key step to provide immunization coverage for individuals exposed to Hepatitis.

Method: The participants of this study were the personnel of ICU ward in Kowsar Hospital of Sanandaj. Their antibody titer was measured; afterwards, a checklist was developed for the vaccination project. Twenty-five of the personnel agreed to participate in the present study. The collected data was analyzed using SPSS 18.

Result: The collected data revealed that 64% of the participants had done the vaccination schedule completely for three times, 20% of them only completed the first and the second phases of the schedule, and 16% of them just did the first phase. The results showed that 81.25% of the participants, who had done the vaccination for three times, indicated antibody titer which was over 200.

Conclusion: Completed vaccine schedule can often guarantee healthy individuals in 95% of cases. It can be concluded that personnel of hospitals, in particular those who work in ICU wards, need to check their antibody titer regularly.

Keywords: Hepatitis, Personnel of ICU ward, Antibody Titer



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Immunology of Mycobacterium Tuberculosis

Oral Presentation

ICIA2018\Immunology of Mycobacterium Tuberculosis\Oral\9873

9873

The analysis of exosomal miRNAs released from the human macrophages after infection with bovis bacillus Calmette-Guerin (BCG)

Esmacil Mortaz¹, Shamila D Alipoor², Payam Tabarsi¹, Mehdi Mirsaeidi³, Johan Garssen⁴, Masoud Movassaghi⁵, Ian M Adcock⁶

1. Clinical Tuberculosis and Epidemiology Research Center, National Research Institute of Tuberculosis and Lung Diseases (NRITLD), Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Institute of Medical Biotechnology, Molecular Medicine Department, National Institute of Genetic Engineering and Biotechnology (NIGEB), Tehran, Iran

3. Division of Pulmonary, Critical Care, Sleep and Allergy, Department of Medicine, University of Miami, FL, USA

4. Nutricia Research Centre for Specialized Nutrition, Utrecht, Netherlands.

5. Department of Pathology and Laboratory Medicine, University of California, Los Angeles (UCLA), Los Angeles, CA, USA

7. Airways Disease Section, National Heart & Lung Institute, Imperial College London, London, UK

Background: Tuberculosis (TB) remains a significant global health concern and its diagnosis is challenging due to the limitations in the specificity and sensitivity of the current diagnostic tests. Exosomes are bioactive 30-100nm vesicles that are produced by most cell types and are found in almost all human body fluids. Exosomal microRNAs (miRNAs) can transfer biological information between cells and tissues and may act as potential biomarkers in many diseases. In this pilot study we assessed the miRNA profile of exosomes released from human monocyte-derived macrophages upon infection with Mycobacterium bovis Bacillus Calmette-Guerin (BCG).

Methods: Human monocytes were obtained from the peripheral blood of 3 healthy subjects and driven to a monocyte-derived macrophage (MDM) phenotype using standard protocols. MDMs were infected with BCG or left uninfected as control. 72h post-infection, exosomes were collected from the cell culture medium, RNA isolated and RNA-seq performed. The raw reads were filtered to eliminate adaptor and primer sequences and the sequences were run against the mature human miRNA sequences available in miRBase. MicroRNAs were identified using an E value <0.01. miRNA network analysis was performed using the DIANA miRNA tool, miRDB and functional KEGG pathway analysis.

Results: Infection of MDMs with BCG leads to the release of several exosomal miRNAs. These included miR-1224, -1293, -425, -4467, -4732, -484, -5094, -6848-6849, -4488 and -96 all of which were predicted to target metabolism and energy production-related pathways.

Conclusions: This study provides evidence for the release of specific exosomal miRNAs from BCG-infected MDMs. These exosomal miRNAs reflect host-pathogen interaction and subversion of host metabolic processes following infection.

Keywords: Mycobacterium, Exosome, miRNA, Macrophage, Biomarker

ICIA2018\Immunology of Mycobacterium Tuberculosis\Oral\9869

9869

miR-1224 Expression is Increased in Human Macrophage After Infection With Bacillus Calmette-Guerin (BCG)

Mohammad Varahram¹, Shamila D. Alipoor^{2,3,4}, Esmail Mortaz², Payam Tabarsi², Majid Marjani², Gert Folkerts⁵, Johan Garssen^{5,6}, Ian M. Adcock^{7,8}

1. Mycobacteriology Research Center (MRC) National Research Institute of Tuberculosis and lung diseases (NRITLD), Shahid Beheshti University of Medical Sciences, Tehran, Iran
2. Clinical Tuberculosis and Epidemiology Research Center, National Research Institute of Tuberculosis and Lung Diseases (NRITLD), Shahid Beheshti University of Medical Sciences, Tehran, Iran
3. Institute of Medical Biotechnology, Molecular Medicine Department, National Institute of Genetic Engineering and Biotechnology (NIGEB), Tehran, Iran
4. Department of Biotechnology, School of Advanced Technologies in Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran
5. Division of Pharmacology, Utrecht Institute for Pharmaceutical Sciences, Faculty of Science, Utrecht University, Utrecht, Netherlands
6. Nutricia Research Centre for Specialized Nutrition, Utrecht, Netherlands
7. Airways Disease Section, National Heart & Lung Institute, Imperial College London, London, UK
8. Priority Research Centre for Healthy Lungs, Hunter Medical Research Institute, The University of Newcastle, Newcastle, New South Wales, Australia

Background: Tuberculosis (TB) remains a major threat to human health. Understanding the strategies mycobacterium takes to overcome immune defense is important to control the infection. miRNAs are master regulators of most pathways in the human body. Infection with mycobacterium affects upon host metabolic pathways to obtain the nutrition for its intracellular survival. In this study, we aimed to investigate the effect of BCG infection on the expression of three miRNAs (miR-1224, -484 and -425), which have been previously demonstrated to be important in infection and metabolic pathways.

Methods: Peripheral blood monocyte derived macrophage (MDM) cultures were prepared and infected with BCG at a multiplicity of infection (MOI) =10 or left uninfected as a control. 72h post-infection, the cultured cells were subjected to RNA extraction, cDNA synthesis and real-time PCR. Expression levels miRNAs were normalized to the levels of U6 snRNA (Rnu6) using the $2^{-\Delta\Delta C_t}$ method.

Result: Infection with BCG resulted in a highly significant increase in miR-1224 expression (24.4 ± 3.8 -fold induction) in human MDMs. The induction of miR-484 (1.8 ± 0.3 -fold increase) and of miR-425 (1.2 ± 0.2 -fold increase) was less increased compared to miR-1224. Mycobacterium tolerates a hostile microenvironment by escaping from lysosomal degradation and providing a lipid-rich niche by trigger with and re-patterning host metabolism.

Conclusion: This study highlighte the potential roles of miRNAs in host responses upon mycobacterium infection.

Keywords: Macrophages, MDM, TB, miR-1224, miR-484, miR-425

ICIA2018\Immunology of Mycobacterium Tuberculosis\Oral\8625

8625

Assessment of the CC and CXC chemokine family's amounts in bronchoalveolar fluid and peripheral blood samples from patients with pulmonary tuberculosis in Mashhad Persian date Farvardin 1391 to 1392

Marzie Motaghy¹, Amin Seyfaddiny², AmirReza AghebatKhayr³, Gholam Hossien HasanShahi⁴

1. Kerman University of Medical Sciences
2. Medical Student, Rafsanjan University of Medical Sciences, Rafsanjan, Iran
3. Medical Student, Rafsanjan University of Medical Sciences, Rafsanjan, Iran
4. Molecular Medicine Research Center, Rafsanjan University of Medical Sciences, Rafsanjan, Iran

Background: Pulmonary tuberculosis is one of the most common infectious diseases, the causative agent of the disease is acid fast bacilli called Mycobacterium tuberculosis. Approximately one-third of the world's population is infected with the bacteria. This disease despite extensive control programs in the past two decades, is still one of the most dangerous infectious diseases as it is estimated that around the world people develop TB disease every 15 minutes and 6 people die of the disease. For prevention of TB, early detection of the disease may be one of the most important strategies.

Methods: This cross-sectional study was conducted on a group of 150 patients with pulmonary tuberculosis, were involved. Peripheral blood and BAL was performed for patients as well as controls carried out in coordination with the Mashhad and 150 healthy individuals were collected and blood was obtained. Demographic data were collected through clinical history and methods, and were matched for age and sex. The chemokine family of CC and CXC in peripheral blood samples and bronchoalveolar lavage samples were measured and the results, along with demographic information by SPSS software and statistical method T-Test and Mann - Whitney were analyzed.

Results: 150 patients were involved in the scheme, were matched for age and sex. The serum level mean were measured every 3 households CC chemokines in peripheral blood and BAL in patients with pulmonary tuberculosis had increased significantly and statistical differences were significant ($P < .0001$). The serum level mean of 4 CXC chemokine family in peripheral blood and BAL in patients with pulmonary tuberculosis in peripheral blood than the control group also showed significant difference except one (The compare of serum level mean of CXCL9 (Mig) in peripheral blood between patients and controls was not significant ($P = 0.092$)).

Conclusion: Our study showed that all CC and CXC chemokine families in the study of BAL (bronchoalveolar lavage) sample in pulmonary tuberculosis patients was significantly associated with the disease, and suggests CCL11 as a new chemokine associated with pulmonary tuberculosis in bronchoalveolar fluid that could serve as a diagnostic marker from alveolar fluid to be used .well as all the items in the peripheral blood of patients with pulmonary tuberculosis in chemokine significantly related to disease and are discussed as tuberculosis diagnostic markers in blood, except CXCL9 (Mig) that local nature of this chemokine probably is production in bronchial epithelial cells, thus making it suggestive tuberculosis marker only in BAL.

Keywords: Pulmonary tuberculosis (TB), CC chemokine, CXC chemokine, Broncho alveolar lavage (BAL)

ICIA2018\Immunology of Mycobacterium Tuberculosis\Oral\7603
7603

Comparing effects of route of immunization on the cellular immune responses induced by Mycobacterium tuberculosis recombinant antigen (ESAT-6/CFP-10)

Mozhdeh Namvarpour¹, Majid Tebianian², Reza Mansouri¹

1. Department of Medical Immunology, Shahid Sadoughi University of Medical Sciences and Services, Yazd, Iran

2. Department of Biotechnology, Razi Vaccine and Serum Research Institute, Agricultural Research Education and Extension Organization (AREEO), Karaj, Iran

Background: Tuberculosis, as one of the main health problems of the present century, affects about 10 million people annually. The BCG vaccine is the only available vaccine against this disease in the recent 75 years and its protective responses can make enormous changes. Therefore, study on development of a new vaccine against tuberculosis is the new topic of many investigations. Route of vaccination is one of the most important factors in creating a good immune response. Recently, intranasal administration route has increased vaccines efficiency against respiratory infections. In this study, the effects of vaccine administration routes including intra-nasal, subcutaneous, and intramuscular routes on creating protective humoral immunity against tuberculosis has been compared and examined.

Methods: Recombinant antigen ESAT-6/CFP-10 was administered by three routes of IM, IN and SC to 8 weeks Balb/C mice with or without adjuvant for three times at two-week intervals. For IM and SC administration routes, adjuvant MF59 and for IN route adjuvant CTB were used. Then, the case-control study was carried out to evaluate the cellular responses elicited against ESAT-6/CFP-10, that were evaluated by mice spleens' immune cells culture proliferation and cytokine assay

Results: The results showed that vaccinated mice with recombinant antigen have higher titers of IFN γ , IL5 and significant increase in immune cells proliferation compared to the control groups. In the comparison between the other routes of injection, although SC and IM routes, were both able to make a good immune responses, IN route could activate high cellular immune responses that had fewer side effects than other aggressive injection routes.

Conclusion: This study emphasize that immunization of mice with recombinant ESAT-6/CFP-10 protein by IN route are durable and protective.

Keywords: Mycobacterium tuberculosis, Intramuscular, Subcutaneous, Intranasal administration

ICIA2018\Immunology of Mycobacterium Tuberculosis\Oral\11084
11084

The Immunomodulatory Changes in Patients with Tuberculosis

Rafal J. Al-Saigh¹, Hussam Al-Humadi²

1. Department of Clinical Laboratory Sciences, Faculty of Pharmacy, University of Babylon, Babylon, Iraq

2. Dean of Faculty of Pharmacy, University of Babylon, Babylon, Iraq

Background: Tuberculosis (TB) is an infectious disease caused by *Mycobacterium tuberculosis*. The aim was to investigate the levels of immunomodulatory markers like interleukin-6 (IL-6), tumor necrotizing factor- α (TNF- α), cell differentiation-4 (CD4) and CD8 levels in those patients with active tuberculosis (TB) disease in comparison with control group.

Method: 41 adults diagnosed with TB were included in comparison to 32 healthy individuals at Babylon health center for pulmonary diseases and TB. Descriptive data for patients and control group were collected by well-trained researcher following a structured questionnaire. In parallel, peripheral blood collected to determine IL-6, TNF- α , CD4 and CD8. Then the assessment for the association between clinical and descriptive data and immunomodulatory markers levels was investigated statistically.

Results: The majority of TB patients were males (56%) and 71% were resident in rural areas; 47% of them were living in middle socio-economic state, moreover, 47% of TB cases had diabetes, furthermore, 51% had chronic obstructive pulmonary diseases, 12% had hypertension and 39% of them had chronic anemia with 47% smokers with no significant difference versus control. Following to that, there was highly increased in IL-6 and TNF- α levels in TB patients versus control ($P < 0.001$), with low CD4 level versus control ($P < 0.001$). While there was no significant change shown in CD8 levels versus control and this might highly be correlated with 30% of abnormal liver function tests among TB patients.

Conclusion: A high proportion of TB patients have low CD4 level mostly associated with active disease. Moreover, the increase of IL-6 and TNF- α levels suggests an inverse impact on CD4 level which closely associated with the outcome of the disease.

Keywords: Tuberculosis, Immunomodulatory markers, IL-6, TNF- α , CD4, CD8.



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Immunology of Rheumatic Diseases

Oral Presentation

ICIA2018\Immunology of Rheumatic Diseases\Oral\7790

7790

Effect of *Lactobacillus rhamnosus* and *Lactobacillus delbrueckii* on the expression levels of miR-181a and miR-155 in the PBMCs of systemic lupus erythematosus patients

ZohrehVahidi¹, Mahmoud Mahmoudi², Zahra RezaeiYazdi³, Maryam Sahebari³, AlirezaEsmaceli⁴, NafisehTabasi⁵, Maryam Rastin^{6*}, Morteza Samadi^{7*}

1. Department of Immunology, Faculty of Medicine, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

2. MD-Ph.D of immunology. Immunology Research Center, BuAli Research Institute, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

3. MD of Rheumatology. Rheumatic Diseases Research Center, Ghaem Hospital, Internal Medicine section, Mashhad University of Medical Sciences

4. Ph.D. Student of Immunology. Immunology Research Center, BuAli Research Institute, Mashhad University of Medical Sciences, Mashhad, Iran

5. MSc of Immunology. Immunology Research Center, BuAli Research Institute, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

6. Ph.D of immunology. Immunology Research Center, BuAli Research Institute, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

7. Reproductive Immunology Research Center, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

Background: Systemic lupus erythematosus (SLE) is a chronic inflammatory autoimmune disease described by the production of autoantibodies and activated immune cells against self-antigens. The expression patterns of some miRNAs in PBMCs were shown to be associated with SLE development and activity. Probiotics showed anti-inflammatory and modulatory effects on the immune responses. In this study, we evaluated the effect of probiotics *Lactobacillus rhamnosus* and *Lactobacillus delbrueckii* on the expression levels of miR-155 and miR-181a in the PBMCs of SLE patients.

Methods: A total of 20 newly diagnosed SLE patients and 20 healthy controls were enrolled in the study. Peripheral blood mononuclear cells (PBMCs) were isolated from whole blood and were treated with probiotics *Lactobacillus rhamnosus* (10^7 Bac/ ml) and *Lactobacillus delbrueckii* (10^5 Bac/ ml) and cultured in RPMI-1640 for 48 hrs. After that, total RNA containing micro-RNAs were extracted. cDNA was synthesized for miRNAs and gene expression of miR-181a and miR-155 was analyzed by real-time PCR method.

Results: Findings showed expression levels of miR-181a and miR-155 in SLE patients were higher than those of healthy controls. Expression of miR-181a and miR-155 were down-regulated in the PBMCs of SLE patients after treatment with *Lactobacillus rhamnosus* (* $P < 0.1$, ** $P < 0.02$) and *Lactobacillus delbrueckii* (* $P < 0.01$, ** $P < 0.001$). Also, Expression of miR-181a and miR-155 were significantly down-regulated in the PBMCs of SLE patients after treatment with a mixture of *Lactobacillus delbrueckii* and *rhamnosus* (* $P < 0.004$, ** $P < 0.03$).

Conclusion: Treatment of PBMCs with *Lactobacillus rhamnosus* and *Lactobacillus delbrueckii* and with a mixture of 2 probiotics, downregulate the expression rate of miR-181a and miR-155 in treated cells in compared with untreated cells. *Lactobacillus rhamnosus* and *Lactobacillus delbrueckii* modulate the expression of miR-181a and miR-155 in SLE patients, in result they should have beneficial in the management of SLE patients.

Keywords: miR-155, miR-181a, systemic lupus erythematosus, autoimmunity, probiotic

ICIA2018\Immunology of Rheumatic Diseases\Oral\8217

8217

The effect of DNA methylation on expression of adhesion molecules In peripheral blood mononuclear cells of patients with limited and diffuse scleroderma

Navid Dashti^{1, 2}, Mohammad Vodjgani¹, Mahdi Mahmoudi², Farhad Gharibdoost², Hoda Kavosi², Ahmad Reza Jamshidi²

1. Department of Immunology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

2. Rheumatology Research Center, Tehran University of Medical Sciences, Tehran, Iran

Background: Scleroderma is an autoimmune rheumatic disease that may result in damages to the skin, endothelium of blood vessels, muscles, and internal organs through inflammation and fibrosis. Products of L-selectin (SELL) and β 2-Integrin (ITGB2) genes participate in several functional pathways of immune system. The aim of this study was to survey the mRNA expression level of SELL and ITGB2 genes as well as methylation status of CpG sites in promoter region of differently expressed gene in PBMCs of patients with limited and diffuse scleroderma.

Methods: PBMCs were isolated from whole blood of 50 scleroderma patients and 30 healthy controls. After the extraction of total RNA and DNA contents from PBMCs, complementary DNA (cDNA) was synthesized. Afterwards, quantitative analysis of SELL and ITGB2 messenger RNA (mRNA) was conducted by real-time polymerase chain reaction (PCR) using the SYBR Green PCR Master Mix. To evaluate the methylation status of CpG sites in the promoter region of gene, nested-PCR products of bisulfite-treated DNA from scleroderma patients and controls were sequenced via Sanger difficult sequencing method.

Results: ITGB2 gene in PBMCs of scleroderma patients was overexpressed significantly in comparison to healthy controls. However, no altered SELL expression was observed. Three CpG sites of 12, 13 and 14 were significantly hypomethylated in patients group, despite overall methylation status of ITGB2 gene promoter revealed no significant difference between study groups. There was no statistically significant correlation between methylation status of ITGB2 promoter and the gene expression in patients.

Conclusion: Regarding to lack of correlation of increased expression of ITGB2 with its promoter hypomethylation in scleroderma patients, our study suggests that upregulation of ITGB2 in PBMCs from scleroderma patients is probably due to another mechanism other than methylation alteration.

Keywords: Scleroderma, SELL, ITGB2, DNA methylation, CpG site

ICIA2018\Immunology of Rheumatic Diseases\Oral\8428

8428

The effect of endothelial reticulum stress and autophagy pathways on IL-23 production of peripheral blood mononuclear cell-derived macrophages in patients with ankylosing spondylitis

Alireza Rezaeiemanesh^{1,2}, Mahdi Mahmoudi³, Ali Akbar Amirzargar^{2,4}, Mahdi Vojdani³, Ahmad Reza Jamshidi³, Mohammad Hossein Nicknam^{2,4}

1. Department of Immunology, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

2. Department of Immunology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

3. Rheumatology Research Center, Tehran University of Medical Sciences, Tehran, Iran

4. Molecular Immunology Research Center, Tehran University of Medical Sciences, Tehran, Iran

Background: Interleukin (IL)-23/IL-17 pathway involves in the pathogenesis of ankylosing spondylitis (AS). The exact mechanism implicated in overexpression of IL-23 and activation of the IL-23/IL-17 axis has not been understood. The current study aimed to evaluate the role of endothelial reticulum stress and autophagy pathways in the IL-23 production of peripheral blood mononuclear cell-derived macrophages in patients with AS.

Methods: Peripheral blood monocyte isolated from 10 HLA-B27⁺, 5 HLA-B27⁻ patients, 10 HLA-B27⁺, and 5 HLA-B27⁻ normal subjects were differentiated to macrophages by macrophage-colony stimulating factor (M-CSF) for 7 days. Selected samples were treated with IFN- γ to up-regulate HLA-B expression prior to stimulation with LPS for 24 hours. Also, thapsigargin and 3-Methyladenine were used for ER stress and unfolded protein response (UPR) induction and autophagy inhibition, respectively. Flowcytometry was used to detect monocyte purity and expression of macrophage markers. Analysis of mRNA expression for HLA-B and B27, UPR-associated genes (BiP, XBP1, PERK, ATF6, and CHOP) and autophagy genes (Atg5, Atg12, and Atg16L) was performed using RT-qPCR. IL-23 was assessed by RT-qPCR and ELISA methods.

Results: Data showed significant overexpression of HLA-B, HLA-B27, UPR genes (BiP, XBP1, CHOP, and PERK) and IL-23 levels in M-CSF-derived macrophages, also after treatment with IFN- γ \pm LPS from AS patients compared to controls. Thapsigargin significantly increased the level of IL-23 in macrophages from AS patients in comparison to healthy macrophages. No variations found in autophagy genes before and after treatment. It is notable that autophagy inhibition with 3-Methyladenine does not affect IL-23 production.

Conclusion: Our data suggest that UPR activation occurs in macrophages either before treatment with IFN- γ \pm LPS or after that in AS patients and is accompanied by overexpression of IL-23. UPR appears has a regulatory role in the IL-23 production.

Keywords: Ankylosing spondylitis, Autophagy, Endothelial reticulum stress, Interleukin-23, Macrophage

ICIA2018\Immunology of Rheumatic Diseases\Oral\8443

8443

Inhibition of ERAP1 represses of HLA-B27 free heavy chains expression and NK cells responses in Ankylosing Spondylitis

Farhad Babaie^{1,2}, Hamed Mohammadi^{1, 2}, Sorayya Salimi^{1, 2}, Alireza Rezaeiemanesh³, Behzad Baradaran^{1, 2}, Zohreh Babaloo^{1, 2}

1. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

2. Department of Immunology, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

3. Department of Immunology, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

Background: Ankylosing spondylitis (AS) is a type of arthritis that is referred to a group of chronic immune mediated inflammatory diseases termed as seronegative spondyloarthro pathies. It typically affects the joints of the spinal and axial skeleton and exhibits typical clinical features and genetic factors such as HLA-B27 and ERAP1. The ERAP1 trim amino acid residues at the N-terminal to optimize their length for binding to MHC class I molecules and can potentially alter surface expression of HLA-B27 free heavy chains (FHCs). We assessed the effects of ERAP1 polymorphism/chemical inhibition on HLA-B27 FHC expression and NK Cells responses in AS.

Methods: To measure the expression levels of cell surface HLA-B27 FHCs in monocyte-derived macrophages from patients with AS carrying rs30187 (K528R) and rs27044 (Q730E) ERAP1 genotypes, we used flowcytometry. ERAP1 inhibited monocyte-derived macrophages were cocultured with AS NK cells. NK cells responses were analyzed by IFN- γ , CD69, and CD107a by flow cytometry.

Results: The expression of surface HLA-B27-(FHCs) in monocyte-derived macrophages from patients with AS was reduced in the AS-protective ERAP1 polymorphisms. ERAP1 inhibition in monocyte-derived macrophages downregulated HLA-B27 FHC surface expression, reduced IFN- γ production, and also CD107a and CD69 expression by AS NK cells.

Conclusion: ERAP1 plays an important role in determining the expression levels of cell surface HLA-B27 FHCs and potentially upregulated NK cells responses in AS by binding to HLA-B27 FHCs.

Keywords: Ankylosing spondylitis, ERAP1, HLA-B27, NK cell.

ICIA2018\Immunology of Rheumatic Diseases\Oral\8484

8484

Inhibition of ERAP1 suppresses free heavy chain expression and CD8+ T cell responses in Ankylosing Spondylitis

Hamed mohammadi^{1,2}, Maryam Hemmatzadeh^{1,2}, Farhad Babaie^{1,2}, Alireza Rezaeiemanesh³, Jafar Majidi^{1,2}, Behzad Baradaran^{1,2}

1. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

2. Department of Immunology, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

3. Department of Immunology, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

Background: Ankylosing spondylitis (AS) is a chronic immune-mediated inflammatory disease that affects both axial and peripheral skeletons and soft tissues. Evidence offers that disease pathogenesis is strongly associated with human leucocyte antigen (HLA)-B27 and endoplasmic reticulum aminopeptidase 1 (ERAP1). ERAP1 is a key aminopeptidase in HLA class I presentation and can potentially alter surface expression of HLA-B27 free heavy chains (FHCs). We evaluated the effects of ERAP1 variations/inhibition on HLA-B27 FHC expression and TCD8 responses in AS.

Methods: To evaluate the expression of FHC cell surface in monocyte-derived macrophages from patients with AS carrying rs30187 and rs27044ERAP1 genotypes, we used flow cytometry. ERAP1 inhibited monocyte-derived macrophages were co-cultured with AS CD8+ T cells. CD8+ T cells responses were evaluated by IFN- γ , CD69, and CD107a by flow cytometry.

Results: The expression of surface FHCs in monocyte-derived macrophages from patients with AS was decreased in the AS-protective ERAP1 variants. ERAP1 inhibition in monocyte-derived macrophages suppressed HLA-B27 FHC surface expression, reduced IFN- γ production, and also CD69 and CD107a expression by AS CD8+ T cells.

Conclusion: ERAP1 activity determines the surface expression of HLA-B27 FHCs and potentially enhance TCD8 responses in AS by binding to HLA-B27 FHCs.

Keywords: Ankylosing spondylitis, ERAP1, TCD8, HLA-B27 free heavy chains

ICIA2018\Immunology of Rheumatic Diseases\Oral\8667

8667

Increased inflammatory Responsiveness of Peripheral Blood Mononuclear Cells (PBMCs) to in vitro NOD2 Ligand Stimulation in Patients with Ankylosing Spondylitis

Negar Vanaki^{1, 2*}, Taghi Golmohammadi^{1*}, Ahmadsreza Jamshidi², Maryam Akhtari², Mahdi Vojdani², Shayan Mostafaei², Shiva Poursani², Nooshin Ahmadzadeh², Mahdi Mahmoudi^{2*}

1. Department of Biochemistry, Faculty of Medicine, Tehran University of Medical Sciences, Tehran, Iran

2. Rheumatology Research Center, Tehran University of Medical Sciences, Tehran, Iran

Background: Ankylosing spondylitis (AS) is a common debilitating rheumatic disease in which the innate immune responses play important role in its pathogenesis. The Interleukin (IL)-23/IL-17 axis related genes have critical role among the factors that predispose the individuals to AS. Nucleotide binding oligomerization domain (NOD)-2 as an innate receptor is critical for IL-23 production in cells. Therefore, we aimed to stimulate NOD2 signaling and study its effects on cytokine production of peripheral blood mononuclear cells (PBMC) in these patients.

Methods: PBMCs from 18 patients with active AS and 18 healthy individuals were separated by ficoll-hypaque density gradient centrifugation and cultured in the presence of muramyl dipeptide (MDP), as NOD2 ligand. Quantitative expression analysis of NOD1, NOD2, RIPK2, SLC15A4, NLRP1, NLRP3, IL23A, IL17A, IL1B and TNFA genes was performed using Real Time PCR. Finally, protein changes of IL23A and IL17A expression were validated using enzyme linked immunosorbent assay (ELISA).

Results: Apart from NOD1 that tend to be downregulated in the controls, all the selected genes showed overexpression in response to MDP in cells from the studied groups. Except RIPK2, all the genes had higher expression changes upon MDP stimulation in the AS population. Overexpression of IL23A and IL17A were confirmed at protein levels using ELISA.

Conclusion: This study indicated that AS PBMCs were hyper-responsive to MDP stimulation. This observation implies an important role of NOD2 in the pathogenesis of inflammatory diseases including AS.

Keywords: Ankylosing spondylitis, NOD2, PBMC

ICIA2018\Immunology of Rheumatic Diseases\Oral\10039
10039

The expression of CCN1 and its specific integrins in lungs of mouse model of systemic sclerosis

P. Safari¹, F. Vafashoar¹, H. Poormoghim², P. Jorbozedar¹, M. Nadjji³, T. Mousavi⁴, N. Mojtavavi¹

1. Immunology department, Iran University of medical sciences, Tehran/Iran

2. Scleroderma study group Firuzgar Hospital, Iran University of medical sciences

3. Institute of Immunology and Infectious disease, Antimicrobial resistance Center

Background: Systemic sclerosis is an autoimmune disease with vascular abnormalities, skin and internal organs fibrosis. This disease results from deposition of extracellular matrix in tissues. Member of CCN family perform their different functions through interactions with their specific integrins and proteoglycans in health and disease. Concerning the role and interaction of CCN1/ CYR61 with its specific integrin receptors in fibrosis exist paucity of information. Therefore we aimed to evaluate the expression of CCN1 and its related receptors in lungs of mouse model of systemic sclerosis.

Method: Mice received bleomycin for 28 days for induction of fibrosis. On days 10, lungs were removed from lethally anesthetized mice for measurement of hydroxyl proline content, histochemistry staining with H& E, Masson trichrome and RNA extraction. Specific primer for CCN1, α_v , β_3 , α_6 , β_1 were designed and qPCR was performed.

Results: Hydroxyproline assay and Masson trichrome staining of mice, indicated fibrosis on day 28 and even more fibrosis on day 35. The overall infiltrated cells to lungs were significantly increased on day 28 qPCR results showed increment of CCN1 and $\alpha_6 \beta_1$ integrin genes on day 10 and their decline to normal level on day 28. The expression of integrin α_v didn't change at any of the assessment days.

Conclusion: The increase of CCN1 and its integrins transcript on day 10 prior to establishment of fibrosis on day 28, may indicate the role of CCN1 as a profibrotic factor in the lung, however further investigation at the level of protein seems to be necessary.

Keywords: CCN1, integrins, systemic sclerosis.



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Immunomodulation and Immunoregulation

Oral Presentation

ICIA2018\Immunomodulation and Immunoregulation\Oral\8189

8189

Generation of Tolerogenic DC by Lactobacillus Rhamnosus and Delbruki in Healthy and SLE Donors

Seyed-Alireza Esmacili^{1,2,3}, Mahmoud Mahmoudi^{1,2}, Fatemeh Mardani^{1,2,3}, Sahar Khorasani^{1,2,3}, Zohreh Vahidi^{1,2,3}, Nafiseh Tabasi¹, Mahdiah khazaei¹, Maryam Rastin^{1,2*}

1. Immunology Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

2. Immunology Department, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

3. Student Research Committee, Mashhad University of Medical Sciences, Mashhad, Iran

Background: Systemic lupus erythematosus (SLE) is a complicated autoimmune disease. Loss of tolerance against self-antigen and excessive inflammatory immune responses happen in SLE patients. Dendritic cells (DC) induce and promote immune responses in innate and adaptive immune system. Tolerogenic DCs regulate priming of immune response and prevent from inflammatory responses. Tolerogenic probiotics could produce regulatory monocyte-derived DC in culture media. The aim of this project was evaluating the effect of two tolerogenic probiotics (Lactobacillus delbrueckii and lactobacillus rhamnosus) for production of the regulatory DC in healthy and SLE donors.

Methods: in order to immature DC (IDC) production, Monocytes from healthy and SLE donors cultured with optimize concentration of GM-CSF and IL-4 for 5 days; followingly in order to mature DC (MDC) production, IDC of SLE and healthy donors treated for 48 h with cytokines plus L. delbrueckii, L. rhamnosus and mix probiotics. FITC-uptake assay performed for uptake of IDC. Surface marker expression and gene expression of IDC and MDC were evaluated by Flowcytometry and Real-Time PCR methods.

Results: we observed the significant reduction in the expression of co-stimulatory molecules (HLA-DR, CD86, and CD80) also CD83, CD1a, and CD14 in probiotics-derived DC of both healthy and SLE donor in comparison with LPS -derived DC. Also in probiotics-derived DC in compared to LPS-derived DC Indoleamine 2, 3-dioxygenase (IDO) and IL10 were increased but was declined the expression of IL12.

Conclusion: our finding revealed that live probiotics could change the phenotype of DC to the regulatory and increase expression of IDO and IL10 during the differentiation process. Therefore, these modulatory cells probably could be used in treatment of SLE patients.

Keywords: Systemic lupus erythematosus (SLE), Probiotics, Dendritic cell(DC), Lactobacillus delbrueckii and lactobacillus rhamnosus

ICIA2018\Immunomodulation and Immunoregulation\Oral\8219

8219

Regulation of Th17/ Treg Balance by Mesenchymal Stem Cell-Conditioned Medium in Mouse Model of Chronic Colitis

Sedigheh Pouya¹, Maryam Heidari¹, Kaveh Baghaei², Hamid Asadzadeh Afgdaei², Afshin Moradi³, Saeed Namaki¹, Mohammad reza Zali³, Seyed Mahmoud Hashemi¹

1. Department of Immunology, School of medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Basic and Molecular Epidemiology of Gastrointestinal Disorder Research Center, Research Institute for Gastroenterology and Liver Disease, Shahid Beheshti University of Medical Science, Tehran, Iran

3. Gastroenterology and Liver Disease Research Center, Research Institute for Gastroenterology and Liver Disease, Shahid Beheshti University of Medical Science, Tehran, Iran

Background: Inflammatory bowel disease (IBD) is a chronic inflammatory disease of the gastrointestinal tract. Among the factors that cause IBD the role of immune response in manifestation and progression of IBD has been strongly emphasized. On the other hand, the role of mesenchymal stem cells (MSC) and mesenchymal stem cell-conditioned medium (MSC-CM) in the regulation of immune responses (inhibition of T lymphocytes proliferation and induction of regulatory T cells) also has been proved. In this study, the effects of MSC-CM on the Th17/ Treg cell balance in the chronic model of colitis have been investigated.

Method: In this study after induction of chronic colitis in female C57BL/6 mice, intraperitoneal injection of MSC-CM (500μl) was done for 6 times. Briefly, for colitis induction, the mice were received dissolved 2% dextran sulfate sodium (DSS) in water for 4 days that followed by only water consumption for 7 days. This step repeated in 3 cycles. After euthanizing the mice the serum and mononuclear cells of mesenteric lymph nodes and spleen were isolated. Then serum and the cells were cultured with and without PHA for 72 hours. At the end of the experiment the levels of IL-10, IL-17 and TGF-β in serum and cell culture supernatant were evaluated (ELISA method).

Results: The results showed that after MSC-CM injection, IL-17 production was reduced. Also, the levels of anti-inflammatory cytokines, IL-10 and TGF-β, were increased both in the serum and in the cell culture supernatant.

Conclusion: According to the results mesenchymal stem cell- conditioned medium injection, can make a balance between Th-17 and regulatory T cells function and can ameliorate inflammatory responses in colitis.

Keywords: Inflammatory bowel disease, Mesenchymal stem cells, Conditioned medium, Dextran sulfate sodium.

ICIA2018\Immunomodulation and Immunoregulation\Oral\8227
8227

Oleoylethanolamide improves inflammation and oxidative stress in obese people. A randomized double-blind, placebo-controlled trial

Payahoo L¹, Khaje-Bishak Y², Asghari M³, Ostadrahimi A⁴

1. Talented student center, Student Research Committee, Nutrition Research Center, Faculty of Nutrition, Tabriz University of Medical Sciences, Tabriz, Iran
2. Student Research Committee, Faculty of Nutrition, Tabriz University of Medical Sciences, Tabriz, Iran
3. Road Traffic Injury Research Center, Tabriz University of Medical Sciences, Tabriz, Iran
4. Nutrition Research Center, Faculty of Nutrition, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Oleoylethanolamide (OEA) is structurally similar to the endocannabinoid system components, but, it does not bind to the cannabinoid receptors. Obesity as a chronic low-grade inflammation disorder has been prevailed dramatically, in recent times. The aim of the present study was to determine the effects of Oleoylethanolamide supplementation on the markers of inflammation and oxidative stress in obese people.

Methods: This randomized, double-blind, placebo-controlled clinical trial was carried out on 60 healthy obese people in 2016 in Tabriz, Iran. Eligible subjects were randomly divided into two interventions (received daily two 125 mg OEA capsules) and control groups (the same amounts starch) and treated for 8 weeks. Blood samples (5 ccs) were taken in fasting state at baseline and at the end of the study. The concentrations of malondialdehyde (MDA) and total antioxidant status (TAS) were measured by spectrophotometer method. High sensitive-C reactive protein (hs-CRP) level measured by Immunoturbidimetry assay using the commercial kits and IL-6 and TNF- α levels were assayed by ELISA method. The differences between groups were assessed by analysis of covariance (ANCOVA) and $p < 0.05$ defined as statistically significant.

Results: The analysis was done on 56 samples that completed the study. OEA decreased significantly serum levels of IL-6 and TNF- α ($p < 0.001$). The levels of hs-CRP decreased significantly in the intervention group ($p = 0.011$), however, ANCOVA test didn't reveal significant amounts. Changes in other variables were not detectable ($p > 0.05$).

Conclusion: Use of Oleoylethanolamide as a complementary and alternative approach could be effective in attenuating inflammation in obesity and related disorders.

Keywords: Inflammation, Endocannabinoids, Obesity, Oleoylethanolamide, Oxidative stress.

ICIA2018\Immunomodulation and Immunoregulation\Oral\8303

8303

Dendritic cell migration to skindraining lymph nodes is controlled by dermatan sulfate and determines adaptive immunity magnitude

Reza Nadafi¹, Jasper J. Koning¹, Henrike Veninga¹, Xanthi N. Stachteas², Tanja Konijn¹, Antonie Zwiers¹, Anders Malmström², Joke M. M. den Haan¹, Reina E. Mebius¹, Marco Maccarana², Rogier M. Reijmers¹

1. Department of Molecular Cell Biology and Immunology, VU University Medical Center, Cancer Center Amsterdam, Amsterdam, the Netherlands

2. Department of Experimental Medical Science, Lund University, Lund, Sweden

Background: For full activation of naïve adaptive lymphocytes in skindraining lymph nodes (LNs), presentation of peptide: MHC complexes by LNresident and skinderived dendritic cells (DCs) that encountered antigens, is an absolute prerequisite. To get to the nearest draining LN upon intradermal immunization, DCs need to migrate from the infection site to the afferent lymphatics, which can only be reached by traversing a collagendense network located in the dermis of the skin through the activity of proteolytic enzymes.

Methods: Using in-vitro experiments, in-vivo mouse model and performing multicolor FACS staining, we could showed impaired DCs migration due to altered collagen fibrillogenesis in skin.

Result: Here, we show that mice with altered collagen fibrillogenesis resulting in thicker collagen fibers in the skin, display a reduced DC migration to the draining LN upon immune challenge. Consequently, the initiation of the cellular and humoral immune response was diminished. Antigen-specific CD8⁺ and CD4⁺ T cells as well as antigen-specific germinal center B cells and serum immunoglobulin levels were significantly decreased.

Conclusion: Hence, we postulate that alterations to the production of extracellular matrix, as seen in various connective tissue disorders, may in the end affect the qualitative outcome of adaptive immunity.

Keywords: skindraining lymph node, adaptive immunity, DCs, dendritic cells, dermatan sulfate epimerase, collagen, antigen.

ICIA2018\Immunomodulation and Immunoregulation\Oral\8347
8347

Investigating the involvement of microRNA-146 family in the pathogenesis of multiple sclerosis using EAE animal model

Farideh Talebi, Samira Ghorbani, Farshid Noorbakhsh

Department of Immunology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

Background: MicroRNAs have recently emerged as a new class of modulators of gene expression. miRNAs are small non-coding RNA molecules which control protein synthesis through translational repression or degradation of mRNA transcripts. Recent studies have shown that dysregulation of immunity-related miRNAs might contribute to autoimmune disorders such as multiple sclerosis. Herein, we investigated the contribution of miR-146 isoforms and their target gene in the pathogenesis of multiple sclerosis using EAE animal model.

Methods: The expression levels of isoforms of miR-146 as well as their potential target genes were measured in the CNS tissue from EAE mice and stimulated lymphocytes by real-time RT-PCR. The miR-146a-5p/b-5p mimic sequences were transfected into T CD4⁺ cells and the role of them were then investigated in T cell differentiation by flowcytometric analysis of intracellular cytokines. Luciferase assays using vector containing the 3'UTR of predicted target was performed to confirm the interaction of miRNA sequences with target genes transcripts.

Results: Expression of both isoforms miR-146 were significantly increased in acute and chronic phases of EAE as well as activated lymphocytes in comparison with control groups whereas the levels their predicted target gene decreased. CD4 T cell differentiation was skewed towards Th17 subset in miR-146a-5p and miR-146b-5p transfected cells. Luciferase assays revealed SMAD4 as direct target of both of miR-146 isoforms and overexpression of miRNA mimic sequences suppressed the expression of this target transcripts in lymphocytes.

Conclusions: Our findings suggest that increased expression of miR-146 isoforms in the CNS tissue, as well as in lymphocyte might be involved in the pathogenesis of multiple sclerosis through altering the process of differentiation of these cells into the inflammatory phenotype. Our findings suggested that targeting and suppression of protective gene in multiple sclerosis such as SMAD-4 by miR-146 is a possible mechanism for miRNA effects.

Keywords: Multiple Sclerosis, miR-146, SMAD4

ICIA2018\Immunomodulation and Immunoregulation\Oral\9749
9749

Association between serum cytokine concentrations and the presence of hypertriglyceridemia

Seyed Reza Mirhafez^{1,2}, Mohammad Tajfard², Amir Avan², Alireza Pashar^{2,3}, Reza Nedaeinia^{4,5}, Malihe Aghasizade², Hafezeh Davari², Mostafa Manian^{6,7}, Adeleh Mahdizadeh⁸, Zahra Meshkat⁹, Ali Movahedi¹, NahidGhaedAmini¹⁰, NahidEskandari¹¹, RasoulSalehi¹², Gordon A. Ferns¹³, Majid Ghayour-Mobarhan¹⁴

1. Department of Basic Medical Sciences, Neyshabur University of Medical Sciences, Neyshabur, Iran.
2. Department of Modern Sciences and Technologies, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran
3. Division of Applied Medicine, Medical School, University of Aberdeen, Foresterhill, Aberdeen, AB25 2ZD, UK
4. Student Research Committee, Department of Medical Biotechnology, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran
5. Department of Microbiology, Deputy of Food and Drug, Isfahan University of Medical Sciences, Isfahan, Iran
6. Immunology Research Center, Iran University of Medical Sciences, Tehran, Iran
7. Neurosciences Research Center, Alzahra Hospital, Isfahan University of Medical Sciences, Isfahan, Iran
8. Cardiovascular Research Center, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran.
9. Antimicrobial Resistance Research Center, Mashhad University of Medical Sciences, Mashhad, Iran
10. Department of Biology, Payame Noor University of Isfahan, Isfahan, Iran;
11. Department of Immunology, Faculty of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran
12. Department of Genetics and Molecular Biology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran.
13. Brighton & Sussex Medical School, Division of Medical Education, Falmer, Brighton, Sussex BN1 9PH, UK.
14. Biochemistry of Nutrition Research Center, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

Background: Hypertriglyceridemia is an established risk factor for coronary-heart-disease. Inflammatory cytokines are known to be important mediators of atherogenesis; however the relationship between the concentrations of specific inflammatory cytokines and the presence of hypertriglyceridemia has not been well established. The purpose of this study was to investigate the relationship between the serum levels of several pro-and anti-inflammatory cytokines and the presence of hypertriglyceridemia. **Methods:** Four hundred and eighty-four subjects with/without established hypertriglyceridemia were recruited. Anthropometric-parameters and biochemical-analysis (including a full fasted lipid profile) were determined. The serum levels of several cytokines and growth factors including IL-1 α , IL-1 β , IL-2, IL-4, IL-6, IL-8, IL-10, TNF- α , MCP-1, IFN- γ , EGF, and VEGF were measured followed by univariate and multivariate analyses. **Results:** Individuals with hypertriglyceridemia had a significantly higher body mass index, total cholesterol and triglyceride, compared to the group without hypertriglyceridemia. Serum levels of MCP-1, TNF- α and IL-8 were significantly higher in subjects with hypertriglyceridemia [e.g., IL-8 from 7.8 ng/L (95%CI: 4.6-18.9) versus 5.7 ng/L (95%CI: 3.6-11.9), $P < 0.05$]. The multivariate analysis showed that the increased serum concentration of TNF- α was independently associated with high-density lipoprotein cholesterol (HDL-C), while the serum levels of IL-8 and MCP-1 were associated with hypertriglyceridemia. **Conclusion:** Subjects with serum triglycerides of ≥ 2.25 mmol/L had an altered cytokine-profile, particularly with respect to serum IL-8, MCP-1 and TNF- α , which might partially account for its adverse clinical-consequences. Further-investigations in a large multi-center setting are warranted to unravel the potential functional-importance of these cytokines in individuals with hypertriglyceridemia.

Keyword: cytokines, hypertriglyceridemia, dyslipidemia, triglyceride

ICIA2018\Immunomodulation and Immunoregulation\Oral\9853
9853

A new module combined therapy using of azoximer bromide and laciumin treatment irritable bowel syndrome

Sheikh Sajjadih Mohammadreza

Department of clinical laboratory immunology and allergy, National medical academy for post graduate education, Kiev, Ukraine

Background: Some evidence suggests that irritable bowel syndrome (IBS) is affected by the immune system. This study focused on the effect of Azoximer bromide in complex with Lacium as a probiotic in compare with classical therapy of IBS.

Methods: Our study included 22 children with IBS with diarrhea treated by probiotic and azoximer bromide (study group), and 19 children treated only by classical therapy (control group). Lymphocytes subset was analyzed by flow cytometry, phagocytic activity was assessed by latex article, IgE and serum cytokine were evaluated by ELISA. The intestinal bacterial microbiota was assessed by medical microbiologic method.

Results: The change of immune status in IBS patient is the holistic description of our articles. After therapy in group study the percentage of all lymphocyte phenotype subsets, phagocytic activity, IgE, cytokine level and intestinal microbiota were near of healthy subjects compared with control group. In control group the mean percentage of CD3 and CD4 were less than in healthy subjects ($P<.05$), the mean percentage of CD8, CD16 and CD22 were more than in healthy subjects ($P<.05$). The mean level of IL-4 and TNF- α were significantly lower when compared with before therapy ($P<.001$). No significant difference between before and after therapy was observed in serum concentration of the mean level of IFN-c, IL-1 and phagocytic activity. Intestinal dysbiosis is corrected after therapy in both groups.

Conclusions: Immunomodulator azoximer bromide when contributed with probiotic can be used to elicit or amplify an immune response in patient having suppressed immune system. Probiotic is a prescription medication for treatment of irritable bowel syndrome and in addition for correct immune status may be prescribed azoximer bromide.

Keywords: Probiotic , Immunomodulator , Irritable bowel disease



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Immunoparasitology

Oral Presentation

ICIA2018\Immunoparasitology\Oral\5318
5318

Immunohistochemical observation of the inflammatory cell infiltration in adventitia of human hydatid cysts

Rasoul Jafari¹, Behnam Sanei², Azar Baradarn³, Hossein Yousofi Darani¹

1. Department of parasitology and mycology, Isfahan university of medical sciences, Isfahan, Iran

2. Department of Surgery, Isfahan university of medical sciences, Isfahan, Iran

3. Department of pathology, Isfahan university of medical sciences, Isfahan, Iran

Background: Human cystic echinococcosis (CE) is a parasitic disease with many immunological aspects which needs to be studied. The aim of the present study was to evaluate the local immune cell infiltration pattern in human CE caused by the *Echinococcus granulosus* using immunohistochemistry (IHC) method.

Methods: In this study 50 surgically removed hydatid cysts samples and their surrounding tissue were collected from patients referred to Al Zahra Hospital, Isfahan, Iran. IHC was performed on the pre-cyst of 44 liver, five lung and one kidney hydatid cysts using anti human CD3, CD19, CD8, CD4, CD68, CD56, Ki-67 and Foxp3 antibodies.

Results: There were positive cells for CD3+ T cells, CD19+ B cells, CD8+ cytotoxic T cells, CD4+ helper T cells, CD68+ macrophages, and Ki-67+ proliferating cells, yet in some cases, there was no positive cells for CD56+ natural killer cells (7 CE patients) and Foxp3+ regulatory T cells (one CE and one chronic hepatitis), and also there was no eosinophils in non-CE samples.

Conclusion: Based on the results of the present study, the presence of regulatory T cells and a considerable number of proliferating B cells maybe the cause of persistence of hydatid cyst in human cases for long period of time.

Keywords: Hydatid cyst, immune cells, Immunohistochemistry

ICIA2018\Immunoparasitology\Oral\6317

6317

Evaluation of serum concentration of AFP marker in toxoplasmosis pregnant women with high level of IgG & IgM toxoplasma antibody by ELISA assay

Hamideh Rahimi¹, Ehssan Shariat bahadori²

1. Department of Immunology of medical University of mazandaran.

2. Department of parasitology University of Tarbiatmodares of Tehran.

Background: Toxoplasmosis is a parasitic disease which may cause some laboratory symptoms in infected individuals. One of the main ways to transmit this organism is placenta to fetus pathway. If this transmission occurs in the 3th month of pregnancy the abortion, central nerve system and ocular disorder will happen. Because of this issue, the precise technique for detection of Toxoplasma Antibody such as IgG and IgM is important, that contain ELISA to detect Toxoplasma Antibody such as IgG and IgM and AFP.

Methods: In this survey the main sample is serum that can be collected from 255 pregnant women infected with toxoplasma gondii in avesina center. Then we detected serum concentration of AFP in toxoplasmosis pregnant women with high level of IgG & IgM toxoplasma antibody by ELISA assay.

Results: The results of this survey showed that, in these total pregnant women the infection by toxoplasma gondii is occurred and 13% of them had high levels of AFP in their serum. The statistical survey was done by SPSS18.

Conclusion: in some pregnant women with high level of IgG & IgM toxoplasma antibody had high levels of AFP in their serum and this index correlates with NTD in their fetus.

Keywords: Toxoplasma gondii, ELISA ,IgG antibody, IgM antibody, AFP

ICIA2018\Immunoparasitology\Oral\7543

7543

Efficacy and mechanism of action of selected antimicrobial peptides on Leishmania major parasite: in vitro and in vivo assaysFarnaz Zahedifard¹, Hyeryon Lee³, Joo Hwan No³, Mona Salimi², Ahmad Asoodeh⁴, Sima Rafati¹*1. Immunotherapy and Leishmania Vaccine Research Department, Pasteur Institute of Iran**2. Physiology & Pharmacology department, Pasteur Institute of Iran**3. Leishmania Research Lab, Pasteur Institute of Korea**4. Departement of Chemistry, Ferdowsi University of Mashhad*

Background: Leishmaniasis as a vector born and neglected disease affects tropical and subtropical areas in the world. The only ideal available drug, Amphotericin B is expensive for majority of infected communities and need patients to be hospitalized. Immurgence of having a new drug in leishmaniasis made researchers to find new candidates. Antimicrobial peptides nowadays are in major concern to battle infectious agents. In this study two different antimicrobial peptides (Brevinin 2R and Jellein) and their Lauric acid conjugated version were studied against L. major in vitro and in vivo. Also peptides' mechanism of action on promastigotes was defined.

Methods: We measured effective concentration 50 (EC50) of antimicrobial peptides on leishmania promastigotes and amastigotes. Hemolysis assay and toxicity in THP1 cells were performed for the peptides. Membrane permeability with SYTOX green and membrane potential changes with bisoxonol fluorescence probes were investigated. SEM views of parasite-treated peptides were captured. Caspase 3 and 7 activation in promastigote were measured. Changes in mitochondrial potential were also studied in promastigotes. Apoptosis induction was also investigated through flow cytometry assay of Annexin/PI dyes. In vivo studies are now going on in Balb/c mouse model of cutaneous leishmaniasis.

Results: Lauric acid-Brevinin(L- Brevinin) had the best effect on L. major promastigotes but simultaneously had toxic effect on THP1 cells and erythrocytes. Jellein destroyed promastigote in higher concentrations. Both peptides can change membrane permeability and potential fast after exposure to promastigotes. Peptides couldn't activate caspases 3 and 7 in promastigotes but they could change mitochondrial membrane potential effectively. Flowcytometry assay showed necrosis versus apoptosis in peptide-treated promastigotes. SEM results vividly showed changes in the size and membrane of parasites in EC50 concentration of peptides (L-Brevinin and Jellein). Infected Balb/c mice treatment is now being done.

Conclusion: L-Brevinin and Jellein both are effective against L.major promastigotes. They damage parasites through membrane disruption.

ICIA2018\Immunoparasitology\Oral\7888

7888

Transcriptome profiling of the *Leishmaniatropica* infected cutaneous lesion

Nasrin Masoudzadeh¹, Amir Mizbani², Yasaman Taslimi¹, Sima Habibzadeh¹, Vahid Mashayekhi³, Hossein Mortazavi⁴, Hosein Malekafzali Ardekani¹, Sima Rafati¹

1. Immunotherapy and Leishmania Vaccine Research Dept., Pasteur Institute of Iran, Tehran, Iran

2. Zurich, Switzerland

3. Cutaneous Leishmaniasis Research Center, Emam Reza Hospital, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

4. Department of Dermatology, Razi Hospital, Tehran University of Medical Sciences, Tehran, Iran

Background: *Leishmania* (L.) *tropica* is the main causative agent of anthroponotic cutaneous leishmaniasis (CL) in Iran. CL caused by L. *tropica* infection show a longer healing time, often resulting in disfiguring scars and are more difficult to treat compared to CL caused by L. *major*. Despite these problems, there is still a lack of data describing the host immune pathways that active in localized L. *tropica* infection in human patients. To address this issue, we applied RNA sequencing technology in skin biopsies to characterize indetail the transcriptome of the human host at the lesion during L. *tropica* infection.

Methods: Skin biopsies were obtained from confirmed L. *tropica* infected patients with acute lesion using a 4-mm punch. The CL diagnosis was based on both clinical examination and PCR. Normal skin samples were taken from volunteers without a history of leishmaniasis. All participantsprovided written informed consent. Total RNA isolated from frozen tissues and sequencing libraries were prepared following Ribosomal RNA depletion. Sequencing of single-end 125-nucleotide sequences was performed on an Illumina HiSeq-2500 platform. Transcript quantification and differential expression analysis were done using Bioconductor package edgeR.

Results: Comparing the transcription profile of the L. *tropica* skin lesions with normal skin identified 5396 differentially regulated genes. Forty-seven of the up-regulated genes encode immunoglobulin fragments, highlighting the remarkable involvement of B cells in the infection site. Gene set enrichment analysis indicated that cytotoxic CD8⁺, inflammasome, B and T cell activationare associated with acute lesions.

Conclusion: We observed prominent role for B cells and immunoglobulins in lesions, suggesting that B cells may be correlated with disease severity rather than protection or cure. Defining the host inflammatory pathways within L. *tropica* lesions will be critical for development of new therapeutic interventions. In addition, dual RNA-seq of host and parasite genes is highly recommended for in-depth investigation.

Keywords: Cutaneous leishmaniasis, *Leishmaniatropica*, RNA sequencing, Transcriptome.

ICIA2018\Immunoparasitology\Oral\8087

8087

The Role of *Toxoplasma gondii* Lysate, Excretory/Secretory Antigens and a Combination of them on the expression of IFN- γ and IL-4 and mice surveillanceAli Fattahi Bafghi^{*1}, Aghdas Mirjalili², Mohammad Kazemi Arababadi³

1. Associate professor, Medical Parasitology and Mycology Department, School of Medicine, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

2. MSc Histologist, Morph-anatomy Department, School of Medicine, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

3. PhD in Immunology, Immunology of Infectious Diseases Research Center, Rafsanjan University of Medical Sciences, Rafsanjan, Iran.

Background: *Toxoplasma gondii*, an obligate intracellular parasitic protozoan, is capable of infecting a wide range of warm-blooded vertebrates. This survey was designed to The Role of *Toxoplasma gondii* Lysate, Excretory/Secretory Antigens and a Combination of them on the expression of IFN- γ and IL-4 and mice surveillance.

Methods: The above-mentioned antigens were prepared, encountered with the isolated murine peritoneal leukocytes and then the IFN- γ and IL-4 levels were measured in the stimulated leukocytes culture supernatants using ELISA technique. Moreover, the effect of these antigens on the mice survival time was examined.

Results: The results of one-way ANOVA showed that there was a significant difference among the testing groups under study (TLAs, ESAs and TLA: ESA) on the basis of IL-4 and IFN- γ concentrations ($P < 0.05$). In addition, the result of Turkey's HSD test showed only a significant difference between the TLA: ESA and TLA groups ($P < 0.05$). The results of two-tailed t-test for pair-wise comparison of the mice survival time in the immunized and challenged groups showed a significant difference between the two testing groups ESAs as well as TLA: ESA and the control groups ($P < 0.05$). The Pearson's correlation coefficient results showed that there was a significant correlation between the mice survival time and the spleen size ($P < 0.01$).

Concussion: This study discussed on the host-parasite interaction with the involvement of IFN- γ and IL-4 and how these interactions possibly help the parasite to establish its life-cycle in nature.

Keywords: IFN- γ , IL-4, *Toxoplasma gondii*, host-parasite interaction.

ICIA2018\Immunoparasitology\Oral\8241
8241

Preparation of Theileria annulata infected Hyalomma anatolicum ticks stabilate for use in efficacy of bovine Theileriosis vaccine

Asghar Afshari¹, Gholamreza Habibi¹, Kasra Esmacilnia¹, Hossein Modirrousta²

1. Department of Parasite vaccine research and production of Razi vaccine and serum research institute, Agriculture Research, Education and Extension Organization (AREEO), Karaj, Iran

2. Department of Wildlife research of Razi vaccine and serum research institute, Agriculture Research, Education and Extension Organization (AREEO), Karaj, Iran

Background: Tropical theileriosis caused by the protozoan parasite Theileria annulata, is one of the most important tick-borne disease that affect cattles, accompanied with economic loses in endemic area such as Iran. Theileriosis is a vector-borne disease that is transmitted by Hyalomma ticks (Acari-ixodidae). Control and prevention of tropical theileriosis are carried out by both acaricides and administration of live attenuated vaccine. Efficacy of vaccine needs to be challenged by live virulent parasite preserved in tick stabilate that infected with acute strain of Theileria annulata. This study was carried out in order to preparing of tick infected stabilate to applying in vaccine efficacy test.

Methods: The experimental theileriosis was achieved by inoculation of previously infected blood sample to a naïve healthy calf. Different stages of Hyalomma anatolicum tick life cycles including eggs, larvae, nymph and adult gained by incubation and feeding on white rabbit in laboratory conditions. The unfed nymph ticks fed on experimentally infected calf. Theileria annulata infection in calf and ticks was detected by PCR using specific primers. The tick stabilate was prepared by grinding infected ticks in normal saline and preserved in deep freeze using 15% of glycerol as cryoprotectant. Bioassay was performed to confirm the infectivity of prepared stabilate by subcutaneous injection to intact susceptible 4 months old calf.

Result: Hyalomma anatolicum ticks successfully reared and complete life cycle lasted about 90 days. Theileria annulata infection was confirmed by clinical signs, microscopic observation of blood smears and specific PCR test in both calf and ticks specimens. Incubation period of disease by injection of blood and stabilate lasted 14 and 8 days, respectively.

Conclusion: Severity of clinical signs and low incubation period of disease caused by injected stabilate revealed its acuteness and so it can be used in vaccine efficacy test and other molecular, serological and challenge studies.

Keywords: Theileria, Tick, Stabilte, Vaccine efficacy

ICIA2018\Immunoparasitology\Oral\8262
8262

Evaluation of Natural Killer Cells Function Exposed to Recombinant *Leishmania major* Lipophosphoglycan-3 Antigens

Sanaz Rasolzadeh Nori^{1,4}, Arash Pourgholaminejad³, Vahid Younesi², Mehdi Yousefi^{1,4}

1. Stem Cell and Regenerative Medicine Institute, Tabriz University of Medical Sciences, Tabriz, Iran

2. Pishtaz Teb Zaman Diagnostics, Tehran, Iran

3. Department of Immunology, School of Medicine, Guilan University of Medical Sciences, Rasht, Iran

4. Department of Immunology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Leishmaniasis is one of the widespread infectious diseases that caused by intracellular parasite, *Leishmania major*. Nowadays, there is not effective vaccination procedure and immune-booster strategy for this disease. Among many antigenic molecules in the structure of *Leishmania major*, lipophosphoglycan-3 (LPG-3) antigen and its fragments are critically immunogenic and play major role in the pathogenicity of the parasite. It has been hypothesized that LPG-3 enhances immune response against Leishmaniasis through natural killer (NK) cells activation and can increase immune response of the host. However, in the current study, we analyzed the effects of LPG-3 in activation of human NK cells.

Methods: Human NK cells were isolated from healthy volunteers (n=10) by magnetic-activated cell sorting (MACS). NK cells purity was checked by Flowcytometry. Then, purified cells were co-incubated with recombinant LPG-3 in presence/absence of TLR-2 inhibitory-monoclonal antibody for 72 hours. Afterward, the gene expression and protein secretion of IFN- γ and TNF- α were measured in treated-NK cells and cell-supernatants by real-time PCR and ELISA methods, respectively.

Results: Our findings indicated that recombinant LPG-3 increases the expression of IFN- γ and TNF- α significantly ($p<0.01$) in NK cells after three days of co-incubation in compare with untreated-NK cells. Also, the concentration of IFN- γ and TNF- α in cell-supernatants increased ($p<0.01$).

Conclusion: We concluded that *Leishmania major* LPG-3 antigen can activate NK cells toward enhancement of inflammation, and enhances immune response against leishmaniasis. Our findings suggest immunization with LPG-3 may enhance innate immune system against leishmaniasis.

Keywords: NK cell, LPG-3, IFN- γ , TNF- α , Innate immunity

ICIA2018\Immunoparasitology\Oral\8316

8316

The Evaluation Effect of fucose-mannose ligand of *Leishmania infantum* on murine peritoneal macrophages in vitro

Namdar Ahmadabad Hasan¹, Shafiei Reza¹, Nezafat Ferizi Maryam¹, Kalantar Kurosh², Hatam GholamReza³, Bakhshi Joibari Fahimeh⁴

1. Department of Pathobiology and Laboratory Sciences, School of Medicine, North Khorasan University of Medical Sciences, Bojnourd, Iran

2. Department of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

3. Department of parasitology and Mycology, school of Medicine, Shiraz University of Medical Sciences

4. Student Research Committee, School of Medicine, North Khorasan University of Medical Sciences, Bojnourd, Iran

Background: Fucose-mannose ligand (FML) is glycoprotein, which is present both on promastigotes and amastigotes of *Leishmania infantum*. It has potential use as a tool for prognosis, diagnosis, and vaccine development of visceral leishmaniasis. In present study we evaluated the effect of FML on murine peritoneal macrophages in vitro.

Methods: We treated macrophages and LPS stimulated macrophages with different concentration of FML purified from *Leishmania infantum* and measured the concentration of TNF- α , IP-10, IL-12p70, and IL-10 production in culture supernatant of these cells by sandwich ELISA. We also measured culture supernatant levels of nitric oxide (NO) by Griess reaction.

Results: Our findings showed that that FML significantly increases NO, IL-10, IL-12p70 and IP-10 production in macrophages, but doesn't have any effect on TNF- α production (*P< 0.05). We also found that FML significantly increases the IL-12p70/IL-10 ratio in culture supernatants of macrophages(*P< 0.05).

Conclusion: We concluded that FML can polarize macrophages to an M1-like phenotype, although our results showed that FML enhances IL-10 production from macrophages and doesn't increase TNF- α production from them.

Keywords: Fucose-mannose ligand, Macrophage, Nitric oxide, Visceral Leishmaniasis

ICIA2018\Immunoparasitology\Oral\8503

8503

Evaluation of stability of the attenuated line of Leishmania major transfected with PSA gene in BALB/c mouseSahar Mashayekhi ¹, Fahime Aeine negin¹, Hamid Daneshvar¹*1. Department of Immunology, Medical School of Kerman University*

Background: Prostate cancer is the second most common cancer among men worldwide. The attenuated leishmania can be a good candidate for live recombinant vaccine to express proteins and a tumor-associated antigen (TAA). In this study, attenuated line of Leishmania major (H-line) was used to express the PSA gene. The aim of this study was to evaluate the expression of the PSA gene in this parasite and to investigate the stability of transfected parasite in BALB / c mice.

Methods: In this study, we cloned PSA gene into attenuated leishmania major. To examine the safety of the transgenic parasite, BALB/c mice were injected subcutaneously with stationary-phase L. major wild, H-line, recombinant promastigotes. The lesion volume was measured weekly. To investigate the ability of promastigotes to infect macrophages and their survival within macrophages, peritoneal macrophages were exposed to stationary-phase promastigotes. The culture slides then were incubated in 5% CO₂, 95% air at 37°C. At the end of the incubation period, the infection status of the macrophages was determined by microscopy.

Results: The PSA gene cloned into leishmania major H-line successfully for first time and we could detect the expression of PSA gene in leishmania major H-line. Safety evaluated by 2 criteria: first, the promastigotes of the recombinant leishmania major could enter but not survive in macrophages, and second, that mice inoculated subcutaneously with attenuated parasites did not develop skin lesions.

Conclusions: These finding represent that leishmania major H-line could be used as a new vector vaccine for expression of TAAs and other proteins.

Keywords: L.major H-line, vector vaccine, prostate cancer, PSA

ICIA2018\Immunoparasitology\Oral\8641
8641

Effect of *Lavandula angustifolia* extract on the antigenemia and WBC responses to acute infection with *Toxoplasma gondii* in mice

Mohammadreza Alijani¹, Yaser Pirali², Rasool Rahimi Junqani³

1. DVM Student of Veterinary Medicine, Shahrekord University, Shahrekord-Iran

2. Professor, Department of Clinical Sciences, Faculty of Veterinary Medicine, Shahrekord University, Shahrekord-Iran

3. DVM Graduated Student of Veterinary Medicine, Shahrekord University, Shahrekord-Iran

Background: Toxoplasmosis, caused by the protozoan parasite *Toxoplasma gondii*, is one of the most common parasitic infections of man and other warm-blooded animals. To control toxoplasmosis, a combination of antifolates, such as pyrimethamine and sulfadiazine, has been used, but a large percentage of undesirable side effects such as hematological toxicity (caused by pyrimethamine), cutaneous rash, leukopenia and thrombocytopenia (caused by sulphonamides) are often reported. The purpose of this study was to determine the effect of *Lavandula angustifolia* extract on the antigenemia and WBC responses of mice to acute infection with *Toxoplasma gondii*.

Methods: 30 female mice were infected by intraperitoneal (IP) injection of 104 tachyzoites of the virulent RH strain, and divided in 3 groups as follows: 1 control group. 2 infected mice that received extract of *Lavandula angustifolia* 50 mg/kg. 3 infected mice that received extract of *Lavandula angustifolia* 100 mg/kg. for 6 days after infection blood sample were collected and toxoplasma antigen were tested by dot-ELISA. Also white blood cell count (neutrophil, lymphocyte, eosinophil and monocyte) were measured. All data were analyzed via ANOVA, with a significant level $P < 0.05$.

Results: We observed that toxoplasmosis cause lymphopenia, neutrophilia and eosinophilia in all groups. Although extract of *Lavandula angustifolia* can decrease total WBC, but the difference observed is not significant ($p > 0.05$). Circulating antigens were detected early on day 2 PI in control group and day 3 PI in treatment groups, and the titers increased from day 4 PI in all groups. In other factors such as body weight, splenomegaly and hepatomegaly there was no significant difference between groups received extract of *Lavandula angustifolia* and the control group ($p > 0.05$). Acutely infected mice died within 7 to 10 days' post-infection as a function of the strain and inoculum size.

Conclusion: According to the results of the present study *Lavandula angustifolia* extract has no effect on *Toxoplasma gondii* infection in mice.

Keywords: *Withania somnifera*, antigenemia, *Toxoplasma gondii*, mice



Immunopharmacology and Medicinal Plants

Oral Presentation

14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Immunopharmacology and Medicinal Plants\Oral\5269
5269

A Comparison of the Immunosuppressive Effects of Silymarin and Cyclosporine A on the Proliferation of Regulatory T Cells and Function

Niloufar Keyhanmehr¹ , Hossein Motedayyen¹ , Nahid Eskandari¹

1. Department of Immunology, Faculty of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

Background: Immunosuppressive agents are necessary to enhance allograft tolerance after transplantation and the treatment of autoimmune disorders. Regulatory T cells (Tregs), a subset of T cells, play a pivotal role in improving allograft tolerance and determining the fate of transplanted organs. Therefore, the specific aim of this study was to investigate the immunomodulatory effects of cyclosporine A (CsA) and silymarin on Treg proliferation and their cytokine production.

Methods: Tregs were isolated from peripheral blood mononuclear cells (PBMCs) from healthy donors and phenotypic characteristics of Tregs were determined by flow cytometry. Tregs were expanded and then cultured with different concentrations of CsA and silymarin. Immunomodulatory effects of CsA and silymarin on the proliferation of Tregs and function were determined after 3 and 5 days of culture.

Results: Our results showed that CsA significantly decreased Treg proliferation in a dose-dependent fashion. However, CsA did not have the ability to induce Treg function through the production of transforming growth factor-beta 1 (TGF- β 1). In contrast, silymarin significantly increased the proliferation of Tregs. A statistically significant increase was also observed in the function of Tregs through TGF- β 1 production.

Conclusion: The results of this study for the first time show that silymarin, unlike CsA, possess the ability to increase the proliferation and function of Tregs and may be beneficial in treatment of autoimmune disorders and improvement of Treg-dependent allograft tolerance after transplantation.

Keywords: Silymarin, Cyclosporin A, Regulatory T cells, TGF- β 1

ICIA2018\Immunopharmacology and Medicinal Plants\Oral\7391
7391

An in vitro study on cytotoxic effects of *Androctonus crassicauda* scorpion venom on K562 cell line

Golnaz Rashidi^{1,2}, Ata Ghadiri¹, Babak Vazirianzade³, Nazanin Fathi^{1,2}, Ali Khodadadi^{*1,2}, Mohammad Rashno^{1,2}, Mohammad Ghasemi^{1,2}

1. Department of Immunology, Faculty of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

2. Cancer Petroleum and Environmental Pollutants Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

Infectious and Tropical Diseases Research Center, Health Research Institute” and “Medical Entomology and Vector Control Department, Public Health Faculty”, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

Background: Today, cancer is posed as one of the major problems in the world. The incidence of cancer and its associated mortality rates are increasing and in spite of diverse medical treatments in this field none of them were deemed thoroughly effective. Therefore using the existing rich resources in the nature for therapeutic purposes is considerable. The objective of the present study was to measure the ability of *Androctonus crassicauda* scorpion venom on inhibition of cell proliferation in K562 cell line derived from human Chronic Myeloid Leukemia.

Methods: Crude venom of *Androctonus crassicauda* scorpion was obtained by electrical stimulation of telson. Bradford assay was employed to measure the protein concentration. K562 cell line was cultured in RPMI-1640 with 10% heat inactivated fetal bovine serum, 100 units/ml penicillin and 100µg/ml streptomycin. Measurement of cell viability and metabolic activity was performed by MTT assay. 5×10^4 cells/well of K562 cells were seeded in a 96-well tissue culture plate. The cells were treated with increasing concentrations of *A. crassicauda* venom from 40 to 160 µg/ml for 24h. The absorbance was read at 570 nm in a micro-plate reader.

Result: Protein concentration of scorpion venom in this study was calculated 0.5mg/ml. This study showed that the scorpion venom inhibits the growth of K562 cells with $IC_{50}=111.46\mu g/ml$.

Conclusion: The results of this study proved that *A. crassicauda* venom induces growth inhibition in K562 cell line. According to other studies as well as the current study, which demonstrated the cytotoxic effect of natural toxins on cancer cells, apoptotic effects induced by *A. crassicauda* in K562 cell line will be dealt with in our future research, which is hoped to be a step in the improvement of cancer treatments using natural products.

Keywords: *Androctonus crassicauda*, Scorpion venom, K562 Cell Line, MTT, IC_{50}

ICIA2018\Immunopharmacology and Medicinal Plants\Oral\7817

7817

The in vitro impact of Glycyrrhizic acid on CD4⁺ T lymphocytes through OX40 receptor in the patients with allergic rhinitis

Saloomeh Fouladi¹, Nahid Eskandari¹, Mohsen Masjedi¹

1. Department of Immunology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

Background: Glycyrrhizic acid(GA) the major bioactive component of glycyrrhiza possess a great variety of pharmacological properties such as anti-inflammatory, anti-allergic and immunomodulatory activities. In the present work, the in vitro anti-allergic effect of GA through OX40 receptor was investigated in patients with allergic rhinitis.

Methods: Purified CD4⁺ T cells from patients with allergic rhinitis (n=12) were activated with anti-CD3/anti-CD28 and anti-OX40 mAb agonist antibody and then treated with different concentration of GA and dexamethasone. Then, cells were incubated (72 hr) for proliferation assay. Protein expression of OX40 in anti-OX40 mAb stimulated CD4⁺ T cells were evaluated by flow cytometry. mRNA expression of the T-bet, GATA3 and forkhead box P3 (Foxp3) transcriptional factors were measured by quantitative polymerase chain reaction.

Results: GA significantly inhibited the augmented T cells proliferation induced with anti-OX40 mAb ($p < 0.05$ - $p < 0.01$ at 100 μ M and 200 μ M GA compared to control). OX40 expression was also significantly decreased ($p < 0.05$). Dexamethasone and GA showed inhibitory effect on T-bet and GATA3 gene expression, but this inhibition was only significant in the case of GATA3 ($p < 0.05$ - $p < 0.01$ at dexamethasone and 200 μ M GA). In contrast, enhanced gene expression of Foxp3 was seen using both of them ($p < 0.05$).

Conclusion: We suggest that GA may have a therapeutic effect on allergic rhinitis partly by regulating the Th1/Th2 balance through suppressing OX40 and increasing the activity of regulatory T cells.

Keywords: Glycyrrhizic acid, OX40, Allergic Rhinitis, Th1/Th2 balance

ICIA2018\Immunopharmacology and Medicinal Plants\Oral\7855

7855

Decline in the expression and release of inflammatory mediators by the extracts from *Satureja hortensis* in macrophagesZahra Farzaneh¹, Kurosh Kalantar¹, Zahra Amirghofran^{1,2}*1. Department of Immunology, Shiraz University of Medical Sciences, Shiraz, Iran**2. Autoimmune Disease Research Center, Shiraz University of Medical Sciences, Shiraz, Iran*

Background: *Satureja hortensis* is a medicinal plant used in traditional medicine for its anti-inflammatory effects. Macrophages are among the main cells involved in the generation of inflammation. This study has investigated the anti-inflammatory effect of *Satureja hortensis* extracts on macrophages by measuring the expression and release of important inflammatory mediators by these cells.

Methods: J774.1 mouse macrophages were stimulated with lipopolysaccharide (LPS) and treated by various extracts of *Satureja hortensis*. We examined the macrophages for nitric oxide (NO) production using the colorimetric assay followed by real time-PCR for gene expressions and ELISA for cytokine levels. Flowcytometry was used for measuring the surface expression of intercellular adhesion molecule (ICAM)-1.

Results: Dichloromethane and hexane extracts more efficiently reduced NO production at non-cytotoxic concentrations compared to the aqueous and butanol extracts. The extracts at 25 µg/ml decreased NO release to less than 43% (dichloromethane) and 53% (hexane) of the control ($p < 0.05$). All concentrations of both extracts decreased gene expression of inducible NO synthase (iNOS) (< 0.44 fold of control), cyclooxygenase (COX)-2 (< 0.29 fold) interleukin (IL)-1 β (< 0.41 fold), IL-6 (< 0.25 fold) and tumor necrosis factor (TNF)- α (< 0.2 fold). At 25 µg/ml, dichloromethane extract reduced the IL-6 protein level to $79.2 \pm 0.84\%$ and the hexane extract to $49.4 \pm 6.8\%$ of the control. The IL-1 β level decreased to $62.5 \pm 14.6\%$ (dichloromethane) and $36.7 \pm 24.3\%$ (hexane). Both extracts reduced ICAM-1 fluorescent intensity of expression in comparison with the control.

Conclusion: The inhibitory effects of *Satureja hortensis* extracts on the expression and release of inflammatory mediators in macrophages suggest the presence of useful anti-inflammatory compound(s) in this plant.

Keywords: *Satureja hortensis*, Anti-inflammatory effect, Macrophages

ICIA2018\Immunopharmacology and Medicinal Plants\Oral\8058

8058

Teucrium polium extract modulates the immune response in Streptozotocin-induced diabetic rats

Hamidreza Safari¹, Mohsen Naseri², Gholamreza Anani Sarab², Tayebeh Nowras¹, Mahdi Abdollahi¹, Zeinab Rostamy¹

1. Student Research Committee, Birjand University of Medical Sciences, Birjand, Iran

2. Infectious Diseases Research Center, Birjand University of Medical Sciences, Birjand, Iran

Background: Diabetes is a chronic metabolic disorder that is an epidemic disease affecting a large proportion of the population around the world. Recent studies have suggested that proinflammatory cytokines play an important role in the pathogenesis of type 1 diabetes (T1DM). This study aimed to investigate the Immunomodulatory effects of Teucrium polium in Streptozotocin-induced diabetic rats.

Methods: In this experimental study, diabetes was induced by streptozotocin in male Wistar rats (65 mg/kg). Rats were divided into the following groups: control, untreated diabetic, 2 Teucrium polium (100, 200 mg/kg/d)–treated diabetic groups. Rat's spleen cells cultured, then cell proliferation and inflammatory cytokine release (interleukin-6 (IL-6), interleukin-1 (IL-1)) from cultured spleen cells measured by MTT and ELISA kit, respectively. Data were analyzed using one-way ANOVA and Tukey post-hoc tests in SPSS software and P-values less than 0.05 were considered statistically significant.

Results: Proliferation of spleen cells in treatment groups was significantly reduced compared with Control group ($P < 0.05$). There was no significant difference in anti-inflammatory activity between the groups.

Conclusion: The result demonstrates that Teucrium polium increased the immune cells proliferation without increasing inflammatory cytokine. Teucrium polium has Immunomodulatory effects, with the goal that the effect of continuous severity due to inflammation can be suppressed.

Keywords: Teucrium polium, STZ, Anti-inflammatory effect, Diabetes.

ICIA2018\Immunopharmacology and Medicinal Plants\Oral\8080
8080

Induction of Apoptosis in Raji Burkitt Lymphoma Cell Line by *Ferula hezarlalehzarica* Extract

Yahya Asemani¹, Maryam Bayat¹, Abbas Azadmehr², Reza Hajiaghachee³, Zahra Amirghofran^{1,4}

1. Immunology Department, Shiraz University of Medical Sciences, Shiraz, Iran

2. Immunology Department, Babol University of Medical Sciences, Babol, Iran

3. Autoimmune Disease Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

4. Medicinal Plants Research Center, Institute of Medicinal Plants, ACECR, Karaj, Iran

Background: *Ferula* plants have recently been considered for their anti-cancer effects. In this study we aimed to determine the anti-tumor and apoptosis-inducing activity of a newly discovered native species, *Ferula hezarlalehzarica* against cancerous cell lines.

Methods: The cytotoxic activity of methanol, hexane, ethyl acetate, butanol and water extracts of *F. hezarlalehzarica* on Raji, EL-4, HepG2, HeLa, and K562 cell lines was examined by the MTT colorimetric assay. Analysis of apoptosis was performed by Annexin V-PE/7-AAD staining using flow cytometry. Changes in the expression of genes associated with apoptosis including Bax, Bcl-2 and Fas were investigated by real-time PCR. The JC-1 lipophilic cationic dye staining was used for assessment of changes in mitochondrial membrane potential (MMP).

Results: The most effective inhibitory activity was observed for the hexane extract of *F. hezarlalehzarica* with IC₅₀ value of 31.6 µg/ml against Raji cell line. A 24-hour analysis of apoptosis in Raji cells showed that this extract induced apoptosis dose-dependently such that at 100 µg/ml of the extract 68.53±1.2% and at 200 µg/ml 96.93±1.45% underwent apoptosis compared to the negative control (5.2±2.02%). Treatment of cells with 75 µg/ml of the hexane extract significantly up-regulated the gene expression levels of pro-apoptotic molecules Bax (1.75±0.31-fold, p<0.01) and Fas (5.02±0.74-fold, p<0.01) and down-regulated the expression level of Bcl-2 (0.23±0.008-fold, p<0.001). The ratio of Bax to Bcl-2 expressions at 75 µg/mL was 7.4±0.2. A significant decrease was detected in MMP at concentrations more than 100 µg/ml (≈0.4, p<0.05) of extract.

Conclusion: The hexane extract of *F. hezarlalehzarica* induced apoptosis in Raji cell line by regulating apoptotic and anti-apoptotic genes as well as inducing MMP depolarization. Therefore this extract has the potential to be considered as an anti-cancer agent against Burkitt lymphoma.

Keywords: Apoptosis, Burkitt Lymphoma, *Ferula hezarlalehzarica*

ICIA2018\Immunopharmacology and Medicinal Plants\Oral\8268

8268

Anti-inflammatory Activity of *Lactobacillus paracasei* on Isoproterenol-Induced Heart Failure in Rat

Bahador Bagheri¹, Zohreh Hesari², Mahdi Barati³, Parviz Kokhaei⁴

1. Cancer Research Center and Department of Pharmacology, Semnan University of Medical Sciences, Semnan, Iran

2. School of Veterinary Medicine, Semnan University, Semnan, Iran

3. Department of Immunology, Mashhad University of Medical Sciences, Mashhad, Iran

4. Cancer Research Center and Department of Immunology, Semnan University of Medical Sciences, Semnan, Iran

Background: A wealth of evidence has associated heart failure with inflammatory cytokines that can predict clinical outcome. The emerging information has suggested that the certain probiotics have immunomodulatory effects in diseases that have linkages with inflammation. This study was performed to determine whether oral probiotic administration could attenuate or prevent the heart failure induced by isoproterenol in animal model.

Methods: *Lactobacillus paracasei* subsp. *paracasei* 8700:2 was used. Adult male Wistar rats (n=40, 185±15g), were randomly assigned to five groups, including the control group, probiotic control (0.25 mg/kg/day, oral), isoproterenol control (5 mg/kg/day, subcutaneous), pretreatment group and treatment group were divided. The groups were studied for 30 days. Serum levels of TNF- α and ANP were measured by ELISA and the standard histopathological examination was performed.

Results: A significant reduction was seen in the levels of TNF- α and ANP in the treatment group compared to the control group and isoproterenol group (p<0.05). Pretreatment with *L. paracasei* subpp.*paracasei* 8700:2 could not prevent the development of heart failure induced by isoproterenol. Histopathological analysis showed a marked attenuation of cardiac necrosis and infiltration of immune cells in treatment group.

Conclusion: This report indicates that the strain of *L. paracasei* attenuates inflammatory responses in rats with heart failure. These effects are mediated by reduction in serum levels of the TNF- α and ANP.

Keywords: heart failure, isoproterenol, TNF- α , ANP, *Lactobacillus paracasei*

ICIA2018\Immunopharmacology and Medicinal Plants\Oral\8282

8282

The alteration of human colon cancer, HT29 cell line, after administration of Kombucha, a fermented microbial productMahsa Alem¹, Somayeh Ahmadi afshar¹, Norouz Delirezsh¹*1. Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran*

Background: Kombucha is an ancient healing supplement of Asian origin. It strengthens the immune system as a natural medicine. Anti-cancer properties of Kombucha and its derivatives have been shown in many in vivo and in-vitro experiments. It can revive the ionic balance in the cell membrane so prevent many of ailments. The current research was conducted to investigate the apoptotic and cytotoxic effects of administration of Kombucha on some functions of the HT29 as human colon cancer cell line.

Methods: The study population consisted, cells were allocated in to different groups such as negative control, treatment with 4 doses (50, 100, 150 and 200 µl/ml) of Kombucha, for 24 h, after cell line culture. All groups were tested via tetrazolium dye (MTT) assay for vitality investigation, natural red (NR) for cell membrane viability investigation, invert microscopy for evaluation of cell's morphology, acridine orange (AO) and propidium iodide (PI) staining for evaluation of necrosis and apoptosis respectively.

Results: The cell membrane viability was decreased in HT29 cell line. MTT assay test showed that Kombucha could make significant changes in the vitality of Kombucha treated colon cancer cell line. PI and AO staining manifested that Kombucha induced apoptosis in lower dose and necrosis in upper dose. These results were dose-dependent and statistically significant compared to the control group.

Conclusion: Based on these results, it appeared that this agent could be a good candidate for further evaluation as effective therapy acting through induction of apoptosis in human colon cancer cell line.

Keywords: HT29-cell line, Human colon cancer, Kombucha

ICIA2018\Immunopharmacology and Medicinal Plants\Oral\8436

8436

Epigallocatechin-3-gallate enhances differentiation in acute promyelocytic leukemia cells via inhibition of PML-RAR α and HDAC1

Maliheh Moradzadeh¹, Abazar Roustazadeh², Saiedeh Erfanian², Alijan Tabarraei³

1. Golestan Rheumatology Research Center, Golestan University of Medical Sciences, Gorgan, Iran

2. Research Center for Non-Communicable Diseases and Biochemistry Department; Department of Advanced Medical Sciences & Technologies, School of Medicine, Jahrom University of Medical Sciences (JUMS), Iran

3. Infectious Diseases Research Center, Golestan University of Medical Sciences, Gorgan, Iran

Background: Introducing all trans retinoic acid (ATRA) for the treatment of acute promyelocytic leukemia (APL) patients could dramatically improve their survival rate. However, resistance to this drug and their toxicity are major problems in the treatment of APL. Earlier studies suggested that in haematopoietic neoplasms, the green tea polyphenol Epigallocatechin gallate (EGCG) induces cell death without adversely affecting healthy cells.

Methods: In the present study, the potential anti-promyelocytic leukemia activity of EGCG and the underlying molecular mechanisms were investigated.

Results: EGCG (100 μ M) significantly inhibited proliferation and induced apoptosis in HL-60 and NB4 cells. The effect was associated with the decreased expressions of MDR proteins ABCB1 and ABCC1, while the expressions of proapoptotic genes CASP3, CASP8, p21 and Bax/Bcl-2 ratio were significantly increased. EGCG at 25 μ M concentration, like ATRA (1 μ M), induced differentiation of leukemic cells towards granulocytic pattern. Furthermore, EGCG suppressed the expression of clinical marker PML/RAR α in NB4 cells, and reduced the expression of HDAC1 in leukemic cells.

Conclusion: The results suggest that EGCG can be considered, alone or in combination with ATRA, for preclinical and also clinical testing in APL patients.

Key words: APL, EGCG, Apoptosis, Differentiation, MDR

ICIA2018\Immunopharmacology and Medicinal Plants\Oral\10043

10043

Evaluation and comparison of Albumin-curcumin Nano particle and curcumin anti-cancer effects on Sk-BR3 cell lineZahra Matluobi¹, Parisa Amir Kalvanagh¹, Zuhair Saraf Hasan¹*1. Immunology Department, Tarbiat Modares University, Tehran, Iran*

Background: The advent of chemotherapy and radiation-resistant cancer cells, as well as severe complications of chemotherapy drugs, has led researchers to try to find solutions to these problems. Among the solutions that are very much considered is the use of some herbal compounds with antioxidant properties as therapeutic supplements along with chemotherapy drugs that reduce the dose of these drugs and thus reduce their side effects. Among the plant antioxidants, curcumin is one of the strongest antioxidants to inhibit the proliferation of cancer cells and induce apoptosis in them. However, curcumin has had very low dissolution in water and its ketone form has been predominant in acidic conditions (the PH of environment surrounding the tumor is acidic). In addition, degradation rate of curcumin by enzyme and its elimination from the bloodstream is very high. So, the antioxidants effects of curcumin in the body have had greatly reduce. To overcome these problems, various compounds of Nano-curcumin have been produced and tested. In this study, curcumin was conjugated with human serum albumin (HSA), and were measured for its effects on breast cancer cell lines (SK-BR3) and were compared with curcumin anti-cancer effects.

Methods: After making of curcumin-albumin Nano drug, anticancer effects of its have been examined by MTT and flow cytometry tests on SK-BR3and PBMC cells.

Results: FTIR graph and TEM picture confirmed that curcumin and albumin have been combined together. The MTT results showed that in first 24 hours none of the two drugs had a significant effect. After 48 hours curcumin and Nano curcumin had led to 14 % and 24% death in Sk-Br3 cells, respectively. Results of 72 hours showed that Nano curcumin had 52 % cytotoxic effect on cells while the cytotoxicity of curcumin on cells was only 24%. None of the drugs had significant toxicity on PBMC. The results of MTT were approved by Flow cytometry.

Conclusions: Nano-curcumin drug increases curcumin solubility in water and its stability in physiological and acidic conditions. These factors had led to a significant increase in the toxicity of curcumin on the SK-BR3 cells without increasing the cytotoxicity of the normal cells.

Keywords: curcumin, Albumin, Nano drug, SK-BR3



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Innate Immunity and Inflammation

Oral Presentation

ICIA2018\Innate Immunity and Inflammation\Oral\7551
7551

IL-17 producing CD4+CD45RO+ T-cells in Atherosclerosis express GITR molecule

Atefe Ghamar Talepoor¹, Negar Behnamfar¹, Mohammad Javad Zibaenezhad², Mehrnoosh Doroudchi¹

1. Memory T-cell Laboratory, Department of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran.

2. Cardiovascular Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Atherosclerosis (AS) is a chronic inflammatory disease of vessel walls associated with infiltration of immune cells which their function is controlled by different co-stimulatory and co-inhibitory receptors. In this study, we investigated the expression of co-inhibitory molecules on the memory and effector T-cells in patients with Atherosclerosis.

Methods: Patients included 9 hypertensive, dyslipidemic, non-diabetic, non-smoker individuals with the diagnosis of coronary artery disease and controls were 8 normotensive, normolipemic, non-diabetic, non-smoker individuals with normal coronary angiography/ insignificant coronary artery disease. PBMCs were separated from the blood and memory T-cell subsets as well as the expression of Glucocorticoid-induced tumor necrosis factor receptor (GITR), Programmed Death-1 (PD-1), IL-17A and IFN- γ were quantified by flowcytometry.

Results: CD4+CD45RO+ memory T-cells and CD4+CD45RO- effector T-cells in patients expressed the highest level of GITR molecule. The IL-17 producing memory CD4+CD45RO+ T-cells were enriched in GITR molecule in the patients group (P=0.03). The increased population of GITR+ effector CD4+CD45RO- T-cells in patients, however, did not produce IL-17 (P=0.03). PD-1 expression on memory T-cells of the patients was higher than the controls and was concomitant with the lack of IFN- γ expression (P=0.05). IFN- γ production by effector T-cells was only seen in the PD-1- population in both groups.

Conclusions: We provide data on the expression of GITR molecule on IL-17 producing memory T-cells in patients with CAD. A population of memory T-cells, which expressed PD-1 and were not producing IFN- γ , also increased in patients' blood. These data suggest the modified phenotype/function of T-cell subsets in the atherosclerotic inflammation.

Keywords: Atherosclerosis, Immunopathology, T cell, inflammation

ICIA2018\Innate Immunity and Inflammation\Oral\7851

7851

Differential pattern of eotaxins (CCL11, CCL24, CCL26) and CCR3 expression in Iranian Parkinson's patients

Hossein Khorramdelazad^{1,2}

1. *Molecular Medicine Research Center, University of Rafsanjan, Rafsanjan, Iran*

2. *Department of Immunology School of Medicine. University of Rafsanjan, Rafsanjan, Iran*

Background: The Parkinson's disease (PD) is defined as the most frequent human neurodegenerative disorder, after Alzheimer's disease. Recent studies indicated that cytokine/chemokine network and their downstream inflammatory responses are remarkably involved in the PD pathogenesis.

Methods: The present study was aimed to examine the expression of eotaxin family members, including CCL11, CCL2, and CCL26 in parallel with their receptor CCR3 in PD patients at both mRNA and protein levels. Blood specimens were isolated from 30 patients suffering from PD and 30 healthy participants as control for analysis in this investigation. Demographic information of patients and controls were also collected via a researcher-designed questionnaire. The enzyme-linked immunosorbent assay (ELISA) was performed for the detection of serum levels of CCL11, CCL24, and CCL26. Flow cytometric and real-time PCR methods were also employed to determine expression of CCR3.

Results: We observed that CCL11 and CCL24 were significantly elevated in serum of PD patients, whereas CCL26 was not significant changed whereby compared to control group ($P < 0.001$). In addition, results obtained from flow cytometry and RT-PCR showed that there was no remarkable change in the expression of the CCR3 in PD patients compared with control group.

Conclusion: According to these findings, it could presumably be reasonable to conclude that CCL11, CLL24 might be play a part in the PD pathogenesis and they can be considered as potential serum-based biomarkers in this disease. Moreover, it is probable that microglial cells are expressing CCR3 more than circulatory eosinophils in response to released eotaxins from astrocytes due to progressed neurotoxicity as well as PD.

Keywords: CCL11, CCL24, CCL26, eotaxin, CCR3, Parkinsonism.

ICIA2018\Innate Immunity and Inflammation\Oral\8154

8154

Macrophage repolarization using PLGA-PEG nanoparticles containing Chrysin: possible application in regenerative medicine

Akram Firouzi Amandi¹, Zohreh Babaloo¹, Younes Pilehvar-Soltanahmad², Leili Aghebati-Maleki³, Farhad Jadidi-Niaragh¹, Mehdi Dadashpour²

1. Immunology Department, Medicine Faculty, Tabriz University of Medical Sciences, Tabriz, Iran

2. Department of Medical Biotechnology, Faculty of Advance Medical Science, Tabriz University of Medical Science, Tabriz, Iran

3. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

Background: The aim of this study was to evaluate the efficiency of using a natural substance, Chrysin, encapsulated in PLGA-PEG nanoparticles (NPs) for the modulation of macrophage polarity from the pro-inflammatory M1 to anti-inflammatory M2 phenotype.

Methods: Characterization of the NPs was monitored using FTIR, DLS and FE-SEM. The effects of Chrysin-encapsulated PLGA-PEG NPs on the viability of LPS/IFN- γ stimulated peritoneal macrophages were determined using MTT assay. The cellular uptake of free Chrysin and nano-formulated Chrysin was measured using confocal microscopy. Also, the expression levels of M1 marker (iNOS and SOCS3) and M2 marker (Arg1 and Fizz) and also pro-inflammatory cytokines were determined by real time PCR

Results: Data showed that the nano-formulated curcumin with spherical shape, an average diameter of 102.5 nm and high cellular uptake was significantly less toxic to peritoneal macrophages. Furthermore, the nano formulated Chrysin effectively indicated a reduction in iNOS-2 and an increase in Arg-1 levels than free Chrysin. The change in macrophage phenotype by Chrysin-encapsulated PLGA-PEG NPs could suppress the inflammation in LPS/IFN- γ stimulated macrophages as verified by a major reduction in pro-inflammatory cytokines.

Conclusion: The results proposed that the Chrysin formulation with PLGA-PEG NPs might be a hopeful platform for the treatment of inflammatory diseases.

Keywords: Chrysin, PLGA-PEG, nanoparticles, macrophage polarity, inflammation

ICIA2018\Innate Immunity and Inflammation\Oral\8185

8185

Reciprocal role of hBD2 and hBD3 on the adaptive immune response by measuring T lymphocyte proliferation in terms of CD4 and CCR6 expression

Sina Taefehshokr^{1*}, Pouriya Dadebighlu², Arezoo Naderi³, Maryam Diansaei¹, Saeed Farhadzadeh¹, Farzin Javid¹, Ebrahim Taghizadeh², Nima Taefehshokr⁴

1. Young Researchers and Elite Club, Tabriz Branch, Islamic Azad University, Tabriz, Iran

2. Faculty of Veterinary Medicine, Tabriz Branch, Islamic Azad University, Tabriz, Iran

3. Faculty of Medicine, Tabriz Branch, Islamic Azad University, Tabriz, Iran

4. Division of Bioscience, Department of Life Sciences, College of Health and Life Sciences, Brunel University, UK

Background: Human β -defensins are small cationic antimicrobial peptides of innate immune system which can act as a barrier against the majority of pathogens, contributing to the host immune defense. The aim of study is to determine whether human β -defensins (hBD2 and hBD3) play a role in development and proliferation of human effector cluster of differentiation 4 (CD4) T cells or not. Furthermore, if enhanced proliferation is observed in the presence of hBD2 and hBD3, these data will demonstrate whether CC chemokine receptor 6 (CCR6) is required to be present for this activity to occur.

Methods: In this study, we examined the effect of hBD2 and hBD3 on CD4⁺ T cell proliferation in CCR6⁺ and CCR6⁻ T cells through co-culture of PBMCs (Peripheral blood mononuclear cells) with anti-CD3 and anti-CD28 stimulation in the presence or absence of hBD2 and hBD3. Proliferation was assessed using flow cytometry.

Results: It was demonstrated that, co-culture with hBD2 and hBD3 led to up-regulation of CD4⁺ T cell proliferation after 72 hours whereas, CD4⁺ T cell proliferation was suppressed after 96 hours. On the other hand, CCR6⁻ and CCR6⁺ T cell proliferation was up-regulated after 72 hours. But, CCR6⁺ only was down-regulated in the second cycle in the presence of hBD3. In contrast, after 96 hours CCR6⁺ and CCR6⁻ T cell proliferation was decreased.

Conclusion: Collectively, our data indicated that human β -defensins (hBD2 and hBD3) play a positive and negative regulatory role in development and proliferation of human effector CD4⁺ T cells which is essential for optimal adaptive immune responses and the control of immunopathology.

Keywords: hBD2, hBD3, T cell proliferation, CD4, CCR6

ICIA2018\Innate Immunity and Inflammation\Oral\8381

8381

Xanthine oxidase is elevated and associated with disease activity in Behçet's disease: study on ALGERIAN PATIENTS

Arezki Chekaoui¹, Houda Belguendouz¹, Karima Lahmar-Belguendouz¹, Djanette Hakem², Fatma Zohra Mazari³, Malika Terrahi³, Chafia Touil-Boukoffa¹

1. Cytokines & NO Synthases" team, LBCM-FSB-USTHB, Algiers, Algeria

2. Internal medicine department, CHU Bab El Oued, Algiers, Algeria

3. Ophtalmology department, CHU Nafissa Hammoud, Algiers, Algeria

Background: Behçet's disease (BD) is a multisystem disease. It stands at the crossroad between autoimmunity and autoinflammatory diseases. Xanthine Oxidase (XOD) is an enzyme that is involved in free radicals production. Several studies have demonstrated that free radicals trigger biological material oxidation leading to severe inflammatory reactions in BD. In the present study, we aimed to assess the XOD activity plasma levels and elucidate its possible association with BD activity in Algerian BD.

Methods: We investigated the Xanthine oxidase activity (XOD) and nitrosative stress marker (Nitric oxide (NO)) in Algerian Behçet's Disease patients (n=78): Active Behçet's disease patients (ABP, n=28) and Inactive Behçet's Disease patients (IBP, n=50) and healthy controls (HC, n=41). Mann-Whitney U test and Pearson correlation test were used for statistical analyses.

Results: In our study, we showed that ABP displayed significantly higher XOD activity levels ($p < 0.0001$, $p = 0.038$) compared to IBP and HC respectively. In contrast, XOD levels in IBP showed no statistical differences ($p = 0.45$) versus HC. Nitric oxide levels significantly increased in ABP ($p < 0.0001$) when compared to IBP and HC and in IBP ($p < 0.0001$) versus HC. XOD significantly and positively correlated with NO ($r = 0.4882$, $p = 0.0002$) in Behçet's disease patients.

Conclusion: We emphasize that Xanthine oxidase activity increased and correlated with the disease severity in BD patients. We suggest exploring further this enzyme (XOD) with the aim to serve as additional marker for assessing BD activity and being an eventual target for therapeutics.

Keywords: Behçet's disease, disease activity, Xanthine oxidase, chronic inflammatory disorder.

ICIA2018\Innate Immunity and Inflammation\Oral\8589

8589

Berberine attenuates high glucose-induced lipotoxicity and lipid droplet accumulation in RAW264.7, involvement of suppression of skewing RAW264.7 toward M1

Maryam Shabani^{*1}, Reyhaneh Babaei Khorzoughi¹

1. Department of Clinical Biochemistry, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

Background: Macrophages are flexible cells and able to play multiple functions in inflammatory conditions. These major components of innate immunity cells secrete various cytokines and a vast number of mediator molecules and CD markers. These processes are tightly regulated and these molecules have both beneficial and detrimental outcome in health. macrophages can acquire specialized phenotypes; M1 (pro-inflammatory phenotype) and M2 (anti-inflammatory phenotype). M1 subset secreted inflammatory mediators such as iNOS, TNF, MCP1, IL-6, and have CD11-c surface marker. In contrast, M2 markers are CD206, CD163, TGF β , Arginase. M1/M2 balance has a critical role in inflammation. Diabetes Mellitus is an inflammatory condition and polarization pattern of macrophage is going toward M1. Berberine is an active plant compound that is extracted from *CoptisChinensis*, which has strong anti-inflammatory properties. This study was conducted to evaluate the anti-inflammatory effect of Berberine.

Method: RAW 264.7 macrophages were stimulated by high glucose concentration, then we have studied effects of Berberine (10 μ M) on high glucose-induced lipotoxicity and macrophage polarization. We have measured high glucose-induced lipogenesis by oil-red O staining. For investigating macrophage polarization, we assessed M1 marker, CD11c, as an inflammatory factor via flowcytometry. Also, inflammatory genes expression include: iNOS, IL-6, and MCP1 were assayed by Real-time PCR.

Results: This study suggests that high glucose induce RAW264.7 lipogenesis. Flow cytometry analyses showed that high glucose skew RAW264.7 toward M1 subset, whereas Berberine attenuated M1 marker up to about 70%. RT-PCR confirmed the anti-inflammatory effect of Berberine through attenuation of inflammatory genes expression ($P < 0.05$).

Conclusion: Reduction of lipogenesis, M1 surface marker and above inflammatory genes approves that Berberine can be used as a therapeutic drug in inflammatory diseases.

Keywords: Inflammation, Berberine, High Glucose, Macrophage polarization

ICIA2018\Innate Immunity and Inflammation\Oral\9847
9847

Increased circulating Chitinase-3-like Protein-1 (YKL-40) concentrations after cardiopulmonary bypass in coronary surgery patients, the effect of Erythropoietin treatment

Mahnoosh Foroughi^{1*}, Zohre Mohammadi^{1,2}, Mostafa Haji molahoseini³, Mahmood Bakhtiyari², Ali Dabbagh⁴

1. Cardiovascular Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

2. Clinical Research and Development Center, Shahid Modarres Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran

3. Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

4. Anesthesiology Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Cardiopulmonary bypass (CPB) and the imposed global myocardial ischemia induce extensive inflammatory reaction. YKL-40 is a rather new biomarker in acute and chronic inflammation and cardiovascular disease. Among patients undergoing cardiac surgery, the serum level of YKL-40 has not been investigated yet. The discrepancy and gap about the cardio protective effects of erythropoietin (Epo) still exist. The aim of this study was to assess the effects of CPB and Epo treatment on serum YKL-40.

Methods: In this pilot trial, 97 patients were admitted for elective coronary surgery with CPB, and randomly allocated to receive one of three treatments: group A (n = 35, infusion of 20000 IU Epo after aortic clamp in 30-45 min), group B (n = 31, the same intervention after anesthesia and before CPB, control group (n = 31).

Results: In comparison between the groups, there was a sharp increase in serum YKL-40 with a 24 h delay after CPB in all groups without significant difference. The increase in serum IL-6 was significant in group B compared with both groups A and control (P=0.001 and P=0.001, respectively). Serum NT-pro BNP reached maximum level after 24 h in all groups and group B was significantly higher than group A (P=0.008). CK-MB was significantly increased in all groups (P<0.001), but this change was less prominent in group B compared to control (P=0.003).

Conclusion: The inflammatory response induced by CPB has a dramatic effect on YKL-40 level. Serum YKL-40 reached peak level one day after CPB. Epo treatment has no effect on postoperative serum YKL-40.

Keywords: Coronary artery bypass, Erythropoietin, Inflammation, IL-6, Reperfusion injury, YKL- 40.



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Monoclonal Antibody Diagnostic and Therapeutic

Oral Presentation

ICIA2018\Monoclonal Antibody Diagnostic and Therapeutic\Oral\7606
7606

Production of a Mouse Monoclonal Antibody against cell-surface antigen of new established breast cancer cell line

Farzaneh Ghaderi Shabankareh ^{1,2}, Abbas Ghaderi ^{1,2}

1. Department of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

2. Shiraz Institute for Cancer Research, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Breast cancer is the most common malignancy and the leading cause of cancer-related death among women worldwide. In Iran, breast cancer accounts for 24.6% of total cancers with the crude incidence rate of about 22.6 per 100,000 females, which even is increasing annually. Murine monoclonal antibodies (MoAbs), reactive with several human tumor-associated antigens, have been produced and characterized in many laboratories. The purpose of these studies has been to generate effective monoclonal antibodies that could be useful in tumor diagnosis and therapy.

Methods: Splenic lymphocytes of BALB/c mice immunized with a new established breast cancer cell line (Pari-ICR cell line, established in Shiraz Institute for Cancer Research) were fused with the mouse myeloma cell line, SP2/0 in the presence of polyethylene glycol, MW 1450. We generated a panel of monoclonal antibodies against the newly established cell line. The hybrid cultures were screened by flowcytometry for the production of antibodies.

Results: Hybridomas that produced antibodies to surface antigens of the immunizing cell line and other breast carcinoma cell lines, but not to human fibroblast cells, mesenchymal stem cells and leucocytes isolated from peripheral blood, were selected and cloned by limiting dilution method. One clone designated 1H3, with IgG2a isotype was further analyzed for specificity by flowcytometry. Monoclonal antibody was affinity purified from mouse ascites fluid for further analysis. The target of monoclonal antibody had a molecular weight of approximately 150 kDa as determined by Western blotting. The functional characteristics and the specific target of the produced monoclonal antibody on immunizing cell line are underway.

Keywords: monoclonal antibody, breast cancer cell line, hybridoma.

ICIA2018\Monoclonal Antibody Diagnostic and Therapeutic\Oral\7620

7620

Generating a Novel Monoclonal Antibody against CD73 as a Potential Diagnostic Marker in Human Carcinomas

Kimia Zandehi¹, Saeideh Milani², Mehrdad Hashemi¹, Javad Mohammadi^{2, 3}, Fatemeh Ghaemimanesh², Ali Ahmad Bayat², Reza Hadavi², Hodjatallah Rabbani²

1. Department of Genetics, Islamic Azad University, Tehran Medical Sciences Branch, Tehran, Iran

2. Antigen and Antibody Engineering Department, Monoclonal Antibody Research Center, Avicenna Research Institute, ACECR, Tehran, Iran

3. Department of Life Science, Faculty of New Science and Technology, University of Tehran, Tehran, Iran

Background: CD73 is an enzyme which catalyzes the hydrolysis of AMP to adenosine. It also has a key regulatory role in cancer cells proliferation, migration and invasion. Over expression of CD73 has been found in many type of cancer. All of these characteristics led scientists to consider CD73 as a favorable marker for diagnosis and assessments of cancer. The aim of this study was to produce a novel and effective monoclonal antibody against CD73 as a diagnostic tool in human epithelial cancer.

Methods: A synthetic peptide derived from CD73 protein was conjugated to Keyhole limpet hemocyanin (KLH) and used for immunization of BALB/c mice. The generated antibody was purified from the supernatant of final hybridoma clone using peptide-affinity chromatography column and its ability to recognize the immunizing peptide was measured by ELISA. The reactivity of the antibody with CD73 protein was then evaluated by Western blot, immunocytochemistry and flowcytometry using different epithelial cancer cell lysates.

Results: Western blot, immunocytochemistry and flowcytometry analysis using produced anti-CD73 monoclonal antibody showed a specific recognition of this protein in different carcinoma cell lines including MCF-7, SW480 and A431.

Conclusion: Our generated mAb has potential to be utilized in three application including W.B., IHC and FACS for detection and assessment of CD73 expression in human carcinomas.

Keywords: Monoclonal Antibody, CD73, cancer, detection

ICIA2018\Monoclonal Antibody Diagnostic and Therapeutic\Oral\8019
 8019

Introduction of the First Antibody Drug Conjugate (ADC) based on anti-Placenta-specific1 (PLAC1) Antibody as a Novel Immunotherapeutic Tool for Prostate Cancer

Mohammad Reza Nejadmoghaddam^{1,2}, Rassoul Dinarvand^{2,3}, Ramin Ghahremanzadeh¹, Roya Ghods^{4,5}, Jafar Mahmoudian², MaryamYousefi¹, Mahboobeh Nazari⁶, Mohammad Hossein Ghahremani², Maryam Abolhasani⁷, Ali Anissian⁸, Morteza Mahmoudi², Amir-Hassan Zarnani^{9,10}

1. Nanobiotechnology Research Center, Avicenna Research Institute, ACECR, Tehran, Iran.
2. Nanotechnology Research Center, Faculty of Pharmacy, Tehran University of Medical Sciences (TUMS), Tehran, Iran
3. Department of Pharmaceutics, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran
4. Oncopathology Research Center, Iran University of Medical Sciences, IUMS, Tehran, Iran
5. Department of Molecular Medicine, Faculty of Advanced Technologies in Medicine, Iran University of Medical Sciences, IUMS, Tehran, Iran.
6. Monoclonal Antibody Research Center, Avicenna Research Institute, ACECR, Tehran, Iran.
7. Department of Pathology, Hasheminejad Kidney Center, Iran University of Medical Sciences, IUMS, Tehran, Iran
8. Veterinary department, Islamic Azad University, Abhar branch, Abhar, Iran
9. Reproductive Immunology Research Center, Avicenna Research Institute, ACECR, Tehran, Iran
10. Immunology Research Center (IRC), Iran University of Medical Sciences, IUMS, Tehran, Iran

Background: Our group has recently reported differential expression of PLAC1 in prostate cancer suggesting the potential usefulness of PLAC1 for targeted immunotherapy of prostate cancer. We hypothesized that taking advantages of cancer cell specificity of anti-PLAC1 antibodies and cytotoxic activity of toxic agents could be a platform for generation of an antibody drug conjugate (ADC) that selectively target prostate tumor cells.

Methods: The expression profile of PLAC1 in prostate cancer cells, LNCaP, PC3 and DU145, was assessed by PCR, Western blotting, and flow cytometry. To prepare ADC, SN-38, a potent anti-topoisomerase I derivative of camptothecin, was first aminated and characterized by FT-IR and NMR spectroscopies. Modified SN-38 was conjugated to carbohydrate residues of a monoclonal anti-PLAC1 antibody through amine linkage and drug-antibody ratio (DAR) was measured by HPLC. Binding properties, affinity and internalization capacity of ADC were analyzed by ELISA, flow cytometry and immunofluorescent staining, respectively. Cytotoxicity of SN38-anti-PLAC1 on prostate cancer cells was studied by fluorimetry and single cell gel electrophoresis. Cytotoxic ex vivo effect of anti-PLAC1-ADC was further assessed by Annexin V apoptosis assay in human primary prostate cancer cells. Off target effects of the anti-PLAC1-ADC was also tested in vivo in mice.

Results: Prostate cancer cells expressed PLAC1 at both gene and protein levels. Anti-PLAC1 binding to the cells induced rapid internalization of the antibody within few minutes which completed within an hour. After SN38 conjugation to antibody, a DAR of about 6 was achieved. Drug conjugation did not negatively affect affinity and binding property of anti-PLAC1 antibody to cell surface antigen. The ADC retained intrinsic antibody activity and showed enhanced and selective cytotoxicity with an IC₅₀ about 15 fold lower compared to free drug. Anti-PLAC1 ADC induced selective apoptosis in human primary prostate cancer cells and cancer cell lines, while exerted no off target effects as judged by histopathology examination of different organs of the injected mice.

Conclusion: Our results showed for the first time the potential application of anti-PLAC1-based ADCs as novel immunotherapeutic modality for patients with prostate cancer.

Keywords: PLAC1, Prostate cancer, Antibody-drug conjugate (ADC), Immunotherapy

ICIA2018\Monoclonal Antibody Diagnostic and Therapeutic\Oral\8086

8086

Designing, cloning, expression and purification of the bispecific antibody which can retarget oncolytic Newcastle Disease virus to carcinoma

Monireh Kazemimanesh¹, Frank Momburg², Mostafa Jarahian³,Vahid Baniasadi⁴, Kayhan Azadmanesh¹

1. Department of Virology, Pasteur Institute of Iran, Tehran, Iran

2. Clinical Cooperation Unit, Applied Tumor Immunity, Antigen Presentation and T/NK Cell Activation Group, German Cancer Research Center(DKFZ), Heidelberg, Germany

3. Department of Toxicology and Chemotherapy, German Cancer Research Center (DKFZ), Heidelberg, Germany.

4. Department of Arboviruses and Viral Hemorrhagic Fevers, Pasteur Institute of Iran, Tehran, Iran

Background: Carcinoma is a type of cancer that develops from epithelial cells. Oncolytic viruses such as Newcastle Disease virus (NDV) are a promising class of drugs that preferentially replicate in cancer cells as well as finally kill the malignant cells. For retargeting NDV to tumor tissue, bispecific antibody (biAb) can be used to bind NDV to tumor cells. Our biAb binds with one arm to the hemagglutinin-neuraminidase (HN) molecule of NDV and with the other arm to the epithelial cell adhesion molecule (EpCAM) which is overexpressed on carcinoma cells.

Methods: The plasmid which encodes the biAb was designed and constructed. Suitable competent E. coli DH5α cells were transformed using the CaCl₂ standard method. A single bacterial colony from a selective plate was used for DNA preparation. Concentration of extracted plasmid was measured by spectrophotometry and confirmed by double digestion with restriction enzymes. This plasmid was stably transfected in HEK 293 cell using lipofectamin LTX-PLUS. The stable cell line was cultured and after 72 hours the medium (supernatant) was collected. For investigating expression of biAb, SDS-PAGE and Western blotting of medium and filtered medium (concentrated using Amicon® Ultra-15 centrifugal filter devices) as well as flow cytometry analysis were performed. Purification of biAb was performed using rProtein A Gravi Trap (GE Healthcare Life Sciences). The ability of purified biAb to attach to HN of different strains of NDV and EpCAM expressed by carcinoma cells was investigated by flow cytometry.

Results: The cloned plasmid which encodes the biAb was confirmed by digestion with restriction enzymes. Expression of biAb was confirmed by flow cytometry analysis. These results show that biAb could attach to HN of different strains of NDV and EpCAM.

Conclusion: This biAb binding to NDV-HN and the pan-epithelial antigen EpCAM, may be used to retarget oncolytic NDV to carcinomas.

Keywords: Bispecific antibody, Carcinoma, Newcastle Disease virus.

ICIA2018\Monoclonal Antibody Diagnostic and Therapeutic\Oral\8184
8184

Contribution of Fc fragment of monoclonal antibodies in tetanus toxin neutralization

Somayeh Ghotloo¹, Mohammad Mehdi Amiri¹, Jalal Khoshnoodi¹, Mahmood Jeddi-Tehrani²,
Forough Golsaz-Shirazi¹, Fazel Shokri^{1,2}

1. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran.

2. Monoclonal Antibody Research Center, Avicenna Research Institute, ACECR, Tehran, Iran

Background: Monoclonal antibodies (MAbs) against neurotoxin of *Clostridium tetani* are considered as a novel source of immunoglobulins for passive immunotherapy of tetanus. Fab and F(ab')₂ may be more effective in neutralizing neurotoxin than intact antibodies due to their faster and higher tissue distribution. In addition, removing Fc portion from the mouse antibodies reduces HAMA response in human. We generated Fab and F(ab')₂ of three mouse MAbs, which previously showed neutralizing activities either individually or in combination and compared their neutralizing activities with those of their intact antibodies.

Methods: Fab and F(ab')₂ fragments were generated by papain and pepsin digestion, respectively. Immunoreactivity of the fragments was compared with their intact forms by ELISA. Toxin neutralizing activity of the antibodies was evaluated in an in vivo toxin neutralization assay in which various amounts of MAbs or their fragments were mixed with 20MLD of the toxin and injected intraperitoneally to BALB/c mice. Human polyclonal immunoglobulins (tetabulin) and their fragments were used as the control.

Results: Enzymatic digestion of MAbs did not affect reactivity of the antigen binding sites. While intact MAbs were able to fully protect the mice even at the lowest dose, none of the mice, which received F(ab')₂ or Fab fragments, survived longer than 14 days even at the highest administered dose. All mice receiving tetabulin or their fragments were fully protected.

Conclusion: Loss of protective activity of MAbs fragments is not due to the attribution of Fc region in toxin neutralization, since the mice receiving tetabulin fragments were fully protected. It may be explained by the removal of steric hindrance of Fc region on the receptor binding site of the toxin. It seems that the polyclonal pool of tetabulin contains some molecules with variable regions recognizing receptor binding site of the toxin leading to toxin neutralization.

Keywords: Tetanus toxin, Monoclonal antibody, Toxin neutralization, Fab, F(ab')₂

ICIA2018\Monoclonal Antibody Diagnostic and Therapeutic\Oral\8211

8211

Development and Characterization of a Camelid Single Domain Antibody Directed to Human CD22 Biomarker

Fatemeh Faraji¹, Nader Tajik², Mahdi Behdani³, Mohammad Ali Shokrgozar⁴, Amir Hassan Zar nani², Fatemeh Shahhosseini⁴, Mahdi Habibi-Anbouhi⁴

1. Department of Immunology, School of Medicine, Iran University of Medical Sciences, Tehran, Iran

2. Immunology Research Center (IRC), Iran University of Medical Sciences, Tehran, Iran

3. Biotechnology Research Center, Venom & Biotherapeutics Molecules Laboratory, Pasteur Institute of Iran, Tehran, Iran

4. National Cell Bank of Iran, Pasteur Institute of Iran, Tehran, Iran

Background: CD22 is a B-cell-specific trans-membrane glycoprotein which is found on the surface of the most B-cells and modulates their function, survival, and apoptosis. Recently, targeting this cell surface biomarker in B-cell malignancies and disorders has attracted a lot of attention. The variable domain of camelid single chain antibodies (VHH, Nanobody) is a form of antibodies with novel properties including small size (15-17 kDa), thermal and chemical stability, high affinity and homology to human antibody sequences.

Methods: In this study, a novel anti-CD22 specific VHH (Nb) has been developed and characterized by the screening of an immunized phage display library and its binding to CD22⁺ B-cells evaluated.

Results: Produced anti-CD22 VHH had a single protein band about 17 kDa of molecular size in western blotting and its binding affinity was approximately 9×10^{-9} M. Also, this product had high specificity and it was able to recognize the natural CD22 antigen in CD22⁺ cell lysate as well as on the cell surface (93%).

Conclusion: This anti-CD22 VHH with both high affinity and specificity recognizes CD22 antigen well and can be used in diagnosis and treatment of B cell disorders and malignancies.

Keywords: CD22, Camelid Single Domain Antibody, VHH

ICIA2018\Monoclonal Antibody Diagnostic and Therapeutic\Oral\8215
8215

A novel method for purification of hyaluronidase from *Streptococcus pyogenes* in one step

Amir Shakouri¹, Jalal Abdolalizadeh²

1. MSc of Biotechnology, Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

2. PhD of Clinical Biochemistry, Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Hyaluronidase contains some classes of catalysts that cleave Hyaluronic Acid that exist in mammalian spermatozoa, venom of scorpions and snakes, Leeches, Bacteria and etc, which can digest hyaluronic acid and is also known as a developer factor. It open the way for other active ingredients found in the saliva of leeches to penetrate into deep tissue. The aim of this study is the purification of Bacterial hyaluronidase with affinity chromatography which packed with Anti- Hyaluronidase proteins that produced in rabbit.

Methods: The rabbit polyclonal antibody was prepared against the hyaluronidase enzyme. Thus the pure enzyme was injected to rabbit and to evaluation of antibody production, ELISA and double Diffusion tests were done. Antibody purification was done by ion exchange chromatography and protein G affinity chromatography. Then the purified antibody was attached to the CNBr-activated Sepharose and finally it used for purification of hyaluronidase from Bacteria. After the purification, purity of fractions was assessed by SDS-PAGE electrophoresis and proved within Western- Blot.

Results: SDS-PAGE analysis showed the purity of protein was up to 98%. Also the single band with a molecular weight of approximately 50 KD is related to *Streptococcus Pyogenes* hyaluronidase. Western- Blot analysis demonstrated that single band was related to hyaluronidase.

Conclusion: Affinity chromatography using produced antibody would be an economical and safe method for purification of hyaluronidase from *Streptococcus Pyogenes*.

Keywords: Hyaluronidase, Immunoaffinity Chromatography, *Streptococcus Pyogenes*, Polyclonal antibody

ICIA2018\Monoclonal Antibody Diagnostic and Therapeutic\Oral\8295

8295

Phage display-derived antibody fragments against conserved regions of VacA toxin of *Helicobacter pylori*

FarnazFahimi^{1,2}, ShamimSarhaddi², Mehdi Fouladi^{1,2}, NaserSamadi², JavidSadeghi⁴, Asal Golchin ¹,
MohammadrezaTohidkia^{1*}, JalehBarar^{1,3}, YadollahOmid^{1,2,3}

1. Research Center for Pharmaceutical Nanotechnology, Biomedicine Institute, Tabriz University of Medical Sciences, Tabriz, Iran

2. School of Advanced Biomedical Sciences, Tabriz University of Medical Sciences, Tabriz, Iran

3. Department of Pharmaceutics, Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran

4. Department of Microbiology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Infection with *helicobacter pylori*, the only carcinogen bacterium, is attributed to gastroduodenal disorders and gastric adenocarcinoma. Due to inaccuracy of common diagnostic methods new detection approaches based on monoclonal antibodies are in priority. Among various toxins assisting pathogenesis of *H. pylori*, VacA is one of the most potential toxins, which is known as the major cause of peptic ulcer in patients.

Methods: Solution phase biopanning and screening were performed in order to isolate scFvs against two conserved regions of VacA. Characterization of scFvs was carried out by ELISA, immunoblotting and SPR. Bioinformatic analysis also performed in order to characterize structural and functional properties of isolated scFvs and antibody-antigene interactions.

Results: After four rounds of panning and screening, positive colonies in scFv ELISA were harvested for plasmid extraction and sequencing. Among those of which, high diversity of the VacA1 and diversity of 2 colonies for VacA2 were determined. Large scale expression and purification were performed for further complementary tests such as SDS-PAGE, western blot and SPR in order to determine affinity and specificity. Docking results done by bioinformatics analysis determined the interaction of the CDRs with the VacA peptide.

Conclusion: In this study, for the first time recombinant antibody fragments were isolated against conserved residues of VacA toxin with high specificity and affinity. New diagnostic approaches based on monoclonal antibody production and combination with drug delivery methods would minimize misinterpretation of the clinical laboratory tests.

Keywords: Antibody, Phage antibody display, scFvs, *Helicobacter pylori*, VacA toxin



Mucosal and Gastrointestinal Immunology

Oral Presentation

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Mucosal and Gastrointestinal Immunology\Oral\5257

5257

Diverse profiles of TLR2, 4, 7 and 9 mRNA in Peripheral blood and biopsy specimens of patients with Celiac Disease

Haniye Ghasiyari^{1,2}, Mohammad Rostami Nejad^{3*}, Davar Amani^{2*}, Kamran Rostami⁴,
Mohammad Amin Pourhosseingholi³, Mohammad Reza Zali³

1. *Basic and Molecular Epidemiology of Gastrointestinal Disorders Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran*

2. *Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran*

3. *Gastroenterology and Liver Diseases Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran*

4. *Department of Gastroenterology Milton Keynes University Hospital, Milton Keynes, UK*

Background: Celiac disease (CD) is an organ-specific autoimmune disease, and both adaptive and innate immunity are involved in its development. The aim of this study was to investigate the TLRs 2,4,7,9 genes expression in both peripheral blood and intestinal mucosa of patients with celiac disease compare to healthy control(HC).

Methods: Blood samples from 120 confirmed CD patients who had been on a gluten-free diet (GFD) at least for 1 year and 120 age- and sex-matched healthy volunteers served as control group were collected during 2016. Also 20 biopsy specimens from persons who had a clinical manifestation despite GFD were randomly taken. Total RNA was isolated using a standard commercial kit. The mRNA expression of TLRs were quantified by relative qPCR with β 2 microglobulin (β 2m) as a reference gene.

Results: TLR4 and TLR9 mRNA were significantly higher expressed in blood samples from GFD-CD compared to the healthy controls ($P=0.02$); but not for TLR2. Furthermore, mRNA expression of TLR2 ($P=0.03$) and TLR4 ($P=0.0003$) expression level was increased in CD biopsy specimens compared to controls, whereas expression of TLR9 mRNA was significantly decreased in CD patients. There was no significant differences expression of TLR7 in biopsy and blood specimens.

Conclusions: It seem that upregulation of TLR4 and TLR9 suggests the contribution of gut microbiota or dysregulation of the immune response to commensal flora in small bowel mucosa in celiac patients.

Keywords: celiac disease, Toll like receptor, innate immunity

ICIA2018\Mucosal and Gastrointestinal Immunology\Oral\5306

5306

Generation of recombinant cytokine secreting strains of *Lactococcus Lactis* to enhance mucosal innate immunity

Sadegh Feizollahzadeh¹, Hossein Khanahmad³, Ilnaz Rahimmanesh³, Mazdak Ganjalikhani-hakemi², Alireza Andalib², MohammadHossein Sanei⁴, Abbas Rezaei²

1. Department of Medical Laboratory Sciences, Paramedical faculty, Urmia University of Medical Sciences, Urmia, Iran

2. Department of Immunology, Faculty of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

3. Department of Molecular Biology and Genetics, Faculty of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

4. Acquired Immunodeficiency Research Center, Isfahan University of Medical Sciences, Isfahan, Iran

Background: Cytokines are secretory mediators of immune system that play a crucial role in innate immunity. Interleukin-18 (IL-18) mainly identified as interferon- γ inducing factor. This cytokine is a pleiotropic cytokine secreted by different types of immune and nonimmune cells. It is involved in defense against viral, intracellular bacterial and protozoan infections and intestinal homeostasis. *Lactococcus lactis* (L. lactis), is a safe bacterium which increasingly used for heterologous protein expression in therapeutic and industrial applications. In the present study, we engineered L. lactis to express murine mature IL-18 in order to use the new straining the enhancement of intestinal innate immunity.

Methods: For a generation of such strain, coding DNA sequence of mature murine IL-18 was optimized for codon usage of L. lactis. It cloned downstream of Nisin system in PNZ8149 plasmid and electroporated to L. lactis NZ3900. Positive clones were selected based on yellow colony formation on Elliker medium and confirmed by polymerase chain reaction (PCR) and restriction digestion. IL-18 production was confirmed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), western blotting and ELISA in lysates and medium.

Results: Transformed bacteria formed yellow colonies on Elliker. PCR with IL-18 specific primers on extracted plasmids were positive. SDS-PAGE on bacterial lysate showed a thick sharp band around 18kD protein band. The identity of the protein band was then verified using IL-18 primary antibodies in western blot test. IL-18 level in bacterial lysate was 3–4 mg/L as detected by specific ELISA kit. Secreted IL-18 in medium was about 6–7 μ g/L.

Conclusion: The successfully generated L. lactis strain that expressed active murine IL-18 can be used to evaluate the possible therapeutic effects of IL-18 on mucosal surfaces.

Keywords: Cytokine, Innate immunity, Interleukin-18, *Lactococcus lactis*

ICIA2018\Mucosal and Gastrointestinal Immunology\Oral\7823
7823

Normal total duodenal intraepithelial lymphocytes and CD3⁺, and CD8⁺ intraepithelial T cells in celiac patients: first report on cell density counts of the intestinal intraepithelial lymphocytes in Isfahan population, Iran

Hadi Hossein-Nataj¹, Mohsen Masjedi², Sayed Mohammad-Hassan Emami Najaffi Dehkordi², Mojgun Mokhtari², Fereshteh Alsahebhosoul²

1. Mazandaran University of Medical Sciences

2. Isfahan University of Medical Sciences

Background: Increased numbers of the duodenal intraepithelial lymphocytes (IEL) are one of the key histological findings in celiac disease (CD). However, this finding may vary from one region to another. There are cases of borderline histology and seronegative CD which might be different from other causes of the duodenal immunological reactions by measuring the mean percentages of iIEL expressing the α/β TCR, γ/δ TCR, CD3, CD4, and CD8 markers. This study aimed to measure the total CD3⁺, CD8⁺ IEL T cells in celiac patients and healthy controls in Isfahan, Iran.

Methods: D2 biopsies from definite CD cases (17 cases) and healthy controls (22 cases) were evaluated. H&E staining and several monoclonal antibodies were used in this study by the immunoperoxidase staining method.

Results: Immunoperoxidase staining method showed that the mean total IEL was 19 and 40 in the control subjects and CD cases, respectively ($P \leq 0.05$). The upper normal limit of the CD3⁺ intraepithelial T cells (mean+2SD of IEL T cells counts in the controls) was 20, while it was 14% for the CD8⁺ T cells. In addition, the H&E staining method showed the cut-off of 34% for IEL in CD.

Conclusion: This study suggests that the total IELs are of > 34%, CD3⁺ IEL > 20% and CD8⁺ IELs > 14% by the immunoperoxidase staining technique. These might be considered as Marsh I among the general population of Isfahan. Since immunoperoxidase staining of IEL by monoclonal antibody has an additional value in CD diagnosis and should be done for interpreting when there is disagreement between the serology and H&E staining methods.

Keywords: Celiac disease, duodenal biopsy, intestinal intraepithelial lymphocytes, normal range.

ICIA2018\Mucosal and Gastrointestinal Immunology\Oral\8254
8254

The up-regulation of NRF-2 and IL-11 in mild Ulcerative colitis

Milad sabzevary-ghahfarokhi¹, Hedayatollah Shirzad²

1. Student Research Committee, Shahrekord University of Medical Sciences, Shahrekord, Iran

2. Cellular and Molecular Research Center, Basic Health Sciences Institute, Shahrekord University of Medical Sciences, Shahrekord, Iran

Background: Ulcerative colitis is an inflammatory disorder of the colon in which the balance of the intestinal immune system is disturbed. Oxidative stress plays crucial roles in the pathogenesis of UC. Our aim was to measure protein level of IL-11, and characterized the expression pattern of the NRF-2 protein in histological tissues.

Methods: 20 patients with mild ulcerative colitis and 21 healthy subjects were enrolled in the study. The IL-11 level in tissues was determined by means of ELISA. NRF-2 protein and mRNA in mucosal tissue were also performed by immunohistochemistry and real-time PCR.

Results: The increased level of IL-11 protein and NRF-2 mRNA were shown in the mild patient compared to control subjects. Furthermore, NRF-2 protein dominantly appears in mucosal epithelium.

Conclusion: the existence of oxidative stress in UC patients induces NRF-2 expression to resist against damages. The NRF-2 factor is a key regulator transcription factor of IL-11 production. Our data indicated that NRF-2 increased in patients leading to the protection of mucosal tissue. It is suggested that IL-11 act as a regenerator factor which may be a point of research in the novel therapeutic approaches for IBD.

Keywords: Oxidative stress; NRF-2; IL-11.

ICIA2018\Mucosal and Gastrointestinal Immunology\Oral\8349
8349

Interleukin-6 gene polymorphism -174 G/C, -572G/C association with susceptibility to Iranian celiac disease population

Zeinab Barartabar¹, Abdolrahim Nikzami¹, Majid Sirati Sabet¹, Vahid Chaleshi³, Elham Aghamohammadi², Mohammad Reza Zali³, Mohammad Rostami Nejad³

1. Biochemistry Department, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Basic and Molecular Epidemiology of Gastrointestinal Disorders Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran

3. Gastroenterology and Liver Diseases Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: polymorphisms of -572G and -174G in the interleukin 6 promoter can effect on the production and secretion of interleukin 6 which may play a role in the inflammation and pathogenesis of celiac disease. The aim of this study was to investigate the relationship between -572G and -174 G polymorphisms with susceptibility to celiac disease in Iranian population.

Methods: A total of 105 patients with celiac disease and 106 healthy individuals were recruited in this study during 2016. DNA of both groups were extracted by available kits and gene polymorphism studied by polymerase chain reaction-restriction fragment length polymorphism method (PCR-RFLP). Also serum levels of Interleukin 6 were evaluated by ELISA method in both groups.

Results: The results showed a significant relationship between polymorphism -572Gin CD patients compared with control subjects in genotype ($p=0.001$) and alleles ($p=0.022$), respectively. But there was no significant relationship between polymorphism -174G and frequency of genotype, but the association of this polymorphism with the frequency of allele ($p=0.034$), age ($p=0.001$) and BMI ($p=0.003$) was seen. On the other hand serum level of interleukin 6 was significantly associated only with rs1800796 ($p=0.000$).

Conclusions: Our results show that (572- rs1800796) polymorphism may play a role in susceptibility to CD in the Iranians patients.

Keywords: Celiac disease, PCR-RFLP, IL-6 gene polymorphism, autoimmune and inflammatory reaction

ICIA2018\Mucosal and Gastrointestinal Immunology\Oral\8615

8615

In vitro activation of Dendritic cells by a novel peptide as gastric cancer antigen-presenting cells

Kaveh Baghaei¹, Ali Asadirad², Bahareh Mohebi, Samaneh Beigi¹, Fatemeh Yarian, Seyed Mahmoud Hashemi², Maryam Nourizadeh³, Hamid Asadzadeh Aghdai¹, Mohammad reza Zali⁴

1. Basic and Molecular Epidemiology of Gastrointestinal Disorder Research Center, Research Institute for Gastroenterology and Liver Disease, Shahid Beheshti University of Medical Science, Tehran, Iran

2. Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

3. Immunology, Asthma and Allergy Research Institute, Children's Medical Center Hospital, Tehran University of Medical Sciences, Tehran, Iran.

4. Gastroenterology and Liver Disease Research Center, Research Institute for Gastroenterology and Liver Disease, Shahid Beheshti University of Medical Science, Tehran, Iran

Background: Dendritic cells (DC) play an important role in forming the quantity and quality of antitumor immunity. Therefore, DC-based peptide delivery of tumor antigen is becoming a potential approach in cancer immunotherapy. Effective antitumor immunity requires the generation and persistence of functional tumor-specific T-cell responses. The use of peptide-based vaccination has been found to be a promising means to induce antitumor T-cell responses. In current study we showed role of novel gastric cancer drive peptide which present by DCs and activate lymphocytes against tumor cells.

Methods: In this approach, autologous DC were generated from their precursors in bone marrow by using conventional cytokine and loaded with specific gastric cancer tumor antigen and then co-cultured with Lymphocyte cells. Immature and mature DCs were confirmed by Flow cytometry. Lymphocyte proliferation was measured by BRDU.

Results: Mature dendritic cells were successfully obtained from bone marrow mononuclear cells. The level of CD markers (CD40, CD80, CD83, CD86) were analyzed by flow cytometry. We could show, designed peptide activated DCs to proliferate lymphocytes.

Conclusion: Our DC-based peptide vaccination can act as an antigen delivery vehicle, resulting in measurable antitumor immunity in cancer settings in preclinical and clinical after in vivo studies.

Keywords: Dendritic cell, Gastric Cancer, Peptide vaccine

ICIA2018\Mucosal and Gastrointestinal Immunology\Oral\8711
8711

Investigation of immunogenic effects of *Syphacia obvelata* in treatment of experimental colitis mouse model

Niloofar Taghipour^{1,2}, Nariman Mosaffa³, Hamid Asadzadeh Aghdaei⁴, Mohammad Rostami Nejad¹

1. Gastroenterology and Liver Diseases Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

2. Department of Biotechnology, School of Advanced Technologies in Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran;

3. Department of Immunology, Faculty of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

4. Basic and Molecular Epidemiology of Gastrointestinal Disorders Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Many evidences have been suggested a reverse relationship between helminths infections and allergic or autoimmune diseases. In inflammatory bowel disease, excessive Th1 and Th17 cell responses to commensal germs or intrinsic antigens cause progressive inflammation in the tissues. The ability of helminth parasite infections to manipulate the immune system of their host has been proposed to suppress the inflammatory response.

Methods: male C57BL / 6 mice were divided into 5 groups: control group, colitis group, Receiving *S. obvelata* group, Preventive group, Curative group. After creating the experimental model, one group fed the *S. obvelata* eggs and then infected with colitis, and the other group was vice versa. A disease activity index score can be assessed to evaluate the clinical progression of colitis. TNF- α , IL-10, IL-17, IFN- γ were evaluated by ELISA in the culture supernatants of MLN and PP cells. The percentage and expression Treg in different study groups were investigated using Real-time PCR and flow cytometry.

Results: Treg cells in the colitis group had a significantly decreased in both sites compared to other groups. Significant increase in the population of Foxp3 + cells in MLN and PP was observed in the groups receiving *S. obvelata* compared to the control and the DSS group. Improvement of symptoms and complications of induced colitis, improvement of histologic inflammation, increased Treg response (IL-10 cytokines) and decreased inflammatory cytokines in both site of mice in the preventive and curative groups were significantly better than the colitis group.

Conclusion: Our results showed that *S. obvelata*, in addition to its role in prevention, has curative potential to reduce inflammatory colitis in mouse models. This is probably due to immunomodulatory effects of *S. obvelata* the production of anti-inflammatory cytokines and inhibition of inflammatory cytokines.

Keywords: Helminth, Colitis, *S. obvelata*, Treg, Foxp3

ICIA2018\Mucosal and Gastrointestinal Immunology\Oral\9906
9906

Evaluation of the expression level of BANCR Long none coding RNA in the Stool samples of patients with colorectal cancer, advance adenoma and healthy individual

Shadi Sadri¹, Mitra Salehi², Ehsan Nazemol hosseini Mojarad³

1. Department of cellular and molecular, North Tehran Branch, Islamic Azad University, Tehran, Iran

2. Department of microbiology, North Tehran Branch, Islamic Azad University, Tehran, Iran

3. Gastroenterology and Liver Diseases Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Colorectal cancer is mostly diagnosed in the late stages with poor prognosis. CRC is mostly derived from adenoma polyps and using of biomarkers such as long non coding RNA, is recommended for early detection. Long none coding RNA (lncRNAs) are key participants of gene regulations in disease and healthy situation. In this study we evaluation of the expression level of BANCR Long none coding RNA in the Stool samples of patients with colorectal cancer, advance adenoma and healthy individual.

Methods: In this study, 20stool samples of CRC patients , 20stool samples of adenoma polyp and 20 stool samples from healthy individual were collected. The clinical information was collected by a questionnaire and clinical reports, mRNA was extracted and cDNA was synthesized. BANCR gene expression was investigated by Real time PCR method and relative quantification. Fold change of genes expression were evaluated by ($2^{-\Delta\Delta Ct}$) method. Data were analyzed by 7500 system SDS version 1.3 and Prism version 5 software.

Results: The data show that BANCR was significantly over expressed in stool samples of CRC patients comparing to patients with adenoma polyp and healthy individual. However, these changes were not linked with tumor pathological characterizations ($p < 0.05$).

Conclusion: It seems that, increasing in BANCR expression is highly linked with tumor development and it may be a reason for malignancy progressing.

Keywords: BANCR Lnc RNA, Colorectal cancer, early diagnosis

ICIA2018\Mucosal and Gastrointestinal Immunology\Oral\9926
9926

Comparing plasma level of miR-106a to Differentiates Crohn's disease from Ulcerative Colitis

Vahid Chaleshi¹, Mohammad Hossein Ghorbani, Nazanin Amiri, Shabnam Shahrokh, Hamid Asadzadeh Aghdai, Mohammad Reza Zali

1. Basic and Molecular Epidemiology of Gastrointestinal Disorders Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Inflammatory bowel disease (IBD) comprising Crohn's disease (CD) and ulcerative colitis (UC). Finding new biomarker for detection and monitoring of IBD is importance in diagnosis, prognosis and treatment. miRNAs are a class of non-coding RNAs which regulate gene expression. To date, the majority reports have identified dysregulated miRNAs in IBD. The aim of this study was to evaluate gene expression of plasma miR-106a between ulcerative colitis and Crohn's disease patients and healthy control.

Methods: A total of 73 patients with IBD (55 UC, 18CD), and 15 healthy controls were enrolled in the study. Colonoscopy in combination with histological finding confirm the IBD diagnosis. Healthy controls showed no IBD symptoms or any gastrointestinal diseases such as celiac disease, polyp and cancer. miRNA isolated from plasma samples using miRNeasy mini Kit (QIAGEN). cDNA was synthesized from miRNA using miScriptII RT Kit (QIAGEN). miR-106a gene expression examined by q-Real Time PCR using miScript SYBR Green PCR Kit (QIAGEN). All kits were used according to manufacturer's instructions.

Results: miR-106a expression was significantly increased in patients with Crohn's disease in proportion of UC patients (P value: 0.0003) and healthy control group (P value: 0.0143). No significant differences were observed between UC patients and control group (P value: 0.7319).

Conclusion: According to our results expression of miR-106a between CD and UC maybe used as a new non-invasive factor for diagnosis and differentiation between IBD type.

Keywords: Ulcerative colitis (UC), Crohn's disease (CD), Inflammatory bowel disease (IBD), miR-106a

ICIA2018\Mucosal and Gastrointestinal Immunology\Oral\9957

9957

CD8 T cells are more frequent than CD4 T cells in patients suffering from chronic rhino sinusitis with polyp

Farhad Seif^{1,2}, Majid Khoshmirsafa¹, Hossein Aazami¹, Monireh Mohsenzadegan³, Babak Ghalehbaghi⁴, Mehdi Shekarabi^{1,2}

1. Immunology Department, School of Medicine, Iran University of Medical Sciences, Tehran, Iran

2. Immunology Research Center, Institute of Immunology and Infectious Diseases, Iran University of Medical Sciences, Tehran, Iran

3. Department of Medical Laboratory Science, Faculty of Allied Medical Sciences, Iran University of Medical Sciences, Tehran, Iran

4. ENT and Head and Neck Research Center and Department, Hazrat Rasoul Akram Hospital, Iran University of Medical Sciences, Tehran, Iran

Background: Chronic Rhinosinusitis (CRS) is a common inflammatory disorder influencing sinonasal mucosa. It is classified into two different subtypes referred to as CRS with nasal polyps (CRSwNP) and CRS without nasal polyps (CRSSNP). Although multiple immune mechanisms are suspected to be involved in CRSwNP etiology, the impact of T cells is not completely determined. The aim of this research was to evaluate the pattern of T cells (CD8+ and CD4+ T cells) in CRSwNP patients.

Methods: In our study, 13 healthy controls and 20 CRSwNP patients enrolled. Sinonasal tissues were investigated to find CD4 and CD8 T cells by immune histochemistry method. The diagnosis of CRS was confirmed by the current European Position Paper on Rhino sinusitis and nasal polyps. The Kruskal-Wallis, Mann-Whitney were used in statistical analyses. Gathered data were analyzed using SPSS V.21 and presented by GraphPad Prism V.6 softwares.

Results: The median (min-max) age of the groups was: 30 (14-51) for CRSwNP, and 29 (18-47) for healthy controls. The frequency of eosinophils was elevated in CRSwNP patients in comparison to controls. The frequency of total inflammatory cells in CRSwNP and control groups was 89 (66.5-127) and 11.5 (19-18.5), respectively. The frequency of CD4+ and CD8+ T cells as well as total inflammatory cells were significantly elevated in CRSwNP patients in comparison to healthy controls. CRSwNP group showed elevated infiltration of CD8+ T cells in sinonasal mucosa ($P < 0.0001$). CRSwNP group showed enhanced infiltration of CD4+ T cells into sinonasal mucosa in comparison to healthy controls ($P < 0.0001$).

Conclusion: The number of CD8+ and CD4+ T cells was increased in CRSwNP patients. Furthermore, the frequency of CD8+ T cells was higher than CD4+ T cells in sinonasal mucosa of CRSwNP patients. Targeted therapeutic approaches may be accessible by evaluating the pattern of T cells. Therefore, further complementary researches on T cells are needed to achieve this objective.

Keywords: Chronic Rhino sinusitis, CRSwNP, T cell, CD4, CD8

ICIA2018\Mucosal and Gastrointestinal Immunology\Oral\10007
10007

Increased expression of IL 17A and IL21 in the small intestine of patients celiac disease

Manizhe Faghih¹, Mohammad Rostami Nejad², Davar Amani¹

1. Department of Immunology, School of medical science, University of Shahid beheshti, Tehran, Iran

2. Department of Gastroenterology and Liver Diseases Research Center, School of medical science, University of Shahid beheshti, Tehran, Iran

Background: Th17 and its related cytokines play an important role in the pathogenesis of celiac disease (CD). Therefore the aim of this study was to evaluate the expression levels of IL-17, IL-21 in treated and untreated celiac disease patients compared to the control group.

Methods: In this study, duodenal biopsied were collected from 60 treated CD patients, 60 healthy control and 8 newly diagnosed celiac patients, during 2016. RNA extracted by commercial kit, cDNA synthesized and specific primer pairs were designed for each gene. The expression of genes was investigated by Real-time PCR technique using SYBER Green method.

Results: 60 treated CD patients (58.3% female; mean age of 38.8 years old), 60 healthy control (55% female; mean age of 36.5 years old) and 8 newly diagnosed celiac patients (75% male; mean age of 43.3 years old) were involved in the present study. The expression levels IL21 were insignificantly increased in the untreated CD group compared with control group. But the expression of IL-17A in the untreated CD patients was significantly higher than the control group ($p = 0.007$).

Conclusion: Our data showed that IL-17A, IL-21 has different expression profile in studied groups. According to the result, the investigated genes can be used as a diagnostic marker for screening patients with celiac disease. However, much more studies are required to confirm our findings and clarify the underlying mechanisms.

Keywords: Celiac Disease, IL-17A, IL-21, Gene Expression



Nanoimmunology

Oral Presentation

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Nanoimmunology\Oral\7528
7528

Nanocurcumin Influences Regulatory T cell Responses in Ankylosing Spondylitis

Rozita Abolhasan^{1,2}, Sanam Dolati^{1,2}, Mehdi Yusefi^{1,2}

1. Stem Cell and Regenerative Medicine Institute, Tabriz University of Medical Sciences, Tabriz, Iran

2. Department of Immunology, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Ankylosing spondylitis is a chronic, inflammatory disease of the axial spine that can appear with chronic back pain and progressive spinal stiffness. Participation of the spine and sacroiliac (SI) joints, peripheral joints, digits, entheses are distinguishing of the disease. Regulatory T cells with suppressive effects on inflammation and autoimmunity have been described to implicate in pathology of AS. This present study aimed to explore the effects of nanocurcumin on the frequency of Treg and their associated cytokines.

Methods: 24 patients with active AS joined in this study that were divided into two groups: 12 patients received 80 mg nanocurcumin every day for 16 weeks and 12 other patients received placebo as control group. First, the peripheral blood mononuclear cells were collected, and the frequency of Treg was analyzed by flow cytometry. Then, FoxP3 gene expression was appraised by using real-time qPCR. In addition, ELISA was carried out to determine the levels of cytokines such as IL-10, TGF- β and IL-6 in the peripheral blood of AS patients.

Results: Our study displayed that nanocurcumin could significantly increase the frequency of peripheral Tregs in AS patients (p value = 0.032). The results of real-time PCR also exhibited that the expression of FoxP3 was significantly increased (p value=0.02 and p value=0.0005, respectively). In addition, nanocurcumin group had higher IL-10 (p value=0.015) and TGF- β (p value=0.043) production and lower IL-6 (p value=0.018) expression than that of placebo group. The changes in the frequency and expression level of cytokines were positively correlated with the disease improvement.

Conclusion: Our results proposed that dysregulation of Tregs in peripheral blood of AS patients impacts the AS progress and nanocurcumin therapy through regulation of Tregs and the expression of the related cytokines can be appreciated in the treatment of ankylosing spondylitis and other autoimmune/ inflammatory diseases.

Keywords: Ankylosing spondylitis, Nanocurcumin, Regulatory T cells, Cytokines

ICIA2018\Nanoimmunology\Oral\7796
 7796

Combination treatment of tumor cells with the STAT3-specific siRNA loaded nanoparticles and BV6

Afshin Nikkhoo^{1,2}, Fatemeh Atyabi^{4,6}, Narges Rostami^{1,2}, Ali Masjedi^{1,2}, Ali Rastegari⁴, Masoomah Baghaei⁶, Vida Hashemi^{7,1,2}, Shima Bastaki^{3,1}, Farhad Jadidi Niaragh^{1,2,5}

1. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

2. Department of Immunology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

3. Department of cellular and molecular biology, faculty of Basic science, Azarbaijan Shahid Madani university, Tabriz, Iran

4. Department of Pharmaceutical Nanotechnology, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran 1714614411, Iran

5. Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

6. Nanotechnology Research Centre, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran

7. Department of basic science, faculty of medicine, Maragheh university of medical Science, Maragheh, Iran

Background: Apoptosis resistance is mediated by IAPs that inhibit caspases. Smac mimetics such as BV6 can bind to and inhibit IAPs. Signaling of STAT3 can enhance cancer cell survival. Therefore, combination therapy by BV6 smac mimetic and STAT3 siRNA-loaded nanoparticles (NPs) can be considered as potent anti-tumor therapy.

Method: STAT3-specific siRNA loaded NPs were generated by ionic gelation method and applied to downregulate the expression of STAT3 in order to increase apoptosis in 4T1 breast cancer cells and B16F10 melanoma cancer cells in combination with BV6smac mimetic. Apoptosis induction was analyzed by MTT assay and expression of apoptosis related genes was assayed by real-time PCR. The expression of angiogenic and metastatic genes was also investigated. The migration of cancer cells was investigated by wound healing assay.

Results: Synthesized NPs had about 100 nm size with a polydispersity index below 0.3 and a zeta potential about 13. NPs efficiently encapsulated siRNA, showed high serum stability, and efficient cellular uptake. Expression of STAT3 was significantly decreased in the tumor cells following treatment with nanoformulation. Combination therapy could significantly induce apoptosis in cancer cells which was associated with downregulation of survival molecules. Moreover, tumor cells exhibited lesser expression of angiogenic and metastatic molecules accompanied with decreased migration capacity after combination therapy.

Conclusion: Treatment with combination of STAT3 siRNA -loaded ChLa NPs and BV6 may be considered as a promising approach for cancer therapy; however, further in vivo investigations are necessary.

Key word: STAT3, BV6, NIK, apoptosis, cancer, Trimethyl chitosan

ICIA2018\Nanoimmunology\Oral\7810

7810

The immunomodulatory effect of nanocurcumin on acute Toxoplasmosis in Balb/c mice

Seyed Mohammad Mahdi Ghahari¹, Abolghasem Ajami², Mehdi Mahdavi³, Majid Sadeghizadeh⁴, Aahmad Reza Esmaeili Rastaghi^{5,*}

1. Department of Immunology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran
2. Department of Immunology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran
- 2- Molecular and Cell Biology Research Center, Mazandaran University of Medical Sciences, Sari, Iran
3. Department of Immunology, Pasteur Institute of Iran, Tehran, Iran
4. Department of Genetics, School of Biological Sciences, Tarbiat Modares University, Tehran, Iran
5. Department of Parasitology, Pasteur Institute of Iran, Tehran, Iran

Background: One of the controversial infections that most of the warm-blood creatures in the world are involved is related to *Toxoplasma gondii*. It is an important obligate intracellular protozoan parasite for immune deficient patient and pregnant women who sometimes are caused a death and abortion in individuals, respectively. Herein, immunomodulatory effect of nanocurcumin was evaluated against Toxoplasmosis killed vaccine model in Balb/c mice.

Methods: 144 Balb/c mice were included in 8 groups and the mice were administered different regimens of vaccine; vac +30, 20 mg/kg of curcumin and nanocurcumin, vac +Freund, killed vac, vac +Alum and PBS via subcutaneous route for three times with two weeks interval. Two weeks after the last immunization, the sera were assessed for total antibody, IgG1 and IgG2a with an optimized ELISA method. Lymphocyte proliferation was evaluated with Brdu method. The splenocytes culture supernatant was analyzed by ELISA for the presence of IL-4, IFN- γ , IL-2 and TNF α and ratio of IFN- γ /IL4, IFN- γ / TNF α and IL-2/IL4 cytokines. Then survival rate were determined at 10 days.

Results: Our results show that nanocurcumin 20 mg/kg significantly increased IFN- γ /IL4 and IFN- γ /TNF α cytokines but contemplative increased IL-2/IL4 cytokines and lymphoproliferative response versus control group. These results demonstrate that this dose of nanocurcumin could elicit strong Th1 immune responses at the lowest level of inflammation and production IgG antibody titers predominance of IgG2a and finally increased survival rate compared with control groups and P-values less than 0.05 was considered to be statistically significant.

Conclusion: This study provides evidence of immunomodulatory effects of nanocurcumin.

Keywords: *Toxoplasma gondii*, nanocurcumin, curcumin, cytokines

ICIA2018\Nanoimmunology\Oral\8500

8500

Enhanced immunological responses with tumor antigen loaded polymeric nanoparticlesSanaz Sheikhzadeh¹, Nowruz Delirezhi¹, Rahim Hobbenaghi²*1. Department of Microbiology, Faculty of Veterinary Medicine, Urmia University**2. Department of Pathology, Faculty of Veterinary Medicine, Urmia University*

Background: Cancer immunotherapy, the treatment that harnesses the patient's immune system to fight cancer, is now emerging as an important addition to conventional therapies. Cancer vaccines are active immunotherapy. Enhancing the immunogenicity of the vaccines requires efficient targeting of antigens and adjuvants to dendritic cells (DCs), that are the most potent antigen presenting cells for naive T cell activation. Nanoparticles (NP) are new vehicles for delivering vaccines. These particles are efficiently taken up by DCs because of their size and particulate structure which resembles pathogens. This study aimed to investigate the efficacy of a new nanoparticle-based cancer vaccine against breast cancer in mouse model.

Methods: Poly (D, L-lactide-co-glycolide) (PLGA) nanoparticles (NPs) containing breast tumor cell lysate were prepared using a double-emulsion solvent evaporation technique. Mice were challenged subcutaneously with 4T1 tumor cells. After induction of palpable tumor, immunotherapy was initiated. The control animals received PBS and the test animals were immunized with PLGA-NPs containing breast tumor cell lysate. One week after the last immunization, mice were euthanized. To evaluate the antitumor effects of vaccine; tumor growth, survival rates and cytokine level were determined.

Results: In mice immunized with PLGA-NPs loaded with tumor cell lysate, significantly reduced tumor growth and the survival rate was higher compared to the control group. The production of IFN- γ in spleen cell culture was increased while the production of IL-4 was decreased.

Conclusion: The goal of this study was to evaluate the antitumor activity of PLGA-based nanoparticle vaccine in a mouse model of breast cancer and so it is useful to develop a new way for the treatment of cancer.

Keywords: Immunotherapy, Nanoparticles, Tumor cell lysate, Cancer vaccine

ICIA2018\Nanoimmunology\Oral\8635

8635

Improvement of Hematologic and Immunologic Findings in Gamma Irradiated Rats Treated with Selenium Nano-Particles and Selenium Selenite

Siamak Asri Rezaei¹, Ali Asghar Tehrani², Arya Sohrabi², Zahra Nori Sabzikar³

1. Department of Clinical Pathology, School of veterinary college, University of Urmia

2. Department of Pathology, School of veterinary college, University of Urmia

3. Department of Clinical Pathology, School of veterinary college, University of Mashhad

Background: The objective of this study was to evaluate the efficacy of the selenium nano-particles and selenium selenite supplementation in boosting the immune response and protecting from oxidative stress in irradiated rats.

Methods: 45 mature male albino rats weighing 150 - 200 g were obtained and divided into 3 groups (control, Nano-Selenium and Selenite selenium). Selenium supplementation at dose 0.1 mg/kg was given by IP injection to rats, two weeks before gamma irradiation and then gamma therapy at the dose levels of 0, 2 and 8 Gy was done for two weeks (twice in a week and each time 20 min). After 14 days, blood samples were taken from heart and CBC were done and serum level of MDA, IL-2, TNF- α and IL-6 were determined.

Results: The results revealed that total body irradiation induced significant decreases in RBCs, WBCs and lymphocytes, as well as tumor necrosis factor alpha (TNF- α) and interleukin 2 (IL-2) also decrease significantly while interleukin 6 (IL-6) and lipid peroxidation marker malondialdehyde (MDA) in serum were elevated. In irradiated animals receiving selenium nano-particles and selenium selenite, values of RBC and WBC significantly increased as compared with the irradiated group, and similarly IL-2, IL-6 and TNF- α were significantly elevated. In contrast MDA levels in rats receiving selenium supplementation was decreased significantly. Also, the results of this study revealed that in immune system protection selenium nanoparticles be more effective than selenium salt.

Conclusion: The curative action of selenium nanoparticles enforcing significant innate response could trigger and augment adaptive immune response, thus protecting immune system from radiation induced damage as well as oxidative stress.

Keywords: Selenium nanoparticles, Gamma Radiation, Rat, Immune System

ICIA2018\Nanoimmunology\Oral\9965
9965

Gelatin nanoparticle based delivery of Urease and Omp31 in mice protects against *Brucella melitensis* 16M infection

Morteza Abkar¹, Saeed Alamian², Naghmeh Sattarahmady¹

1. *Nanomedicine and nanobiology research center, Shiraz University of Medical sciences, Shiraz, Iran*
2. *Brucellosis department, Razi vaccine and serum research institute, Karaj, Iran*

Background: *Brucella*(B) species are causative agents of brucellosis, a worldwide zoonotic illness causing Malta fever in humans and abortion in domestic animals. Due to the serious economic and medical consequences of the disease, many efforts are to prevent infection of domestic animals through vaccines.

Methods: In this work, we examined the vaccine potential of Gelatin nanoparticles formulation of Urease (Gelatin/Urease) and Omp31 (Gelatin/Omp31) against brucellosis. Gelatin/Urease and Gelatin/Omp31 nanoparticles were separately and in combination administered orally. Particle size and loading efficiency of the nanoparticles were determined. Antibody detection, cytokine measurement, lymphocyte proliferation assay and protection assay were performed. Finally, immunized mice were challenged with the virulent *B. melitensis*16M.

Results: All immunized mice elicited titers of specific immunoglobulin G (IgG). According to cytokine assay and antibody isotypes, oral administration with all vaccine formulations induced a mixed Th1-Th17 immune response. In lymphocyte proliferation assay, spleen cells from all-immunized mice showed a strong recall proliferative response. The combination of Gelatin/Urease and Gelatin/Omp31 nanoparticles conferred protection against *B. melitensis* infection equivalent to that of vaccine strain *B. melitensis* Rev.1. In comparison to Gelatin/Omp31 and Gelatin/Urease nanoparticles alone, combination of Gelatin/Omp31 and Gelatin/Urease nanoparticles induced only a low increase in protection level.

Conclusion: Altogether these results indicate that Gelatin/Omp31 or Gelatin/Urease nanoparticles could be useful candidates for the development of subunit vaccines against brucellosis. Furthermore, Gelatin nanoparticles are a suitable delivery system for orally-administered *Brucella* antigens.

Keywords: Omp31, Gelatin, Urease, Oral Vaccine

ICIA2018\Nanoimmunology\Oral\9968
9968

Subcutaneous vaccination with aluminium hydroxide/Omp31 and chitosan/Omp31 nanoparticles induces protection against *Brucella melitensis* infection in BALB/c mice

Morteza Abkar¹, Saeed Alamian², Naghmeh Sattarahmady¹

1. Nanomedicine and nanobiology research center, Shiraz University of Medical sciences, Shiraz, Iran

2. Brucellosis department, Razi vaccine and serum research institute, Karaj, Iran

Background: Vaccination has a high impact on the control and prevention of infectious illnesses. Subunit vaccines, such as recombinant proteins, are promising vaccine candidates because they are nonpathogenic, a virulent, well defined, nonviable and safer for manipulators, and economical. Since recombinant vaccines have low immunogenicity, application of an efficient antigen delivery system and adjuvant is very important. The use of the nanoparticle mediated delivery system is an effective strategy for site specific delivery of antigens.

Methods: In this study, we compared the potential of chitosan and aluminium hydroxide nanoparticles (NPs) to assess their ability in the development of new vaccines against brucellosis. Aluminium hydroxide and chitosan NPs were prepared, loaded with the protein antigen *B. melitensis* Omp31 and characterized. The immunostimulatory capacity of these vaccine delivery systems was examined in vivo.

Results: DLS and SEM images showed that most of NPs had a mean size distribution less than 100 nm. The amount of Omp31 loaded onto chitosan and aluminium hydroxide NPs were about 61% and 74%. According to the antibody subclasses and cytokine profile, subcutaneous immunization by both chitosan/Omp31 and aluminium hydroxide/Omp31 NPs induced T helper 1 (Th1) and Th1–Th2 immune responses, respectively. In comparison to aluminium hydroxide/Omp31 NPs, chitosan/Omp31 NPs induced only a low increase in protection level.

Conclusion: Altogether, our results indicate that both chitosan and aluminium hydroxide NPs are a potent delivery system for subcutaneous vaccination against brucellosis.

Keywords: Brucellosis, Omp31, Chitosan, Aluminium hydroxide.



Oral Immunology

Oral Presentation

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Oral Immunology\Oral\7637

7637

Assessment of microRNA-146a in aggressive periodontitis and its association with disease severity

Somayeh Ghotloo¹, Hossein Motedayyen^{2,3}, Mandana Sattari³, Mahmood Saffari^{1,4}, Davar Amani³

1. Department of Laboratory Medicine, Kashan University of Medical Sciences, Kashan, Iran

2. Department of immunology, Isfahan University of Medical sciences, Isfahan, Iran

3. Department of Immunology, Faculty of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

4. Department of Microbiology and Laboratory Medicine, Kashan University of Medical Sciences, Kashan, Iran

Background: MicroRNA-146a (miR-146a) is a small non-coding RNA that plays a critical role in negative regulation of innate immune response and its dysregulation has been associated with several inflammatory disorders. Aggressive periodontitis is an almost less common but severe form of periodontitis in which chronic inflammation leads to rapid destruction of tissues surrounding the teeth. This study was performed to evaluate miR-146a expression level in gingival tissues of patients with aggressive periodontitis and its association with disease severity.

Methods: Gingival samples of 18 patients with aggressive periodontitis and 10 healthy subjects were collected and level of miR-146a and its targets including TNF- α , IL-1 β and IL-6 was assessed using real-time PCR assay. Clinical parameters including probing depth (PD) and clinical attachment level (CAL) were measured and their correlations with miR-146a level were determined.

Results: Our results demonstrated an elevation in expression level of miR-146a in aggressive periodontitis compared to healthy control (p value < 0.001) which was directly associated with the disease severity (p value < 0.05). Over-expression of miR-146a was accompanied by the reduction of pro-inflammatory cytokines levels.

Conclusion: Positive association between miR-146a level and clinical parameter in aggressive periodontitis suggests that miR-146a may contribute to the pathology of the disease and serves as an indicator of disease severity. Elevation of miR-146a in these patients may occur in response to the bacterial components such as LPS leading to the control of pro-inflammatory cytokines level through a negative feedback loop.

Keywords: Aggressive periodontitis, MicroRNA -146a, Pro-inflammatory Cytokines

ICIA2018\Oral Immunology\Oral\7706

7706

Correlation Between Expression of Gingival TLR5, TLR2, RANKL in Tissues and Chronic Periodontal DiseasesAli Saeedi Boroujeni¹, Mandana Sattari², Samaneh Delavari¹, Ali Khodadadi¹*1. Department of Immunology, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran**2. Department of Immunology, Shahid Behashti University of Medical Sciences, Tehran, Iran*

Background: Periodontal diseases are the most common chronic infections in human. Inflammatory responses play an important role in periodontal disease. TLRs play a key role in induction of inflammatory responses by periodontal cells. In addition to TLR4, LPS of *Porphyromonas gingivalis* can stimulate TLR2 and its fimbria can stimulate TLR5, so the aim of this study was to evaluate the gingival gene expression of TLR2 and TLR5 and also RANKL as the most important factor in bone resorption, in periodontal diseases.

Methods: Gingival samples were collected from 20 individuals with clinically healthy gingival; 25 patients with moderate to severe chronic periodontitis after homogenizing tissue samples, RNA extraction and cDNA synthesis were done. Then, Real-time PCR technique was used for evaluating gene expression.

Results: There were no significant difference between the expression of TLR2 (0.209), TLR5 (0.553) and RANKL (0.597) in control group and chronic periodontitis group. There was also significant correlation between TLR2 and TLR5 gene expression ($P \approx 0.000$).

Conclusion: According to the results, no difference was seen between chronic periodontitis and control groups in terms of inflammatory responses.

Keywords: Periodontitis, Inflammation, RANKL, TLR

ICIA2018\Oral Immunology\Oral\8313

8313

Correlation between TLR5 expression and asymptomatic irreversible pulpitis

Shideh Mehr Mofakham*, Mandana Sattari, Hengameh Akhavan, Sarira Shah Navaz, Aram Mohammadi

Shahid Beheshti University of Medical Sciences, Tehran, IRAN

Background: Innate immunity in the pulp uses receptors to recognize molecular patterns common to microbes to initiate inflammation in the dental pulp, which eventually leads to irreversible damage and even necrosis of dental pulps. Since most of studies have been done on the roles of TLR2 and TLR4 in asymptomatic pulpitis, so the purpose of the present study was to compare TLR5 expression between asymptomatic irreversible pulpitis and normal pulps.

Methods: Pulps were taken from 20 teeth with asymptomatic irreversible pulpitis and 20 from intact premolars scheduled for extraction as control group. RNAs were collected from homogenized pulps and were subjected Reverse Transcriptase, producing cDNA, which is more stable. Forward and reverse primers were designed using Beacon designer software. Quantitative real time PCR were done by using SYBR Green. Statistical analysis was made by using REST software.

Results: The fold change of TLR5 expression in asymptomatic pulpitis was 1.510 but there is not any significant difference between asymptomatic pulpitis and normal pulp ($P \approx 0.852$).

Conclusion: It is concluded that with the progression of dental caries, despite the probable prevalence of some motile bacteria that are labeled with flagellum, but absence of a difference between a healthy pulp and an asymptomatic pulpitis can be a reflection of domination of non-motile bacteria in the pulp.

ICIA2018\Oral Immunology\Oral\8328
8328

The assessment of correlation between salivary matrix metalloproteinase-8, 20 concentration and severe early childhood caries S-ECC

Mina Biria¹, Mandana Sattari², Atieh Ehsani³

1. Department of Pediatric Dentistry, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Department of Immunology, Shahid Beheshti University of Medical Sciences, Tehran, Iran

3. Department of Pediatric Dentistry, Ghazvin University of Medical Sciences, Ghazvin, Iran

Background: Dental caries is a transmissible bacterial disease process caused by acids from bacterial metabolism diffusing into enamel and dentin and dissolving the minerals. Collagen degradations considered to be initiated by endogenous proteolytic enzymes, mainly collagenolytic matrix metalloproteinases (MMPs). Hence the aim of this research was to evaluate the relationship between severe early childhood caries and salivary MMP-8 (or salivary collagenase-2) and MMP-20 (or enamelysin) concentration.

Method: 50 children aged 36-72 months were included in this study. They were divided into two groups: caries-free (control) and severe early childhood caries (S-ECC), with 25 children in each group. Standard clinical examinations were performed, and around 1ml of unstimulated and whole expectorated saliva was collected. 3 months after dental restorations, children in S-ECC group were re-sampled and analyzed for salivary concentrations of MMP-8, MMP-20 using an ELISA assay.

Results: Before the treatment, subjects in S-ECC group presented with elevated levels of MMP-8 compared with subjects in control group ($p < 0.001$), but salivary concentration of MMP-20 in 2 groups was not statistically significant ($p = 0.189$). The difference between salivary concentrations of MMP-8 and MMP-20 before and after 3 months in S-ECC group were statistically significant (0.698 vs 0.331, $P < 0.001$; 3.801 vs 3.438, $P = 0.024$, respectively).

Conclusion: Our data reveal that subjects with severe early childhood caries have elevated levels of salivary MMP-8 compared to subjects with no caries lesions. 3 months after dental restorations, the concentration of MMP-8 and MMP-20 decreased significantly.

Keywords: Dental caries, Matrix Metalloproteinase 8, Matrix Metalloproteinase 20, saliva

ICIA2018\Oral Immunology\Oral\8444
8444

Comparison of gene expression of different isoforms of osteopontin in superficial dental caries

Aram Mohammadi¹, Mandana Sattari¹

1. Department of Immunology, School of medicine, Shahid Beheshti University of Medical Sciences

Background: Osteopontin (OPN), a kind of human genum glycoprotein, plays an important role in immune system modulation. OPN can activate osteoclasts, thus causing resorption. Also it may have a protective function against polymicrobial endodontic infection. Because different isoforms of OPN might have diverse roles, so the purpose of the present study was to compare gene expression of different isoforms of osteopontin in superficial dental caries and normal teeth.

Methods: Pulp samples were taken from 20 teeth with superficial dental caries as the case group and 20 pulpal samples from intact premolars scheduled for extraction as control group. RNAs were collected from homogenized pulps and were subjected Reverse Transcriptase, producing cDNA, which is more stable. Forward and reverse primers were designed using Beacon designer software. Finally, quantitative real time PCR were done in order to estimate the expression of different genes (Isoforms of osteopontin and GAPDH as housekeeping gene). Statistical analysis was made by REST software.

Results: There was lower expression of OPN2 in superficial caries but there were no significant difference between superficial dental caries and normal teeth regarding expression of OPN, OPN3 and OPN5.

Conclusion: It is concluded that probably OPN2 or variant 2 of OPN has a protective role against pulpal inflammation and there is a decrease in its expression by appearance and subsequent progression of dental caries.

Keywords: osteopontin, dental caries, pulp.

ICIA2018\Oral Immunology\Oral\8473

8473

Study of the effects of the dental pulp stem cells (DPSCs) on proliferation of peripheral blood mononuclear cells(PBMCs)Nikoo Hossein-khan nazer¹, Arash Khojasteh², Mandana Sattari¹, Seyed Mahmoud Masiha Hashemi¹*1. Department of Immunology, School of medicine, University of Shahid Beheshti**2. Department of Oral and Maxillofacial Surgery Shahid Beheshti University of Medical Sciences*

Background: Dental pulp stem cells (DPSCs) are multipotent mesenchymal stem cells that have the ability to self-renew and differentiate into a variety of cell types including osteoblasts, chondroblasts and adipocytes. Recent studies demonstrate that these cells exhibit immunomodulatory and anti-inflammatory effects on immune system. DPSCs capable of regulating immune cells which may provide a foundation for clinical use of DPSCs. In this study we determined immunoregulatory effects of DPSCs on peripheral blood mononuclear cell (PBMCs) proliferation using trans well co-culture system.

Methods: Dental pulp stem cells extracted from impacted third molars. Cells characterized for differentiation potential to adipogenic and osteogenic lineage and expression of mesenchymal stem cells markers. For transwell co-culture experiment, Stimulated and non-stimulated PBMCs from allogeneic donor cultured in different ratios with or without DPSCs (1:1, 1:5) for 72 hours. After incubation period proliferation of PBMCs evaluated using a cell proliferation BrdU ELISA kit.

Results: The BrdU results showed that Dental pulp stem cells reduced allogeneic PBMC proliferation. DPSC could inhibit stimulated and non-stimulated PBMC, 72 hours after incubation. This inhibiting property was more remarkable in 1:1 (MSC: PBMC) ratio.

Conclusion: This study demonstrated that Dental pulp stem cells exert anti proliferative effects on proliferation of allogeneic immune cells due to secreting soluble factors.

Keywords: PBMC, DPSC, co-culture

ICIA2018\Oral Immunology\Oral\8554
8554

HSP70 and CD184 are not important in tissue destruction in aggressive periodontitis

Sarira Shahnava¹, Mandana Sattari¹, Aram Mohammadi¹, Fatemeh Sarlati², Farshid Yeganeh¹

1. Department of Immunology School of medicine Shahid Beheshti University of medical sciences

2. Department of Periodontal School of Dental University of Islamic Azad

Background: Aggressive periodontitis is severe disease that its progression is fast and leads to bone destruction. The aggressive periodontal disease also divides two subtype groups that called local aggressive periodontal (LAP) and generalized aggressive periodontal (GAP). In other hand, CD184 is a chemokine receptor which is involved in the trafficking of leukocytes in to and out of extravascular tissues. CD184 may serve a homeostatic role to prevent excessive inflammation; alternatively, CD184 may be exploited by *Porphyromonas gingivalis* for suppressing innate immunity. However, the interaction of *P. gingivalis* with CD184 impairs antimicrobial host defense and it may be one mechanisms of immune evasion. Also, HSP70 family constitutes the most conserved and well-known class of HSPs. HSP70 has some regulatory and cytoprotective functions. So the aim of this study was to evaluate the relationship between expression of CD184 and HSP70 and clinical parameters of tissue destruction [such as CAL (Clinical attachment loss) and PD (Pocket depth)].

Methods: For this purpose, gingival tissue samples were collected from 20 individuals with clinically healthy gingiva and 25 patients with aggressive periodontitis. After homogenizing the gingival tissues, RNA was extracted by RNA isolation kit. After synthesis of cDNA, expression of CD184, HSP70 and beta-actin (as housekeeping gene) was evaluated by Real-time PCR.

Results: There is not any significant difference regarding expression of CD184 or HSP70 between aggressive periodontitis and healthy groups. In addition there is not any significant correlation neither between CD184 expression and CAL or PD nor HSP70 expression and CAL or PD ($P>0.05$).

Conclusion: Based on the results, we cannot suggest any significant role for CD184 in periodontal tissue destruction. Regarding no correlation between HSP70 and clinical parameters we cannot contribute any protective role for HSP70 in aggressive periodontitis. Of course more studies are needed in order to define the precise roles of CD184 and HSP70 in aggressive periodontitis.

Keywords: aggressive periodontitis, CD184, CAL, HSP70, PD

ICIA2018\Oral Immunology\Oral\9862

9862

Chemotactic and proliferative responses of human gingival fibroblasts (HGFs) to insulin-like growth factor-1 (IGF-1)Rokhsareh Sadeghi¹, Christopher A McCulloch²*1. Assistant Professor, Department of Periodontics, Faculty of Dentistry, Shahed University, Tehran, Iran**2. D.D.S., Ph.D., F.R.C.D. (C), Canada Research Chair in Matrix Dynamics, Professor, Matrix Dynamics Group, Faculty of Dentistry, University of Toronto, Toronto, Canada*

Background: It has been demonstrated the Insulin-like Growth factor-1 (IGF-1) has mitogenic and chemotactic effects on various cell lines. The aim of the present study was to assess the chemotactic and proliferative response of human gingival fibroblasts (HGFs) to IGF-1.

Methods: Gingival fibroblasts were obtained from the gingival tissue of medically and periodontally healthy patients. Chemotaxis assay was performed in modified Boyden chambers using Transwell® permeable inserts. The lower chamber of each well was filled with growth medium containing IGF-1 at different concentrations (0, 50, 100, 500, 1000 ng/ml). The cell suspension was placed on the top of the filters. Cell migration was assessed after 4 hours of incubation by counting the number of cells on the bottom surface of membranes using confocal microscopy. The proliferative effect of IGF-1 was evaluated by direct cell count. HGFs were plated in 24-well plates in growth medium supplemented with IGF-1 concentrations of 0, 50, 100, 200, 500, and 1000 ng/ml. The relative proliferation rate of cells was evaluated after 3, 5 and 7 days, by counting the nuclei of cells using Epifluorescence light microscope.

Results: IGF at a concentration of 50 ng and 100 ng showed a significant chemotactic effect on HGFs compared to the negative control ($P=0.0221$, $P=0.0009$, respectively). Regarding proliferation assay, no statistically significant differences were seen between the groups containing IGF-1 and negative control at any time point ($P>0.05$).

Conclusion: While HGFs showed a dose-dependent chemotactic response to IGF-1, they did not demonstrate any significant proliferative reaction to this growth factor.

Keywords: human gingival fibroblast, insulin-like growth factor, chemotaxis, proliferation



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Psychoneuroimmunology and Immunoendocrinology

Oral Presentation

ICIA2018\Psychoneuroimmunology and Immunoendocrinology\Poster\7476
7476

Dysregulation of peripheral Th17 cells in elderly patients with ischemic stroke

Farnaz Hajizadeh^{1,2}, Sanam Dolati^{1,2}, Mehdi Yousefi^{1,2}

1. Stem Cell and Regenerative Medicine Institute, Tabriz University of Medical Sciences, Tabriz, Iran

2. Department of Immunology, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Ischemic stroke causes brain injury with activation of an inflammatory response that can contribute to clinical impairment. The involvement of IL-17A and IL-17-secreting T cell subsets in IS patients has not been verified. In the present study, we studied Th17 frequency, cytokine secretion, expression of transcription factors related to Th17 which is compared between IS patients and control group.

Methods: 30 patients with IS and 30 persons as control group were enrolled in this study. Blood samples were collected from IS patients at 24 hours, 5 and 10 days after stroke. Th17 was detected among PBMCs using flowcytometric analysis, SYBR Green method of polymerase chain reaction (PCR) was employed for the analysis of ROR γ t mRNA levels, Quantification of IL-17 was carried out with the TaqMan Real-Time PCR. Secretion of IL-17, was evaluated in the supernatant acquired from cultured PBMCs using ELISA.

Results: The proportion of Th17 cells were significantly elevated in IS patients, ROR γ t mRNA expression levels were significantly increased in IS patients after 1 and 5 days compared to the controls 2.06 ± 72 and 1.77 ± 0.5 ($p < 0.0001$), IL-17A levels were increased in IS patients at 1, 5 and 10 days after stroke compared with the controls from 118 ± 35.9 to 171 ± 34 ($p = 0.0003$), 147 ± 24.84 ($p = 0.005$) and 146 ± 27.57 ($p = 0.005$), respectively.

Conclusion: These studies suggest that the increase in proportion of Th17 cells might contribute to the pathogenesis of IS.

Keywords: Ischemic Stroke, Th17, Cytokines, Inflammation

ICIA2018\Psychoneuroimmunology and Immunoendocrinology\Oral\7479
7479

Frequency and function of regulatory T cells in elderly patients with ischemic stroke

Farzaneh Ghorbani^{1,2}, Sanam Dolati^{1,2}, Mehdi Yousefi^{1,2}

1. Stem Cell and Regenerative Medicine Institute, Tabriz University of Medical Sciences, Tabriz, Iran

2. Department of Immunology, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

Background: CD4+CD25+regulatory T (Treg) cells play important role in pathogenesis of ischemic stroke (IS). In the current study, we studied Treg cells frequency and determined the levels of their signature transcription factors and cytokines in the peripheral blood of IS patients which is compared between IS patients and control group.

Methods: 30 patients with IS and 30 individuals as control group were enrolled in this study. The frequency of Treg lymphocytes, the expression of transcription factor (FOXP3) and the serum levels of associated cytokine (TGF- β) were assessed by flow cytometry, Real-time PCR and ELISA, respectively.

Results: The proportion of Treg cells were significantly reduced in patients compared with controls at 1 and 5 days after stroke, from 10.17 ± 3.12 to 6.83 ± 1.85 ($p < 0.0001$) and 8.47 ± 2.06 ($p = 0.016$), respectively. FoxP3 transcripts were significantly downregulated in IS patients 1 and 5 days after stroke compared to controls and TGF- β levels were diminished in IS patients at 1, 5 and 10 days after stroke compared with the controls from 155.7 ± 32.9 to 95.5 ± 28.6 ($p = 0.001$), 102 ± 25.4 ($p = 0.002$) and 122 ± 32.57 ($p = 0.08$), respectively.

Conclusion: This study suggests that the reduction of Treg cells contribute to the pathogenesis of IS. Also reduction in levels of TGF- β and FOXP3 attributed to that; inflammatory reaction was initiated to decrease and anti-inflammatory function of Tregs was started to increase subsequently 10 days after stroke.

Keywords: Ischemic Stroke, Treg, TGF- β , FOXP3

ICIA2018\Psychoneuroimmunology and Immunoendocrinology\Oral\7880
7880

Cytokine profile in autistic patients

Mohammad Mahdi Eftekharian¹, Mohammad Taheri^{2,3}

1. Neurophysiology Research Center, Hamadan University of Medical Sciences, Hamadan, Iran
2. Department of Medical Genetics, Shahid Beheshti University of Medical Sciences, Tehran, Iran
3. Urogenital Stem Cell Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: The etiology of Autism Spectrum Disorders (ASDs) as severe neurodevelopmental ailments is not known. However, several evidences point to dysregulation of immune system as an underlying cause of ASD.

Methods: In the present study we evaluated the mRNA expression levels of TNF- α , TGF- β , IFN- γ , CXCL8, IL-1 β , IL-2, IL-4, IL-6, IL-17 in whole blood samples of 30 ASD patients and 41 age and sex-matched healthy subjects with means of real-time PCR.

Results: TNF- α , IL-6 and IL-17 have been shown to be significantly up-regulated in ASD patients compared with healthy subjects ($P < 0.0001$, $P = 0.001$ and $P < 0.0001$ respectively). IL-2 has been shown to be significantly down-regulated in total ASD patients ($P < 0.0001$). No significant difference has been found in expression levels of other cytokines between patients and healthy subjects.

Conclusion: The present study provides further evidences for dysregulation of immune response in ASD patients.

Keywords: autism, TNF- α , TGF- β , IFN- γ , CXCL8, IL-1 β , IL-2, IL-4, IL-6, IL-17

ICIA2018\Psychoneuroimmunology and Immunoendocrinology\Oral\8642

8642

Association of major depressive disorder with activation of NLRP3-inflammasome and Total Antioxidant Capacity

Abdolghader Taene¹, Ghazaleh Khalili Tanha², Gholamreza AnaniSarab³

1. Student Research Committee, Birjand University of Medical Sciences, Birjand, Iran

2. MSc of cellular and molecular biology, Cellular and Molecular Research Center, Birjand university of medical science, Birjand, Iran.

3. Cellular and Molecular Research Center, Birjand University of Medical Sciences, Birjand, Iran

Background: Depression is one of the most common, expensive, and devastating psychiatric disorder. Approximately, 16% of males and 25% of females experience depression during their lifetime course. Despite several studies, the pathogenic mechanism of major depressive disorder (MDD) have been poorly understood. It seems that changes in the expression of neurotransmitters, the hypothalamus-adrenal-pituitary (HPA) axis, oxidative stress and immune dysregulation can be a major factor of MDD. The present study examined the hypothesis to determine whether NLRP3-inflammasome is activated in peripheral bloods mononuclear cells from MDD patients. Furthermore, we analyzed the differentiation of Total Antioxidant Capacity (TAC) between MDD patients and healthy groups.

Methods: Twenty MDD patients without treatment with a diagnosis of major depression, twenty patients treated with antidepressant drugs and twenty healthy volunteers were enrolled in this study. Samples were collected from Department of psychiatry Emam Reza Hospital, in Birjand. The inflammasome activation was studied by SYBR green real time PCR and TAC of samples were measured by using the determination of ferric reducing ability power (FRAP). The differences in mean expression of NLRP3 and Caspase 1 gene as well as the amount of TCA between groups were analyzed.

Results: We found increased NLRP3 ($P=0.023$) and caspase-1 ($P=0.026$) gene expression relative to B-actin in bloods mononuclear cells of MDD patients respect to healthy controls. Interestingly, the amount of NLRP3 and caspase-1 gene expression reduced in treated group. Also we found out that serum TCA level in healthy group were significantly higher than both untreated and treated patient groups ($P<0/005$).

Conclusion: We conclude that NLRP3-inflammasome is activated in MDD patients and antidepressant drugs reduce NLRP3-inflammasome in MDD patients. Also reduced levels of TAC were observed in MDD patients.

Keywords: Major depressive disorder, Total Antioxidant Capacity, NLRP3-inflammasome

ICIA2018\Psychoneuroimmunology and Immunoendocrinology\Oral\9815
9815

The Effects of D-aspartate in Experimental Autoimmune Encephalomyelitis

Yaser Panahi¹, Mahdi Goudarzvand², Kosar Abouhamzeh³, Antimo D'Aniello⁴, Gholamreza Azizi^{3*}

1. North Khorasan University of Medical Sciences, Bojnurd, Iran. Vector-born Disease Research Center, Bojnurd, Iran
2. Department of Physiology and Pharmacology, Faculty of Medicine, Alborz University of Medical Sciences, Karaj, Iran
3. Non-Communicable Diseases Research Center, Alborz University of Medical Sciences, Karaj, Iran
4. Laboratory of Neurobiology, Zoological Station of Naples "Anton Dohrn", Villa Comunale, Napoli, Italy

Background: Experimental autoimmune encephalomyelitis (EAE) is a widely used model for MS. In the present research, our aim was to test the effects of D-aspartate (D-Asp) on remyelination in EAE.

Methods: In this study the EAE was induce on C57BL/6 mice and treated by D-Asp. On days 20, brains and cerebellums of mice were removed for histological analysis. Quantitative determination of D-Asp was performed using liquid chromatography-tandem mass spectrometry.

Results: Our findings demonstrated that IP injection of D-Asp had beneficial effects on EAE severity than orally gavage.

Conclusion: IP injection of D-Asp had more effects on remyelination than orally received EAE mice.

Keywords: Experimental autoimmune encephalomyelitis, D-aspartate



Reproductive Immunology

Oral Presentation

14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Reproductive Immunology\Oral\6337
6337

Immunomodulatory effects of human amniotic epithelial cells on cytokine profile of naïve CD₄⁺ T cells from women with unexplained recurrent spontaneous abortion

Hossein Motedayyen^{1*}, Nader Tajik², Amir Hassan Zarnani², Abbas Rezaei¹

1. Department of Immunology, Faculty of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

2. Immunology Research Center (IRC), Iran University of Medical Sciences, Tehran, Iran

Background: Immune imbalance at the maternal-fetal interface plays a fundamental role in the pathogenesis of unexplained recurrent spontaneous abortion (URSA). Human amniotic epithelial cells (hAECs) possess pregnancy-friendly immunomodulatory effects. Here, we investigated how cytokine profile of naïve T cells from USRA patients is affected by hAECs.

Methods: Naïve CD4⁺ T cells were isolated from 18 patients with URSA using immunomagnetic separation method. hAECs were obtained from healthy women during elective cesarean deliveries. hAECs were co-cultured at different ratios (1:1, 1:2, 1:5, 1:10) with naïve CD4⁺ T cells (4×10^5) in 24-well plates and stimulated with anti-CD3/CD28A. After 3 and 6 days, the co-culture supernatants were collected and the level of IFN- γ , IL-17A and IL-4 were analyzed using an Enzyme-linked immunosorbent assay (ELISA) kit. In this experiment, hAECs and unstimulated naïve T cells cultured alone served as negative controls, while naïve CD4⁺ T cells cultured alone and stimulated with anti-CD3/CD28A were considered as positive control.

Results: Our results showed that a statistically significant reduction in the level of IFN- γ and IL-17A was observed for all hAECs: naïve T cells ratios after 3 and 6 days of co-culture ($p < 0.05$ -0.0001). Moreover, the level of IL-4 was significantly higher at all hAECs: naïve T cells ratios after 3 days ($p < 0.01$ -0.0001). This significant increase in IL-4 level was not observed on day 6 for any of the hAECs: naïve T cells ratios.

Conclusion: These findings suggest that hAECs have immunomodulatory effects on cytokine profile of naïve T cells from URSA patients through stimulating Th2 cytokine (IL-4) production and decreasing Th1 and Th17 cytokines (IFN- γ , IL-17A) production.

Keywords: Immunomodulatory effects, Amniotic Epithelial Cells, Naïve T cells, Recurrent Spontaneous Abortion.

ICIA2018\Reproductive Immunology\Oral\7494
7494

Th17 and regulatory T cells frequency in RIF's patients treated with IVIG

Zahra Pourmoghdam¹, Majid Ahmadi¹, Mehdi Yousefi^{1, 2}

1. Stem cells and Regenerative Medicine Institute, Tabriz University of Medical Sciences, Tabriz, Iran

2. Department of Immunology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Failure to achieve successful implantation and clinical pregnancy after transferring at least three good quality embryos in to normal uterus is called recurrent implantation failure (RIF). Several studies have shown the importance of local mother's immune system in the success or failure of implantation. Recent evidence suggests the role of immunotherapy in the treatment of RIF's patients; among them intravenous immunoglobulin (IVIG) has become very important in recent years, with the hope that it will increase the chances of success in in vitro fertilization (IVF). The aim of this study is assessment of immune cells (Th17 and Treg) and their cytokines after IVIG therapy in women with RIF.

Methods: 72 patients with immunological abnormalities (elevated Th1/Th2 ratio and natural killer cells frequency) selected and divided into two groups. 40 patients as the IVIG group received aspirin, heparin and IVIG. 32 patients as control group received just aspirin and heparin. 400 mg/kg of IVIG was administered intravenously 2 days prior to embryo transfer. Then Th17 and Treg frequency was evaluated by flow cytometry. RT-PCR and ELISA were used to measure the Th17 and Treg cytokines expression and secretion. ROR γ t and FOXP3 transcription factors expression were evaluated by RT-PCR.

Results: After IVIG treatment elevated Treg frequency ($p=0.186$) and FOXP3 expression ($p=0.0004$) were detected. TGF β and IL-10 expression ($p=0.0038$, $p=0.0058$ respectively) and secretion ($p=0.0156$, $p=0.0413$ respectively) were significantly increased. In comparison with control group no Significant differences were seen in IL-17 and IL-23 expression and secretion as well as Th17 frequency; only ROR γ t expression was decreased significantly ($p=0.0218$).

Conclusion: Our study results suggest that treatment with IVIG can be very helpful in successful implantation and pregnancy after IVF by affecting immune regulatory system. So it can be used as a therapeutic option in infertility following IVF.

Keywords: Intravenous immunoglobulin, Recurrent implantation failure, Regulatory T cell, T helper 17 cell.

ICIA2018\Reproductive Immunology\Oral\7512

7512

The frequency and function of Follicular T helper in preeclampsiaHaniehheydarlou^{1,2}, Majid Ahmadi^{1,2}, Mehdi Yousefi^{1,2}*1. Stem Cell and Regenerative Medicine Institute, Tabriz University of Medical Sciences, Tabriz, Iran**2. Department of Immunology, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran*

Background: Pre-eclampsia is a relatively common multisystem disorder of human, affecting about 3% of pregnancies. It develops in several stages only the last being the clinical illness and causes different maternal and fetal problems. Studies have revealed that immunological changes in the placental microenvironment may have an important role in the initiation of preeclampsia. Whereas, another study show that there is a relationship between TFH and its cytokine (IL-21) in immunological infertility so it may happen through the generation of some antibodies.

Methods: 25 preeclampsia women and their control group were assessed at the Alzahra Hospital of Tabriz University of Medical Sciences, Tabriz, Iran. PBMCs were separated from the blood and then RNA was extracted. Afterwards, the frequency of TFH cells was evaluated by flow cytometry and using fluorescent antibodies against CD4, CXCR5 and ICOS markers. The expression of CXCR5 and Bcl6 genes were measured. The TFH related cytokines including IL-6 and IL-21 were assayed by ELISA technology.

Results: TFH cells with the CD4⁺ CXCR5⁺ ICOS⁺ phenotypes showed a significant increase in preeclamptic patients compared to normal pregnant women. The expression of CXCR5 and BCL6 were increased. In addition, IL-6 and IL-21 secreted from TFH cells up regulated in patients group. Correlation analysis showed a strong relationship between the TFH CD4⁺ CXCR5⁺ ICOS⁺ and pathogenesis of preeclampsia.

Conclusions: These data indicate that CD4⁺ CXCR5⁺ BCL6⁺ TFH cells may participate in the pathogenesis of preeclampsia.

Keyword: cytokine, Follicular T helper, Preeclampsia.

ICIA2018\Reproductive Immunology\Oral\7846
7846

Difference in the seminal plasma protein expression in unexplained infertile men with successful and unsuccessful in vitro fertilization outcome

Zahra Kanannejad¹, Behrouz Gharehi Fard^{1,2}

1. Department of Immunology, Shiraz University of Medical Sciences, Shiraz, Iran

2. Infertility Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Unexplained male infertility (UMI) is a condition in which routine semen analysis fails to detect subcellular sperm dysfunctions. In the present research, a comparative proteomics study of seminal plasma (SP) was conducted in men with unexplained infertility whose female partners undergone in vitro fertilization (IVF) treatment to find differences in the SP protein profile.

Methods: Five UMI men with successful and eight with unsuccessful IVF outcome enrolled in this study. Two-dimensional polyacrylamide gel electrophoresis (2D-PAGE) technique was used for protein separation. The differentially expressed proteins were identified using Mass Spectrometry technique.

Results and conclusion: indicated that at least three different protein spots, including clusterin, epididymal secretory protein E1, and prostate specific antigen, were differentially expressed in the successful group as compared with the unsuccessful couple (more than 1.5 fold change, $P < 0.05$). Considering the role of all three identified proteins in the sperm quality, the results of the present study introduced these proteins as new biomarkers for success of IVF in UMI couples.

Keywords: Unexplained male infertility, in vitro fertilization, seminal plasma, Proteomics.

ICIA2018\Reproductive Immunology\Oral\8103

8103

Decidual microenvironments from resorbed and non-resorbed fetuses show divers effects on dendritic cells maturation state and function

Eskandarian Maryam, Yaftiyan Atefeh, Jahangiri Amir Hossein, Moazzeni Seyyed Mohammad

Immunology Department, Faculty of Medical science, Tarbiat Modares University, Tehran, Iran

Background: Dendritic cells (DCs) constitute a system of antigen presenting cells that are key regulators of immune responses. DCs can acquire immunogenic or tolerogenic properties depending to their microenvironment. In this study we aimed to determine the effects of decidual cells from resorbed and non-resorbed fetuses from abortion prone mice on DC functions.

Methods: DCs were differentiated from mouse bone marrow cells in the presence of DC differentiation cytokines, GM-CSF and IL-4. The DCs were then co-cultured with decidual cells from resorbed and non-resorbed fetuses and their immunophenotype and pinocytotic activity were evaluated using the flow cytometric analysis. The obtained DCs were also pulsed with paternal antigens and used for immunization studies. The antigen specific T cell responses were determined in T cells derived from lymph node of immunized animals.

Result: Our finding revealed that treatment of dendritic cells with decidual cells from resorbed fetuses significantly increased the MHC-II, CD86 and CD40 expression by DCs compared to DCs treated with decidual cells from non-resorbed fetuses. A remarkable reduction in the endocytic capacity were also observed in iDCs that were co-cultured with decidua of resorbed fetuses. Meanwhile, the ability of DCs to induce lymphocyte proliferation significantly increased following their treatment with decidua of resorbed fetuses.

Conclusion: We concluded that in decidua of resorbed fetuses an immune potentiating microenvironment dominancy could be observed which can modulate the DCs phenotype and functions, leading to triggering of an undesirable immune response that is associated with fetal rejection.

Keyword: dendritic cells, decidua, microenvironment

ICIA2018\Reproductive Immunology\Oral\8104

8104

The immunosuppressive activity of decidual cells protect the semiallogenic fetus via modulation of dendritic cells functions

Eskandarian Maryam, Yaftiyan Atefeh, Jahangiri Amir Hossein, Mohseni Mahboobeh, Moazzeni Seyyed Mohammad

Immunology Department, Faculty of Medical Science, Tarbiat Modares University, Tehran, Iran

Background: Dendritic cells (DCs) represent a system of antigen presenting cells with a dual property of triggering an immune response or promote cell tolerance, depending on the cellular and molecular context of their microenvironment. This study was done to investigate the immunomodulatory properties of decidual cell from normal pregnancy on DCs.

Methods: Immature dendritic cells (iDCs) were prepared from mouse bone marrow progenitors in the presences of IL-4 and GM-CSF cytokines. iDCs were pulsed with paternal antigens. In some cultures of DCs, decidual cells were added during the antigen pulsing process. The immunophenotype and antigen uptake properties of DCs were measured using the flow cytometry. The antigen specific T cell stimulation potency of DCs was also determined using the LTT assay.

Results: Our results demonstrated that decidual cells from pregnant mice had inhibitory effect on DCs maturation and function, a significant reduction in the expression of co-stimulatory molecules (MHC-II, CD80 and CD86) and induction of antigen specific T cell proliferative response by DCs were observed. The Ag uptake capability of DCs were remained unchanged after treatment with decidua of the pregnant mice.

Conclusion: It seems that decidual microenvironment modulate the function of DCs which result in the induction of immunologic tolerance to the semi allogeneic fetus at the feto-maternal interface.

Keyword: BM-derived DC, Decidua, Microenvironment

ICIA2018\Reproductive Immunology\Oral\8159

8159

Menstrual blood-derived stem cells modulate functional features of natural killer cells

Mohammad Reza Shokri¹, Mahmood Bozorgmehr^{2,3}, Alireza Ghanavatinejad⁴, Mehdi Aleahmad⁴, Shohreh Nikoo⁵, Maryam Tavakoli², Somaieh Kazemnejad⁶, Reza Falak^{1,5}, Fazel Shokri^{4,7}, Amir Hassan Zarnani^{2,5}

1. Department of Immunology, School of medicine, Iran University of Medical Sciences, Tehran, Iran
2. Reproductive Immunology Research Center, Avicenna Research Institute, ACECR, Tehran, Iran
3. Oncopathology Research Center, Iran University of Medical Sciences, Tehran, Iran
4. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran
5. Immunology research center (IRC), Iran University of Medical Sciences, Tehran, Iran
6. Reproductive Biotechnology Research Center, Avicenna Research Institute, ACECR, Tehran, Iran
7. Department of Hybridoma, Monoclonal Antibody Research Center, Avicenna Research Institute, ACECR, Tehran, Iran

Background: Up to now, however, there is a great debate on how functional features of uterine NK cells are regulated during normal pregnancy. We hypothesized that endometrial stem cells could potentially attenuate bona fide cytotoxic features of NK cells.

Methods: Menstrual blood stem cells (MenSCs), as surrogates for endometrial stem cells, and Bone marrow mesenchymal stem cells (BMSCs) were isolated and fully characterized. The effects of MenSCs in a co-culture system on various features of NK cells including proliferation, K-562 cytotoxicity, and expression of cytotoxic markers were determined by CFSE, Calcein-AM and flow cytometry, respectively. Secretome of MenSCs was determined by growth factor array. IDO activity in MenSCs upon treatment with IFN- γ was determined by a colorimetric method. The involvement of IL-6, IL-10 and, TGF- β in MenSCs-modulated NK cell proliferation was assessed by neutralizing antibodies.

Results: MenSCs exhibited multi-lineage differentiation potential and expressed markers associated with mesenchymal and embryonic origins. Contrary to BMSCs, MenSCs induced proliferation of NK cells, while IFN- γ /IL-1 β pre-treated MenSCs significantly inhibited NK cell proliferation. Of 42 growth factors tested, BMSCs produced higher levels of IGFBP 1-4, VEGF-A, b-NGF and M-CSF compared to MenSCs. Both MSCs showed a high activity of IDO upon IFN- γ treatment. The Anti-proliferative potential of IFN- γ /IL-1 β -pretreated MenSCs was mediated through IL-6 and TGF- β . Similar to BMSCs, MenSCs impaired the cytotoxic activity of NK cells on K-562 cells, consistent with the lower expression of perforin and granzymes A/B. Interestingly, MenSCs were found to be prone to NK-mediated lysis in an MHC-independent manner.

Conclusion: Our findings may imply that NK cell dysregulation in such pregnancy-related disorders as infertility and miscarriage may not be considered as an intrinsic defect, rather stemmed from dysfunction of endometrial stromal stem cells.

Keywords: Natural killer cells, Endometrial stem cells, Menstrual blood stem cells, Bone marrow mesenchymal stem cells, Mesenchymal stem cells, Cytotoxicity

ICIA2018\Reproductive Immunology\Oral\8195

8195

Pro-inflammatory Milieu Makes Menstrual blood stem cells a Potent inducer of Function Regulatory T cells

Mehdi Aleahmad¹, Mahmood Bozorgmehr^{2,3}, Shohreh Nikoo⁴, Mohammad-Reza Shokri⁵,
Alireza Ghanavatinejad¹, Maryam Tavakoli², Somaieh Kazemnejad⁶, Fazel Shokri^{1,7}, Amir Hassan Zarnani^{2,4}

1. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran
2. Reproductive Immunology Research Center, Avicenna Research Institute, ACECR, Tehran, Iran
3. Oncopathology Research Center, Iran University of Medical Sciences, Tehran Iran
4. Immunology research center (IRC), Iran university of medical sciences, Tehran, Iran
5. Department of Immunology, School of medicine, Iran University of Medical Sciences, Tehran, Iran
6. Reproductive Biotechnology Research Center, Avicenna Research Institute, ACECR, Tehran, Iran
7. Department of Hybridoma, Monoclonal Antibody Research Center, Avicenna Research Institute, ACECR, Tehran, Iran

Background: Regulatory T cells (Treg) are fundamental for successful pregnancy. There are several mechanisms proposed so far for induction of Tregs at the feto-maternal interface. However, the potential role of endometrial stem cells (ESCs) in this context remained to be elucidated.

Methods: Menstrual blood stem cells (MenSCs), as surrogate for endometrial stem cells, were isolated and evaluated for their immunophenotypic characteristics and multi-lineage differentiation capacity. Effect of untreated and IFN- γ /IL-1 β -pre-treated MenSCs on CD4+CD25+FOXP3+ Treg generation, proliferation of CD3/CD28-activated CD4+ T cells and IL-10 and IFN- γ production was studied by flow cytometry. Capacity of MenSCs to produce functional IDO was assessed by a colorimetric assay. Involvement of IDO, PGE2, IL-6, IL-10 and TGF β in MenSCs-modulated T cell proliferation and generation of Tregs was investigated by blocking antibodies. Functional capacity of MenSCs-induced Tregs was studied in an allogeneic MLR system. Bone marrow-derived stem cells (BMSCs) were used as positive control in some settings.

Results: Unlike BMSCs, MenSCs induced proliferation of T CD4⁺ lymphocytes in a dose dependent manner, while pre-treatment with IFN- γ /IL-1 β reversed proliferation promoting effect of MenSCs through induction of PGE2. Both mesenchymal stem cell types showed high IDO activity following IFN- γ treatment. Untreated MenSCs inhibited generation of Tregs, while treatment of MenSCs with pro-inflammatory cytokines resulted in a significantly higher numbers of Tregs mainly through IL-6, IL-10 and TGF- β production. Tregs generated in the presence of MenSCs were found functionally active and significantly inhibited MLR response in a dose dependent manner.

Conclusion: The results of this study imply that pro-inflammatory microenvironment, which is established upon blastocyst implantation; may trigger endometrial stem cells to induce generation of T cells with a regulatory phenotype and function.

Keywords: Menstrual blood stem cells, Tregs, Pregnancy, MLR inhibition

ICIA2018\Reproductive Immunology\Oral\8478
8478

Adipose-derived mesenchymal stem cells could alter cytokine profile of natural killer cells in abortion prone mice

Fatemeh Rezaei, Seyed Mohammad Moazzeni

Immunology Department, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

Background: Cytokine production by NK cells at the feto-maternal interface play a pivotal role in the maintenance of normal pregnancy and the imbalance of cytokine profile is associated with recurrent spontaneous abortion (RSA). MSCs are shown to have immunomodulatory effects on NK cells and their cytokine production. The purpose of this study is to evaluate the impact of MSCs therapy of RSA mouse model on NK cells cytokine profile.

Methods: Adipose derived MSCs were injected (IP) at day 4 of gestation to female CBA/J mice following their mating with DBA/2 male. In RSA control group PBS was injected and CBA/J × BALB/c mating was also used as normal pregnancy control. Decidual cells were isolated at gd12.5 and the production of TGF- β , IL-4, IL-10, and IFN- γ by uNK cells were examined using the flow cytometry.

Results: Our results showed that the administration of MSCs could alter the cytokine profile of uNK cells in abortion- prone mice. We demonstrated that MSCs therapy caused the downregulation of IFN- γ and upregulation of IL-4 and IL-10 production by uNK cells at the feto-maternal interface and skewed cytokine production toward a Th2 bias. Also MSCs didn't have any effect on TGF- β production.

Conclusion: This finding indicated that MSCs could modulate the uterine NK cells cytokines profile from Th1 predominance to Th2 bias and thereby help to maintain tolerogenic microenvironment at the feto-maternal interface.

Keywords: Recurrent Spontaneous Abortion, Mesenchymal stem cell, NK cells, cell therapy, cytokine expression.

ICIA2018\Reproductive Immunology\Oral\9964
9964

MSC therapy reduces complement precipitation at the feto-maternal interface of abortion prone mice

Shahab Shahgaldi¹, Amir Salek Farrokhi¹, Seyed Mohammad Moazzeni¹

1. Department of Immunology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

Background: Recurrent spontaneous abortion, which is quite prevalent among pregnant women, is mainly due to immunological factors and disorders. Complement is involved in a wide and diverse variety of immune and non-immune operations, including the development of miscarriages and has emerged as a common event in recurrent pregnancy loss. Mesenchymal stem cells (MSCs), which possess unique in vivo immunosuppressive properties, reduce the abortion rate through modulation of immune responses. In the present study, the quantity of complement C3 deposition is determined at the feto-maternal interface of abortion prone mice (CBA/J mated DBA/2 males).

Methods: MSCs were derived from abdominal fat (AT-MSCs) of CBA/J mice. On the 4.5th day of gestation, the test group (CBA/J × DBA/2) received an IP injection of 1×10^6 of AT-MSCs while the control (CBA/J × DBA/2) and normal pregnancy (CBA/J × BALB/c) groups received an IP injection of PBS. On the 13.5th day of gestation, tissue samples (placentae and deciduae of pregnant mice) were analyzed to measure complement C3 deposition using immunohistochemistry.

Results: The resorption rate was significantly reduced following MSC therapy as shown in our previous studies. Furthermore, immunohistochemical analysis of complement deposition demonstrated that MSC administration considerably diminished complement C3 deposition at the feto-maternal interface of abortion prone mice.

Conclusion: In the present study, we showed for the first time that adoptive transfer of MSCs reduced abortion rate in abortion-prone mice by adjusting the levels of complement C3 deposition. Our results suggested that diminished complement deposition induced by MSCs could contribute to better pregnancy outcomes.

Keywords: Mesenchymal stem cell, Recurrent spontaneous abortion, Cell therapy, Complement deposition, Complement C3, Pregnancy.

ICIA2018\Reproductive Immunology\Oral\9976
9976

The regulatory influence of 1,25-Dihydroxy Vitamin D₃ on VEGF related angiogenesis in patients with endometriosis

Zahra Moradi^{1,2}, Sepideh Khodaverdi³, Mahdi Shekarabi¹, Tahereh Mohammadi², Ali Akbar Delbandi^{1,2}

1. Immunology Research Center, Immunology and Infectious Disease Institute, IRAN University of Medical Sciences

2. Immunology Department, School of Medicine, IRAN University of Medical Sciences

3. Endometriosis Research Center, IRAN University of Medical Sciences

Background: Endometriosis is a chronic inflammatory disease in women, which characterized by the growth of endometrial tissue outside the uterus. Angiogenesis is one of the most important mechanisms in the pathogenesis of endometriosis. The aim of this study was to evaluate the regulatory effects of vitamin D₃ on the VEGF expression in ectopic (EESCs) and eutopic (EuESCs) endometrial stromal cells, peritoneal fluid and blood mononuclear cells in patients with endometriosis compared to nonendometriotic controls.

Methods: Endometrial ectopic and eutopic stromal cells were isolated by enzymatic digestion from 20 endometriotic patients and 10 non endometriotic controls. Also, the peritoneal fluid and peripheral blood mononuclear cells were separated by ficoll method from 10 patients and 10 controls. The VEGF gene and protein expression levels were measured by Real-Time PCR and ELISA, respectively. In the following, the cells were treated with optimal concentration of 1,25(OH)₂D₃ in three different times, and the regulatory effects of this vitamin were evaluated in gene and protein expression levels of VEGF in all mentioned cells. Subsequently, the effects of 1, 25(OH)₂D₃ on proliferation, expression and production of VEGF in HUVEC cells was examined.

Results: EESCs showed significantly higher VEGF expression than EuESCs and control endometrial stromal cells (P<0.0001). In addition, peritoneal fluid mononuclear cells in patient group produced higher levels of VEGF protein compared to controls (P<0.01). Considerably, Vitamin D₃ reduced the expression levels of VEGF in EESCs and peritoneal fluid mononuclear cells (P<0.001, P<0.01, respectively). Also 1, 25(OH)₂D₃ caused inhibition of proliferation and VEGF expression at the gene and protein level in HUVEC cells in vitro.

Conclusion: Our results showed that 1, 25(OH)₂D₃ is an appropriate candidate for reduction of angiogenesis in endometriosis by regulating the production of angiogenesis- related inflammatory factors.

Keywords: Endometriosis, 1,25(OH)₂D₃, Stromal cells, VEGF



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Research and Development

Oral Presentation

ICIA2018\Research and Development\Oral\5320

5320

Apigenin: a flavonoid to manage the expression of P53 gene in gastric and bladder cancer

Fereshteh Abdolmaleki¹, Effat Farzanehnia²

1. Student of Master in medical biotechnology School of Paramedical Sciences, Qazvin University of Medical Sciences

2. Student of Master in medical biotechnology School of Paramedical Sciences, Qazvin University of Medical Sciences

Background: This study is aimed to evaluate the effect of Apigenin on the expression of P53 in gastric and bladder cancers (MKN45 and EJ138 cell lines)

Methods: The cell lines were cultured in enriched medium with FBS and kept in CO₂ incubator to proliferate. The cells were treated with different concentration of apigenin (0 to 100 μM) and exposed to MTT solution to determine their cell viability. After detecting the IC₅₀ of Apigenin, the cell was treated with the chosen concentration of Apigenin (0 to 60 μM) and their expression in both levels of mRNA and protein were studied through real-time qPCR and Western blotting technique respectively, also MSP technique for DNA methylation and apoptosis assay were done for better understanding the action of the gene. All experiments were done triplicate and for three periods of times (24, 48, and 72 h).

Result: MTT assay showed that there is no cytotoxic effect on cells up to 30 μM of Apigenin. The result of real-time qPCR and western blotting presented that the expression of P53 has been developed after exposing with apigenin dose and time dependently so that in 60 μM of apigenin and after 72 h we had the most expression of P53 in both level of mRNA and protein. Also, the result of MSP approved our hypothesis base on the methylation of P53 promoter and the result of apoptosis assay by Hoechst 33258 kit demonstrated that apigenin could promote the apoptosis of treated cells.

Conclusion: In both gastric and bladder cancer (MKN45 and EJ138 cell lines) P53 gene was methylated so its expression has been reduced. Our study displayed that apigenin could increase P53 expression to induce apoptosis and suppression of cancer cell. This flavonoid has not been done in this both cell lines and we hoped that it could be used as a breakthrough in cancer therapy.

Keywords: Apigenin, P53 gene, gastric cancer, bladder cancer

ICIA2018\Research and Development\Oral\6310

6310

Investigation and Suggestion of Allergenicity Reduction from Osmotin/thaumatin-like superfamily

Esmail Ebrahimie^{1,3,4,5}, Ali Niazi¹, Zarrin Minuchehr², Bijan Bambai², Najaf Allahyari Fard², Nassim rahmani¹

1. *Institute of Biotechnology, Shiraz University, Shiraz, Iran*

2. *National Institute of Genetic Engineering and Biotechnology (NIGEB), Tehran, Iran*

3. *School of Medicine, The University of Adelaide, Adelaide, Australia*

4. *Division of Information Technology, Engineering and the Environment, School of Information Technology and Mathematical Sciences, University of South Australia, Adelaide, Australia*

5. *School of Biological Sciences, Faculty of Science and Engineering, Flinders University, Adelaide, Australia*

Background: Allergy is a widespread disorder that can lead to a systemic anaphylactic shock. Underlying asthma has been associated with increased risk of severe reactions and even death caused by food allergy. The prevalence and impact of allergic diseases are increasing worldwide. Osmotin/thaumatin-like superfamily is one of the most important plant allergen superfamily that is found in apple, sweet cherry, grape, kiwi and etc. In this research, we tried to suggest the solution for reducing or removing Allergenicity of this great group.

Methods: we collected all plant allergens from the allergen databases and prepared Osmotin/thaumatin-like superfamily, and then the pattern was obtained by using PROSITE. In addition, we find all submitted epitopes of Osmotin/thaumatin-like superfamily members by using SDAP and BIOPEP databases. Afterward with comparing the pattern and epitopes sequences; we selected similar amino acids and changed two of them with amino acids belonging to the same families to omit the Allergenicity and provide changing in epitope sequence.

Results: K83 amino acid from allergen epitope changed to H83, the only amino acid which could be replaced, and G84 amino acid from allergen epitope changed to D84, E84, S84, T84, and Y84. Before and after mutations, positions of allergen epitopes binding with heavy and light chains of IgE were analyzed and ΔH (the change in enthalpy) for these reactions was calculated.

Conclusion: We determined that replacing of Lys with His and Gly with Thr, reduce the possibility of docking between allergen epitope and IgE, therefore reduction of allergenicity was expected.

Keywords: Plant allergen, Allergenicity, Osmotin/thaumatin-like, IgE

ICIA2018\Research and Development\Oral\7462

7462

Effects of extremely low frequency electromagnetic fields with various intensities on immune system of ratsMahsa Sobhani^{1,2}, Mohammad Mahdi Eftekharian^{1,3}, Ghasem Solgi^{1,4}, Safoora Nikzad⁵, Alireza Zamani^{1,4}*1. Department of Immunology, School of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran**2. Student Research Committee, Hamadan University of Medical Sciences, Hamadan, Iran**3. Neurophysiology Research Center, Hamadan University of Medical Sciences, Hamadan, Iran**4. Molecular Immunology Research Group, Research Center for Molecular Medicine, Hamadan University of Medical Sciences, Hamadan, Iran**5. Department of Medical Physics, School of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran*

Background: Some epidemiologic studies have shown that extremely low frequency electromagnetic fields (ELF-EMF) affect human health as well as immune system cells and organs. Therefore, it will be reasonable to test whether ELF-EMF can change the level of cytokines like interferon-gamma (IFN- γ), interleukin-4 (IL-4) and expression of T-box transcription factor (T-bet) and GATA binding protein 3 (GATA-3) that regulates TH1/TH2 balance. Aim of this study was to evaluate the effects of ELF-EMF with various intensities on immune system function.

Methods: The experiments were performed on eighty two-month-old-male wistar (202.5 ± 7.5 gram) rats, which were divided to 5 groups and subjected to 50 Hz ELF-EMF with the flux densities of 1, 100, 500, and 2000 μ T, 2h/day for two months. The controls were placed in identical chamber without ELF-EMF. In order to activate the immune system, human serum albumin (HSA) was injected by intraperitoneal route to the all groups on days 31, 44, and 58. The relative expression of T-bet and GATA-3 in spleen and thymus were evaluated with RT-qPCR. The serum level of IFN- γ and IL-4 were measured by enzyme-linked immunosorbent assay (ELISA).

Results: Under our experimental conditions, there was no significant differences between the mean mass of rats and spleens, whereas, the weights of thymuses were significantly decreased in ELF-EMF exposed group compared with controls ($p < 0.05$). Our experiments showed that ELF-EMF significantly decrease the expression of T-bet and GATA-3 in spleen ($p < 0.05$), whereas, there were no significant changes on genes expression in thymuses. There were significantly decrease in concentrations of IFN- γ and IL-4 before stimulation of the immune system ($p < 0.01$, $p < 0.001$, respectively), but the serum levels of this cytokines were not significantly changed after stimulation.

Conclusion: It can be concluded that ELF-EMF exposure could altered the immune system function in the way suppression of immune system response.

Keywords: Cytokine, Extremely Low Frequency Electromagnetic Fields, Spleen, Thymus, T-Lymphocytes.

ICIA2018\Research and Development\Oral\7738
7738

Natural killer cells expansion for adoptive immunotherapy: comparison of two isolation methods, three cytokines, IL-2, IL-15, or IL-18 and impact on NK cytotoxicity

Anahita Najafalizadeh¹, Nafiseh Esmacil¹, Farzaneh Peighambarzadeh¹, Fariborz Mokarian¹, Abbas Rezaei¹, Mazdak Ganjalikhani Hakemi¹

1. Department of Immunology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran
2. Cancer Prevention Research Center, Isfahan University of Medical Sciences, Isfahan, Iran

Background: NK cells are progressively considered as medical tools for cancer immunotherapy. The development of applicable methods to generate a large number of functional NK cells is a crucial step to maximize the potential of this approach.

Methods: In this article, NK cells were isolated from PBMCs using 2 different methods. One method with MagniSort Human NK cell Enrichment kit, and the second method isolation with antibodies and complement (cytotoxic method). Purified NK cells were activated in vitro by IL-2 and IL-15 for 14 days and we used IL-18 on the day 14 for another 48 hours to increase the cytotoxicity. Finally, the HL-60 (Human promyelocytic leukemia cells) cell line was used as a target to assess NK cytotoxic activity.

Result: The purity of NK cells was 86% and 92% by cytotoxic method and MACS method, respectively. NK cells expanded 60-100 folds in day 14. The expanded NK cells were highly cytotoxic (90%) and they were more than 80% viable. Interestingly, the cytotoxic activity of NK cells was decreased in presence of IL-18.

Conclusion: Here we present a simple approach that enables the isolation of NK cells without any beads and magnets using mouse complement with anti-CD3 and anti-CD19 and compared it with the routine isolation method, using magnetic cell sorting. This simplified and efficient method for NK cells isolation and activation could be used in future clinical trials.

Keywords: Immunotherapy, NK cells, AML, Interleukin 2, Interleukin 15, Interleukin 18

ICIA2018\Research and Development\Oral\8152

8152

High Yield Magnetosome Production by the *Magnetospirillum Gryphiswaldense* MSR-1 After Optimizing The Oxygen, Iron, and Carbon Sources and Concentrations in Culture Medium

Leila Hatami Giklou Jajan¹, Masoud Ghorbani², Seyed Fazlollah Mousavi, Seyed Nezamodin Hosseini, Behzad Ghareyazie, Behrooz Vaziri, Mohsen Abolhassani

1. Pasteur Institute of Iran

2. Pasteur Institute of Iran, Research and Production Complex, Department of Viral Vaccines, Kilometer 25 Autobahn Karaj, Alborz, Iran

Background: Magnetosome production yield of the *Magnetospirillum gryphiswaldense* strain MSR-1 was significantly increased after optimization of the oxygen, iron, carbon and nitrogen sources and concentrations on current culture medium without the need to the genetic manipulation.

Results: The optimum concentration of Sodium-L-lactate (25Mm), NaNO₃ (40 Mm), ferric citrate (200 µM) and dissolved oxygen (2-5%) in the culture medium resulted a higher magnetosome yield than the current culture medium. An approximately four times higher magnetosome yield of 186.67 mg/liter was reached in batch cultures after optimization of the growth medium along with a significant increase of bacterial culture. The iron uptake was also increased significantly after optimization of the medium. A more than 3-fold magnetic response of bacteria to the magnetic field was significantly higher than the other test groups. Based on these findings, it was concluded that alteration in growth medium and optimizing the parameters could increase the magnetosome production yield without any gene manipulations within the bacteria genome. This could be of great importance in the production of industrial magnetosome products.

Conclusion: This optimization technic cannot only reduce the culture time period but can reduce the cost of production. Refinement of this method will enable the further increase of magnetosome yield.

ICIA2018\Research and Development\Oral\8298

8298

Comparison of the Effects of Multimeric and Natural Soluble CD40L on the Antibody Production from B cell LineBahare Shokoohian¹, Fatemeh Yari¹, Mahshoid Mohammadipour¹, Zohreh Sharifi¹*1. Blood Transfusion Research Center, High Institute for Research and Education in Transfusion Medicine, Tehran, Iran*

Background: CD40L is a member of TNF family that plays an important role in innate and adaptive immunity. Since this protein plays its role through clustering the receptors, CD40, and also the natural soluble form of CD40L is not stable, it seems that multimeric form of CD40L can have a stronger effect than the natural trimeric form in activation of B lymphocytes and IgG production.

Methods: dodecameric (four-trimeric) form of CD40L was designed in silico by replacing the extra-cellular domain of CD40L with lectin domain of surfactant protein-D (SP-D). The fusion protein (SPD-CD40L) was expressed by HEK293 cell line. SPD-CD40L and natural soluble CD40L from platelet concentrate were purified by affinity chromatography. Specificity and molecular weight of both proteins were determined by ELISA and western-blot. Daudi cells were encountered natural and dodecameric CD40L, amounts of produced IgG were investigated by sandwich ELISA. The data were analyzed by the statistical tests of ANOVA, Tukey HSD and Wilcoxon.

Results: The specificity of the recombinant CD40L was confirmed. IgG production from Daudi cells increased due to exposure with both natural and recombinant CD40L. This increase was higher in exposure to recombinant CD40L than the natural protein and the difference in IgG increase after exposure to natural and recombinant CD40L was considered statistically significant.

Conclusion: recombinant dodecameric soluble CD40L protein (SPD-CD40L) can mimic the orientation of membrane-bound CD40L and thus create a stronger message than the natural soluble trimeric molecule in B-lymphocytes because of the spatial orientation of the molecule.

Keywords: CD40L, surfactant protein-D, IgG, B lymphocyte cell line

ICIA2018\Research and Development\Oral\9856
9856

Development of a novels TRAIL-conjugate with improved half-life

Reza Falak^{1,2}, SepidehMohseni¹, Mohammad DavoodzadehGholami¹, Elaheh Safari^{1,2}, Gholamalikardar³

1. Immunology Research Center, Iran University of Medical Sciences, Tehran, Iran

2. Department of Immunology, School of Medicine, Iran University of Medical Sciences, Tehran, Iran

3. Asthma, Allergy and Immunology Research Center, Tehran University of Medical Sciences, Tehran, Iran

Background: TRAIL is a potent inducer of cell death and may selectively eradicate a wide variety of human cancer cells without damaging normal ones. However, recombinant TRAIL has short half-life, necessitating its repetitive injection for maintenance of effective therapeutic concentrations, thus limiting its clinical applicability.

Methods: In this study, we designed a TRAIL fused with an albumin binding peptide (ABP) and evaluated its serum half-life. TRAIL and ABP-TRAIL were cloned in pET28a (+) and transformed into bacterial host. To compare the serum half-lives of the TRAIL and ABP-TRAIL in blood circulation, samples were collected after administration of a single dose in Balb/c mice and TRAIL was measured at periodic intervals by ELISA.

Results: ABP-TRAIL not only retained similar bioactivity as TRAIL in vitro, it also conferred 3.3 fold prolonged half-life in mice.

Conclusion: It seems that ABP-TRAIL is a suitable substitute for TRAIL, of course, if it can pass trial steps.

Keywords: ABP, ABP-TRAIL, Albumin, Half-life, TRAIL.

ICIA2018\Research and Development\Oral\9890

9890

Improvement of CM11 a Antimicrobial Peptide Using D-diastereomer Amino Acid Substitution

Somayyeh Ghadimi, Kazem Ahmadi, Mahdi Fasihi-ramandi

Molecular biology research center, system biology and poisonings institute, Baqiyatallah University of medical sciences

Background: In order to improve the antibacterial activity of CM11, a cecropinA-melittin hybrid, we have substituted some of L-amino acids with their D-counterparts in the CM11 sequence.

Methods: Here structure – activity relationship studies of these anti-microbial peptides (AMPs), by means of in silico analyses of AMPs structures as well as in vitro evaluation of hemolytic and cytotoxicity assays, antimicrobial activity and cell selectivity were performed.

Results: Our results showed that different structural characters of L- and D-CM11 affected its biological behavior and reduced the hemolysis and cytotoxicity of D-CM11 in compare with L-CM11. Furthermore, D-CM11 had an improved antimicrobial activity and therapeutic index (TI) towards some of the gram-negative pathogenic bacteria such as *S. sonnei*, *E. coli*, *S. marcescens*, *A. baumannii*, *P. vulgaris*, *P. aeruginosa*, *B. melitensis*, *B. abortus*, *E. cloacae* that isolated from clinical samples in microdilution assays. These observations indicate that the antimicrobial activity of this amphipathic peptide (D-CM11) was not chiral specific.

Conclusion: Besides increasing our knowledge on the structure-function of CM11 and its D-enantiomer analogue, this study revealed that D amino acid substitution could preserve strong anti-bacterial activity of D-CM11 mostly due to direct peptide–membranes electrostatic interactions, without giving a toxic effect towards eukaryotic cells.

Keywords: Antimicrobial peptide, CM11, Cytotoxicity, D-amino acid, Diastereomer.

ICIA2018\Research and Development\Oral\9929
9929

Production of functional recombinant human GM-CSF (rhGM-CSF) under specific promoter in *Escherichia coli*

Mahmoud Vahidi¹, Bahram Kazemi^{2,3}, Mahmood Bozorgmehr⁴, Mojgan Bandehpour^{1,2,3}

1. Department of Biotechnology, School of Advanced Technologies in Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Cellular and Molecular Biology Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

3. Department of Biotechnology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

4. Oncopathology Research Center, Iran University of Medical Sciences, Tehran, Iran

Background: Granulocyte-macrophage colony-stimulating factor (GM-CSF) is an important hematopoietic growth factor that stimulates multipotent progenitor cells and it also stimulates the differentiation of myeloid leukemic cells. Recently, some of the cytokines as other recombinant proteins could be produced using the recombinant DNA technology. Current study is aimed to clone and production of the Human GM-CSF under specific promoter in *Escherichia coli* and assessed its biological functions.

Methods: The hGM-csf gene construct was artificially synthesized; subsequently it was subcloned into the pcDNA3.1 (+) vector in HindIII restriction enzyme sites. Recombinant DNA was transferred and expressed in BL21 cells. rhGM-CSF protein was evaluated by SDS-PAGE and Western blotting. It was purified by Ni-NTA affinity chromatography. The purified protein concentration and also confirmation were determined by ELISA. MTT assay was applied to evaluate the biological activity of rhGM-CSF on the Erythroleukemic cell line proliferation.

Results: The hGM-csf gene was successfully cloned and transformed into expression *E. coli* cells. As a result, a specific band was observed both on the SDS-PAGE and nitrocellulose membrane after Western blotting. The purified protein concentration was equal to 100 pg/ml. MTT assay showed that exposed cells with rhGM-CSF were proliferated in a dose dependent manner.

Conclusion: The GM-CSF gene with specific promoter was successfully cloned in the prokaryotic system and transcription was carried out by T7 RNA polymerase. Therefore, mass production of GM-CSF can be a great help in clinical trials and research studies. Additionally, prokaryotic system, which used in current work, is less costly and less time-consuming.

Keywords: Human GM-CSF, Cloning, Prokaryotic system, MTT assay

ICIA2018\Research and Development\Oral\11094
11094

Transfection of human beta-globin gene into the AAVS1 locus of K562 cell line using CRISPR/Cas9 technology

Elahe Alikhani, Mahin Nikogoftar, Amir Ali Hamidieh, Mehdi Shamsara

1. Iran Blood Transfusion Organization , Tehran, Iran
2. National Institute of Genetic Engineering and Biotechnology, Tehran, Iran
3. Tehran University of Medical Sciences, Tehran, Iran

Background: Beta thalassemia is a kind of hereditary illness, which is due to an impairment in production of globin chain. There is a need to find a suitable treatment for this disease. CRISPR-Cas9 is a new promising genome editing technology having high efficiency in gene correction.

Method: In this study, we aimed to insert the human beta-globin gene specifically into the AAVS1 locus using AAVS1 Transgene knock-in vector kit (Origen). At first, the gene was amplified with PCR and cloned into the pTG19-T. So then subcloned into the pAAVS1-puro-DNR. For the second part of our study, the K562 cells were cultured. For determining the minimal fatal concentration of puromycin, the cells were exposed to the different dose of the antibiotic. We transfected this cells with both vectors (Pcas-Guide-AAVS1 and the recombinant PAAVS1-Puro-DNR) (Bio-Rad)and PE-GFPas control. The transfected cells were treated with Puromycin for two weeks. For evaluating the insertion; we extract the DNA from cells and confirmed the insertion of the gene by PCR. In the end, we evaluated the expression of inserted segment by RT-PCR.

Results: The Beta-globin gene was cloned into the PAAVS1-Puro-DNR. We confirmed our results by both digestion and sequencing. For determining the minimal fatal dose of puromycin, we exposed the cells with the different dose of antibiotics. The appropriate was 1 µg/mL. We transfected the cells with different protocols of electroporation. For evaluating the insertion, we amplified 700 bp segment with PCR from transfected cells. We confirmed the expression of inserted segment into the cells by amplified 200 bp with RT-PCR.

Conclusion: Our results showed that K562 is appropriate for studies of blood disease especially beta thalassemia because they are derived from blood cells and do not express the beta-globin gene naturally. One of the biggest problems of these cells is transfection and manipulating of them because of being suspended.

Keywords: Beta thalassemia, CRISPR-Cas9, K562 cell



Stem Cells Based Immunotherapy

Oral Presentation

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Stem Cells Based Immunotherapy\Oral\7727
7727

Effects of Adipose-derived mesenchymal stem cells injection on immune responses to *Leishmania major* in BALB/c mice

Elham Zangane¹, Sara Soudi¹, Ahmad Zavaran Hosseini¹

1. Department of Immunology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

Background: Mesenchymal stem cells have immunomodulatory property. MSCs response to infection through migration to infection site and modulation of immune cells that terminate to augmentation or suppression of inflammation. Recently MSCs are used in cell therapy of infection disease. In the correct study, we considered AD-MSCs immunomodulatory effect on *L.major* infection in BALB/c mice.

Methods: BALB/c mice were footpad injected by *L.major*, one group was received AD-MSCs at different interval post infection intravenously (i.v), second group received DMEM in i.v routs. Three time post challenge three mice of each group were sacrificed, spleen and lymph node cells were under secondary PHA/LPS and no-Antigen stimulation for IFN- γ , IL-4, TNF- α , IL-10 and nitric oxide (NO) measurement. Spleen parasite burden were determined by limiting dilution.

Results: Intravenous injection of AD-MSCs to *L.major* infected BAB/c mice have stimulated increase in IFN- γ /IL-4 and TNF- α /IL-10 ratio in spleen and lymph node of infected mice at all study times. In addition, more NO production were detected in splenocyte of this group, and the significant less spleen parasite load were observed. As expected non-MSCs treated mice were demonstrated lower IFN- γ /IL-4 and TNF- α /IL-10 ratio and higher spleen parasite burden.

Conclusion: Immunomodulatory properties of AD-MSCs can direct immune response to the benefit of IFN- γ /IL-4 increase and better control of parasite burden in lymph node tissues against *L.major* infection. Although increase in TNF- α /IL-10 ratio and NO production did not happened at protection level, the results showed the AD-MSCs power in their regulation, finally it seems that AD-MSCs do not have complete protective proportion alone against Leishmaniasis and needs some alterations considered on a candidate in immunotherapy methods.

Keyword: Mesenchymal stem cells, *Leishmania major*, BALB/c

ICIA2018\Stem Cells Based Immunotherapy\Oral\7808

7808

Entosis –like phenomenon and abrogation of MHC class I molecules (H-2D^b) in mouse primary bone marrow mesenchymal stem cells by CRISPR/Cas9 approach

AmrollahMostafazadeh^{1,2*}, BehnamKalali², Karin Mink², Martin Skerhut², Ramona Nitui², RaphaelaSemper², Ahmed Sedegh², ThureAdler³, Markus Gerhard²

1. Cellular and Molecular Biology Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, Iran

2. Institute for Medical Microbiology, Immunology and Heigen, Munich, Germany

3. Institute for Stem Cell Research, Helmholtz Center Munich, Germany

Background: The glycoproteins encoded by the major histocompatibility complex (MHC) are a major barrier in allogenic bone marrow transplantation. In this study, mouse mesenchymal stem cells (MSCs) were used for the first time to examine the application of the Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR/Cas9) approach in MHC de-mismatching.

Methods: MSCs were isolated from femur bone marrow of C57bl/6 mouse, cultured in RPMI-1640-10%FBS, and characterized by immunophenotyping using anti-stem cell antigen – I (Sca-I) and -CD45 antibody, and then transfected with px330 plasmids harbouring the Cas9- and H-2D^bspecific guide RNA- encoding genes or with empty px330 plasmids or remained as untransfected cells. Then GFP (reporter gene) positive cells were sorted by FACS. After 8 days of incubation at 37°C in CO₂ incubator, the sorted cells were stained with anti mouse MHC-I (H2D^bmolecules) FITC conjugated antibody, and then analyzed by flowcytometry.

Results: 97.5% of the MSCs transfected with px330 empty plasmid, and 98.2% of untransfected cells were H2D^b antigen positive. This value for px330-H2D^bsg RNA plasmids transfected cells was 31.5%. It means that the efficacy of CRISPR/Cas9 technique in abrogation of MHC –I gene expression was 68%. Moreover, during the bone marrow cells culture we provided some morphological evidence indicating that the Sca-I⁺ CD45⁺ stem cells can be invaded by several small bone marrow cells (more than 25 cells), implicating anentosis-like phenomenon.

Conclusion: CRISPR/Cas9 approach is an efficient tool in MHC abrogation. Independent to a complex cytokines and growth factors and necessity to a smaller panel of antibody for immunephenotyping, MCSs can be considered as an affordable source for MHC engineering study by CRISPR/Cas9 approach. Moreover this study for the first time provided some evidence for invasion of many cells to one mesenchymal stem cell.

Keywords: CRISPR/Cas9, mesenchymal stem cell, Major histocompatibility complex, Entosis

ICIA2018\Stem Cells Based Immunotherapy\Oral\8194

8194

The in-vitro Immunomodulatory Effects of Exosomes Derived from Tumor Cell Line or Mesenchymal Stem Cell on the Functions of Splenocytes Cells isolated from a Mouse Model of Experimental Autoimmune Encephalomyelitis

AnwarFathollahi¹, Farshid Yeganeh¹, Mostafa Haji Molla Hoseini¹, Seyed Mahmoud Hashemi¹

1. Department of Medical Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Immunomodulation is a shared feature of mesenchymal stem cells (MSCs) and tumors. Exosomes are 40–100nm extracellular vesicles released from cells. Previous studies showed that MSCs and tumors utilize the exosomes to modulate immune responses. This preliminary study was performed to find out that whether exosomes derived from MSCs or 4T1 tumor cell line enable to modulate cytokine productions and gene expressions corresponding to T helper1 (Th1) and Th17, in an Experimental Autoimmune Encephalomyelitis (EAE) mouse model.

Methods: The splenocytes isolated from EAE mice re-stimulated with disease-inducing peptide (MOG35-55, 150 μ /ml), in the presence or absence of exosomes from either 4T1 (15 μ g/ml) or MSCs (15 μ g/ml). Forty-eight hours later, the cells were harvested, and the relative expression levels of master regulators of T cell sub-types, including T-bet, GATA-3, ROR γ t, and FOXP3 were determined using Real-Time RT-PCR assay. Additionally, the alteration of ELF4 mRNA levels was evaluated. The levels of IL-10, IFN- γ , TGF- β , and IL-17 in the supernatants of splenocytes culture were measured by ELISA 72 hours after the same treatments.

Results: The concentrations of IL-17 and IFN- γ in the supernatants of splenocytes culture stimulated with MOG alone, MOG+4T1 exosomes or MOG+MSC were not different ($P > 0.05$). However, both Exosomes from 4T1 and MSCs increased the levels of ROR γ t expression (1.633 and 1.365 fold, respectively). Exosomes treatment enhanced the expression of FOXP3, nonetheless, the production of IL-10 and TGF- β were increased only in splenocytes treated with 4T1 exosome ($P=0.02$ and $P=0.021$ respectively). Comparison between cells that treated with 4T1 or MSCs exosome revealed that 4T1 exosome up-regulate ELF4 expression, (1.621 fold), a transcription factor which prevents Th17 differentiation.

Conclusion: MSCs and 4T1 exosomes modulate Th1/Th17 responses through FOXP3 up-regulation, however, 4T1 exosome has superiority for immunomodulation due to enhancement of ELF-4 transcription and increases IL-10 and TGF- β production.

Keywords: 4T1 cell line, Exosome, Immunomodulation, Mesenchymal stem cell

ICIA2018\Stem Cells Based Immunotherapy\Oral\8276

8276

Histological study of Stem cell-based therapies to promote angiogenesis in ischemic vessel disease by tissue engineering method

Amani Saeede¹, Shahrooz Rasoul¹, Mortaz Esmacil^{2,3,4}, Hobbenaghi Rahim⁵, Mohammadi Rahim⁶, Khoshfetrat Ali Baradar⁷

1. Department of Basic Sciences, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

2. Clinical Tuberculosis and Epidemiology Research Center, National Research Institute of Tuberculosis and Lung Diseases (NRITLD), Shahid Beheshti University of Medical Sciences, Tehran, Iran

3. Department of Immunology, Faculty of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

4. Division of Pharmacology, Utrecht Institute for Pharmaceutical Sciences, Faculty of Science, Utrecht University, Utrecht, The Netherlands

5. Department of Pathobiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

6. Department of Surgery and Diagnostic Imaging, Faculty of Veterinary Medicine, Urmia University, Tabriz, Iran

7. Department of Chemical Engineering, Sahand University of Technology, Tabriz, 14 51335-1996, Iran

Background: Stem cell therapy is novel approach to control or treatment of tissue ischemia associated with peripheral arterial disease. Progenitor cells derived from bone marrow or pluripotent stem cells have shown therapeutic value for restoring tissue function by boosting angiogenesis. The most common technique for stem cells is an injection of cells into specific areas. Therapeutic angiogenesis using bioengineered tissues composed of hydrogen scaffold and hematopoietic stem cell like mast cell and mesenchymal cell were adopted to assess their ability to induce vascular network formation in transected femoral artery location and improve functional recovery in ischemic hind limb in rat.

Methods: Thirty male white Wistar rats weighing approximately 200-250g were divided into three experimental groups (n = 6), randomly: In ischemia group (Ischemia) the femoral artery and was transected and hind limb of animals became ischemic, in hydrogel such as group (Hydro) in the ischemic animals the transected location was immersed with 50 μ L hydrogel solution and in cell transplanted group (Hydro/Mast cells), in the ischemic animals the transected location was immersed with 50 μ L Hydrogel mixed with 50 μ L PBS containing 10×10^6 mast cells combination.

Results: Analyses of capillary density and histomorphometrical analysis were performed on day 14 after surgery. Mean number of blood vessels in groups with MC, MSc and mixture of them indicated a significant difference compared to other experimental groups ($P < 0.05$). The mean number of medium and large blood vessels in HYDRO/MC group was significantly more than placebo groups ($P < 0.05$). Immunohistochemistry indicated more positive immunoreactivity to CD34 protein in Hydro/Mast cells.

Conclusion: Our finding show, Infected locally Mast cells and Mesenchymal cells are a Novel approach for therapeutic angiogenesis that may be able to control ischemic vessel disease and provide improved long-term neovascularization in cases of peripheral arterial diseases.

Key words: Mast cells, Mesenchymal cell, Histology, Tissue engineering, Angiogenesis, Rat

ICIA2018\Stem Cells Based Immunotherapy\Oral\8440

8440

The potential benefits of activated platelet-rich plasma on neuroprotection of Human Umbilical Cord Blood Mesenchymal Stem Cells in Mouse Model of Experimental Autoimmune Encephalomyelitis

Maryam Kheirandish¹, Ghazaleh Bordbari², Sharareh Bordbari¹, Afsaneh Jahangirian¹

1. Blood Transfusion Research Center, High Institute for Research and Education in Transfusion Medicine, Tehran, Iran

2. Dept of Immunology, Faculty of Medical Sciences, Esfahan University

Background: Current clinical research supports the immunomodulatory and neuroprotective effects of administered Mesenchymal Stem Cells (MSCs) in Multiple Sclerosis. The therapeutic efficacy of expanded hUCB-MSCs in aPRP was evaluated in EAE model in comparison with ordinary cultured hUCB-MSCs.

Methods: hUCB-MSCs were collected from the umbilical cord blood. MSCs were cultured in DMEM-low glucose with 10% FBS or 10% aPRP. After inducing EAE in mice, the hUCB-MSCs were transplanted and disease progress was evaluated. Then, the mice were sacrificed, and histological analyses of CNS tissue were done.

Results: The comparison of weight and disease severity between groups showed that there is significant difference between experimental group which received cultured hUCB-MSCs in medium containing aPRP and other groups ($P < 0.05$).

Conclusions: Our findings demonstrated that aPRP supplemented medium not only improves the proliferation but also maintains functional activity of UCBMSCs and it can be a suitable substitution of FBS (Fetal Bovine Serum) in clinical purposes.

Keywords: Experimental allergic encephalomyelitis, Activated platelet-rich plasma (aPRP), Human Umbilical Cord Blood-derived Mesenchymal Stem Cells (hUCB-MSCs), Multiple Sclerosis.

ICIA2018\Stem Cells Based Immunotherapy\Oral\8585
8585

Induction of cytotoxic T lymphocyte responses using dendritic cells transfected with total mRNA of cancer stem cells isolated from patients with gastric adenocarcinoma

Mehran Gholamin¹, Vahid Bagheri², Mohammad Reza Abbaszadegan³

1. Department of Laboratory Sciences, School of Paramedical Sciences, Mashhad University of Medical Sciences, Mashhad, Iran
2. Human Genetic Division, Immunology Research Center, Bu-Ali Research Institute, Mashhad
3. Medical Genetics Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

Background: Cancer stem cells (CSCs) are responsible for tumor initiation, invasion, metastasis, relapse, and resistance to anticancer therapies. Therefore, targeting CSCs, especially using immunotherapy is very critical for cancer treatment. The present report is an evaluation of the immunotherapeutic potential of autologous dendritic cells (DCs) pulsed with total mRNA from gastric CSCs (GCSCs) in vitro for vaccination in GC.

Methods: Tumor samples were enzymatically dissociated. Cells were cultivated in serum-free media under low-adherent conditions. Sphere-forming cells were evaluated by flow cytometry using CD44 and CD54 antibodies. The mRNA levels of CD44 and its isoforms and stemness-related genes in sphere-forming cells were quantitated by comparative real-time PCR. To assess the tumorigenicity, sphere-forming cells injected into nude mice. Monocytes-derived DCs were transfected with autologous total mRNA from GCSCs. After co-culturing DCs with peripheral blood lymphocytes, expression of INF- γ gene and anti-GCSCs activity were investigated in primed lymphocytes.

Results: Primary spheres were observed in cell suspension obtained from four specimens from 36 GC cases. GCSCs expressed surface markers of CD44 and CD54. RT-PCR results indicated that genes of CD44, CD44-v3, -v6 and -v8-10 and stemness factors OCT4, SOX2, SALL4, and Cripto-1 were upregulated in GCSCs. GCSCs were also able to generate tumors in mice. mRNA level of IFN- γ gene increased in stimulated lymphocytes with DCs transfected with total mRNA of GCSCs (6-9 folds). Comparing cytotoxicity assay of primed lymphocytes with antigens of GCSCs isolated from four patients with normal tissue antigens and mock DCs, there was a significant difference in lysis of GCSCs.

Conclusion: The sphere-forming ability is beneficial for the proliferation and enrichment of GCSCs. Considering adverse autoimmune events or other side effects, DCs pulsed with total mRNA from sphere-forming cells that lyse GCSCs by primed lymphocytes in vitro may be utilized as a promising therapeutic vaccination in GC patients in future.

Keywords: gastric cancer, cancer stem cells, sphere-forming cells, dendritic cells

ICIA2018\Stem Cells Based Immunotherapy\Oral\9781
9781

Assessment of phenotype and function of Tregs in an allograft islets and fat derived mesenchymal stem cells composite transplant for experimental autoimmune type 1 diabetes

Bahare Niknam^{1,2}, Jamal Mohammadi Ayenehdeh^{1,3}, Nader Tajik¹, Seyed Mahmoud Hashemi⁴

1. Immunology Research Center, Iran University of Medical Sciences, Tehran, Iran

2. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

3. Department of Immunology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

4. Department Medical Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Islet transplantation is one of the most promising therapeutic approaches in Type 1 Diabetes. In order to improve the viability and function of islet transplantation, it has been proposed to associate pancreatic islets with Adipose Tissue-derived Mesenchymal Stem Cells (AT-MSCs).

Methods: Allograft islets from BALB/c mice co-embedded with syngenic AT-MSCs from abdominal fat tissue of C57BL/6 in hydrogelic composite and delivered in to the peritoneal cavity of Streptozotocin (STZ) induced diabetic C57BL/6 mice. Five groups consist of Control, Hydrogel alone, Gel+MSC, Gel+Islet and Gel+MSC+Islet, delivered into the peritoneal cavity. 32 days after transplantation, Mononuclear cells from the Mesenteric lymph nodes (MLNs) and spleen were analysis For Treg and intracellular cytokine assay by flowcytometry.

Results: Analyses showed that AT-MSCs co-transplanted with allograft significantly increased Treg($P<0.05$) in MLNs but there is no significant variation among the groups in spleen. According to the MFI for IL-10 and TGF- β 1 in MLNs analysis, the IL-10 and TGF- β 1 significantly increased in Gel+MSC and Gel+MSC+Islet groups, in comparison with the other corresponding groups ($P<0.05$). However, in spleen, TGF- β 1 significantly increased in Gel+MSC and Gel+MSC+Islet groups but variation in IL-10 was not significant.

Conclusion: these result shows that AT-MSCs Can promote islet survival and function and can induced functional Treg cells. Moreover, local MSCs transplantation also exert immunomodulatory effects on infiltrating and resident immune cells.

Keywords: Type 1 Diabetes, Mesenchymal stem cells, Treg

ICIA2018\Stem Cells Based Immunotherapy\Oral\10025

10025

Preliminary Results of a Phase I Trial in Patients with Primary Malignant Gliomas using Transfected Mesenchymal Stem Cell with Herpes Simplex-Thymidine Kinase/Ganciclovir Complex

Saeed Oraee Yazdani, Ali Reza Zali

Functional Neurosurgery Research Center, Shohada Tajrish Neurosurgical Comprehensive Center of Excellence, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: The management of patients with glioblastoma remains challenging. Despite aggressive therapies, median survival for malignant gliomas is less than 15 months. Mesenchymal stem cells (MSCs) possess tumor-tropic properties and consequently have been used to deliver therapeutic agents for cancer treatment.

Methods: In this study, six patients with newly diagnosed malignant glioma received autologous bone marrow MSC transfected by Herpes Simplex-Thymidine Kinase (HSV-tk) atvector particles through tumor bed injection 2 weeks after surgery and after confirmation of the pathology. They received ganciclovir as prodrug 14 days after tumor resection. All patients were treated by chemoradiation through or after completion of medication associated gene therapy. They have been following up every six months after intervention with history taking, filling questionnaire, physical examination and imaging. The first case enrolled in the study 15 months ago.

Results: The average time of follow up for patients are 12 months. One patient unfortunately passed away due to deep vein thrombosis and consequent pulmonary emboli that was unrelated to intervention. The rest of patients are living without any evidence of radiological or clinical recurrence. Local or systemic toxicity or increased brain edema did not occur in this study.

Conclusion: These data show that the MSC/HSV-tk/GC complex at the dose used in this study is safe. Long- term follow up and comparing with control group are needed to assess the efficacy of this combination for glioblastoma multiform treatment.



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Tolerance and Autoimmunity

Oral Presentation

ICIA2018\Tolerance and Autoimmunity\Oral\7401

7401

Chitin Micro Particles Induce Regulated T_H1 Response via IL-10 /IFN- γ AxisFatemeh Mami¹, Farshid Yeganeh¹, Elnaz Farahani², Ali Anisian³, Mostafa Haji MollaHoseini¹*1. Department of Immunology, Faculty of Medicine, ShahidBeheshti University of Medical Science, Tehran, Iran**2. Department of Biology, Genetics, Cellular and Molecular group, School of Basic Science, Science and Research Branch of Islamic Azad University, Tehran, Iran**3. Department of Veterinary Pathology, Islamic Azad University, Abhar branch, Abha , Iran*

Background: Our previous studies have shown marked protection among chitin microparticle treated *Leishmania* mice via regulated IL-10/IFN- γ response. Since new contrasting studies are reported on the induction of IL-10 and IFN- γ via Chitin Microparticles (CMPs) during immune stimulation, the present study, once more, examined the inconsistent responses to regarding immunologic response of chitin.

Methods: To verify whether CMPs could indeed up-regulate IL-10 /IFN- γ axis, spleen cells isolated from the MOG35-55-induced experimental allergic encephalomyelitis mice were cultured in the presence of MOG peptide and/or CMPs. The effects of CMPs on IFN- γ , IL-10, and IL-17 production were measured using ELISA. Moreover, T-bet, GATA-3, and ROR γ t expressions (real-time PCR) were investigated.

Results: MOG alone stimulated the production of IFN- γ but not IL-10. MOG/chitin stimulation resulted in a significant increase of IFN- γ and IL-10 levels. Additionally, the expression of T-bet, but not GATA-3, was increased in the CMP-treated spleen cells.

Conclusions: CMP induces GATA-3 independent IL-10 production and promotes T-bet dependent IFN- γ regulation. These results, together with our previous data, may contribute to clarify the adjuvant effect, which has been attributed to CMPs.

Keywords: Chitin; EAE; Immunomodulation

ICIA2018\Tolerance and Autoimmunity\Oral\7445

7445

Lymph node stromal cells control B cell response directed against self-antigen

Reza Nadafi¹, Rogier M. Reijmers¹, Martje Erkelens¹, Eelco D Keuning¹, Lisa G M van Baarsen^{2,3},
Reina E Mebius¹

1. Department of Molecular Cell Biology and Immunology, VU Medical Center, Amsterdam, Netherlands

2. Amsterdam Rheumatology & immunology Center (ARC), Academic Medical Center, University of Amsterdam, Amsterdam, Netherlands

3. Department of Experimental Immunology, Academic Medical Center, University of Amsterdam, Amsterdam, Netherlands

Background: Lymph nodes play a critical role in the induction of immune responses by promoting the interaction between antigen-presenting cells and antigen-reactive lymphocytes. Previous studies have reported that this interaction is additionally controlled by lymph node stromal cells (LNSCs). There is much evidence to support the hypothesis that LNSCs can mediate deletional tolerance of autoreactive CD8⁺ T cells, through presentation of self-antigens in major histocompatibility complex I (MHC class I). However, the properties of LNSCs in controlling autoreactive CD4⁺ T cells through presentation of self-antigens in MHC class II has only been partially addressed by showing the LNSCs dependent maintenance of Tregulatory cells. Here we further determine whether LNSCs can control T follicular helper cells (Tfh) as they play a critical role in the development of humoral immunity by controlling the formation and cellular reactions that occur in germinal centres B cells.

Methods: Here we show that LNSCs control the formation of Tfh cells directed against self-antigen in an antigen dependent manner both in-vitro, by co-culture of LNSCs and CD4 T cells, and in-vivo by lymph node transplantation.

Results: In addition, controlling the Tfh cells directed against self-antigen significantly reduced B cells response for the same self-antigen. The data provides a mechanism by which LNSCs are central to prevent autoantibody production by B cells through the presentation of self- antigens in MHC-class II.

Conclusion: These findings have implications for opportunities to modulate humoral immunity at different stages of autoimmunity.

Keywords: Lymph node stromal cells, T follicular helper, T regulator, B cells, Tolerance, Autoimmunity

ICIA2018\Tolerance and Autoimmunity\Oral\7630
7630

Up-regulation of co-inhibitory genes (PD-1 and Tim-3) by AD-MSCs: implication for immunotherapy of EAE

Adel Mohammadzadeh¹

1. Department of Immunology and Genetics, Faculty of Medicine, Urmia University of Medical Sciences, Urmia, Iran.

Background: Experimental autoimmune encephalomyelitis (EAE) is characterized by central nerve system inflammation that causes axonal myelin sheath damage by autoreactive T cells particularly Th1 and Th17. Impaired hemostasis of immunosystem is the One of the key characteristics of T cells dysfunctionality in this context.

Adipose derived –Mesenchymal stem cells (AD-MSCs) are unique stem cells, as they showed several immunomodulatory features, which have been representing a promising cell-based therapy for autoimmune diseases. Although several mechanisms are shown to be engaged in the process of Immunomodulatory function by MSCs including production of anti-inflammatory cytokines such as TGF- β , it is notable that the main mechanisms are not elucidated yet.

Method: after 72h co-culture of AD-MSCs and PBMC from the spleen of C57BL6 mice, to investigate the mechanisms through which MSCs might modulate T cells, the relative expression of the important co-inhibitory genes i.e. PD-1 (Program cell Death-1) and Tim-3 (T-cell immunoglobulin and mucin-domain containing-3) were analyzed on peripheral blood mononuclear cells (PBMCs), by using quantitative real time-PCR (qRT-PCR).

Results and conclusion: Our results demonstrate that MSCs contribute to T-cell down-regulation in the context of gene modulation of both PD-1 (pv= 0.0012) and Tim-3 (pv= 0.027) pathways and suggest that the understanding of the roles and interactions between MSCs and Immune cells is highly relevant for the development of immunomodulatory drugs and the discovery of biomarkers predictive of therapeutic response.

Keywords: EAE, PD-1, Tim-3, MSCs

ICIA2018\Tolerance and Autoimmunity\Oral\7799
7799

One adjuvant with two different functions: Acylated and deacylated quillaja saponin-21 adjuvants have opposite roles when utilized for immunization of C57BL/6 mice model with MOG₃₅₋₅₅ peptide

Amir Reza Heidari¹, Maryam Rastin², Reza Nosratabadi³, Mahmoud Mahmoudi²

1. Department of Immunology and Allergy, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran
2. Immunology Research Center, Department of Immunology and Allergy, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran
3. Department of Immunology, Immunology of Infectious Diseases Research Center, Rafsanjan University of Medical Sciences, Rafsanjan, Iran

Background: A major of patients with multiple sclerosis (MS) suffer from central neuropathic pain (CNP). Currently, due to interference with motor disability, there are rare experimental autoimmune encephalomyelitis (EAE) models for assessing pain behaviors over the disease course. To address this issue, complete Freund's adjuvant (CFA) was replaced with an acylated triterpene glycoside saponin adjuvant named quillaja saponin-21 (QS-21) in EAE mouse model for CNP development. Deacylated form of QS-21 named QT-0101 has been suggested to have an immunomodulatory effect. Thus, QT-0101 was also used as vaccine adjuvant to understand if it can modulate immune system against myelin oligodendrocyte glycoprotein (MOG₃₅₋₅₅) antigen.

Methods: Thirty two, 5-7 weeks old female C57BL/6 mice were divided into four groups. Except for negative control (PBS) group, other three groups received MOG₃₅₋₅₅ emulsified with CFA, QS-21 and QT-0101 adjuvants, respectively. Thermal hyperalgesia as a CNP clinical manifestation through hot plate test and clinical signs were assessed for two month post immunization (p.i). On day 21 and day 60 p.i mice were sacrificed and TCD4⁺, TCD8⁺, IL-17⁺, IL-4⁺ and CD25⁺/FoxP3⁺ cells in total splenocytes population by flow cytometry technique were evaluated.

Results: Unlike MOG+QT-0101 group, EAE was established in MOG+QS-21 and MOG+CFA groups as mild relapsing-remitting and monophasic models, respectively. Thermal hyperalgesia developed in the bilateral hindpaws on the onset of clinical symptoms in MOG+CFA and MOG+QS-21 groups and it was maintained until study completion in MOG+QS-21 group. TCD4⁺, TCD8⁺ and IL-17⁺ cells population in MOG+QS-21 and MOG+CFA groups, IL-4⁺ and CD25⁺/Foxp3⁺ cells population in MOG+QT-0101 group increased significantly (P<0.05) compared to PBS group.

Conclusion: QS-21 could be a suitable adjuvant in CNP model establishment for future therapeutic researches in MS disease, while QT-0101 seems to have potential for vaccine adjuvant with immunomodulatory property.

Keywords: Multiple sclerosis, Experimental autoimmune encephalomyelitis, Central neuropathic pain, Quillaja saponin-21, QT-0101

ICIA2018\Tolerance and Autoimmunity\Oral\7827

7827

MiR-10a increase expression of ROR γ t and induce proinflammatory T helper 17 cellsZohreh Bolandi^{1,2}, Hossein Ghanbarian^{1,2}, Sara Soudi³, Seyed Mahmoud Hashemi^{4*}

1. Department of Biotechnology, School of Advanced Technologies in Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Cellular and Molecular Biology Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

3. Department of Immunology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

4. Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: MicroRNAs (miRNAs) are endogenous, short non-coding RNA molecules (18–24 nucleotides) that regulate target gene expression by inhibiting translation and increasing mRNA degradation. Recently, miRNAs have been revealed to play a major role in T cell activation, proliferation and differentiation, therefore, by promoting the development of inflammatory T cells including the T helper 17, miRNAs can involve in autoimmune disorders. The aim of this study was to investigate the role of miR-10a in the regulation of ROR γ t expression, the key transcription factor in the generation of Th-17 cells, in mouse splenocytes population.

Methods: Splenocytes were isolated from 6-8 weeks old C57Bl/6 female mice by mincing the spleen with a syringe plunger followed by lysis of erythrocytes with 2mL of lysis buffer. Splenocytes were stimulated with anti-CD3/CD28mAbs. Next day, splenocytes were transfected with miR-10a mimic using Lipofectamine 2000 according to the manufacturer's protocol. After 48 hrs, cells were harvested and washed twice with PBS. RNA was extracted and Real-time PCR was performed to quantify the relative expression levels of miR-10a, ROR γ t, and IL-17 in transfected cells compared to nontransfected control.

Results: To overexpress of miR-10a in mouse splenocytes, miR-10a mimic was transfected via Lipofectamine 2000 into the splenocytes. Total RNA was extracted and Real-time PCR assay showed that miR-10a was significantly overexpressed in transfected cells and following the overexpression of the miRNA, the expression of ROR γ t and IL-17 increased in mouse splenocytes.

Conclusion: Here, we indicated that miR-10a could induce ROR γ t transcription factor expression that have an important role in Th-17 differentiation. It can be concluded that the overexpression of miR-10a leads to T cell differentiation into Th-17 cells and producing the IL-17, an important proinflammatory cytokine that has an important role in the induction of various autoimmune diseases.

Keywords: Splenocytes, miR-10a, Th-17, ROR γ t.

ICIA2018\Tolerance and Autoimmunity\Oral\8236

8236

Induction of Regulatory T cell by Monocyte Derived DC after Differentiation with Tolerogenic Probiotics in SLE Patients

Seyed Alireza Esmaeili^{1,2,3}, Mahmoud Mahmoudi^{1,2}, Fatemeh Mardani^{1,2,3}, Sahar Khorasani^{1,2,3}, Zohreh Vahidi^{1,2,3}, Nafiseh Tabasi¹, Mahdieh khazaei¹, Maryam Rastin^{1,2}

1. Immunology Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

2. Immunology Department, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

3. Student Research Committee, Mashhad University of Medical Sciences, Mashhad, Iran

Background: Systemic lupus erythematosus (SLE) is a main autoimmune disease that uncontrolled inflammation lead to tissue damage, therefore control of excessive inflammation is necessary. Dendritic cells (DC) are determinative cell to shape direction of regulatory or inflammatory immune response. Inhibitory DC shift naïve T cell toward the Regulatory T cell (Treg) that could decrease inflammation by production of regulatory cytokines, also help to keep immune hemostasis and return of inflammatory T cells population balance. The aim of this study was induction of the regulatory T cell in co-culture with regulatory DC after producing trough tolerogenic probiotics in in-vitro treatment.

Methods: Monocytes of SLE patient cultured with GM-CSF and IL-4 cytokines for 5 days, then immature DC cultured with Lactobacillus delbrueckii and lactobacillus rhamnosus along with same concentration of both cytokines for two days in order to maturation of DC. The Phenotypes of generated DC examined by Flowcytometry and Real-time PCR. Naive CD4⁺ T cell isolated through negative isolation kit. Generated regulatory DC and Naive CD4⁺ Tcell were cultured in RPMI media for 5 days to induction of Treg cells and finally properties of regulatory T cell evaluated by Flowcytometry and Real-Time PCR methods.

Results: We observed the generated regulatory DC showed reduction in expression of the surface co-stimulatory molecules and elevation of anti-inflammatory cytokines and enzyme genes. Following, we could show significant increase of CD4⁺CD25⁺CD127⁻ FOXP3⁺ cell populations and increase of FOXP3⁺ transcription factor and TGF- β cytokine but not IL10.

Conclusion: Our finding revealed that probiotic-induced regulatory DC could induce regulatory T cells and related inhibitory cytokine. Therefore, in vivo production of Treg cells trough regulatory DCs could be beneficial strategy for prevention or treatment of SLE disorders.

Keywords: Systemic Lupus Erythematosus (SLE), Probiotics, Dendritic Cell (DC), Lactobacillus, Regulatory T Cells (Treg)

ICIA2018\Tolerance and Autoimmunity\Oral\8264
8264

MicroRNA-92a drives Th1 responses in the experimental autoimmune encephalomyelitis

Nahid Rezaei¹, Farideh Talebi², Samira Ghorbani², Abbas Rezaei¹, Farshid Noorbakhsh^{2*},
Mazdak Ganjalikhani Hakemi¹

1. Department of Immunology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

2. Department of Immunology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

Background: IFN- γ -producing Th1 cells are the main players in several autoimmune diseases, including multiple sclerosis (MS) and its animal model, experimental autoimmune encephalomyelitis (EAE). Despite the fact that, Th1 fate is determined by specific transcription factors, the role of microRNAs is dispensable. MicroRNAs dysregulation have been linked to the progress of a number of autoimmune disease including MS. However, little is known about the role of miR-92a, one of the most upregulated miRNAs in MS, in the development of autoimmunity. Based on the remarkable change in CNS of patients with MS, miR-92a was selected for subsequent experiments to ascertain the possible association between the overexpression of miR-92a and immunopathogenesis of MS.

Methods: The expression level of miR-92a was assessed in the CNS tissues and splenocytes from mice with EAE using real-time RT-PCR. Then, the possibility that miR-92a is involved in the Th1 polarization was investigated by flow cytometric analysis. Additionally, the expression level of related targets was explored.

Results: Enhanced miR-92a expression in mouse samples with EAE at the peak of disease was accompanied with reduced expression level of TSC1 and DUSP10. Over expression of miR-92a in splenocytes led to more Th1 differentiation than that in cells transfected with negative control.

Conclusion: Our data discovered that upregulation of miR-92a by ex vivo transfection leads to elevated Th1 differentiation due to silencing of TSC1 and dusp10. Hence, regulation of miR-92a expression, which in turn modulates Th1 responses, may be a potential new therapeutic target in MS.

Keywords: EAE, miR-92a, Th1

ICIA2018\Tolerance and Autoimmunity\Oral\8273

8273

The Effect of Lactobacillus Probiotics on Frequency of Th1 and Th17 Cells in Mouse Model of SLE Induced by Pristane

Fatemeh Mardani^{1, 2}, Seyed Alireza Esmacili^{1, 2}, Sahar Khorasani^{1, 2}, Fahimeh Lavi^{1, 2}, Nafiseh Tabasi³, Mahmoud Mahmoudi¹, Maryam Rastin¹

1. Immunology Research Center, BuAli Research Institute, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

2. Student Research Committee, Mashhad University of Medical Sciences, Mashhad, Iran

3. Immunology Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

Background: Systemic lupus erythematosus (SLE) is an important autoimmune disease and uncontrolled inflammation engaged multiple organs. Th17 cells are the major inducer of inflammation in SLE and IL-17 cytokine is the main Th17 cell mediator that responsible in recruitment and activation of inflammatory immune cells therefore, suppressing these cells is a major therapeutic strategy in SLE. Tolerogenic probiotics are able to modulate the activated immune system and decrease the level of inflammation in SLE and other autoimmune diseases. The aim of this project is evaluation the effects of lactobacillus rhamnosus and delbrueckii on Th17 and their mediators in Pristane induced Balb/C mice.

Methods: SLE induced in mice by injection of Pristane to establishment of disease. Animals divided into two groups including pretreatment and treatment. probiotics and prednisolone receiving groups feeds daily for sixth month, then the level of Anti-nuclear (ANA) , Anti – dsDNA and anti RNP antibodies measured by immunofluorescent and ELISA methods in mice sera. After six months the mice killed and Frequency of Th1 and Th17 cells evaluated by flow cytometry method as well as The expression of ROR γ t and IL-17 measured by real-time PCR. Also INF- γ , IL-17 and IL-10 cytokines measured by ELISA techniques.

Results: We observed reduction the level of serum ANA, Anti-dsDNA and Anti-RNP and also delay onset SLE disease along with decrease of mass of lipogranuloma in probiotic and prednisolone receiving groups in bothpretreatment and treatment mice. Treatment with Probiotics and prednisolone decreased the frequency of both Th1 and Th17 cells as well as the amount of INF- γ , IL-17 cytokines and ROR γ t transcript while no change in the level of IL-10 has seen.

Conclusion: The results of current study showed that feeding with lactobacillus rhamnosus and delbrueckii can reduce the frequency of Th1/Th17 cells and suppress production of IFN gamma and IL-17 in mouse model of SLE, and therefor using probiotics may be an alternative way for alleviating the inflammation in SLE and possibly other TH1/Th17 mediated immune disorders.

Keywords: SLE, Lactobacillus, Tolerogenic Probiotics, Th1/Th17, Animal Model

ICIA2018\Tolerance and Autoimmunity\Oral\8597
8597

Neat-1 LncRNA regulates inflammation and T cell differentiation in an animal model of multiple sclerosis

Farimah Masoumi¹, Samira Ghorbani¹, Farideh Talebi¹, Farshid Noorbakhsh^{*1}

1. Immunology Department, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

Background: Despite growing evidence that Long noncoding RNAs (lncRNAs) can widely take part in autoimmune diseases, our knowledge of Multiple Sclerosis-related lncRNAs remains limited. Herein, we investigated the potential role of Neat-1 lncRNA, in the context of autoimmune neuroinflammation.

Methods: The expression level of Neat-1 was measured in the CNS tissues from mice with experimental autoimmune encephalomyelitis (EAE). Expression analyses were also performed in mitogen and antigen-stimulated splenocytes, as well as activated macrophages using real-time RT-PCR. To examine the role of Neat-1 in macrophages polarization, Neat-1 siRNA was transfected into primary macrophages followed by M1/M2 macrophage polarization. Then M1/M2 genes expression were evaluated. Also, the role of the Neat-1 in T cell differentiation was investigated by transfection of CD4⁺ T cells with Neat-1 siRNA, followed by flow cytometric analysis of intracellular cytokines. Moreover, effect of Neat-1 downregulation on T cells proliferation was investigated using CFSE staining.

Results: Expression of Neat-1 was significantly increased in the spinal cords of EAE mice at days 15 and 25 post disease induction. Splenocytes stimulated with MOG peptide or anti-CD3/anti-CD28 antibodies showed significant upregulation of Neat-1, whereas expression of Neat-1 in stimulated bone marrow-derived macrophages was reduced. Neat-1 down regulation increased proinflammatory gene expression in macrophages and polarization toward M1 phenotype. Also, Neat-1 downregulation in activated lymphocytes shifted the pattern of T cell differentiation towards Th1 cells. However, T cell proliferation remained unchanged following Neat-1 downregulation.

Conclusion: Our data highlight the anti-inflammatory actions of Neat-1 in the context of autoimmune neuroinflammation. Neat-1 influence differentiation of T cells and activation of macrophages, providing potential therapeutic options for controlling inflammation in MS.

Keywords: LncRNA, Neuroinflammation, Multiple sclerosis, T cell differentiation



14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

Transplantation

Oral Presentation

ICIA2018\Transplantation\Oral\5252

5252

Down-regulation of inflammatory signaling pathways despite up-regulation of Toll-like receptors; the effects of corticosteroid therapy in brain-dead kidney donors, a double-blind, randomized, controlled trial

Reza Jafari¹, Reza Aflatoonian², Reza Falak¹, Gholamreza Pourmand³, SanazDehghani³, Mojgan Mortazavi⁴, Adeleh Adelipour³, Abbas Rezaei⁵, Nader Tajik¹

1. Immunology Research Center (IRC), Iran University of Medical Sciences, Tehran, Iran

2. Department of Endocrinology and Female Infertility at Reproductive Biomedicine Research Center, Royan Institute for Reproductive Biomedicine, ACECR, Tehran, Iran

3. Urology Research Center, Tehran University of Medical Sciences, Tehran, Iran

4. Isfahan Kidney disease research center, Isfahan University of Medical Sciences

5. Department of Immunology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

Background: The brain death of a potential organ donor induces a systemic inflammatory response, resulting in inferior organ quality and function. Our study aimed to evaluate the effects of methylprednisolone (MPN) therapy on pattern recognition receptor (PRR) signaling in potential brain-dead (BD) kidney donors.

Methods: To evaluate the effects of MPN therapy on PRR signaling in BD kidney donors we performed a prospective randomized treatment-versus-control study. Fifty-one potential kidney donors were randomly divided into three groups: brain-dead donors (BDDs) who received 15mg/kg/d of methylprednisolone (group T1, n=17), BDDs who received 15mg/kg/d of MPN at the time of filling consent for kidney donation and 100mg/2h until kidney harvest (group T2, n=17), and normal donors as controls n=17. Gene expression for Toll-like receptors (TLRs) 1-9 and their signaling pathway molecules including MYD88, TRIF, NF-KB1, IRAK, IRF3, and IRF7, as well as the inflammatory cytokines RANTES, IL-1 β , TNF- α , IL-6, CXCL8, IL-18, IFN- α , and IFN- β was determined by PCR array. Due to the crucial role of TLRs 2 and 4 in pattern recognition, surface expression of these molecules was analyzed by flow cytometry. Plasma levels of inflammatory cytokines were measured by immunoassay. Finally, serum creatinine and cystatin C were measured in 100 kidney recipients one week and one, three, and six months after transplant.

Results: Polymerase chain reaction (PCR) array gene expression revealed greater expression of TLRs and signaling molecules in group T1 than in the controls. Surface expression of TLRs 2 and 4 were significantly greater in group T2 than in group T1 ($P<0.05$). Plasma concentrations of inflammatory cytokines were significantly greater in group T1 than in controls ($P<0.05$). The recipients that received kidneys from group T1 had significantly higher levels of creatinine and cystatin C than the recipients of kidneys from both group T1 and controls ($P<0.05$).

Conclusion: Administration of MPN to BDDs at specified periods until kidney harvest resulted in less systemic inflammation in the BDDs and improved renal function in kidney graft recipients compared with common MPN therapy.

Keywords: Innate immunity, Inflammation, Brain-dead donors, Toll-like receptors

ICIA2018\Transplantation\Oral\7396

7396

Increased expression of TGF- β , SMAD3 and miR-21 in renal transplant recipients that developed allograft dysfunction: a cohort study

Farshad Foroughi¹, Pedram Ahmadpoor², Morteza Hosseinzadeh³, Sanaz Keshavarzshahbaz^{4,5}, Mehri Barabadi⁴, Mir Saeed Yekaninejad⁶, Mina Ghorbanpour⁶, Ehsan Soltaninejad⁴, Farshid noorbakhsh⁴, Aliakbar Amirzargar^{4,7}

1. Department of Immunology, School of Medicine, Qazvin University of Medical Sciences, Qazvin, Iran
2. Department of Nephrology, Shahid Labbafinejad Medical Center and Chronic Kidney Disease Research Center, Shahid Beheshti
3. University of Medical Sciences, Tehran, Iran
4. Department of Immunology, School of Medicine, Ilam University of Medical Sciences, Ilam, Iran
5. Department of Immunology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran
6. Department of Immunology, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran
7. Department of Epidemiology and Biostatistics, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran
8. Molecular immunology research center, Tehran university of medical sciences, Tehran, Iran

Background: Chronic allograft nephropathy is the most important factor in long term survival of kidney transplantation. Leukocyte infiltration into the graft and profibrotic gene expression have pivotal effects on kidney transplantation outcome. The present study sought to determine the expression of sequential TGF β , SMAD3, miR-21 and miR-29 gene expression in human renal allograft.

Methods: Blood samples from 52 consecutive renal transplant patients were evaluated at the time of transplantation and at two times (90 and 180 days) after transplantation to analyze the gene expression of TGF β , SMAD3, miR-21 and miR-29 by real time PCR. A total of 30 biopsies, including protocol biopsy (n=24) and cause biopsy (n= 6), were investigated according to the Banff criteria.

Results: The gene expression of TGF β , SMAD3 and miR-21 were found to be significantly higher in graft dysfunction than well-functioning grafts ($p<0.001$, $p=0.02$, $p<0.001$), while the expression of miR-29 was lower significantly in graft dysfunction ($p<0.001$). Receiver Operating Characteristic (ROC) Curve Analysis showed that the calculated AUC was 0.77 and 0.83 at the third month for TGF β and miR-21 respectively ($P=0.001$, $p<0.001$). Multiple logistic regression analysis showed that an increase in TGF β and miR-21 expression leads to higher risk of graft dysfunction (OR=1.96, OR=4.19).

Conclusion: During renal transplantation, TGF β , Smad3 and miR-21 gene expression increased in graft dysfunction subjects. Performing such measurements by using non-invasive techniques that include real-time PCR, a comparison of their results with the pathology findings and the patient's clinical condition may assist with the discovery of appropriate cellular and molecular markers for patients with kidney transplants. These findings may prospectively predict allograft dysfunction and help elucidate the underlying pathogenic mechanisms.

Keywords: kidney transplantation, graft dysfunction, miRNA, TGF β

ICIA2018\Transplantation\Oral\7503

7503

Urinary Kidney injury molecule-1 (KIM-1) mRNA analysis in kidney transplant recipientsSanaz Keshavarz Shahbaz^{1,2*}, Mehri Barabadi¹, Farshad Foroughi³, Aliakbar Amirzargar^{2,4}*1. Department of Immunology, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran**2. Department of Immunology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran**3. Cellular and Molecular Research Center, Qazvin University of Medical Sciences, Qazvin, Iran**4. Molecular Immunology Research Center, Tehran University of Medical Sciences, Tehran, Iran*

Background: The Occurrence of acute rejection episode as the outcome of renal transplantation is so difficult to predict, even with an allograft biopsy. Kidney injury molecule-1 (KIM-1), a transmembrane glycoprotein demonstrates the considerable expression in dedifferentiated cells within damaged regions of the proximal tubule but it does not express in normal renal tissue. Hence, we hypothesized that KIM-1 might serve as biomarkers for predicting early acute rejection after kidney transplantation.

Methods: In this study, 30 kidney transplant recipients were studied. Urine samples at 3 times (1 day, 3 months and 6 months after transplantation) were taken. RNA extraction, cDNA synthesis and mRNA expression of KIM-1 gene by RT-PCR were performed.

Results: The KIM-1 gene expression in urine samples at three and six months after transplant increased significantly in comparison with stable graft function patients ($p = 0.01$).

Conclusion: These data suggests that urinary KIM-1 mRNA could be used as promising markers for prediction of early acute kidney allograft rejection.

Keywords: KIM-1, Renal Transplant recipient, acute renal allografts Rejection.

ICIA2018\Transplantation\Oral\9857

9857

Natural killer cells exhibit an activated phenotype in peripheral blood mononuclear cells of renal allograft rejection recipients: a preliminary study

Azita Talebzadeh Bonakdar¹, Mohammad Mahdi Mohammadi¹, Afshin Namdar², Pedram Ahmadpoor³, Davood Jafari², Arron Mungela Foma⁴, Mir Saeed Yekaninejad⁵, Ali Akbar Amirzargar^{2,4}

1. Department of Immunology, School of Medicine, Kerman University of Medical Sciences, Kerman, Iran

2. Department of Immunology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

3. Chronic kidney disease research center, Labbafinejad Medical Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

4. Molecular Immunology Research Center, Tehran University of Medical Sciences, Tehran, Iran

5. Department of Epidemiology and Biostatistics, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

Background: A growing body of evidence has revealed the role of innate immune cells in transplantation; however, the nature of natural killer (NK) cells involvement in rejection is still elusive. The purpose of the current study is to determine the impact of NK cell activities in acute and chronic renal transplant rejection.

Methods: This preliminary case-control study comprised 63 participants, of which 19 were patients with kidney allograft rejection (8 patients with acute rejection (AR) and 11 patients with chronic rejection (CR)) and 44 were the control groups, comprising 22 patients who received kidney transplant with well-functioning graft (WFG) and also 22 healthy subjects. In addition to the frequency of NK cells, the intracellular IFN- γ production and surface expression of CD107a as cytotoxic activity were measured using flow cytometry.

Results: Our results revealed a significant increase in CD107a expression ($P=0.021$) in patients with AR in comparison with WFG. Moreover, there was a significant rise in the production of IFN- γ in patients with CR when compared to WFG group ($P=0.003$). Finally, a decrease in the frequency of NK cells in rejection patients was observed compared with control groups; however, this reduction was not statistically significant.

Conclusion: These findings suggest that the increase in NK cell cytotoxicity is correlated with rejection in patients with kidney transplantation and might be consider as a predictive marker in prevalence of graft rejection.

Keywords: Kidney transplant rejection, Natural killer cell, CD107a, IFN- γ , Cytotoxicity.



Vaccine

Oral Presentation

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Vaccine\Oral\7540

7540

Plant base production of virulence factors from Diarrheagenic Escherichia coli; An insight into its immunological evaluation in animal model

Fatemeh shojaei jeshvaghani¹, Jafar Amani², Mahyat Jafari¹, Ali Hatef Salmanian¹

1. Department of Agricultural Biotechnology, National Institute of Genetic Engineering and Biotechnology (NIGEB), Tehran, Iran

2. Applied Microbiology Research Center, Baqiyatallah Medical Science University, Tehran, Iran

Background: The most important infectious agents responsible for human enteric disorders include several viruses and bacterial, such as Diarrheagenic Escherichia coli. Among this group, EHEC and ETEC cause the largest number of diarrheal cases. In humans, EHEC infections result in bloody or non-bloody diarrhea, which may be complicated by haemorrhagic colitis and haemolytic uraemic syndrome (HUS). Infection by ETEC is accompanied by a non-inflammatory watery diarrhea. We hypothesized that the chimeric recombinant forms of the antigenic moiety of these bacteria, which delivered via edible route, could reduce colonization of infectious agent.

Methods: In our study a synthetic gene composed of cfa, eae, stx and It (SICL) attached by linkers was constructed and codon optimized for expression in plant host. The chimeric gene was cloned in the binary vector pBI1400 containing fatty acid elongase (FAE) promoter as a seed-specific promoter. The gene was then transferred to canola plants by agrobacterium mediated protocol. Three mice groups were used in the immunization program, one group fed by whole seed of transgenic canola plant, the second immunized with combination of oral and parenteral administration and the last group fed by nontransgenic canola plant as a control. Blood samples were taken from mice after last immunization for analysis of SICL specific IgG and IgA antibodies.

Results: The chimeric protein showed considerable percentage of total soluble protein in transgenic canola seed. Our studies demonstrate that oral feeding alone or combination with injection can equally develop SICL specific immune response. No antibody response was detected in the mice that were immunized with nontransgenic plant.

Conclusion: We demonstrated that the application of transgenic plants containing recombinant protein could act as an effective tool for the production of a vaccine candidate. Furthermore, the oral delivery of antigen could protect the animal model against bacterial challenge.

Keywords: Diarrheagenic Escherichia coli, transgenic plants, vaccine candidate, immune response.

ICIA2018\Vaccine\Oral\7658

7658

Th1-biased immune responses elicited by Plasmodium falciparum thrombospondin related adhesive protein and poly (I: C) adjuvant in BALB/c mice

Maryam Ameri Torzani^{1,2}, Akram Abouie Mehrizi^{1*}, Sedigheh Zakeri¹, Laleh Babaekhou²

1. Malaria and Vector research Group, Biotechnology Research Center, Pasteur Institute of Iran, Tehran, Iran

2. Department of Biology, Islamshahr Branch, Islamic Azad University, Islamshahr, Iran

Background: Plasmodium falciparum thrombospondin related adhesive protein (PfTRAP) is a leading malaria vaccine candidate antigen on sporozoite. However, the recombinant malaria vaccine candidate antigens are poor immunogenic and additional appropriate immune stimulants, such as an efficient adjuvant, are highly needed to enhance protective Th1 type immune responses. In this regard, polyinosinic: polycytidylic acid (poly (I: C)), was considered as the potential adjuvant and the immunogenicity of PfTRAP formulated with this adjuvant was evaluated in BALB/c mice.

Methods: The pftrap gene was codon optimized based on the E. coli codon usages and cloned in NdeI-XhoI sites of pET24a plasmid. The recombinant PfTRAP antigen was expressed in E. coli BL21 (DE3) and purified by using Ni-NTA agarose. Three test groups of BALB/c mice were immunized with rPfTRAP antigen alone or in combination with poly (I:C) or CFA/IFA adjuvants. The control groups received PBS1× or corresponding adjuvants without antigen. Immunization was carried out three times, 2 weeks intervals via subcutaneous rout. Anti-PfTRAP IgG and IgG subclasses were measured 10 days after each immunization.

Results: rPfTRAP antigen formulated with poly (I: C) elicited high levels of anti-rPfTRAP IgG antibodies comparable to CFA/IFA adjuvant after the second boost. In addition, rPfTRAP formulated with poly (I: C) elicited the higher ratio of IgG2a/IgG1, and IgG2b/IgG1 than the mice immunized with this antigen emulsified in CFA/IFA.

Conclusion: The present investigation revealed that the rPfTRAP delivery in poly (I: C) had potential to modulate Th1 immune responses against rPfTRAP antigen.

Keywords: Plasmodium falciparum, thrombospondin related adhesive protein, Adjuvant, malaria vaccine

ICIA2018\Vaccine\Oral\7661

7661

Evaluation of immune responses to novel *Plasmodium vivax* circum sporozoite based vaccine candidates in combination with second generation adjuvants in C57BL/6 mice

Samaneh H. Shabani^{1,2}, Sedigheh Zakeri^{1*}, Akram A. Mehrizi¹, Yousef Mortazavi²,
Navid D. Djadid¹

1. *Pasteur Institute, Iran*

2. *Zanjan University of Medical Sciences, Iran*

Background: The majority of malaria vaccine studies have mainly focused on *Plasmodium falciparum*, but *Plasmodium vivax* have been ignored. However, *P. vivax* is able to develop hypnozoites, which increases the morbidity caused by a single infection and severe malaria. Additionally, a vaccine targeting of *P. vivax* represents a necessary tool when considering the elimination/eradication goal. Many efforts have been conducted to design a vaccine based on circum sporozoite protein (CSP). Because of limited immunogenicity and less effectiveness of protein-based vaccines, these types of vaccines require adjuvant to induce a protective and long-lasting immune response. In this investigation, we evaluate the immunological potency of two newly designed PvCSP based vaccines in combination with a novel adjuvant system (AS).

Methods: Both CS127 and CS712 constructs include N- and C-terminal parts and a truncated region containing repeat sequences with different arrangement from both PvCSVK210 and PvCSVK247 subtypes. After expression, purification, desalt and concentration, constructs were formulated with novel AS (NLX, CpG and QS21). 6-8 weeks female C57BL/6 mice were immunized with 3 boosts with 2 week interval. Humoral responses include specific antibodies and subclasses against PvCSP, titration and avidity of antibodies, and cellular responses includes lymphocyte proliferation assay and cytokine profiles were evaluated with ELISA, MTT and cytokine assay, respectively.

Results: Our results show that both constructs are highly immunogenic in C57BL/6 mice. Both candidates in combination with AS induce high levels of antibody against PvCSP with high titration and avidity of Th1 related antibodies. Analysis of the induced T cells highlighted different cytokine profile with significant secretion of IFN- γ and Th1 responses.

Conclusion: These results need further clinical investigation of these two candidates in primate models to reach the added value in both immunogenicity and protective efficacy.

Keywords: *Plasmodium vivax*, malaria; adjuvant system, circum sporozoite protein

ICIA2018\Vaccine\Oral\7856

7856

A novel endogenous adjuvant for development of therapeutic vaccine against HIV-1 infection**Alireza Milani¹, Azam Bolhassani¹***1. Department of Hepatitis and AIDs, Pasteur Institute of Iran, Tehran, Iran*

Background: Vaccination is currently considered as an additional therapeutic approach to stimulate HIV-specific immune responses. In order to development of an effective vaccine against HIV-1 infection, novel vaccine strategies are required. Promising results of DNA-based, protein-based and heterologous prime/boost-based strategies showed their influence on eliciting both humoral and cellular immune responses. Recently, the endogenous adjuvants such as heat shock proteins (HSPs) have been suggested effectively to induce antigen-specific humoral and cellular immune responses.

Methods: The recombinant HIV-1 Nef, and Nef fused to Hsp27 DNA and also protein constructs were generated in eukaryotic and prokaryotic expression systems, respectively. Mice were immunized with these constructs based on three strategies of DNA/DNA, Protein/protein and DNA/Protein. To determine the induction of immune response, sera and splenocytes were analyzed for humoral and cellular responses, respectively.

Results: Analysis of the immune responses indicated that the Hsp27-Nef fusion protein significantly increased the Nef-specific T cell responses. Indeed, this regimen induced high levels of IgG2a and IFN- γ directed toward Th1 responses and also Granzyme B secretion compared to other immunization strategies. Moreover, the immunostimulatory properties of Freund's adjuvant were significantly less than Hsp27 in different immunization strategies.

Conclusion: These data demonstrated that the use of Hsp27 in protein-based strategy could improve HIV-1 Nef-specific B- and T-cell immune responses as a promising HIV-1 vaccine candidate in future.

Keywords: Therapeutic HIV vaccine, Nef, Endogenous adjuvant, Small heat shock protein 27

ICIA2018\Vaccine\Oral\8042

8042

Characterization and evaluation of DNA plasmids coding for immunogenic salivary proteins of Phlebotomus sergentias vaccine candidates against Leishmania tropica in a BALB/c mice model

Elham Gholami^{1,2}, Tahereh Taheri², Negar seyed², Sima Habibzadeh², Yasaman Taslimi²,
Mohammad Ali Mazloomi¹, Shaden Kamhawi³, Jesus G Valenzuela³, Fabiano Oliveira³, Sima Rafati²

1. School of Advanced Technologies in Medicine, Tehran University of Medical Sciences, Tehran, Iran

2. Department of Immunotherapy and Leishmania Vaccine Research, Pasteur Institute of Iran, Tehran, Iran (the Islamic Republic of)

3. Vector Molecular Biology Section, Laboratory of Malaria and Vector Research, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Rockville, Maryland, United States of America

Background: In the leishmaniasis, phlebotomine (Ph.) sand flies transmit Leishmania to a mammalian host through depositing the parasite in the presence of their saliva in the skin during feeding. Cutaneous Leishmaniasis is caused by *L. tropica* and *L. major* in Iran, being mostly transferred to the host with *Ph. sergenti* and *ph. papatasi*, respectively. There is no human vaccine for this disease. According to the reports on animal experiments, a pre-exposure to sand fly saliva results in protection against leishmaniasis. Furthermore, it has been reported that *Ph. papatasi* salivary protein SP15 (PpSP15) provides protection against *L. major* infection. Here, we have analyzed the immune response to distinct salivary proteins of *Ph. sergenti* in the BALB/c mice model.

Methods and Results: Results of immunization experiments using 14 DNA plasmids coding for *Ph. sergenti* salivary proteins show that PsSP9 (14.05-kDa) induces protection against *L. tropica* infection. Forty-eight hours after *Ph. sergenti* salivary gland homogenate (SGH) injection, PsSP9 immunized mice exhibited a DTH response and a low antibody response against SGHs, and was associated with a high ratio of IFN γ /IL5 expression levels in the draining Lymph node. These mice exhibited a small ear swelling as well as a low parasite load after *L. tropica* infection. IFN- γ induction after *Ph. sergenti* SGH exposure could be partially responsible for the provided protection in the mice immunized with PsSP9, via anti-Leishmania immunity priming and/or direct parasite killing.

Conclusion: Our work suggests that PsSP9, a member of the PpSP15 family, is a salivary protein that can provide protection against *L. tropica* infection and it also suggests that this family of proteins in *Ph. sergenti*, *Ph. duboscqi* and *Ph. papatasi* may have similar immunogenic and protective properties against different species of Leishmania. Our data imply the beneficial effect of applying arthropod saliva components in vaccination strategies for diseases caused by vector-borne pathogens.

Keywords: Phlebotomus sergenti, Salivary Gland antigen, Leishmania tropica, DTH, vector-borne diseases, vaccine

ICIA2018\Vaccine\Oral\8416

8416

Investigation of the cellular and humoral immune responses to recombinant fragments of filamentous hemagglutinin and pertactin of *Bordetella pertussis* in BALB/c mice

Peyman Bakhshaei¹, Mohammad Hossein Kazemi^{1*}, Maryam Golar¹, Sara Abdolmaleki¹, Roya Khosravi Eghbal¹, Jalal Khoshnoodi¹, Mohammad Ali Judaki¹, Vahid Salimi², Masoumeh Douraghi³, Mahmood Jeddi Tehrani⁴, Fazel Shokri^{1,4}

1. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

2. Department of Virology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

3. Department of Microbiology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

4. Monoclonal Antibody Research Center, Avicenna Research Institute, ACECR, Tehran, Iran

Background: The immune response to acellular vaccines has been generally associated with a shift towards the Th2 profile. In the present study the cellular and humoral immune responses of BALB/c mice to recombinant fragments of filamentous hemagglutinin (FHA), pertactin (PRN) and native pertussis-toxin (PT) has been investigated through measuring IFN- γ , IL-4 and IL-17 cytokines and also specific antibody response to antigens following immunization with different formulations of the antigens.

Methods: Two and four overlapping recombinant fragments of PRN and FHA, respectively, were produced in E-coli and purified by affinity chromatography, while PT was purified from bacterial suspension of *Bordetella pertussis*. The purity and concentration of antigens were determined by SDS-PAGE and BCA assays, respectively. BALB/c mice were immunized twice with different combinations of antigens together with CpG or alum as adjuvant. Immunized mice were subsequently aerosol challenged with *B. pertussis*. Bacterial growth was assessed in Broncho-alveolar lavage (BAL) samples by CFU-assay. The levels of cytokines in supernatants of antigen-stimulated splenocytes and also the antigen-specific antibody titers in serum were measured by ELISA.

Results: The levels of IFN- γ , IL-4 and IL-17 cytokines in supernatants of all immunized mice were significantly high. The level of IFN- γ was higher in response to CpG formulated antigens. A substantial specific antibody response was observed to both PRN and FHA antigens in serum of all immunized mice, particularly those immunized with all antigens. No bacterial colony was observed in culture of BALs of vaccinated mice.

Conclusion: It seems that all combinations of our recombinant FHA and PRN antigens could effectively induce Th1, Th2 and Th17 immune responses as well as specific antibody response leading to bacterial clearance from lung of immunized mice. Formulation of the antigens with CpG induced substantially higher Th1 response.

Keywords: acellular pertussis vaccine, antibody response, cytokine response, filamentous hemagglutinin, pertactin, *Bordetella pertussis*

ICIA2018\Vaccine\Oral\8582

8582

**Generating recombinant *Lactococcus lactis* expressing SP15 of *Phlebotomus papatasi*
As a live nonpathogenic vaccine**

E. Davarpanah^{1,2}, N. Seyed², J. G. Valenzuela³, S. Rafati², R. Safaralizadeh¹, T. Taheri²

1. Department of Biology, Faculty of Natural Sciences, University of Tabriz, Tabriz, Iran.

2. Department of Immunotherapy and Leishmania Vaccine Research, Pasteur Institute of Iran, Tehran, Iran.

3. Vector Molecular Biology Section, LMVR, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Rockville, Maryland, United States of America

Background: Cutaneous leishmaniasis is a major public health problem that is transmitted by biting of *Phlebotomus papatasi* (*P. papatasi*) sandfly. There is no effective drug therapy and vaccination methods against leishmaniasis. *Lactococcus lactis* (*L. lactis*) is a nonpathogenic gram-positive bacteria normally used in dairy industry. We used *L. lactis* as a live host for expression of SP15 protein as a cell wall secretory component from *P. papatasi* saliva.

Methods: Two genes SP15-EGFP and EGFP were synthesized after codon optimization for heterologous gene expression in *L. lactis*. Both genes were cloned in a specific vector in downstream of signal peptides PrtP that is responsible for expression as cell wall secretory. The protein expression was confirmed by western blotting. Then, different groups of BALB/c mice were immunized with recombinant *L. lactis*. After challenging of immunized mice with *L. major* plus Salivary Gland Homogenate (SGH), footpad swelling, cellular and humoral immune response and also nitric oxide were measured. Furthermore, parasite load in lymph nodes (LN) performed using real-time PCR at different time points.

Results: DNA fragments were codon optimized based on *L. lactis* genome that has 35.24% GC content, synthesized and subcloned in pNZ8121 to express secretory SP15 protein on bacterial cell wall. Immunized mice with recombinant *L. lactis*-SP15+EGFP have shown less swelling in the infected footpad and parasite load in LNs in comparison with immunized mice with wild-type or non vaccinated groups ($P<0.05$).

Discussion: Recombinant *L. Lactis* expressing SP15 could act as potential vaccine candidate in order to combat and control against *L. major* infection in highly susceptible BALB/c mice.

Keywords: *Phlebotomus papatasi*, SP15, *Leishmania major*, *Lactococcus lactis*

ICIA2018\Vaccine\Oral\9998

9998

Vaccination with synthetic peptides from outer membrane protein A increase antibody titers against *Acinetobacter baumannii*

Kobra Mehdinejadani¹, Ali Hashemi², Mojgan Bandehpour³, Seyed Amir Jalali¹, Hoda Rahmani¹, Shayesteh Mehdinejadani⁴, Vahedeh Hosseini³, Nariman Mosaffa¹

1. Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Department of Microbiology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

3. Department of Biotechnology, School of Advanced Technologies in Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

4. Department of Anatomy, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

Background: *Acinetobacter baumannii* is an opportunistic pathogen and known to cause healthcare-associated infections, especially in ICUs of hospitals. New methods to prevent and treat such infections are a critical need. We have previously reported that five peptides of outer membrane protein A are theoretically immunogenic using bioinformatics tools. In the present study, we report which peptides can induce strong immune responses in mice.

Methods: Mice were vaccinated with various peptides of OmpA plus aluminum hydroxide (Al (OH) 3) adjuvant. Impact of peptides on induction of antibody production was defined.

Results: The present results indicated that Anti-peptide IgG titers related two peptide were higher in comparison with other peptides and groups of control. In fact, immunization with immunogenic peptide vaccine elicits a robust antibody response.

Conclusion: As there are no reports on the immunogenic peptides of OmpA, results presented here, will help to select such peptides as vaccine candidate. These results inform continued development of the vaccine based on peptide against *A. baumannii*, and also are of general importance in that they induce strong immune responses. It may be an effective approach for preventing infection by *A. baumannii*.

Keywords: *Acinetobacter baumannii*, Vaccine, outer membrane protein A, Peptide



Veterinary Immunology

Oral Presentation

14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Veterinary Immunology\Oral\7660

7660

Bovine Enterotoxigenic Escherichia coli Antigen 43 as a potential vaccine for calf colibacillosisReyhaneh Ghorbanpour¹, Gholamreza Nikbakht Brujeni¹*1. Department of Microbiology and Immunology, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran*

Background: Enterotoxigenic Escherichia coli (ETEC) is the most common bacterial cause of diarrhea in human and farm animals. In spite of the global importance of ETEC infections, there is no vaccine available for widespread protection. Recent studies are directed toward recognizing genomic conserved antigens of E.coli, including auto transporter proteins. Antigen 43 (Ag43) is an auto transporter protein that causes autoaggregation and biofilm formation, with a low frequency in commensal strains. It is among a few genes that distinguish ETEC pathotypes from nonpathogenic strains, which are also important in development of efficient vaccine.

Methods: In the current study, a passenger domain of Ag43 in a bovine ETEC strain was amplified and cloned into a pET-28a (+) plasmid. The recombinant Ag43 was purified via a Ni-NTA agarose column and verified by western blotting. Female BALB/c mice were immunized and their infant mice were challenged by the same strain to evaluate the protective effect.

Results: The results implied a significant increase of specific antibody levels due to vaccination of mice by recombinant Ag43 in compared with control group. In addition, the protective effect, by measuring daily percentage of survived/ challenged infant mice, was observed post challenge by 10^2 cfu of homologous bacteria in compared with control group.

Conclusion: According to importance of introduction the novel non classical antigens as vaccine candidates against ETEC in farm animals, here it was indicated that the passenger domain of Ag43 protein can be potentially valuable target in development of bovine colibacillosis vaccine.

Keywords: Enterotoxigenic E. coli (ETEC), Autotransporter, Antigen 43, Vaccine

ICIA2018\Veterinary Immunology\Oral\7705

7705

Production of monoclonal antibody (MAb) against p24 protein of bovine leukemia virus (BLV)

Sanaz Zarasvand Nia¹, Masoud Reza Seyfi Abad Shapouri², Sepide Salehipour Bavarsad¹, Zohre Najafi¹, Reza Pasandideh³

1. DVM Graduated from Faculty of Veterinary Medicine, Shahid Chamran University of Ahvaz, Ahvaz, Iran
2. Professor Department of Pathobiology, Faculty of Veterinary Medicine, Shahid Chamran University of Ahvaz, Ahvaz, Iran
3. PhD Graduated of Animal Genetics and Breeding, Ramin Agriculture and Natural Resources University of Khuzestan, Mollasani, Iran

Background: Bovine leukemia virus (BLV) is the cause of Enzootic bovine leukosis (EBL), the most common neoplastic disease of cattle. Most of the BLV infected cows do not show any clinical signs and as a carrier, have the potential to transfer the virus to other animals through infected lymphocytes, lifetime. BLV infections cause significant economic losses to the livestock industry and a considerable cost is spent for its eradication and control. Since, there is no vaccine or effective treatment against BLV infections, effective control and eradication could be possible by rapid identification of infected cows and culling or separating them.

Methods: Balb/c mice were immunized by the recombinant p24 antigen and the mouse showing the highest titer of anti-p24 antibodies by indirect ELISA was selected for donating spleen for fusion. Spleen cells were fused with SP2/0 myeloma cells using polyethylene glycol. Hybridoma cells were selected in HAT medium and screened by indirect ELISA. The positive clones were subjected to 2 times subcloning. In order to verify the immunogenicity of epitopes against which the monoclonal antibodies have been produced, the possibility for inhibition of monoclonal antibodies reaction by a polyclonal antibody against BLV (the positive control of a commercial ELISA kit) was assessed in a competitive ELISA.

Results: Based on our results, a mAb against p24 was produced and reacted successfully with natural antigens.

Conclusion: With regards to the antigenicity of p24, the specific monoclonal antibodies produced against it may be suitable for developing a competitive ELISA.

Keywords: Monoclonal antibody, p24 protein, bovine leukemia virus

ICIA2018\Veterinary Immunology\Oral\8213

8213

Determination of CD marker expression profiles of Theileria annulata schizont infected cell lines in different passage numbers using RT-PCR analysisGholamreza Habibi¹, Hossein Modirrousta², Asghar Afshari¹

1. Department of Parasite vaccine research and production of Razi vaccine and serum research institute, Agriculture Research, Education and Extension Organization (AREEO), Karaj, Iran

2. Department of Wildlife research of Razi vaccine and serum research institute, Agriculture Research, Education and Extension Organization (AREEO), Karaj, Iran

Background: Bovine theileriosis caused by *Theileria annulata* is an important haemoprotozoan vector-borne disease in Iran. The attenuated vaccine cell line was prepared for induction of immunity after 260 passages in cell culture media. Vaccine seed characterization was performed by molecular and biological methods and now we are going to determine the cell markers of Theileriosis vaccine local cell line. However, CD (cluster of differentiation) molecules are cell surface markers which are very useful for the identification and characterization of cells and the different subpopulations of leukocytes. They are usually recognized by specific antibodies by flow cytometry and immunohistochemistry. Here we emphasize on the application of RT-PCR for defining the cell markers of the bovine leukocyte vaccine cell line with high passage number in comparison with a newly prepared low passage cell line.

Methods: The vaccine cell line at high passage number and a new prepared cell line in low passage number were grown in cell culture medium and the harvested cells were subjected to RNA isolation and further RT-PCR by specific primers for a panel of bovine CD markers (CD4, 5, 11a, 14, 43, 45 and 79a).

Results: The PCR was set up for 7 CDs and was used for the two mentioned above cell lines. The comparison of two sets of CDs expression profiles showed the marker expression for the T cells, dendritic cells, monocytes and macrophage on both examined cell lines but the low passage cell line showed exclusively the CD79a marker for B cells.

Conclusion: The present work is the first study for determination of *Theileriaannulata* schizont infected cell line markers for cell identity test by RT-PCR and showed us the technique might be an alternative assay for flow cytometry.

Keywords: Theileria, RT-PCR, CD, cell line.

ICIA2018\Veterinary Immunology\Oral\8364
8364

In vivo and In vitro Evaluation of Calf Lung Surfactant Extract on Pro-Inflammatory and Anti-Inflammatory Cytokine Changes

Zohre Eftekhari¹, Mohammad Reza Mokhber Dezfouli², Mahiar Jeloudari Mamaghani³, Babak Beikzadeh⁴

1. Department of Quality Control, Research and Production Complex, Pasteur Institute of Iran
2. Department of Clinical Sciences, Faculty of Veterinary Medicine, University of Tehran
3. Department of Drug Substance Production, Noargen Biopharmatech Company
4. Department of microbiology and Immunology, faculty of veterinary medicine, University of Tehran

Background: Inflammatory reactions in the lung pathophysiologic conditions are a critical problem in the treatment process, which in some cases lead to death, particularly in neonate. Exogenous lung surfactant has been considered a candidate to cure inflammation in the lungs. Hence, the efficacy of this substance has been examined for changes in inflammatory and anti-inflammatory cytokine profile.

Methods: Calf lung surfactant extract (CLSE) was obtained by water- salt extraction of freshly slaughtered calves' isolated minced. Finally, obtained active pharmaceutical ingredients formulated by adding dipalmitoylphosphatidylcholine and palmitic acid as excipients and suspension was made by adding NaCl. The quality of finish product determined by hydrophilic interaction liquid chromatography coupled to electrospray ionization–tandem mass spectrometry. The New Zealand White Rabbits as appropriate animal model were treated with formulated CLSE, and then blood samples collected and the level and gene expression of IL-10, IL-6, IL-1 β , IFN- γ and TGF- β were assessment before and after surfactant treatment for 30 days by ELISA and Real time PCR in isolated peripheral blood mononuclear cell, respectively.

Results: The results indicate that IFN- γ and TGF- β increased at 24, 48 and 72 h which were statistically significant compared to baseline. While, IL-6 and IL-1 β in also started to decrease over time in response to surfactant which these changes were significant. The results showed that the gene expression of IL-6, IL-10 and IL-1 β decreased during the time after exposure to the surfactant. The gene expression of IFN- γ and TGF- β increased due to surfactant therapy which reached its maximum expression after 7 days.

Conclusion: This study suggests that CLSE could contribute in reducing pathology effects of respiratory distress syndrome in neonates which can be used as auxiliary and protective drug in respiratory diseases.

Keywords: Exogenous surfactant, Inflammatory, Cytokine profile, Calf lung surfactants

ICIA2018\Veterinary Immunology\Oral\8464
8464

Cellular immune responses of vaccinated chickens with fowl pox vaccine

Mohammad Azizi¹, Bahman Khalesi², EsmailAsli^{1,3}, Ali RezaFarahani⁴, Ali Mohammad Behroozikhah⁵, Abbas Alamian²

1. Department of Microbiology, Faculty of Sciences, Islamic Azad University, Karaj, Iran

2. Department of Poultry Vaccines Production, Razi Vaccine and Serum Research Institute, AREEO, Karaj, Iran

3. Immunology Research Group, Razi Vaccine and Serum Research Institute, AREEO, Karaj, Iran

4. Department of Immunology, Faculty of Medicine, Iran University of Medical sciences, Tehran, Iran

5. Department of Brucella Vaccines Production, Razi Vaccine and Serum Research Institute, AREEO, Karaj, Iran

Background: Fowlpox is a disease of chickens with world-wide distribution. The disease has an economic impact on the poultry industry. This study provides more information that will facilitate poultry industry to a better knowledge of the role of fowl-pox in the poultry immune system. This study was to evaluate the cellular immune responses of vaccinated chickens with fowl pox vaccine.

Methods: Three groups of certain specific pathogens free (SPF) chickens were (21day old, n=40 per group) were used. One group served as then negative control (PBS) and the other two groups were inoculated with either the local FP vaccine or commercial vaccine. Inoculations of all chickens were done by wing-web puncture with a double needle. Blood samples of each group were collected at weeks 1, 2, 3, 4, 5, and 6 after post-vaccination. Peripheral blood mononuclear cells (PBMC) were isolated of each blood sample using Ficoll–Hypaque density gradient centrifugation. The percentages of CD3⁺, CD3⁺CD4⁺, and CD3⁺CD8⁺T-lymphocytes were analyzed by flowcytometry.

Results: The results of this study showed that vaccinated chickens elevated one week later swelling formation (“take”) at the site of vaccination. A maximum elevation of “takes” at day 7 post vaccination were observed. The groups of vaccinated chickens had higher T-cell proliferation responses than the control group. The percentages of CD3⁺, CD3⁺CD4⁺, and CD3⁺CD8⁺T- lymphocytes were increased (P<0.05) in vaccinated chickens with commercial and local FP vaccines. There were no significant differences between both groups of vaccines.

Conclusion: This study revealed that the protective immunity may be associated with increased cellular immunity, which has been interpreted as enhancing the proliferation of T cells and increasing CD4⁺to CD8⁺ratios due to vaccination of the FP vaccine. This suggests that increased the induction of the immune responses by predominantly a Th1-type response.

Keywords: Fowlpox vaccine, Flowcytometry, Chickens, CD4/CD8T-Lymphocytes

ICIA2018\Veterinary Immunology\Oral\8644

8644

Production, Purification and evaluation of Specific Immunoglobulin Y against *Streptococcus iniae* and *Lactococcus garvieae* to prevent streptococcosis in Rainbow trout

Mojtaba Mirzaei¹, Firouz Ebrahimi², Narjes Hasan Fakhrabadi³

1. Doctor of veterinary medicine student, department of pathobiology, school of veterinary medicine, university of Zabol, Zabol, I.R. Iran

2. Department of Biology research center, Faculty of Basic Sciences Imam Hosain University, Tehran - I.R. Iran

3. Doctor of veterinary medicine student, department of pathobiology, school of veterinary medicine, university of Zabol, Zabol, I.R. Iran

Background: The use of IgY technology has been used to control many microbial infections. *Streptococcus iniae* and *Lactococcus garvieae* are two important pathogens in wild and breeding fish, especially salmon, in the world and in Iran, and have caused many economic losses. The purpose of this study was to produce, purify and evaluate IgY antibodies against these bacteria.

Methods: immunization of hens with prepared vaccine of these bacteria in different groups performed, IgG against each antigen in hen serums were detected, IgY was purified from the yolk using PEG 6000 and was dialyzed against distilled water. Product was evaluated by SDS PAGE method. IgY was then evaluated by indirect ELISA. To evaluate the effectiveness of antibodies to prevent pathogenicity of bacteria, the challenge test was carried out in rainbow trout in 6 groups. The first group received only bacteria, the second group received bacteria and IgY (IP rout), the third group received IgY for 7 days (PO rout), the fourth group received non-specific IgY and the bacterium (IP). The fifth received only IgY (IP) and no bacterium was prescribed for the sixth group, this test was performed for each bacterium and each group was evaluated in 2 replicates.

Results: The results of SDS PAGE showed that the produced product contains IgY, and the results of indirect ELISA confirmed that in a concentration of 38 ng of IgY in each well can detect antigens. The results of challenge showed that mortality rate in group1 was 8.3%, and in the group 3, was 58.3% and in group 4 was 83.3% However, in the positive control group, which received only bacteria, mortality was 83.3%.

Conclusion: use of IgY technology can be used to reduce pathogenicity of *Streptococcus* in rainbow trout, so it is readily available and at low cost.

Keywords: Immunoglobulin Y technology, in direct ELISA, *Streptococcus*, rainbow trout

ICIA2018\Veterinary Immunology\Oral\8682
8682

Production of recombinant human polyclonal antibody Fab fragment antivenom specific for Iranian viper *Echiscarinatus* in Rosetta-g bacterium

Mohammad Hasan Motedayen¹, Gholamreza Nikbakht Brojeni², Hosein Zolfagharian¹, Mohammad Mahdy Ranjbar³

1. Department of Immunization and Plasma Production, Razi Vaccine and Serum Research Institute, Agricultural Research, Education and Extension Organization (AREEO), Tehran, Iran

2. Department of Microbiology and Immunology, School of Veterinary Medicine, University of Tehran, Tehran, Iran

3. Department of virology, Razi Vaccine and Serum Research Institute, Agricultural Research, Education and Extension Organization (AREEO), Tehran, Iran

Background: Snake bite poisoning is a serious threat in many of tropical and subtropical countries including in Iran. The best acceptable treatment of envenomated humans is the use of antivenoms. However, present horse-derived antivenoms, despite their unique effects on the treatment of snake victims, they were not fully perfect hence, it needs more improvements.

Methods: In this study, recombinant human Fab fragment antivenom was produced in Rosetta-g bacteria, using a gene library, which was constructed in the previous study. The product Fab was purified in several steps, desalted and LPS-depleted, using ammonium sulfate solution, dialysis against PBS and Triton X-114 solution, respectively. Subsequently, the product was initially confirmed by the SDS-PAGE test and ELISA assay, respectively. Finally, neutralization potency of the product was investigated in laboratory Syrian mice.

Results: Results showed corresponding reduced bands to Fab fragment with the molecular weight of about 28 kDa, at a concentration of 3.1 mg/ml and significant differences of product antivenom with control groups ($P < 0.05$) in ELISA assay. The neutralization potency of present product against *E. carinatus* venom was found to be about 7LD₅₀/ml (54.6 µg/ml) when tested on mice.

Conclusion: Based on the results obtained in the present study, it seems that the product Fab fragment antivenom has the ability to neutralize the *in vivo* biological activity of the Iranian viper *E. carinatus* snake venom. However, more studies require reaching a suitable concentration of antivenom fragment.

Keywords: *Echiscarinatus*, Fab fragment, gene library, antivenom, polyclonal



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Allergy and Immunotherapy of Allergic Disease

Poster Discussion

ICIA2018\Allergy and immunotherapy\Poster Discussion\6331
 6331

Use of a Murine Model to Investigate Potential Allergens Associated to *Ailanthus Altissima* Pollen Allergy

Fateme Mousavi¹, Youcef Shahali², Gholamali Kardar³, Ahmad Majd^{4, 5}, Farrokh Ghahremaninejad⁴, Zahra Pourpak³.

1. Space Biology and Environment center, Aerospace Research Institute, Ministry of Science Research and Technology, Tehran, Iran

2. Razi Vaccine and Serum Research Institute, Agricultural Research, Education and Extension organization (AREEO), Karaj, Iran

3. Immunology, Asthma and Allergy Research Institute, Tehran University of Medical Sciences, Tehran, Iran

4. Department of Plant Sciences, Faculty of Biological Sciences, Kharazmi University, Tehran, Iran

5. Department of Biology, Faculty of Biological Sciences, Islamic Azad University, North Tehran Branch, Tehran, Iran

Background: The murine models that have been widely used in the study of allergy desensitized mice can produce IgE and IgG1 in response to the injection of antigen/adjuvant combination. A.altissima pollen (AAP) has been recently reported as an emerging aeroallergen in Iran. So far, several AAP candidate allergens have been identified by the screening of allergen-specific IgE in human sera from AAP sensitized patients. In the present study, a murine model has been used to investigate AAP specific IgE binding proteins by an immunoproteomics approach.

Methods: Fresh pollens were collected during the 2014 AAP season from male trees planted in an urban green space of Tehran, Iran. The pollen proteins were extracted in phosphate-buffered saline (PBS). Thirty male BALB/c mice were randomly divided into two groups of AP extract sensitized and sham that respectively received AAP PBS extract and a PBS control by intraperitoneal injections at regular intervals. The optimized AAP protein extracts were analyzed using 2D-gel electrophoresis and were subsequently confronted to pooled sera of sensitized mice.

Results: Two-D gel electrophoresis of AAP extract allowed the separation of 125 protein spots distributed in a wide range of pI and molecular masses. Two-DE immunoblotting using pooled sera of sensitized mice led to the detection of 14 IgE reactive spots with molecular masses ranging from 12 to 40-42 KDa, including one very acidic, three acidic, one neutral, eight basic and one very basic protein spots.

Conclusion: The results do not correlate with our previous analyses using human AAP-sensitized sera. These findings reflect some differences in the sIgE reactivity to allergenic proteins in animal models. Since the sensitization process in mice has not been naturally developed but rather induced through the multiple injections of high doses of antigen in combination with adjuvants.

Keywords: Murine model, Allergy, Pollen, *Ailanthus altissima*

ICIA2018\Allergy and immunotherapy\Poster Discussion\7874

7874

Design and Evaluation of Sal k 1 Peptide Based Vaccine and Comparison with Recombinant Allergen and Russian thistle Extract

Saeideh Tabesh^{1,2}, Fanuel Songwe^{2,3}, Mohammad Reza Fazlollahi², Mir Saeed Yekaninejad⁴, Gholam Ali Kardar^{2,3*}, Seyed Alireza Razavi^{1,*}

1. Department of Immunology, School of Public Health, Tehran University of Medical Sciences

2. Immunology Asthma & Allergy Research Institute, Tehran University of Medical Sciences

3. Department of Medical Biotechnology, School of Advanced Technologies in Medicine, Tehran University of Medical Sciences-International Campus (IC-TUMS)

4. Department of epidemiology and biostatistics, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

Background: The Salsola kali pollen is one of the most important causes of allergic rhinitis in the desert and semi-desert area. Its main allergen, Sal K 1, is responsible for more than 90% of the sensitivity of the plant. Allergic rhinitis occupies a large part of the daily life of patients, also the economic consequence of this disease has the large burden on the community. Although the immunotherapy is the only treatment that can modulate the immune system, but given the low efficacy of this method, it is necessary to find a more effective and alternative therapeutic way by these molecular and bioinformatics tools.

Methods: In this study, Sal K 1-KLH vaccine was prepared on the basis of B-cell epitopes. In the next step, the specific IgG and IgE against this allergen or its recombinant protein, Sal K 1, were measured by ELISA in immunized BALB/c mice sera. Also, the inhibition of IgE by mouse IgG was evaluated using an inhibitory ELISA. The IgE reactivity designed vaccine was evaluated by dot blot test and compared with the recombinant protein and Salsola kali extract. Finally, proliferation of T lymphocyte was evaluated in cell culture of patients' PBMCs by MTT assay.

Results: Bioinformatics analysis revealed that peptide based vaccine had different conformational structure than its normal allergens; also had the characteristics of B-lymphocyte epitope. Vaccination with vaccine, produced high levels of protective IgG in mice, which could inhibit the binding of patients IgE to recombinant proteins. The results show designed vaccine, unlike the recombinant protein and extract, did not induce T-cell lymphocytes response also it showed decreased IgE reactivity.

Conclusion: It can be concluded that the designed vaccine will be safe for patients and can be considered as an effective therapeutic candidate in specific immunotherapy.

Keywords: Allergen, Allergen-specific immunotherapy, B epitope, Sal k 1.

ICIA2018\Allergy and immunotherapy\Poster Discussion\8095

8095

Survey the Effect of Saffron Supplementation on The Antibody Titer to Heat-Shock Protein (HSP)70, hsCRP and Spirometry Test in Patients with Mild and Moderate Persistent Allergic AsthmaMarzie Zilaei^{1,2}, Farhad Abolnezhadian³, Seyed Ahmad Hosseini^{*1,2}, Mohammad Ghasemi⁴

1. Nutrition & Metabolic Diseases Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.

2. Nutrition Department, Faculty of Paramedicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

3. Clinical Immunology and Allergy, Allergy Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

4. Immunology Department, Faculty of medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

Background: Asthma is a heterogeneous disease, which usually associated with chronic airway inflammation. The anti-heat shock protein (anti-HSP) 70 is a novel risk factor for asthma. The aim of the present study was to survey the effect of saffron supplementation on anti-HSP70, high-sensitivity C-reactive protein (hs-CRP) and spirometry test in patients with allergic asthma.

Methods: In our clinical trial study, subjects (N= 76, 30 women and 46 men, 18–65 years old) with mild and moderate allergic asthma were randomized into two groups: a group of patients who received two capsules of saffron (100mg/d) and a control group who received two capsules of placebo for 8 weeks. Anti- HSP70, hs-CRP, eosinophil and basophils and spirometry test were determined in patients before (week 0) and after (week 8) intervention. SPSS software (version 16.0; Inc, Chicago, IL) was used for data analysis.

Results: Results showed that saffron in comparison with placebo significantly reduced the hs-CRP (p=0.001), anti-HSP70 (p=0.000), eosinophils (p=0.005) and basophils (p=0.01) concentrations. In spirometry test, FEV1/FVC ratio increased in both saffron and placebo groups (p=0.01). Frequency of the shortness of breath and use of salbutamol spray in saffron group (no in placebo group) decreased significantly (p=0.00).

Conclusion: Results of the present study suggested that saffron supplementation in patients with allergic asthma decreased significantly anti-HSPs 70, hs-CRP and inflammatory cells and improved some clinical symptoms of asthma.

Keywords: Asthma, Anti-HSP70, Inflammation, Saffron.

ICIA2018\Allergy and immunotherapy\Poster Discussion\8620

8620

Environmental Condition and Allergenic Airborne Pollen Counts of Chenopodiaceae/Amaranthaceae Families in Tehran, IRAN

Maryam Sharif Shoushtari^{1,2}, Ahmad Majd³, Mostafa Moin^{2,4}, Fariborz Zandieh⁵, Masoud Movahedi ^{2,4}, Zahra Pourpak^{2,4}

1. Young Researchers and Elites club, Science and Research Branch, Islamic Azad University, Tehran, Iran
2. Immunology, Asthma and Allergy Research Institute, Tehran University of Medical Sciences, Tehran, Iran
3. Department of Plant Biology, faculty of biology science, Islamic Azad University, North Tehran Branch, Iran
4. Department of Immunology and Allergy, Children Medical Center, Tehran University of Medical Sciences, Tehran, Iran
5. Bahrami Children Hospital, Tehran University of Medical Sciences, Department of Immunology, Asthma and Allergy, Tehran, Iran

Background: Pollen grains of Chenopodiaceae/Amaranthaceae family members cause allergic respiratory system diseases in summers especially in areas with arid climates. The objective was to evaluate the association between Chenopodiaceae/Amaranthaceae pollen dynamics in the atmosphere of Tehran and relationships with meteorological factors for eight years.

Methods: The districts of Tehran were located in northern, southern, eastern, western and central regions. Airborne pollen data were collected from January 2010 to December 2017 by two air samplers; volumetric methods employing a Burkard grains/m³ and gravitational method, Durham sampler grains/m², to identify and determine pollen concentration in the air and separated in Chenopodiaceae/Amaranthaceae family. Then, relationship between the meteorological parameters including the mean values of temperature, sunshine hours, relative humidity, wind speed and raining was investigated.

Results: Chenopodiaceae/Amaranthaceae pollen season lasted in Tehran from mid-May until the mid-November. For the five area of Tehran, consisting of 19 genera types were identified in the eight years. Pollen production per square meter was higher in the center and south and lower in the east areas of Tehran. Highest pollen counts were obtained during Mid-August and lowest in the June. The comparison of Chenopodiaceae/Amaranthaceae pollen counts with a Durham sampler showed the automatic counter a better correlation than Durham sampler but this sampler when pollen dispersal was high and pollen identify is better than automatic counter sampler. The correlations obtained between daily pollen counts and meteorological parameters showed that the airborne presence of this pollen type is associated positively with temperature and sunshine hours and negatively with rainfall and relative humidity.

Conclusions: The Chenopodiaceae/Amaranthaceae in weeds pollen calendar and its association with meteorological factors depend mainly on daily temperature, sunshine hours which may help draw the attention of physicians and allergic patients every days with high Chenopodiaceae/Amaranthaceae pollen counts. This is an important relationship to consider while interpreting individual clinical trials.

Keywords: Chenopodiaceae/Amaranthaceae, Pollen counts, Pollen monitoring, Pollinosis, Tehran

ICIA2018\Allergy and immunotherapy\Poster Discussion\8627
8627

CD63 and CD203c Are Strongly Expressed on The Basophils of Patients With Chronic Urticaria and Are Novel Biomarker for Diagnosis of This Disease

Erfan babaei¹, Rasoul Nasiri kalmazi², Ali jalili², Javid Morad Abbasi³

1. Department of Immunology, Faculty of Medicine, Kurdistan University of Medical Sciences, Sanandaj, Iran

2. Cancer and Immunology Research Center, Kurdistan University of Medical Sciences, Sanandaj, Iran

3. Asthma and Allergy Center, Tehran Medical Sciences, Branch of Academic Center for Education, Culture and Research (ACECR), Tehran, Iran

Background: Chronic urticaria (CU) is a skin disorder that considered appearing wheals with itch and pruritus anywhere on the body for more than 6 weeks. Auto reactivity of the CU-patients sera can be established in 30% to 50% of patients, depending on the population, through the autologous serum skin test (ASST). The ASST has only moderate specificity as a marker for functional autoantibodies against the high-affinity immunoglobulin E (IgE)-receptor FcεRIα. There is a need for a strong laboratory test for the diagnosis of CU, which can persistently and objectively support the anamnestic diagnosis. In the current study we aimed to examine the expression levels of both CD63 and CD203c on basophils of patients with CU.

Methods: The study included 45 patients diagnosed with CU, according to the EACCI/GA (2) LEN/EDF/WAO guidelines, Peripheral blood of patients (25 ASST+ CU and 20 ASST- CU) and 25 healthy subjects were tested for expression of activation markers including CD63 and CD203c.

Results: we observed that the number of basophils expressing CD63 and CD203 was significantly higher (both $p < 0.05$) in CU-patients than the normal individuals. In addition, the levels of CD63/CD203+ in patient with positive ASST test were lower than negative ASST test group. However, it was not significant ($p > 0.05$).

Conclusion: Based on this study the flowcytometric analysis for enumerating basophils expressing CD63 and CD203c is a suitable diagnostic tool for patients with CU. However, further studies enrolling more patients are necessary.

Keywords: Chronic Urticaria, CD63, CD203c, ASST

ICIA2018\Allergy and immunotherapy\Poster Discussion\8643

8643

The Modulatory Effects of Estrogen and Progesterone on Expression of Th9-related Transcription Factors and Secretion of IL-9 from PBMCs of Post-Menopausal Asthmatic Patients

Leila Nejatbakhsh Samimi¹, Majid Khoshmirsafa¹, Morteza Fallahpour², Seyed Ali Javad Mousavi³, Rasool Baharloo¹, Paria Jorbozehdar¹, Reza Falak¹

1. Department of Immunology, School of medicine, Iran University of Medical Sciences, Tehran, Iran

2. Department of Clinical Allergy and Immunology, Hazrate-Rasool Akram General Hospital, Iran University of Medical Sciences, Tehran, Iran

3. Department of Internal Medicine, Hazrate-Rasool Akram General Hospital, Iran University of Medical Sciences, Tehran, Iran

Background: IL-9-secreting CD4⁺T cells (Th9), defined as subset of T cells which play a crucial role in development of allergic asthma. The goal of the present study was to determine the effects of 17 β -estradiol (E2) and progesterone (P4) in singular or in combination form on expression of IRF-4, PU.1 and BATF as effective transcription factors on differentiation of IL-9 producing cells in mild to moderate asthmatic patients versus non-asthmatic healthy controls.

Methods: Peripheral blood mononuclear cells (PBMCs) were isolated and cultivated in the presence or absence of phytohemagglutinin (PHA) in parallel to E2 or P4. In brief, 10⁻⁸ M of E2 or 10⁻⁶ M of P4, equivalent to serum levels observed during hormone replacement therapy (HRT), or a combination of both hormones were added on PBMCs in presence or absence of 1% PHA. Following mRNA isolation and cDNA synthesis, real-time PCR was used to evaluate the expression level of IRF-4, PU.1 and BATF. We also measured the concentration of IL-9 as the related cytokine in cell culture supernatants by ELISA.

Results: The median (IQR) IL-9 secretion was 3.2(2.1-5.7) and 1.77(1.3-3) in asthmatic patients and control group for E2 treatment, respectively (Pv=0.0087). In addition, the median (IQR) IL-9 secretion was 2.4(1.6-3) and 1.5(1.1-1.77) in asthmatic patients and control group for E2+P4 treatment, respectively (Pv=0.009). BATF expression in PBMCs was significantly increased in the presence of E2 or the combination of E2+P4 in asthmatic patients compared with the control group. Expression of PU.1 and IRF-4 were also increased; however, they were not statistically significant.

Conclusion: Treating PBMCs with estrogen alone or in combination of estrogen and progesterone (as an in vitro example of HRT) could derive differentiation of PBMCs toward IL-9 producing cells which might exacerbate the clinical symptoms of the allergic asthma.

Keywords: Allergic asthma, PBMCs, Estradiol, IL-9, Progesterone

ICIA2018\Allergy and immunotherapy\Poster Discussion\9895

9895

Relationship between Migraine and Food AllergySafoora pordel¹, Shokrollah Farrokhi¹*1. Department of Immunology, Asthma and Allergy, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Research Institute, Bushehr University of Medical Sciences, Bushehr, Iran*

Background: The factors triggering the onset of migraine attacks are not completely understood. Foods and food additives could induce migraine attacks. The aim of this study was to evaluate the relationship between food allergy and migraine attacks based on skin prick test (SPT) reactivity.

Methods: Fifty-one known patients with migraine disease were enrolled. The extracts of standardized food allergen were used in the skin test. Then, the patients with positive SPT were avoided for food allergen for three months. The quality of life (QOL) of the patients was assessed by using SF-26 questionnaire.

Results: A significant difference was observed between positive SPT reactivity to shellfish ($P=0.002$) as well as tomato ($P=0.038$) and migraine attacks. The most frequent food allergens were peach (45%), strawberry (41.1%) and egg yolk (39.2%), respectively. Moreover, after allergen avoidance, the number of migraine attacks was reduced and QOL was improved for the patients.

Conclusion: There was a significant relationship between migraine attacks and food allergy. Therefore, this study suggested that early diagnosis of food allergens and their avoidance may improve QOL of migraine patients.

Keywords: Food Allergy, Migraine, Skin prick Test

ICIA2018\Allergy and immunotherapy\Poster Discussion\9898
9898

The Efficacy of Autologous Plasma Therapy in Patients with Chronic Urticaria

Safoora pordel¹, Shokrollah Farrokhi¹

1. Department of Immunology, Asthma and Allergy, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Research Institute, Bushehr University of Medical Sciences, Bushehr, Iran

Background: Chronic urticaria is an allergic skin disease characterized by severe pruritic wheal persisting at least for a period of 6 weeks. The aim of this study was to evaluate the efficacy of plasma therapy in treatment of idiopathic chronic urticaria.

Methods: In this cross sectional study, the quality of life (QOL) of 30 patients with idiopathic chronic urticaria (autoimmune and spontaneous) was evaluated before and after autologous plasma therapy by using standard questionnaires (DLQI and TSS).

Results: The findings showed that the means of DLQI and TSS scores in the patients after plasma therapy was significantly reduced. This indicates an improvement of the QOL in the patients. Moreover, it was found that the plasma therapy was significantly effective in both sexes.

Conclusion: The results of the present study suggest that autologous plasma therapy could be a suitable method for the treatment of patients with idiopathic chronic urticaria.

Keywords: Chronic urticaria, Plasma therapy, DLQI and TSS scores, Quality of life

ICIA2018\Allergy and immunotherapy\Poster Discussion\10002
10002

Evaluating Immunomodulatory Effects of Vitamin D on IL-17 Expression in Peripheral Blood Mononuclear Cells of Patients with Asthma

Mohsenfarhang zargar¹, Saeedeh Erfanin², Maliheh Moradzadeh³, Bahareh Razeghi Haghighi¹

1. Student of laboratory sciences, Student Research committee, Jahrom University of Medical Sciences, Jahrom, Iran

2. MSc Department of Advanced Medical Sciences and Technologies, School of Medicine, Jahrom University of Medical Sciences (JUMS), Jahrom, Iran

3. PhD Golestan Rheumatology Research Center, Golestan University of Medical Sciences, Gorgan, Iran

Background: Asthma is a form of hypersensitivity type 1 in respiratory airway that many immunity factors are involved in its development and progression. Laboratory evidence indicates that the anti-inflammatory metabolites of vitamin, 25 D-dihydroxy vitamin D3 and similar compounds can be performed by pairing with vitamin D receptors and caused down regulation or translation of cytokines production such as IL2, IL12, TNF -alpha, INF- γ , agent stimulating macrophages, granulocyte colony-giver and IL-1B and IL-6. Therefore, given the importance of cellular immunity in order to increase the pathogenesis of the disease, the aim of present study is a direct effect of vitamin D on cell blood mononuclear mainly T lymphocytes in medium patients with asthma and control groups.

Methods: 10 cc heparinized peripheral blood was taken from 20 patients with asthma and 20 healthy controls. The isolated peripheral blood mononuclear cells were treated with 10^{-7} and 10^{-6} M of vitamin D and dexamethasone, for 24-hour culture. After RNA extracted from 1 million cells, cDNA synthesis and gene expression Real-time PCR was used for IL-17gene.

Results: The results of this study indicated the overexpression of IL-17 in patients with asthma compared to the control group, although this difference is not significant ($P = 0.061$). In this study, vitamin D3 and dexamethasone at different modes to no treatment group of patients with asthma were significantly different ($P < 0.05$) but treatments 10^{-7} and 10^{-6} of vitamin D3 and dexamethasone did not show a significant difference ($P > 0.05$). Vitamin D3 and dexamethasone different modes compared to no treatment in healthy subjects, in none of the cases did not show significant differences ($P > 0.05$).

Conclusion: The effect of vitamin D treatment as well as on other routes should also be evaluated TBET gene to an effective step in understanding the molecular mechanisms of diseases in order to achieve the objectives described to be removed.

Keywords: IL-17, Vit D, Asthma



Cancer Immunology

Poster Discussion

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Cancer Immunology\Poster Discussion\7482

7482

MiRNA-143 Replacement Therapy Harnesses the Proliferation and Migratrion of Colorectal Cancer ells *in vitro*

Leila Karimi¹ , Dariush Shanehbandi¹ , Behzad Mansoori¹ , Ali Mohammadi¹ , Milad Asadi¹ , Behzad Baradaran¹

1. Immunology Research Center , Tabriz University of Medical Sciences , Tabriz , Iran

Background: Colorectal cancer (CRC) is rated as the third and second most common cancer in men and women. CRC is the fourth most common cause of cancer related mortality worldwide and accounts for ~9% of all cancer cases. Abbarent expression of microRNAs (miRNAs) is correlated with numerous malignancies such as CRC. Restoration of tumor supressor miRNAs seems to be an effectual strategy for treatment of the relating malignancies. MiR-143 is a conspicuous example of tumor supressor miRNAs which its down-regulation has been reported in CRC. This miRNA is involved in negative regulation of metastasis-related genes including KRAS, c-Myc and MMP-9 as well as anti-apoptotic Bcl2.

Methods: In the present study, the lowered expression of miR-143 in SW-480 CRC cell line was confirmed by quantitative Real Time PCR (qRT-PCR). Subsequently, miR-143 replacement was performed by introducing the pCMV-miR-143 vector into the model cells. After the restoration of miR-143 expression, changes in proliferation and migration of the cells were assessed by MTT and scratch tests, respectively. Furthermore, relative expression of KRAS, c-Myc, MMP-9 and Bcl2 as putative targets of miR-143 were investigated by qRT-PCR. DAPI staining was utilized to detect apoptosis occurrence in the transfected cells. Moreover, the ratio of Bax to Bcl2 expression, as a potential indicator of the response to apoptosis induction, was compared between the study groups.

Results: MTT assay revealed a decrease in the viability of miR-143 transfected cells. In addition, a reduced migratory potential for the transfected cells was indicated by scratch test. DAPI staining confirmed the occurrence of apoptosis in the mentioned cells. According to qRT-PCR results, expression of Bcl2, KRAS, MMP-9 and c-Myc mRNAs was significantly decreased in the miR-143 grafted cells compared to the controls. Furthermore, Bax/Bcl2 ratio showed a significant increase in the miR-143 overexpressing cells.

Conclusion: MiR-143 replacement can be an effective strategy for the treatment of colorectal cancer and harnessing its invasive properties.

Keywords: miR-143, Colorectal cancer, Metastasis, microRNA replacement therapy, Apoptosis

ICIA2018\Cancer Immunology\Poster Discussion\7549

7549

Association of IL-27 rs153109 and rs17855750 Polymorphisms with Risk and Response to Therapy in Acute Lymphoblastic Leukemia

Alireza Ghavami¹, Farahnaz Bodaghkhan², Saeedreza Vazirian³, Zahra Amirghofran⁴

1. Department of Immunology and Autoimmune Diseases Research Center, Shiraz University of Medical Sciences

2. Pathology Lab Section, Shiraz University of Medical Science Namazi Hospital

3. Physician of Fars Oil Industry, MD N.I.O.P.D.C

4. Professor of Immunology, Autoimmune Disease Research Center, Shiraz University of Medical Sciences

Background: Interleukin-27 is a cytokine with important anti-cancer activity. This study has evaluated the effects of IL-27 rs153109 and rs17855750 single nucleotide polymorphisms (SNPs) on risk of acute lymphoblastic leukemia (ALL) development, as well as their impact on prognosis and patient survival.

Methods: A total of 200 patients and 210 healthy subjects were genotyped by polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP). There were 40 new cases that had received no chemotherapy at enrollment. We used the samples from these patients for IL-27 serum measurements by enzyme-linked immunosorbent assay (ELISA). The laboratory and clinical characteristics recorded at presentation were sex, age, white blood cell counts, platelet counts, immunophenotype, hemoglobin (Hb) level, percentage of blast in the bone marrow and peripheral blood and extra-medullary involvement (EMI). During the period of follow-up, response to therapy was evaluated by measuring complete remission (CR) rate, CR duration (CRD) and overall survival. Statistical analyses were accomplished using SPSS version 23 for Windows. P-values less than 0.05 were considered statistically significant.

Results: We observed a higher frequency of rs153109 AG and rs17855750 TG genotypes and allele G in patients compared to controls ($p < 0.001$). Combined G variant genotypes (AG+GG and TG+GG) also conferred significantly greater risk of ALL. There was a significant correlation between the genotypes of both SNPs with event-free survival (EFS). Patients with GG genotypes of both SNPs and those of rs153109 AG and rs17855750 TG had a shorter EFS than patients with rs153109 AA and rs17855750 TT genotypes ($p \leq 0.035$). Combined G variant genotypes for both SNPs showed poorer response to therapy in all patients ($p < 0.027$) as well as B-ALL (rs153109, $p < 0.001$) and T-ALL (rs153109, $p = 0.048$) patients. In multivariate analysis, rs153109 combined G variant genotype was associated with shorter EFS (relative risk=9.7, $p = 0.026$). Among those who relapsed, 87.1% had the rs153109 AG genotype and 77.4% had the rs17855750 TG genotype ($p < 0.01$). Patients had higher IL-27 serum levels compared to controls, but this did not differ between genotypes.

Conclusions: The association of IL-27 rs153109 and rs17855750 polymorphisms with risk of ALL development and their impact on EFS suggested an important role for this cytokine in biology and response to ALL therapy.

Keywords: Interleukin-27, Acute lymphoblastic leukemia, polymorphism

ICIA2018\Cancer Immunology\Poster Discussion\7571

7571

Blimp-1 Expression as an Exhaustion Transcription Factor in Chronic Lymphocytic Leukemia

Saeid Taghiloo¹, Esmaeil Allahmoradi¹, Mohsen Tehrani^{1,2}, Ghasem Janbabaei³, Ramin Shekarriz³, Hossein Asgarian Omran^{1,4*}

1. Department of Immunology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

2. Molecular and Cell-Biology Research Center, Mazandaran University of Medical Sciences, Sari, Iran

3. Gastrointestinal Cancer Research Center, Mazandaran University of Medical Sciences, Sari, Iran

4. Immunogenetics Research Center, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

Background: Based on the tumor immune-surveillance theory, various components of the immune system constantly survey the body for any malignant cell proliferation and eliminate them or slow their growth. In the course of chronic cancer and infections, T cells become “exhausted” with poor effector function characterized by the expression of multiple immune checkpoint inhibitory receptors and transcription factors. Here, we investigated the expression of Blimp-1, a transcription factor involving in T-cell exhaustion, in patients with chronic lymphocytic leukemia (CLL).

Methods: Peripheral blood mononuclear cells were collected from 25 untreated CLL patients and 15 sex- and age-matched normal subjects. CLL patients were clinically classified according to the Rai staging system. The relative expression of Blimp-1 mRNA was determined by quantitative Real time Polymerase Chain Reaction (qRT-PCR) after normalization with β -actin.

Results: Expression of Blimp-1 mRNA was much higher in CLL patients than in normal controls ($p=0.001$). Moreover, Blimp-1 was higher expressed in patients with advanced clinical stages of CLL compared to those with early stages of the disease ($p=0.01$). Interestingly, the Blimp-1 expression was correlated with the frequencies of exhausted Tim-3⁺/PD-1⁺/CD4⁺ and Tim-3⁺/PD-1⁺/CD8⁺ T cells in CLL patients.

Conclusions: Our study introduces Blimp-1 as a potential transcription regulator of T cell exhaustion in CLL patients. Better understanding of underlying mechanisms of T cell exhaustion in CLL could be helpful in finding new therapeutic strategies for immunotherapy.

Keywords: Exhausted T cell, Blimp-1, Chronic Lymphocytic Leukemia

ICIA2018\Cancer Immunology\Poster Discussion\7794
7794

SDF-1 α Reduces Human NK cell Cytotoxicity against K562 Cells *in vitro*

Alireza Mardomi¹, Saeid Abediankenari^{1*}

1. Immunogenetics Research Center, Mazandaran University of Medical Sciences, Sari, Iran

Background: Stromal cell derived factor-1 alpha (SDF-1 α) belonging to CXC family of chemokines is up-regulated in a variety of malignancies. The expression of SDF-1 α has been associated with poor prognosis and invasiveness of these neoplasms. It was not clear whether the described poor prognosis attributed to SDF-1 α is due to immunosuppression or not. Evidence on the effect of SDF-1 α in shaping the immune response is controversial. Therefore, the aim of present study was to investigate the effect of SDF-1 α on the cytotoxicity of NK cells, as important effector cells in tumor immune surveillance.

Methods: Human NK cells were freshly isolated from 5 healthy donors using MACSxpress system (Miltenyi) and cultured 24 hours in the presence or absence of recombinant human SDF-1 α (100 nM). A CD107a degranulation assay was conducted by the exposure of NK cells to K562 cells at 2:1 effector to target ratio for 4 hours. The percentage of CD107a positive cells was assessed by FACS. Effect of SDF-1 α was also examined on the relative mRNA expression of NKG2A and NKG2D in NK cells.

Results: Our results demonstrated that SDF-1 α decreased the degranulation of NK cells significantly ($p=0.04$). However, blockage of the SDF-1 α receptor, CXCR4 using AMD3100 did not recover the SDF-1 α induced CD107a down-regulation. The mRNA content of inhibitory and activating receptors, NKG2A and NKG2D as indicators of NK cell activity were both down-regulated in SDF-1 α treated group ($p=0.04$ and $p<0.01$ respectively). Moreover, AMD3100 recovered the NKG2A mRNA level to that of the un-treated cells.

Conclusion: The present study revealed that SDF-1 α , as one of the soluble factors up-regulated in numerous malignancies, has a negative impact on NK cell activity and probably, is involved in tumor-induced immune-suppression. Therefore microenvironment manipulations targeting SDF-1 α may reinforce cancer therapies by disturbing one of the immune-suppressive components existing in the tumor microenvironment.

Keyword: NK cell, SDF-1, Tumor microenvironment

ICIA2018\Cancer Immunology\Poster Discussion\7974

7974

Upregulation of Immune Checkpoint Inhibitors, PD1-PDLs, in Patients with Bladder CancerAli Ariaifar^{1,2,3}, SomayehRezaeifard¹, MaisamGhaedi³, ShahriarZeighami², Zahra Faghih¹

1. Shiraz Institute for Cancer Research, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran
2. Urology-Oncology Research Center, Shiraz University of Medical Sciences, Shiraz, Iran
3. Department of Urology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Malignant cells escape from immune responses through different mechanisms including inhibition of tumor infiltrating lymphocytes by expression of inhibitory molecules. A major hallmark of T-cell exhaustion is enhanced expression of multiple immune checkpoints i.e. PD-1. In the present study, we investigated expression of one inhibitory checkpoint, PD1, and its ligands, PD-L1 and PD-L2, on tumor cells and tumor-infiltrating lymphocytes in bladder cancer (BLC) tissue as well as tumor draining lymph nodes (TDLNs).

Methods: Cells were obtained from 37 tissues and 48 TDLNs were obtained from 61 untreated patients with BLC. The tumor cells were counted and stained for CD45, PD-1, PDL-1 and PD-L2. 7-AAD vital dye was used to discriminate live from dead cells. The cells were then acquired on four-color flow cytometer and analyzed with FlowJo software.

Results: Our results indicated that in TDLNs of BLC patients, the prevalence of total CD45⁺ expressing PD1 was significantly higher in LN⁺ patients and patients with higher stage tumor in comparison to the ones with free lymph nodes and lower stage tumor ($P<0.05$). PD1 expressing CD45^{low} and CD45^{hi} cells were also significantly increased in patients with higher stage tumor ($P<0.05$). Moreover, PD1 and PD-L2 expressing CD45^{hi} cells elevated in LN⁺ patients as well ($P<0.05$). Whilst in tumor tissues, just PD-L1 expressing CD45^{hi} cells were increased in patients with higher stage and higher grade tumors ($P<0.05$).

Conclusion: Elevated frequency of PD1 expressing CD45⁺ cells in TDLNs of patients with advanced diseases implied that immune cells in draining lymph nodes underwent exhaustion or over-activation. This exhaustion leads to decreased production of their effector cytokine and cytolytic activities and hence the failure of cancer elimination, tumor progression and metastasis. These results emphasize the importance of these molecules in cancer biology and designing new anti-cancer treatments.

Keywords: Bladder Cancer, Lymph node, PD1, PD-L1, PD-L2

ICIA2018\Cancer Immunology\Poster Discussion\8016

8016

Evaluation of Potential Therapeutic Effect of Anti-Siglec-F Antibody in a Mouse Dual Breast Cancer and Allergic Asthma Model

Sima Shahmohammadi Farid^{1,2}, Nazanin Mojtavavi³, Ali Anissian⁴, Alireza Razavi¹, Amir Hassan Zarnani^{5,6}

1. Department of Pathobiology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

2. Hematology & Oncology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

3. Department of Immunology, School of Medicine, Iran University of Medical Sciences, IUMS, Tehran, Iran.

4. Veterinary department, agricultural faculty, Islamic Azad University, Abhar branch, Abhar, Iran

5. Reproductive Immunology Research Center, Avicenna Research Institute, ACECR, Tehran, Iran

6. Immunology Research Center (IRC), Iran University of Medical Sciences, IUMS, Tehran, Iran

Background: Eosinophils have long been known to infiltrate tumors. Eosinophils from allergic and non-allergic patients have differential effects on induction of apoptosis in tumor cells. Here, we explore the potential immunotherapeutic effect of anti-Siglec-F (sialic acid-binding immunoglobulin-like lectin F), a mouse eosinophil-specific antibody, on the course of breast cancer in a mouse model of allergic asthma.

Methods: We generated a dual mouse model of allergic asthma and breast cancer and injected mice with anti-Siglec-F antibody to explore the role of eosinophils in breast cancer pathogenesis and metastasis in an asthmatic condition. Chronic and acute allergic asthma models were generated by timeline injection and nebulization of ovalbumin (OVA). Breast cancer was induced by subcutaneous injection of 4T1 cell line. Eight different groups of mice were used in this study.

Results: In cancer group, anti-Siglec-F Ab caused a significant increase of IL-4 and IL-5 level and a significant reduction of IL-10 level compared with cancer group receiving vehicle. In asthma group, anti-Siglec-F Ab significantly increased the level of IL-4, IL-5 and IL-12 and diminished the level of IL-6 and IL-10. In asthma and cancer group, Siglec F Ab leads to increased level of IL-4, -5, -6, -10 and -12 cytokines. Interestingly, such intervention significantly decreased OVA specific IgE level in sera of both asthmatic and asthmatic/cancer mice. Pathological evaluation of lungs from animals received antibody showed extensive metastasis in cancer group which was associated with 25% mortality in mice; however antibody administration in asthma/cancer mice significantly decreased metastatic foci.

Conclusion: Our data showed that, compared to cancerous condition, eosinophils behave differently in the pathogenesis of breast cancer when asthmatic condition has already been established raising the possibility of dual influence of this cell type on cancer prognosis depending upon microenvironment.

Keywords: Allergy, Breast cancer, Eosinophil, Immunotherapy, Metastasis

ICIA2018\Cancer Immunology\Poster Discussion\8047

8047

Proteomic Pattern of Mesenchymal Stem Cells in the Patients with Malignant and Benign Salivary Gland Tumors

Mohammad Reza Haghshenas¹, Zahra Mojtahedi¹, Mahboobeh Razmkhah¹, Nasrollah Erfani¹, Bijan Khademi², Mohammad Javad Ashraf³, Abbas Ghaderi^{1,4}

1. Shiraz Institute for Cancer Research, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

2. Otolaryngology Research Center, Department of Otolaryngology, Shiraz University of Medical Sciences, Shiraz, Iran

3. Department of Pathology, Khalili Hospital, Shiraz University of Medical Sciences, Shiraz, Iran

4. Department of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

Background: To improve the knowledge about molecular mechanisms involved in pathogenesis of tumors, this study aimed to investigate MSCs proteome pattern in the patients with malignant and benign salivary gland tumors (SGTs).

Methods: Mesenchymal and non-mesenchymal fluorochrome antibodies and flow cytometric analysis were used to verify MSCs obtained from malignant (mucoepidermoid carcinoma) and benign (pleomorphic adenoma) tumor tissues, and two-dimensional polyacrylamid gel electrophoresis (2DE) coupled with matrix assisted laser desorption ionization tandem time-of flight mass spectrometry (MALDI-TOF-TOF) was done to identify differentially expressed proteins.

Results: Our result showed that both MSCs were strongly positive for mesenchymal markers such as CD90, CD73, CD105, CD166, and CD44, but they were negative or weakly positive for non- mesenchymal markers such as CD14, CD34, and CD45. Our result also showed that the expression pattern of cytosolic platelet-activating factor acetyl hydrolase type IB subunit beta (PAFAH1B), S-formyl glutathione hydrolase (FGH), type II trans glutaminase (TG2), FK506 binding protein 9 (FKBP9), and Annexin A4 (Anxa4) was significantly higher among MSCs isolated from malignant tissues, while the expression pattern of *keratin, type II cytoskeletal 7* (CK-7), *heat shock protein 70* (Hsp70) was higher in MSCs isolated from the benign tissues.

Conclusion: Our result introduces differential expression of Anxa4 as well as four enzymatic proteins related to malignant transformation, cancer metabolism, and cancer development in the tumor microenvironment of patients with malignant SGTs. The data suggest that MSCs may serve as a desirable therapeutic target in SGTs treatment.

Keywords: Salivary gland tumors, Mesenchymal stem cells, Proteomic pattern

ICIA2018\Cancer Immunology\Poster Discussion\8198

8198

Carcinoma-Associated Fibroblasts Are a Common Finding in the Microenvironment of HPV-Positive Oropharyngeal Squamous Cell Carcinoma

Samira Derakhshan¹, Sedigheh Rahrotaban¹, Alireza Abdollahi², Farzad Yazdani³

1. Department of Oral and Maxillofacial Pathology, School of Dentistry, Tehran University of Medical Science, Tehran, Iran

2. Department of Pathology, Imam Hospital Complex, Cancer institute, Tehran University of Medical Sciences, Tehran, Iran

3. Department of Pathology, Amiralam Hospital, Tehran University of Medical Sciences, Tehran, Iran

Background: The important role of the human papillomavirus (HPV) is widely established in oropharyngeal squamous cell carcinoma (OPSCC). The behavior of a OPSCCs especially induced by HPV might be influenced by the tissue microenvironment and its changes according to the tumor nature. Recognition of the role of the tumor microenvironment on the behavior of neoplastic cells has led to employment the microenvironment to use as therapeutic target. Carcinoma-associated fibroblasts (CAFs), the most abundant cells in the tumor microenvironment, show wide-spread expression of alpha-smooth muscle actin (a-SMA). We focused on CAFs, its presence in oropharyngeal squamous cell carcinoma and the relationship with HPV for the first time. **Methods:** Expression of a-SMA protein in CAFs of the tumor microenvironment of the 44 formalin fixed paraffin-embedded tissue blocks from the primary tumor of OPSCC evaluated by immunohistochemistry between HPV-positive and HPV-negative tumors separated by nested polymerase chain reaction.

Results: In 44 samples 23 HPV positive cases were detected. Statistically there were significant differences between histopathologic grade, percent and final score of a-SMA and HPV expression. Significant difference between HPV expression and inflammation, intensity and clinical parameters was not identified in the present study.

Conclusion: Our results indicate that CAFs are a common finding in the microenvironment of HPV-positive oropharyngeal squamous cell carcinoma and associated with higher histopathologic grade. Therapeutic strategies to employ CAF-mediated drugs need to be considered and evaluated more for treatment of HPV-positive oropharyngeal squamous cell carcinoma.

Keywords: Carcinoma-associated fibroblasts, HPV, Oropharynx, Squamous cell carcinoma

ICIA2018\Cancer Immunology\Poster Discussion\8263

8263

Docosahexaenoic acid and Eicosapentaenoic Acid Induce Cell Death in Peripheral Blood Mononuclear Cells of Multiple Myeloma Patients but Not Healthy Individuals

Milad Moloudizargari¹, Esmail Mortaz^{1,2,3}, Frank A. Redegeld³

1. Department of Immunology, Faculty of Medicine, ShahidBeheshti University of Medical Sciences, Tehran, Iran

2. Clinical Tuberculosis and Epidemiology Research Center, National Research Institute of Tuberculosis and Lung Diseases (NRITLD), ShahidBeheshti University of Medical Sciences, Tehran, Iran

3. Division of Pharmacology, Faculty of Science, Utrecht Institute for Pharmaceutical Sciences, Utrecht University, Utrecht, Netherlands

Background: Poly-unsaturated Fatty Acids (PUFAs) have been shown to have anti-cancer effects in both solid and non-solid tumors. Eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) are among the most studied PUFAs. We have shown in a previous study that EPA and DHA induce apoptotic effects in multiple myeloma cell lines via altering the expression of a wide array of genes involved in oxidative stress as well as NF- κ B inhibition, increasing mitochondrial perturbation, and caspase 3 activation.

Methods: The aim of the present study was to further evaluate the apoptotic effect of these two fatty acids in primary peripheral blood mononuclear cells (PBMCs) obtained from untreated patients (new cases) with confirmed symptomatic multiple myeloma.

Results: The results showed that EPA at the concentration of 100 μ M and DHA at both concentrations of 50 and 100 μ M can induce potent apoptotic effects (up to 44, 76 and 98%, respectively) in the PBMCs of patients with multiple myeloma ($P \leq 0.05$), while they have no effect on the PBMCs isolated from healthy individuals ($P > 0.05$). The observed effect was concentration and time-dependent and 72h treatment with DHA at a concentration of 100 μ M had the strongest effect ($P \leq 0.01$).

Conclusion: These findings confirm the results of our previous study and may suggest a promising future for the application of PUFAs as an adjunctive therapy in the treatment of multiple myeloma with fewer side effects. Further studies are warranted to define the underlying mechanisms shedding light on the possibility of exploiting apoptotic effects of PUFAs in cancer therapy.

Keywords: Docosahexaenoic acid, Eicosapentaenoic Acid, Multiple Myeloma

ICIA2018\Cancer Immunology\Poster Discussion\8441

8441

Genetic variants of PD-1 co-inhibitory molecule (PD1.3 and PD1.5) in patients with basal cell carcinoma

Maryam Fakhmi¹, Maryam Abedi¹, Zahra Faghih¹, Mohamad reza Haghshenas¹, Faride Jowkar³, Nasrollah Erfani^{1,2}

1. Shiraz Institute for Cancer Research, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

2. Departments of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

3. Department of Dermatology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Basal Cell Carcinoma (BCC) is the most common type of skin cancer. About 2% of people above 60 years suffer from this type of cancer. *PD-1* is an inhibitory molecule belonging to immunoglobulin gene super family which strongly inhibits immune system responses after interaction with its ligands. The aim of present study was to investigate the association of two variants of *PD-1* (*PD1.3G/A* and *PD1.5C/T*) with susceptibility and/or progression of BCC in a population from Southern Iran.

Methods: 168 pathologically confirmed BCC patients (mean age: 61.24±13.09) as well as 167 age-sex-matched healthy individuals with no family history of cancer or autoimmune diseases were recruited. Genomic DNA was extracted from the white blood cells, and genotyping was performed using Polymerase Chain Reaction following by Restriction Fragment Length Polymorphism (PCR-RFLP).

Results: No deviation was observed from Hardy Weinberg Equilibration as confirmed by Arlequin 3.1 software package. Statistical analysis revealed the frequency of AA genotype and A allele of *PD1.3* polymorphism to be significantly lower in the patients comparing to controls (0% vs. 4.19% and 9.38% vs. 17.97% respectively, $P=0.01$). Conversely, the frequency of GG genotype at this position was observed to be significantly increased in the patients comparing to controls (81.25% vs. 68.26% respectively, $P<0.01$). However, no statically significant difference were found in the genotypes' and alleles' frequencies of *PD1.5C/T* between two investigated groups ($P>0.05$). No association was also observed between these genotypes and clinic-pathological parameters in the patients ($P>0.05$). Additionally, the haplotype frequencies were not significantly different between patients and control group ($P>0.05$).

Conclusion: Our results collectively suggest that AA genotypes and A allele at *PD1.3* position may play a protective role, whereas GG genotype at this position may increase the risk of BBC of the skin in Iranian population.

Keywords: Basal Cell Carcinoma, Genotypes, Haplotypes, *PD-1*.

ICIA2018\Cancer Immunology\Poster Discussion\8584

8584

CD39/CD73 expressing B cells in breast cancer draining lymph nodesSahar Shariati^{1,2}, Fereshteh Mehdipour², Morteza Samadi¹, Abdol-Rasoul Talei³, Abbas Ghaderi^{2,4}*1. Department of Immunology, School of Medicine, ShahidSadoughi University of Medical Sciences, Yazd, Iran**2. Shiraz Institute for Cancer Research, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran**3. Breast Diseases Research Center, Shiraz University of Medical Sciences, Shiraz, Iran**4. Department of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran*

Background: CD39 and CD73 are two ectoenzymes which contribute to immune suppression by converting ATP to AMP and AMP to adenosine, respectively. These two molecules are expressed by different cell types including B cells. Some subsets of regulatory B cells were reported to express CD39 and/or CD73. As B cells are a major component of the breast tumor draining lymph nodes (TDLNs) which can influence the anti-tumor immunity, the expression of these molecules by B cells in TDLNs of breast cancer were investigated.

Methods: 20 axillary lymph nodes were studied. Mononuclear cells were isolated and stained with anti-CD19, anti-CD39 and anti-CD73 or their isotype matched antibodies and subjected to flowcytometry.

Results: We found that 94.4 ± 3.1 % and 83.5 ± 7.1 % of B cells were CD39⁺ and CD73⁺, respectively. Regarding the CD39 and CD73 expression we could detect four distinct B cell subsets in the breast TDLNs: Majority of B cells (79.4 ± 7.5 %) were CD39⁺CD73⁺ while 14.5 ± 5.8 %, 3.5 ± 2 % and 2.1 ± 1.8 % of CD19⁺ cells had CD39⁺CD73⁻, CD39⁻CD73⁺ and CD39⁻CD73⁻ phenotypes, respectively. The frequencies of these B cell subpopulations did not show significant differences in the metastatic and non-metastatic lymph nodes, however the percentage of CD39⁻CD73⁻ B cells had a non-significant increasing trend in metastatic lymph nodes ($P=0.081$). Moreover, none of these B cell subsets had significant association with disease parameters such as stage, grade, or the tumor size; However, the frequency of CD39⁻CD73⁺ B cells was significantly lower in stage III compared with stage II ($P=0.020$) and showed significant reverse correlation with the number of involved lymph nodes ($R=0.5$, $P=0.014$).

Conclusion: According to CD39 and CD73 expression, we could detect different B cell subpopulations in the TDLNs of breast cancer. CD39⁻CD73⁺ B cells showed negative associations with poor prognosticators such as stage and the number of involved lymph nodes.

Keywords: Breast cancer, Tumor draining lymph nodes, B cells, CD39, CD73

ICIA2018\Cancer Immunology\Poster Discussion\9783

9783

SiRNA-Mediated Silencing of Snail-1 Induces Apoptosis in Human Breast Cancer Cell Line

Seyed mohammad hossein Musavi shenas¹, Amin kamrani¹, Yunes Panahi^{2*}

1. Tabriz University of Medical Sciences

2. Pharmacotherapy Department, School of Pharmacy, Baqiyatallah University of Medical Sciences, Tehran, Iran

Background: Snail-1 known as one of important Transcription factor is a mediator of survival and cell migration, and expression is raised in numerous cancer types. Snail-1 gene may show a role in recurrence of several cancers including breast cancer by down-regulating E-cadherin, inducing an epithelial to mesenchymal transition and its related miRNA. The aim of this study was to investigate the effect of a specific Snail-1 siRNA on apoptosis and alter EMT related miRNA of MDA-MB-468 (breast cancer) cells.

Methods: The cells were transfected with siRNAs using transfection reagent. The cytotoxic effects of Snail-1 siRNA, on breast cancer cells were determined using MTT assay. Relative Snail-1 mRNA levels were measured by QRT-PCR, respectively. Apoptosis was measured by TUNEL test based on labeling of DNA strand breaks. We also evaluated miR-29b, E-cadherin, vimentin and mmp9 expression by QRT-PCR. To determine alteration in miRNA expression and gene expression involved in EMT.

Results: Snail-1 siRNA significantly reduced mRNA expression levels in a 48 hour after transfection at the concentration of 80 PM in breast cancer cells. We also showed that the silencing of Snail-1 led to the induction of apoptosis and mmp9, vimentin and miR-29b depression as well as E-cadherin overexpression has been shown in snail-1 suppressed group in MDA-MB-468 cells in vitro.

Conclusions: These results propose that Snail-1 might play an important role in the progression of breast cancer, and be a potential therapeutic target for trigger apoptosis and suppression EMT related miRNA and gene in breast cancer.

Keywords: microRNA, Breast cancer

ICIA2018\Cancer Immunology\Poster Discussion\10061
10061

Mesenchymal stem cells derived from stage II of human breast tumor tissue promote the proliferation of the PBMCs: An in vitro assay

Koushansineh sepehr¹, Alireza Razavi^{1*}, Zuhair Mohammad Hassan², AbdolReza Fazel³,
Majid Mossahebi Mohammadi⁴, Seyed Mahmoud Hashemi^{5*}

1. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran
2. Department of Immunology, Faculty of Medical Sciences, TarbiatModares University, Tehran, Iran
3. Cancer research center, Golestan University of Medical Sciences, Gorgan, Iran
4. Sarem Cell Research Center (SCRC), Sarem Women's Hospital, Tehran, Iran
5. Department of Immunology, School of Medicine, ShahidBeheshti University of Medical Sciences, Tehran, Iran

Background: Mesenchymal stem cells (MSCs), as a subpopulation of stromal cells found in the tumor microenvironment, are critical in promoting tumor progression and possess immunomodulatory properties. Therefore, the purpose of this study was to isolate MSCs from primary human breast tumor tissue, and to study the effect of soluble factor of MSC on the peripheral blood mononuclear cell (PBMC) proliferation.

Methods: The tumor tissues (n=8) obtain from patients with pathological stage II of breast tumor. MSCs were isolated by explant culture method and identified. The Mitogen-induced PBMC was treated with 25 and 50% MSC conditioned medium (CM) for 72h, and changes in proliferation evaluated by BrdU ELISA kit.

Results: We successfully isolated and identified MSCs from primary breast tumor tissues. Flow cytometry analysis demonstrated that isolated cells were positive for, CD73, CD44, CD29, CD105 and CD90 but negative for, CD11b, CD45, CD34, and HLADR. In addition, cells possessed the capability of multipotential differentiation into osteoblasts and adipocytes.

The results of the BrdU ELISA assay showed that 50% CM concentrations significantly increased the proliferation of PBMCs.

Conclusion: Collectively, our study showed that MSCs were confirmed to exist in stage II of human breast tumor and significantly promote the proliferation of PBMCs in vitro.

Keywords: Breast Cancer, Mesenchymal Stem Cells, Isolation, Immunomodulatory



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Cancer Immunotherapy

Poster Discussion

ICIA2018\Cancer Immunotherapy\Poster Discussion\5233

5233

Evaluation of The Effect of TIM-3 Suppression by miR-498 and Its Effect on Apoptosis and Proliferation Rate of HL-60 Cell Line

Yaser Moghaddam¹, Mazdak Ganjalikhani Hakemi¹, Alireza Andalib¹, Mohammadreza Sharifi²,
Vida Homayouni¹, Maryam Mhammad Ganji¹

1. Department of Immunology, Faculty of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

2. Department of Genetics and Molecular Biology, School of Medicine, Isfahan University of Medical Science, Isfahan, Iran

Background: Acute Myeloid Leukemia (AML) is a Cancer of hematopoietic stem cells with a rapid progression. TIM-3 is expressed on leukemic stem cells (LSCs) in most types of AML and might have a positive effect on maintenance of malignant phenotype. MicroRNAs play important roles in either cancer progression or suppression. In this study were evaluated, the inhibitory effect of miR-498 on TIM-3 expression and its impact on proliferation and survival of HL-60 cell line.

Methods: Firstly, the probable inhibitory effect of miR-498 on TIM-3 expression was predicted. HL-60 cells were cultured and expression of TIM-3 was induced on the cells using phorbolmyristate acetate. The cells were transfected with miR-498 and expression level of TIM-3 was measured using with q-RT-PCR and flow cytometry methods. In addition, the effect of suppression of TIM-3 expression in HL-60 cell line was analyzed with apoptosis and cell proliferation assays.

Results: Bioinformatics analyses predicted that miR-498 has high ability to silence TIM-3 gene expression. Our experiments confirmed that miR-498 was able to strongly silence TIM-3 expression (68% silencing) in HL-60 cell line ($P < 0.002$). Also, the cells with suppressed expression of TIM-3 had a lower proliferation and higher apoptosis rates.

Conclusion: Based on our results, the miR-498 can effectively suppress TIM-3 expression in the AML cell line. TIM-3 suppression, in turn, inhibits malignant cell proliferation and induces its apoptosis. Collectively, suppression of TIM-3 by miR-498 can be considered as a potential powerful way for treatment of AML.

Keywords: Tim-3, mir-498, miRNA

ICIA2018\Cancer Immunotherapy\Poster Discussion\7567

7567

Effect of different intensities of extremely low frequency electromagnetic fields on expression of STAT6, c-Maf and ROR α genes in spleen and thymus

Hanie Mahaki,^{1,2} Naghi Jabarivasal,³ Khosro Sardanian,^{1,2} Alireza Zamani^{1,2*}

1. Department of Immunology, School of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

2. Molecular Immunology Research Group, Research Center for Molecular Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

3. Department of Medical Physics, Hamadan University of Medical Sciences, Hamadan, Iran

Background: The Thymus and the spleen play an important role in the immune system. Some environmental factors such as extremely low frequency electromagnetic fields (ELF-EMFs) have deleterious effects on them. The present study has been undertaken with the aim to investigate the role of long-term exposure of ELF-EMFs in different intensities on expression of signal transducer and activator of transcription 6 (STAT6), transcription factor Maf (c-Maf) and retinoid-related orphan receptor alpha (ROR α) in the thymus and the spleen of rats.

Methods: The adult male rats were exposed to ELF-EMFs in intensities of 1, 100, 500 and 2000 μ T at frequency of 50 Hz for 2 h daily for up to 60 days. **Results:** Under our experimental conditions, there were no differences observed in the body and spleen weights. Furthermore, the thymus weight was decreased at intensities of 500 and 2000 μ T compared with control. Our experimental results in the spleen showed that, ROR α gene expression was declined in low intensities of 1 and 100 μ T, while no changes were observed after ELF-EMFs exposure on expression levels of c-Maf and STAT6 genes. In addition, mean total levels of ROR α and c-Maf genes among groups were higher in the thymus compare with the spleen. **Conclusion:** The overall mean levels of ROR α and c-Maf gene expressions were higher among the groups in the thymus than the spleen.

Keywords: Signal transducer and activator of transcription 6; Transcription factor Maf; Retinoid-related orphan receptor alpha; Extremely low frequency of electromagnetic fields

ICIA2018\Cancer Immunotherapy\Poster Discussion\52470

5247

Investigating Molecular Mechanism Regulating Egr2 Expression and Its Role in the Immune Response

Sina Taefehshokr^{1*}, Su Ling Li², Amin Oveisi³, YasharAzari Key¹, Sahand Fattahi³,
MohaddesehGhezelchek Khamsiyan³, Nima Taefehshokr²

1. Young Researchers and Elite Club, Tabriz Branch, Islamic Azad University, Tabriz, Iran

2. Division of Bioscience, Department of Life Sciences, College of Health and Life Sciences, Brunel University, UK

3. Faculty of Veterinary Medicine, Tabriz Branch, Islamic Azad University, Tabriz, Iran

Background: The immune system is evolved to defend the body against pathogens and is composed of thousands of complicated and intertwined approaches, which are highly controlled by regulatory processes such as transcription and repression of cellular genes. Early growth response gene (Egr2) is important for maintaining immune stability and it has a vital role in controlling inflammation and preventing the development of autoimmune diseases. The aim of this study was to investigate the molecular mechanism regulating Egr2 expression in the adaptive immunity and to address the vital function of Egr2 in regulating the immune checkpoint markers under melanoma tumor model to obtain effective anti-tumor therapy.

Methods: To investigate the effect of Interferon gamma (IFN γ) and Interleukin 6 (IL-6) on Signal transducer and activator of transcription 1 (STAT1) and STAT3 mediated Egr2 induction, green fluorescent protein (GFP)-Egr2 knock-in cluster of differentiation 4 (CD4⁺)T cells stimulated with IFN γ and IL-6 in the presence of STAT1 inhibitor (Fludarabine) and STAT3 inhibitor (Stattic), assessed for Egr2 expression. Moreover, to pinpoint the function of Egr2 under tumor microenvironment, Egr2/3 KO and GFP Egr2 Knock-in mice ranging from 8-10 weeks of age were used in this study. For experiments mice received 1×10^6 B16 melanoma tumor cells via subcutaneous injection on both left and right flank.

Results: In this study, we have found that Egr2 expression is regulated by antigens and cytokines, including IFN γ and IL-6. Furthermore, it is shown that Egr2 can be significantly expressed in tumor infiltrating lymphocytes (TILs) as we observed in a mouse melanoma tumor model.

Conclusion: Collectively our results demonstrate that Egr2 is an intrinsic regulator in the immune system and tumor microenvironment. It can work as an immune checkpoint in adaptive immunity and tumor condition, suggesting its essential role in effective immune response and anti-tumor therapy.

Keywords: Egr2, IFN γ , IL-6, Immune response

ICIA2018\Cancer Immunotherapy\Poster Discussion\6354

6354

Melatonin as an efficient adjuvant for cancer therapeutic DNA vaccines

Sanaz Baghban Rahimi¹, Amir Ghaemi^{2,3}

1. Department of Microbiology, Golestan University of Medical Sciences, Gorgan, Iran

2. Department of Virology, Pasteur Institute of Iran, Tehran, Iran

3. Department of Microbiology, Infectious Diseases Research Center, Golestan University of Medical Sciences, Gorgan, Iran

Background: The aim of this study is to investigate the effects of melatonin as an adjuvant in E7 DNA vaccines in order to control HPV associated tumors and evaluate effective therapeutic cancer vaccines.

Methods: The anti-tumor efficacy of an HPV-16 E7 DNA vaccine adjuvanted with melatonin (MLT) was evaluated in a C57BL/6 mouse bearing TC-1 tumor model. Post vaccination tumor growth and survival rate of tumor-bearing mice was assessed. The specific lymphocyte proliferation response was measured in control and vaccinated mice by MTT assay. IFN- γ secretion in splenocyte cultures was assessed by ELISA and the antitumor effects were estimated by tumor growth curves.

Results: Our results showed that subcutaneous administration of C57BL/6 mice with a DNA vaccine and MLT dose dependency adjuvant significantly induced strong IFN- γ responses. The high level of E7-specific T-cell proliferation was found in the adjuvant-vaccine group. Furthermore, we found a considerable drop in the tumor volume after immunization mice with MLT plus E7 DNA vaccine and the survival time of TC-1 tumor bearing mice was also significantly prolonged.

Conclusion: These data indicate that melatonin will be used as an effective adjuvant in E7 DNA vaccines and also in other gene vaccines and raised hopes that clinically safe cancer therapeutic vaccines could be designed.

Keywords: Melatonin, HPV, DNA vaccines, Tumor

ICIA2018\Cancer Immunotherapy\Poster Discussion\8541
8541

NK cells Reactivation by Combined Blockade of Tim-3 and PD-1 Inhibitory Receptors for Immunotherapy of Patients with Chronic Lymphocytic Leukemia

Mojgan Astaneh¹, Hadi Hossein Nattaj¹, Ramin Shekariz², Hossein Asgarian Omran^{1,3*}

1. Department of Immunology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

2. Gastrointestinal Cancer Research Center, Mazandaran University of Medical Sciences, Sari, Iran

3. Immunogenetics Research Center, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

Background: Recent studies showed that programmed death-1 (PD-1) and T cell immunoglobulin and mucin domain containing molecule-3 (Tim-3) immune checkpoints receptors are more expressed on cytotoxic T lymphocytes and NK cells in some chronic conditions such as viral infections and malignancies. The exhausted phenotype induced by these receptors makes immune cells inactive as they have impaired cytotoxic activities, degranulation and cytokines secretion. Since the exhaustion process is reversible, in the current study, combined blockade of PD-1 and Tim-3 were tested to restore the functional properties of NK cells in chronic lymphocytic leukemia (CLL).

Method: Fresh peripheral blood mononuclear cells were isolated from 15 patients with CLL and NK cells were positively selected by anti-CD56 microbeads via magnetic cell sorting method. Following treatment of purified NK cells with blocking monoclonal antibodies against PD-1 and Tim-3, their functional activities were assessed in comparison to treatment with isotype matched control antibodies. Induction of apoptosis in K562 cell line was determined by Annexin V-PI flow cytometry method. Degranulation activity of isolated NK cells was evaluated by CD107a degranulation assay and the concentrations of IFN- γ and TNF- α were measured in culture supernatant by ELISA.

Results: The results showed improvement in functional properties of isolated NK cells from CLL patients following blocking of PD-1 and Tim-3 with specific monoclonal antibodies. NK cells treated with blocking antibodies represented more apoptosis induction of K562 cells, increasing in degranulation activity and more production of IFN- γ and TNF- α compared to treatment with control antibodies.

Conclusion: Our results suggested that exhausted NK cells in CLL patients can be reactivated by co-blockade of PD-1 and Tim-3 inhibitory receptors. This strategy could be a potential therapeutic approach for immunotherapy of CLL.

Keywords: Chronic lymphocytic leukemia, Tim-3, PD-1, Checkpoint Inhibitors, Exhaustion, NK

ICIA2018\Cancer Immunotherapy\Poster Discussion\8607
8607

Overexpression of MicroRNA-146a-5p Inhibits Migration of Human MKN-45 Gastric Cancer Cell Line via Down-regulating of Metastasis-related Target Genes

NavidShomali^{1,2}, NaghmehShirafkan^{1,2}, BehzadBaradaran¹

1. Immunology Research Center (IRC), Tabriz University of Medical Sciences, Tabriz, Iran

2. Student Research Committee, Tabriz University of Medical Sciences, Tabriz, Iran

Background: The second cause of cancer-related death worldwide is gastric cancer. MicroRNAs are important post-transcriptional regulators that acts as negative regulators of gene expression. They do this role through base-pairing with the 3-UTR (3-untranslated region) region of their target genes. MiR-146a is one of the important microRNAs that plays a role in pathogenesis of wide ranges of human disorders. Though evidences have shown miR-146a as a strong metastasis and invasion inhibitor of gastric cancer, but the molecular mechanisms and the related genes are mainly unknown.

Methods: After miR-146a transfection, quantitative RT-PCR was performed to detect expression level of miR-146a and related target genes including L1CAM, ROCK1, MMP9, HMGA2, CXCR4 and vimentin in GC cell line. Wound-healing assay was performed to measure MKN-45 cells migration after miR-146a transfection. MTT and Annexin V staining were used to determine the role of miR-146a in the regulation of MKN-45 cell viability and apoptosis induction, correspondingly.

Results: We showed that miR-146a has low expression level in MKN-45 gastric cancer cell line. in the same way, we showed increased expression of miR-146a can suppress MKN-45 metastasis capability. Effect of miR-146a replacement on target genes showed that except for the CXCR4, which was not meaningful in the result, other related target genes showed reduced expression. The results of miR-146a replacement on cell viability and apoptosis showed it had not a significant effect on cell viability and apoptosis induction.

Conclusion: Our results indicated that miR-146a, may act as tumor suppressor microRNA in MKN 45 cell line. The freshly identified miR-146a target genes in gastric cancer offers a novel insight into the biological basis of migration in gastric cancer and offers miR-146a as a potential therapeutic approach to suppress gastric cancer migration.

Keywords: Gastric cancer, microRNA, Metastasis, Cell viability, Apoptosis

ICIA2018\Cancer Immunotherapy\Poster Discussion\8640
8640

Polarized Mesenchymal Stem Cells-Microvesicles with Whole Tumor Lysates Pulsed DCs Promote T Cells Cytotoxicity against Glial Cancer Cells

RahmaniKukia Nasim¹, Abbasi Ardeshir², Delirez Nowruz³

1. Division of Immunology, Department of Microbiology, Veterinary Faculty, Urmia University, Urmia, Iran

2. Division of Immunology, TarbiatModares University of Medical Sciences, Tehran, Iran

3. Associate professor, Division of Immunology, Department of Microbiology, Veterinary Faculty, Urmia University, Urmia, Iran

Background: Cancer immunotherapy using dendritic cells (DCs) and peptides is considered as a new promising treatment modality and capable of inducing an anti-cancer response. Mesenchymal Stem Cells (MSCs) migrate to tumor microenvironments and contribute many bioactive factors such as microvesicles (MVs). The MSC-MVs as native or engineered stem cells have similar protective and reparative properties as their cellular counterparts in tissue repair and possibly anti-cancer therapy. Recent studies indicate that polarization of MSCs with Lipopolysaccharide (LPS) can reduce tumor growth and metastasis.

Methods: Peripheral blood monocyte derived DCs were isolated by heart puncture of rats and incubated with GM-CSF, IL-4 and addition of LPS at day 6 to induce maturation of the DCs. MSC-MVs were purified by differential ultracentrifugation after isolation, proliferation and treatment of MSCs with LPS. Autologous T cells were incubated with whole tumor lysate pulsed DCs in the presence of polarized MSC-MVs. Delicate T cells were co-cultured with B₉₂ glial cancer cells in 96-well plates at final volume of 200 µl of medium at an effector:target ratio of 100:1 to evaluate specific cytotoxic activity.

Results: The flow cytometric analysis of cytotoxic T cells showed that using polarized MSC-MVs with tumor lysate induce glial cancer cells death. These results represented that early and late apoptosis was stimulated in tumor cells in compare to controls and were responsible for cells death. Also, the specific lysis of tumor cells was promoted by autologous tumor specific T cell responses *in vitro* in the presence of polarized MSC- MVs.

Conclusion: It appears that an alternative strategy for effective anticancer responses may be the use of polarized MSC- MVs and tumor lysate to elicit T cell responses in terms of proliferation and cytotoxicity.

Keywords: Polarized MSC- MVs, Cancer immunotherapy, Tumor lysate, Cytotoxic T cells, Glial cells, Apoptosis

ICIA2018\Cancer Immunotherapy\Poster Discussion\8650

8650

Antitumor Effects of Liposomes-encapsulated CTLA-4 Immunotherapy Combined with Doxil Chemotherapy in the B16 Large Established Tumors of Melanoma

Seyedamir Jalali¹, Razie Alibeigi²

1. Immunology Department, Medical School, Shahid Beheshti university of medical sciences

2. Immunology Department, Medical School, Mashhad university of medical sciences

Background: Cancer therapy is designed to specifically integrate distinct treatment modalities in the most effective way to achieve the highest cure rate. Chemotherapy has pleiotropic effects on the immune system. The vast majority of chemotherapeutic agents are myelosuppressive and therefore could deplete the very cells responsive for antitumor immunity. In the presented studies, we describe synergy between CTLA-4 blockade, Liposomes-encapsulated CTLA-4 blockade antibody and chemotherapeutic agents, Doxil, in large established tumors in the mice models of melanoma and we were shown the effect of timing immunotherapeutic agent (CTLA-4 blockade antibody) and chemotherapeutic agent (Doxil) on antitumor responses.

Methods: Liposomes were prepared by thin lipid film hydration plus extrusion. The stock chloroform solution of ingredients For PEGylated liposomes, mPEG2000-DSPE, and cholesterol. The lipid film was hydrated with 10 mM phosphate-buffered saline containing 1 mg/ml Anti CTLA-4 monoclonal antibody. Ten days after tumor challenges, mice with large established tumors were randomized into eight different treatment groups of five mice each as follows: 1) free CTLA-4 blockade antibody, 2) CTLA-4 PEG-liposomes, 3) Doxil, 4) Concomitant CTLA-PEG-liposomes and Doxil, 5) concomitant free CTLA-4 blockade antibody and Doxil, 6) First Doxil so free CTLA-4 blockade antibody, 7) free CTLA-4 blockade antibody, 8) PBS and mice were administered intravenously with 100 µg of free or liposomal forms of CTLA-4 blockade antibody and/or single dose of 5 mg/kg Doxil. Flow cytometer analysis of drainage lymph node, tumor infiltrated lymphocytes and spleen were liposomal CTLA-4 blockade antibody with Doxil group (6.17%, 26.1% and 5.5%, 27% respectively).

Result: Our results demonstrating that the optimal timing of CTLA-4 blockade antibody is before Doxil administration in compare with first Doxil CTLA-4 blockade antibody and concomitant administration of Doxil and CTLA-4 blockade antibody

Conclusion: In summary, our findings give strong preclinical rationale for clinical evaluation of Doxil.

Keywords: Doxil, Anti CTLA-4, Chemoimmunotherapy

ICIA2018\Cancer Immunotherapy\Poster Discussion\9793
9793

Accumulative strategy of FoxP3-specific recombinant vaccine and tumor-specific DC vaccine against melanoma in mice

NedaMousaviNiri¹, Maryam Naseroleslami², NosratollahZarghami³, Nasimmohammad kheshtchin⁴,
Jamshid Hadjati⁴

1. Department of Medical Biotechnology, School of Advanced Medical Sciences, Tehran Medical Sciences branch, Azad Islamic university, Tehran, Iran

2. Department of Cellular and Molecular Biology, School of Advanced Medical Sciences, Tehran Medical Sciences branch, Azad Islamic university, Tehran, Iran

3. Department of Medical Biotechnology, School of Advanced Medical Sciences and Department of Clinical Biochemistry, Radiopharmacy Lab, Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

4. Department of Immunology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

Background: Treg apparently have a substantial role in suppression of anti-tumor immune responses. Application of recombinant FoxP3-Fc vaccine inducing anti-Treg immune responses accompanied with tumor antigen expressing dendritic cell therapy is a hopeful approach in anti-tumor treatments.

Methods: Instead of anti Treg vaccine, FoxP3-Fc fusion construct was applied to produce DNA vaccine and its respective recombinant protein. Afterward they were subcutaneously injected into C57/B6 mice followed by melanoma tumor induction (two weeks after last vaccination) and dendritic cell-based anti-tumor immunotherapy. The improvement of immunotherapy results in mice that received anti-Treg vaccination was evaluated by different immune assays compared to other groups.

Results: combinational therapy by dendritic cells and anti-Treg vaccine showed increment in immunological results compared to sole vaccination of anti Treg. Proliferation and IFN- γ secretion of Tumor specific lymphocytes were augmented significantly in dendritic cell and anti-Treg vaccinated group of mice. Tumor growth and survival as well indicated more efficient results in the same group of treatment.

Conclusion: Due to the prospering immune protection of FoxP3-Fc (IgG) vaccine against Treg and its impression in improvement of dendritic cell-based immunotherapy, there would be a good prospect for applied strategy and other means of immunotherapy.

Keywords: Treg, FoxP3, Dendritic cells, Melanoma, Immunotherapy

ICIA2018\Cancer Immunotherapy\Poster Discussion\10012

10012

Recombinant HER2 Protein Vaccination Induces Cellular Immune Response and Tumor Growth Inhibition in Syngeneic Breast Cancer Mice Model

Fateme Sadri Ardalani^{1,2}, Moslem Ahmadi^{1,2}, Mohammad Mehdi Amiri¹, Mahmood Jeddi Tehrani², Fazel Shokri^{1,2}, Mahdi Shabani^{3,2}

1. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

2. Monoclonal Antibody Research Center, Avicenna Research Institute, ACECR, Tehran, Iran

3. Department of Immunology, School of Medicine, ShahidBeheshti University of Medical Sciences, Tehran, Iran

Background: Cellular and protective immune responses against the full HER2 extracellular domain (fHER2-ECD) recombinant protein were investigated in breast cancer mice model.

Methods: Balb/C mice were immunized with produced recombinant eukaryotic fHER2-ECD protein in combination with or without CpG to investigate their potential for induction of HER2 specific cell-mediated immune responses and tumor growth inhibition in vivo.

Results: Immunization resulted in induction of antibody response and induced secretion of IL4 and particularly IFN γ and IL17 cytokines. Challenging of immunized mice with stable 4T1-HER2 transfected cells resulted in partial but significant tumor growth inhibition in the immunized mice (about 8% tumor free in with and without CpG groups) particularly those immunized with fHER2-ECD together with CPG.

Conclusion: The fHER2-ECD along with CpG was a successful formula to induce IFN γ production. The fHER2-ECD induces tumor growth inhibition in immunized mice compared to control groups.

Keywords: Breast Cancer, Cytokine, HER2, Protein immunization, Tumor challenge



Computational Immunology and Systems Biology

Poster Discussion

14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Computational Immunology and Systems Biology\Poster Discussion\7986
7986

Detection of Blood cells' Genes Involved in Responding Sepsis by Weighted Gene Co-expression Network Analysis

Rasoul Godini¹, Fatemeh Hadi¹, Hossein Fallahi¹

1. Department of Biology, Faculty of Science, Razi University

Background: Sepsis, as a systemic inflammation disorder is known as a clinical syndrome caused by infection. Sepsis is a life threatening condition and early recognition and proper treatment is vital to reduce mortality. In here, we use Weighted Gene Co-expression Network Analysis (WGCNA) to detect which genes are most important in Dendritic Cells (DCs) and Peripheral Blood Mononuclear Cells (PBMCs) responding to sepsis.

Methods: In order to do this, we used free available dataset GSE49758. This data contains more than 80 microarray analysis from samples of two cell types, DCs and PBMCs. After preprocessing the matrices for genes with variation, outliers were removed. Then WGCNA analysis was performed to detect associated genes to disease symptoms. Soft threshold of 0.8 was used to detect modules. Hub genes in each module detected and their expression values determined.

Results: Our results showed that there were 4 related modules with the sepsis in each cell type. Searching for the correlations between genes' module membership and significance for the sepsis, we found that two modules with a correlation = 0.44 (p-value = 3.7e-10) and correlation = 0.47 (p-value = 1.6e-31) that are related to immune and inflammatory response in DCs and PBMCs, respectively. Looking for hub genes among members of these modules, 185 in DCs and 549 PBMCs modules, ADORA2A and CD163 in DCs and VIPR1 and LTB in PBMCs were detected as hub genes, which were also differentially expressed. We found 27 and 99 differentially expressed genes in modules regarding to DCs and PBMCs, respectively, however, only a few of them was hub differentially expressed genes, as well.

Conclusion: Results of this study can help not only to find markers to detect sepsis and possibly proper treatments, but also shed lights upon molecular information of sepsis.

Keywords: Dendritic cells, PBM cells, Sepsis, WGCNA

ICIA2018\Computational Immunology and Systems Biology\Poster Discussion\8300
8300

***In silico* Design of Dodecameric Human Soluble CD40L Using Surfactant Protein-D**

Bahare Shokoohian¹, Mahshoid Mohammadipour¹, Fatemeh Yari¹

1. Blood Transfusion Research Center, High Institute for Research and Education in Transfusion Medicine, Tehran, Iran

Background: CD40L is a membrane protein and a member of TNF family that plays an important role in cell signaling in innate and adaptive immunity. Since this protein plays its role through clustering the receptors, CD40, on the target cell surface and also natural soluble form of CD40L is not stable, it is expected that the multimeric form of CD40L can have a stronger effect than the natural trimeric form.

Methods: To design the dodecameric form (four-trimer) of CD40L, aminoacid sequences of CD40L and surfactant protein D (SPD), were obtained from the NCBI protein data bank. To identify consensus sequence, alignment was performed by ClustalW method. Extracellular domain of CD40L was replaced with lectin domain of SP-D and primary construct with the length of 472 aminoacid was built. Physicochemical characteristics and stability of chimeric protein, SPD-CD40L, were determined by ProtParam software. To determine protein localization in eukaryotic expression host, HEK293, PSORTII software was used. Secondary and tertiary structures of the SPD-CD40L were predicted using PSIPRED server and I-TASSER software respectively. Validity of the predicted 3D structure was evaluated by the Ramachandran plot drawn by RAMPAGE. To obtain nucleotidic sequence and optimization of sequence for expression in eukaryotic cells, OPTIMIZER server was used.

Results: Molecular weight was measured 49145.4 Da and the half-life were greater than 30 hours in eukaryotic cells. SPD-CD40L was classified as stable. PSORTII results indicated that protein is secretory. Tertiary structure prediction revealed distinct folding of two parts of SPD-CD40L. RAMPAGE results showed that 95.3% residues clustered in favored region, therefore formation of desired structure between the predicted structures is probable.

Conclusion: We designed the stable soluble multi-trimeric structure of CD40L using SPD that seems to increase clustering of CD40 on the target cells surface and mimics the membranous CD40L orientation.

Keywords: CD40L, Multimeric protein, Surfactant protein D, TNF family

ICIA2018\Computational Immunology and Systems Biology\Poster Discussion\8451
8451

Assessment the Structure and Function of The Model of Cintredekinbesudotox Based on *In-silico* Investigation

Fatemeh Samiezadeh Mashadi¹, Bahareh Hasani Sabzevar², Nazanin Gholampour Feroji²,
Mohammad Rastegar Moghadam^{1,2}, Aliakbar Haddad Mashadrizesh³, Mehran Akbari¹

1. Pharmaceutical Research Group, Khayyam Bioeconomy Institute, Mashhad, Iran

2. Structural biology and Bioinformatics Research Group, Khayyam Bioeconomy Institute, Mashhad, Iran

3. Cell and Molecular Biotechnology Research Group, Institute of Biotechnology, Ferdowsi University of Mashhad, Mashhad, Iran

Background: Immunotoxins are effective and promising drugs for target therapy of cancer, which in turn nowadays has a special prospective in the healthcare system. In this regard, IL-13-PE38QQR which named as Cintredekinbesudotox is one of them which has developed for Glioma healing up to level 2 therapies. Therefore, recognizing the structure and function of this drug could be provide approach for its production based on knowledge, as well as its optimization for achieve higher levels of therapies which are considered in this study based on in-silico biology.

Methods: In this study, NCBI and RCSB databanks were used to achievement to the protein sequences and 3D structures of this drug. Modeller software were employed to assembling the fragment sequences of drug via homology modeling method. Moreover, ERRAT, and RAMPAGE programs were used to determine the quality of the structures of the models. Evaluation the thermostability of the model of this drug in quasi-physiological condition was carried out by GROMACS using GROMOS force fields. Finally, the functionality features of this drug were assayed by their affinity to IL13-R as corresponding antigen as well as immunogenicity properties by HADDOCK and IEDB programs, respectively. Visualizing the structures and interactions was performed via The PYMOL software.

Results: Surveying the sequence of IL-13-PE38QQR has led to reveal domains of IL-2 and *Pseudomonas* Exotoxin A in its context. Modeling and assembling of these fragments with flexible linker led to the production 10 models with various quality and structure. The best model showed stability structure under quasi-physiological circumstance with reasonable immunogenicity. Moreover, its affinity to corresponding antigen compared to negative control was confirmed.

Conclusion: Generally, the results of this study has led to introduced a model of IL-13-PE38QQR immunotoxin, which is suitable in the structure and function in quasi-physiological condition, so can using for its production.

Keywords: Immunotoxin, Cintredekinbesudotox, Glioma, Databases, GROMACS

ICIA2018\Computational Immunology and Systems Biology\Poster Discussion\9866
9866

Common Gene Networks Contribute in Hashimoto and Urticaria Disorders

Nazanin Atieh Kazemi sefat¹, Saeed Talebi²

1. Department of Medical Immunology, Faculty of Medical Sciences, Tarbiat Modares University (TMU), Tehran, Iran

2. Department of Medical Genetics and Molecular biology, Iran University of Medical Sciences (IUMS), Tehran, Iran

Background: Urticaria is common in some patients with Hashimoto thyroiditis. In both disorders the immune system function is affected. In this study we have determined involved common gene networks in these disorders according to the gene ontologies (GO) and molecular pathways.

Methods: A comprehensive literature search was done to find important suggested genes in hashimoto thyroiditis or urticaria. In order to determine the gene networks that were involved, network based enrichment analysis for KEGG and NCI pathway, Biological process, Molecular function, Cellular component was done (on common genes in these disorders) using “EnrichNet” Online tool. Pathways or processes by 5 or more affected genes and significant fisher exact test ($p < 0.05$) were analyzed.

Results: We have found 144 important genes contribute in urticaria and 132 genes involved in hashimoto thyroiditis. 28 genes were common in both disorders. In KEGG pathways most genes were contributed in “Cytokine-cytokine receptor interaction. In both NCI pathways and GO biological processes, IL-12 mediated signaling pathway covered most genes. Gene products typically localized to both extracellular space and plasma membrane in GO cellular component. Gene products with protein binding and cytokine activity were more common in analysis with GO molecular function. IFNG, IL18, IL2 and IL4 were most important cytokine genes and IL2RA and IL2RB were the most common cytokine receptor genes in these pathways.

Conclusion: Cytokine-cytokine receptor gene network specifically the interaction between Interleukin 2 and Interleukin 2 receptor in IL12-mediated signaling pathway may contribute in both urticaria and hashimoto thyroiditis.

Keywords: Enrichment Analysis, Urticaria, Hashimoto thyroiditis

ICIA2018\Computational Immunology and Systems Biology\Poster Discussion\9986
9986

Accurate Prediction Approach Result in Increased Probabilities of Victory: has-miR-27 is a New Potential Autoimmune-Deregulated miRNA

Vahedeh Hosseini¹, Abdolhakim Hosseini², Kobra Mehdinejadani³, Shayesteh Mehdinejadani⁴, Vahid Kia⁵, Samira Mohammadi Yeganeh¹

1. Department of Biotechnology, School of Advanced Technologies in Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Department of Biotechnology, School of Advanced Technologies in Medicine,, Hamedan University of Medical Science, Hamedan, Iran

3. Department of Immunology, Faculty of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

4. Department of Anatomy, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran.

5. Zanzan University of Medical Sciences, Zanzan, Iran

Background: Th17 cells, a subset of CD4⁺ T helper cells, are attracting considerable interest due to their critical role in the pathogenesis of different autoimmune disorders. MicroRNAs are a category of small non-coding RNA that function in a variety of cellular processes by silencing the target mRNA. Several positive and negative regulators govern the Th17 differentiation. Dysregulation of miRNA expression is associated with different autoimmune diseases. The aim of our research was to expand current knowledge of miRNA role as a Th17 differentiation regulator.

Methods: Our proposed search direction for predicting an appropriate miRNA that regulate Th17 differentiation is as follows: systematic literature mining; search in miRvestigator; predict the target genes of the selected miRNA in TargetScan and miRWalk, and final approval in Gene Expression Omnibus databases.

Results: We carried out a systematic literature mining for autoimmune-deregulated mRNAs of Th17 differentiation regulators. Then, using miRvestigator toll, we predicted a miRNA that can target over-represented sequence motif of submitted genes (negative regulators of Th17 differentiation). Among all suggested miRNAs, hsa-miR-27 can target over-represented sequence motif of submitted genes (*P-value*= 1.5e-05). Further bioinformatics studies were performed by TargetScan toll and integrative miRWalk database. To identify the expression pattern of has-miR-27 in Th17 differentiation, microarray dataset GSE75909 was downloaded from the Gene Expression Omnibus database and this data was analyzed by GEO2R to determine differentially expressed miRNAs (DEMs). The result of microarray data shows that has-miR-27 can promote the differentiation of Th17.

Conclusion: Target predictions of has-miR-27 and microarray data indicate its interaction with negative regulators of Th17 differentiation. Therefore, it can be reasonably assumed that inhibitors of hsa-miR-27 can be used as a treatment for autoimmune diseases.

Keywords: miRNA, Th17 cells, autoimmune disease

Diagnostic Methods in Immunology

Poster Discussion

ICIA2018\Diagnostic Methods in Immunology\Poster Discussion\7573

7573

A Novel Sampling and Diagnostic Method for Cutaneous Leishmaniasis

Yasaman Taslimi¹, Pardis Sadeghipour¹, Sima Habibzadeh¹, Vahid Mashayekhi², Hossien Mortazavi³, Housein Malekafzali Ardakani¹, Ingrid Muller⁴, Pascale Kropf^{4*}, Sima Rafati^{1*}

1. Department of Immunotherapy and Leishmania Vaccine Research, Pasteur Institute of Iran, Tehran, Iran

2. Skin center, Mashhad Medical School, Mashhad, Iran

3. Razi Skin Center, Tehran Medical School, Tehran, Iran

4. Section of Immunology, Department of Medicine, Imperial College London, London, United Kingdom

Background: *Leishmania major* and *tropica* are causative agents for CL in Iran causing either permanent lesions or scars in the skin. Accurate diagnosis of CL is crucial in order to the use of chemotherapy with highly toxic drugs such as glucantime. The gold standard for CL diagnosis involves invasive sampling and recognition of parasite in smears and cultures. Albeit these techniques are highly specific but they are not sensitive enough and require technical expertise. Additionally these approaches run the risks of secondary infection and considerable discomfort, especially for children and sensitive body areas. In this report, we present a novel rapid (>1minute), user-friendly and painless sampling method (tape-stripping) of skin to collect DNA from the lesion to perform accurate non-invasive diagnosis method.

Methods: More than 100 patients suspected of CL, 20 fungal-infected lesions and 50 areas of normal skin were examined. DNA isolated from the skin tape-stripping samples, were amplified using kDNA1 minicircle primer. In parallel, DNA was subjected to ITS1 PCR-RFLP for species identification. Standard reference strains of *L. major* and *L. tropica* were used as positive controls.

Results: Among 80 acute cases, half of them were diagnosed as *L. tropica* the rest diagnosed as *L. major*, additionally among 30 chronic CL samples, more than half of them were diagnosed as *L. tropica*, 11 samples as *L. major*. Among 8 healed cases, 7 cases were diagnosed as *L. tropica* and one sample as *L. major*.

Conclusion: In this study, we obtained DNA from around 170 CL patients and healthy control for PCR performance using of non-invasive diagnostic sampling. We confirmed the *Leishmania* infection as well as the species in both acute and chronic form. Our next step is to design a point of care diagnosis method for CL based on mentioned non-invasive sampling method.

Keywords: Cutaneous Leishmaniasis, Non-invasive Diagnosis, Skin-tape Stripping

ICIA2018\Diagnostic Methods in Immunology\Poster Discussion\7697

7697

Detecting Epithelial Cell Adhesion Molecule in Human Carcinoma Cell Lines Using a Developed Polyclonal Antibody

Mehrzaad Ghari¹, Saeideh Milani², Gholamreza Javadi¹, Ali Ahmad Bayat², Reza Hadavi², Hodjattallah Rabbani², Fatemeh Ghaemimanesh²

1. Department of Biology, Science and Research Branch, Islamic Azad University, Tehran, Iran

2. Antigen and Antibody Engineering Department, Monoclonal Antibody Research Center, Avicenna Research Institute, ACECR, Tehran, Iran

Background: Epithelial cell adhesion molecule (EpCAM) is a transmembrane glycoprotein mediating epithelial-specific intercellular cell adhesion. EpCAM is also involved in cell signaling, migration, proliferation and differentiation. The overexpression of EpCAM in various types of human carcinoma makes this molecule to be considered as an interesting diagnostic marker. Therefore, the aim of this study was to produce a specific polyclonal antibody (pAb) against EpCAM to be used as a diagnostic tool in human carcinomas.

Methods: A white New Zealand rabbit was immunized using keyhole limpet hemocyanin (KLH)-conjugated peptide derived from EpCAM molecule. The antibody was purified from immunized rabbit sera using a peptide- affinity chromatography column. The capability of pAb in detection of EpCAM was investigated in different protein read-out systems.

Results: Our results showed that anti-EpCAM pAb specifically detects EpCAM in various human carcinoma cell lines of MCF7, A549, SW480 and A431 by ICC. The IHC experiments showed the expression of EpCAM in human prostate and bladder carcinoma tissues.

Conclusion: The developed pAb might be used as a diagnostic tool in human carcinomas in ICC and IHC applications.

Keywords: Polyclonal Antibody, EpCAM, Cancer, Detection

ICIA2018\Diagnostic Methods in Immunology\Poster Discussion\8099
8099

Determination of *Helicobacter pylori* Antigen in Stool Samples and Comparison with Rapid Urease Test in Patients Suspected of Helicobacter Infection

Reza Marvinam¹, HosseinHadinedoushan¹, Mahdi Dehghanmanshadi¹, FatemeZare¹

1. Department of Immunology, Reproductive Immunology Research Center, ShahidSadoughi University of Medical Sciences, Yazd, Iran

Background: *Helicobacter pylori* (*H. Pylori*) is the microorganism that infects nearly half of world's population. There are several invasive and non-invasive methods for diagnosis of infection. The main objective of this study was to evaluate antigen of *H. Pylori* in feces with *H. pylori* stool antigen (HpSA) test and comparison with rapid urease test (RUT) in the patients suspected to be infected.

Methods: one hundred thirty-seven subjects (56 males, 81 females) were recruited from those patients undergoing a gastrointestinal endoscopic examination in the endoscopy units of Shahid Sadoughi University of medical sciences. One biopsy specimen was obtained from the stomach and each biopsy specimen per subject was tested for the presence of urease using the commercially available CLO test. Stool specimens were taken concurrently with the endoscopic examination and tested by Enzyme-linked immunosorbent assay (ELISA) method for presence of HpSA. In this study, RUT was considered as a gold standard test.

Results: The mean age was 40.4 ± 1.12 years. 13.3% of samples were shown HpSA-positive and negative RUT and 12.9% were shown HpSA-negative and positive RUT. Sensitivity and specificity of HpSA test was 86.6% and 87.1%, respectively. Positive and negative predictive values and accuracy were 89%, 84.3%, 86.2%, respectively.

Conclusion: Our findings showed that the stool enzyme immunoassay for *H. pylori* is a useful method for the primary diagnosis of *H. pylori* in the patients suspected to be infected.

Keywords: ELISA, *H. Pylori*, HpSA, Rapid Urease Test

ICIA2018\Diagnostic Methods in Immunology\Poster Discussion\8570
8570

Optimization of HER2-based ELISA Immunoassay Using Trastuzumab

Maryam Ahmadzadeh^{1,2}, Elham Mohit¹, Leila Nematollahi³, Farzaneh Farshdari⁴

1. Department of Pharmaceutical Biotechnology, School of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Student's Research Committee, Shahid Beheshti University of Medical Sciences, Tehran, Iran

3. Biotechnology Research center, Pasteur Institute of Iran, Tehran, Iran

4. Department of Biotechnology, Faculty of Advanced Sciences & Technology, Pharmaceutical Sciences Branch, Islamic azad University, Tehran- Iran (IAUPS)

Background: Enzyme-linked immunosorbent assay (ELISA) is a sensitive, specific, reproducible and fast method for detection of antigen-antibody reactions. Currently, ELISA is the most common immunoassay approach used to evaluate the biological activity of recombinant whole or fragment of antibody. However, due to the complex nature of immunoassays that involve multiple assay steps using multiple reagents, many factors affect the ELISA performance and should be optimized.

Trastuzumab, a humanized monoclonal antibody directed against the epidermal growth factor receptor 2 (HER2), is used for the treatment of breast cancer patients who overexpress the HER2 receptor. Herein, commercial Trastuzumab was used as a positive control in developing HER2- based ELISA immunoassay for anti-HER2 antibodies/antibody fragments targeting the same epitope as Trastuzumab. The aim of this study was to develop an acceptable HER2-based ELISA procedure using trastuzumab as positive control. Therefore, different factors such as antigen concentration, the type of blocking agent and sample concentration were optimized.

Methods: Accordingly, serial dilution of HER2 antigen (7.8 to 500 ng/ μ L), skim milk 5% and BSA (1 & 3 %) two type of blocking agent and different concentration of trastuzumab (0, 1, 10, 100 ng/ μ L) were analyzed. Anti-human IgG (whole molecule) peroxidase-conjugate (1:5000) and 3, 3', 5, 5'-Tetramethylbenzidine (TMB) were used as detection antibody and substrate, respectively. The reaction was stopped using sulphuric acid and absorbance was recorded at a wavelength of 450 nm.

Results: The lowest concentration of HER2 antigen which has the ability to detect trastuzumab was 7.8 ng/ μ L. Skim milk 5% demonstrated minimal non-specific background and was selected as the optimum blocking agent. Also, the optimal concentration of trastuzumab was between 10 and 100 ng/ μ L which fallen within the linear portion of the curve (absorbance against concentration of trastuzumab).

Conclusion: It was demonstrated that the type of blocking agent is an important factor affecting the performance of the HER2-based assay. The result of this study led to the improvement of the HER2-base ELISA assay and can be used for developing the ELISA method for other anti-HER2 antibodies.

Keyword: ELISA, Trastuzumab, HER2 antigen, Optimization.

ICIA2018\Diagnostic Methods in Immunology\Poster Discussion\8622

8622

Assessment of Serum Level of Stromal Derived Factor-1 α (SDF-1 α) in Serum of Patients with Sepsis at Admission and Discharge in Ali-ebne-Abitaleb at 2013

Marzie Motaghy¹, Mohammad Kamali², Gholamhossein Hasanshahi³

1. Postgraduate student of Hematology, Kerman University of Medical Sciences, Kerman, Iran

2. Medical Student, Rafsanjan University of Medical Sciences, Rafsanjan, Iran

3. Molecular Medicine Research Center, Rafsanjan University of Medical Sciences, Rafsanjan, Iran

Background: Sepsis is a major cause of death and the most common cause of death in the intensive care unit at a rate of 30 to 70 percent. SDF-1 α as an ELR negative CXC chemokine is effective in Call of inflammatory cells in the blood and the role of chemokine activity by a process known as angiogenesis. This study compared the levels of these chemokines in patients with sepsis at admission and discharge based on age and gender studies.

Methods: Fifty four sepsis patients who were enrolled to the study Peripheral blood samples were collected from patients in parallel with demographic data. Methods: SDF-1 α (CXCL12) levels in the samples were measured by ELISA. SPSS, T-test and Chi-square and ANOVA were utilized for data analysis.

Results: The 54 patients participating in the project, 41 were male (75/9%) and 13 females (24/07%) respectively. The minimum age of participants was 51 years and a maximum of 90 years in the project and the mean age was 69/13 years. The mean values of serum chemokines in patients with severe sepsis in the peripheral blood at the time of hospital discharge was significantly reduced and the differences were statistically significant differences ($P < 0/0001$). Average concentrations at admission and discharge were 222/780 and 58/980 respectively. Compared to the overall mean concentration of serum chemokines in patients with severe sepsis at admission by sex and age in peripheral blood concentrations were approximately equal in terms of differences not statistically significant ($P = 0/564$, $P = 0/818$). Compared to the overall mean concentration of serum chemokines in peripheral blood of patients with sepsis at the time of discharge based on sex, age and concentration was almost the same in terms of the differences was not statistically significant ($P = 0/193$ and $P = 0/816$).

Conclusion: According to the results of our study showed that the concentration of this chemokine was not directly associated with age and sex. Finally, our study suggest that SDF-1 α (CXCL12) could possibly use as an early diagnosis marker for sepsis and perhaps other systemic inflammatory syndrome as a new treatment.

Keywords: Sepsis, SDF-1 α , Inflammatory cells.

ICIA2018\Diagnostic Methods in Immunology\Poster Discussion\9750
9750

The Exact Method for Determining the Dose and Type of Chemotherapy Drug

Amirhossein Alimohammadian¹, Negin Maleki²

1. Member of the brilliant talent of young and elite peoples youth club of Ardabil Islamic Azad University, Iran
2. Department of medical laboratory, Ardabil branch, Islamic Azad University, Ardabil, Iran

Background: One of the ways to cure various types of cancer and even after chemotherapy. All types of chemotherapy drugs are somehow inhibitors of cancer cells, but the main problem is that no chemotherapy drug has any significant effect on any type of cancer cell and There is no general method for testing the type and dosage of chemotherapy and the oncologist, based on the type of cancer and its origin, is always using an experiment with an error test, despite the multiplicity of chemotherapy drugs. Especially in malignant tumors, the passage of time with the initial test and error is very unfortunate and irreparable.

Methods: An ELISA plate containing tissue and cell in a suitable medium from RPM or any other medium (for breast cancer cells, prostate cancer, esophageal stomach and leukemia and ...) and a supplement plate containing a chemotherapy drug that was placed on an ELISA plate after 24 hours. Ultimately, an NBT resuscitation spectrometer was detected in aqueous ampullazer with specific wavelengths controlled by the drug and determined by regimen of the drug.

Results: This method makes it possible to customize the chemotherapy of each patient and monitor the stages of treatment, predict and control possible complications, and prescribe appropriate drugs for the continuation of treatment.

Conclusion: With this method, it is possible to diagnose the effect of the drug that is requested by the oncologist and the physician, and calculated by the laboratory.

Keywords: Cancer, Chemotherapy, NBT, Diagnose

ICIA2018\Diagnostic Methods in Immunology\Poster Discussion\9818

9818

Urinary levels of Endothelin-1, Monocyte Chemotactic Peptide-1 and N-acetyl- Glucosaminidase as Prognostic Markers for Severity of Obstruction in Hydronephrotic Neonates

Fatemeh Ezzatifar^{1,2}, Hamid Mohamad Jafari¹, Alireza Rafiei¹

1. *Molecular and Cell biology Research Center , Faculty of Medicine , Mazandaran University of Medical Sciences , Sari , Iran*

2. *Department of Immunology Mazandaran University of Medical Sciences, Sari, Iran*

3. *Student Research Committee, Mazandaran University of Medical Sciences, Sari, Iran*

Background: Antenatal hydronephrosis (AH) is found in 0.5% of neonates, the majority of them had obstructive entity such as UOJO or UVJO. In order to diagnose severity of obstruction, we required invasive testing and exposure of the neonate to radiation. The urinary levels of Endothelin-1(ET-1), monocyte chemotactic peptide-1(MCP-1) and N-acetyl- glucosaminidase (NAG) and their ratios to creatinine were assessed in severely hydronephrotic newborns and were compared to those with milder forms and those without obstruction.

Methods: The neonates with a history of prenatal hydronephrosis were enrolled in three groups based on imaging study results, group 1 the neonates with severe forms of obstruction who need surgical intervention because of functional impairment and group 2 those with milder forms of obstruction without functional impairment. We measured the random urinary levels of ET-1, MCP-1 and NAG, their ratios to creatinine and cut-off values were calculated and finally ROC curve was drawn.

Results: Fifty-nine neonates were enrolled in three groups; 24 patients with severe obstruction (Group 1), 18 neonates with milder forms of obstruction (group 2), and 17 neonates without any obstruction (group 3). The urinary ET-1, NAG, MCP-1 and normalization to creatinine, ET-1/Cr and NAG/Cr values were not significantly different between group 1 and 2 patients but urinary MCP-1/Cr ratio was significantly higher in severely obstructive patients than those with milder obstruction. To compare groups 1 and 2 a cut-off value of ET-1/Cr was measured as 0.5709 ng/mg (sensitivity [sens] 75 %, specificity [spec] 67 %) , of MCP-1/Cr was 0.927 pg/ml (sens 77 %, spec 72%) and the cut-off value of NAG/Cr ratio was 1.1913 (sens 62 %, spec 67 %) .

Conclusions: Evaluation of urinary levels of MCP-1/Cr ratios may help us to differentiate the newborns with severe obstructive hydronephrosis that are candidates of surgical intervention.

Keywords: Antenatal hydronephrosis, Obstruction, Neonates, Endothelin-1

ICIA2018\Diagnostic Methods in Immunology\Poster Discussion\9877

9877

Determination of CD4, CD8 and IL-8 Level in Serum and BAL Fluid of Patients with Pulmonary AnthracosisAtefeh Abedini¹; Arda Kiani²; Mehdi Ramazanpour³; Hamidreza Jamaati¹; Esmail Mortaz⁴; Fatemeh Razavi¹*1. Chronic Respiratory Diseases Research Center, National Research Institute of Tuberculosis and Lung Diseases (NRITLD), Shahid Beheshti University of Medical Sciences, Tehran, Iran**2. Tracheal Diseases Research Center, National Research Institute of Tuberculosis and Lung Diseases (NRITLD), Shahid Beheshti University of Medical Sciences, Tehran, Iran**3. Department of Pulmonology, Qom University of Medical Sciences, Qom, Iran**4. Clinical Tuberculosis and Epidemiology Research Center, National Research Institute for Tuberculosis and Lung Diseases (NRITLD), Shahid Beheshti University of Medical Sciences, Tehran, Iran*

Background: Anthracosis is the black pigmentation of bronchial mucosa. This disease is caused by the deposition of carbon particles, silica, quartz, etc. In the mucosa, sub mucosa, and inside the macrophages and in some cases have been reported with pulmonary tuberculosis (TB). The aim of the present study was to compare serum levels and fluid BAL of CD4/CD8 ratio and IL-8 as diagnosis and prognosis biomarkers for anthracosis.

Methods: 60 patients who referred to Masih Daneshvari Hospital, 30 anthracosis patients that confirmed with bronchoscopy were included as the cases. 5 ml of BAL sample and 5 ml of blood sample containing anticoagulants were taken from these individuals and sent to the Immunology Laboratory for analysis. Patients without anthracosis on bronchoscopy who were suspected to have tuberculosis were included in the control group.

Results: Thirty cases and 30 controls were included in the study. Among them, 60% were women and 40% were men. There were no significant differences in CD4/CD8 ratios in anthracosis compared to control group and no significant differences was observed in IL-8 as well. Pulmonary tuberculosis was confirmed in 88.9% of patients with anthracosis, which was significantly higher than control group.

Conclusion: These results suggest that changes in serum levels and BAL fluid of CD4/CD8 and IL-8 do not play an important role in the diagnosis or pathogenesis for anthracosis. Also due to the strong association of anthracosis and pulmonary tuberculosis, TB should be considered in patients with anthracosis, which in turn can lead to the early diagnosis and treatment of their patients.

Keywords: Anthracosis, CD4/CD8, IL-8, Pulmonary tuberculosis

ICIA2018\Diagnostic Methods in Immunology\Poster Discussion\9971
9971

Study of STING Gene Expression in Patients with Multiple Sclerosis in ChaharmahalvaBakhtiyari in 2015

KambizYousofzadeh Eshkevari¹, NahidJivad, HoseinTeymoori

1. Cellular and molecular research center, Shahrekord University of medical sciences, Shahrekord, Iran

Background: Multiple Sclerosis (MS) is an autoimmune disease which begins by exposure to a pathogen in genetically susceptible individuals. There is still no specific approved gene or collection of genes for MS. STING (Stimulator of interferon genes) also known as transmembrane protein 173 (TMEM173) for induction of type I interferon. In this study, STING gene expression levels in MS patients were measured.

Methods: This study is a comparative cross-sectional study. Patients were selected from the entire ChaharmahalvaBakhtiyari Province. 25 new diagnosed, 25 treated by interferone beta-1 α , 25 treated by interferone beta-1 β MS patients as well as 25 healthy individuals who referred to Kashani Hospital located in Shahrekord city in Iran, were participated in the study. Gene expression levels were determined by Real-Time PCR Device with using the $\Delta\Delta C_t$ method. The data were analyzed with statistical software GraphPad Prism version 5 and SPSS version 22.

Results: STING gene expression in newly diagnosed patients was more than the control group ($P<0.0001$) and in patients treated with interferon was lower than the control group ($P<0.0001$).

Conclusion: This study showed that gene expression of STING can be used to diagnose and treat MS patients by using Interferon beta-1 α and Interferon beta-1 β , especially among high-risk patients (such as first degree relatives). It is also possible to use this method to better determination of prognosis of disease.

Keywords: Multiple Sclerosis, STING Gen Expression, Interferon

ICIA2018\Diagnostic Methods in Immunology\Poster Discussion\10004
 10004

Urinary Levels of Lipoarabinomannan Do not Correlate with Active Tuberculosis in HIV-infected Patients Even in Those with Advanced Immunosuppression

EsmaeilMortaz^{1, 2, 3}, MiladMoloudizargari⁴, MajidMarjani¹, Afshin Moniri¹, Parissa Farnia⁵, Mohammad Varhram⁵, Johan Garssen^{3,6}, Mehdi KazempourDizaji⁵, Ian M. Adcock⁷, Payam Tabarsi¹

1. Clinical Tuberculosis and Epidemiology Research Center, National Research Institute for Tuberculosis and Lung Disease (NRITLD), Shahid Beheshti University of Medical Sciences, Tehran, Iran,

2. Department of Immunology, school of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran,

3. Division of Pharmacology, Utrecht Institute for Pharmaceutical Sciences, Faculty of Science, Utrecht University, Utrecht, Utrecht, Netherlands ,

4. Department of Immunology, school of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: The diagnosis of active tuberculosis (TB) in HIV-positive patients remains a global challenge due to paucibacillary presentations, low sensitivity with sputum based diagnostics, and high rates of extra-pulmonary TB. These issues have led to the pursuit of non-sputum based diagnostics, including the development of a commercially available ELISA-based urinary assay as well as a Lateral Flow (LF)-LAM assay for the detection of mycobacterial cell wall glycolipid lipoarabinomannan (LAM). We examined the specificity and sensitivity of urinary LAM ELISA in the diagnosis of TB in Iranian HIV-positive patients undergoing evaluation for TB. We also assessed whether there was an association between urinary LAM levels and total CD4+T cell count in HIV-TB co-infected patients.

Methods: Urine and blood samples were collected from HIV-positive patients with suspected TB infection. Urinary LAM levels were measured by ELISA and CD4+T cell counts were measured using a flow cytometer. Urinary LAM levels were compared between groups to determine the sensitivity and specificity of the test.

Results: 98 HIV-infected patients were eligible to be included in the study and were able to provide urine samples for LAM analyses. No significant association was found between the levels of urinary LAM in HIV-infected patients and the presence of TB infection. The urinary LAM test to discriminate between HIV-TB co-infected patients and HIV patients without TB had a sensitivity and specificity of 52% and 34%, respectively.

Conclusion: Our results do not support the use of urinary LAM ELISA test for the diagnosis of TB in HIV-positive patients due to low specificity even in patients with severe immunosuppression.

Keywords: LAM, TB, HIV, *Mycobacterium Tuberculosis*

ICIA2018\Diagnostic Methods in Immunology\Poster Discussion\10005

10005

CD4 to CD8 Ratio as a Valuable Diagnostic Test in Iranian Pulmonary Sarcoidosis Patients

ArdaKiani*¹, Atefeh Abedini², Kimia Taghavi², Mihaan Pourabdollah Toutkaboni², Esmaeil Mortaz³, Mehdi Kazempour Dizaji⁴

1. Tracheal Diseases Research Center, National Research Institute of Tuberculosis and Lung Diseases (NRIT-LD), Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Chronic Respiratory Diseases Research Center, National Research Institute of Tuberculosis and Lung Disease (NRITLD), Shahid Beheshti University of Medical Sciences, Tehran, Iran

3. Department of Immunology, Faculty of Medicine, Clinical Tuberculosis and Epidemiology Research Center, National Research Institute of Tuberculosis and Lung Diseases (NRITLD), Shahid Beheshti University of Medical Sciences, Tehran, Iran

4. Mycobacteriology Research Center, National Research Institute of Tuberculosis and Lung Diseases (NRIT-LD), Masih Daneshvari Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Sarcoidosis is a multi-organ disease with noncaseating granulomas that are comprised of T-helper/inducer (CD4⁺) lymphocytes and scant cytotoxic (CD8⁺) T-lymphocytes. This study is the first report from Iran that evaluates the ability of CD4:CD8 to detect sarcoidosis from other interstitial lung diseases (ILDs) in a large population.

Methods: Fifty patients with pulmonary sarcoidosis and 50 patients with non-sarcoidosis interstitial lung diseases (nsILDs) were included in the present study. Bronchoalveolar lavage (BAL) was performed using a flexible fiber optic bronchoscopy. After that, the samples were examined by flow cytometry method.

Results: The non-sarcoid group consisted of 50 patients in eight subgroups of the non-sarcoidosis interstitial lung diseases. As expected, CD4/CD8 ratio was significantly higher in sarcoidosis patients than non-sarcoidosis patients ($p < 0.001$). The best cut-off point in detecting sarcoidosis from other interstitial lung diseases was 1.1, which had a sensitivity of 92% and aspecificity of 80%.

Conclusion: Bronchoalveolar Lavage flow cytometry is a safe and fast test and its use as the first stage of diagnosis of sarcoidosis confirms the disease in 92% of patients (sensitivity to the current study). Hence, only a few patients need to undergo an aggressive procedure.

Keywords: Sarcoidosis, Bronchoalveolar lavage, lymphocyte, CD4⁺:CD8⁺ ratio



Immune Cell Therapy

Poster Discussion

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Immune Cell Therapy\Poster Discussion\8079

8079

Leukemia Inhibitory Factor (LIF) Modulates the Maturation of Murine Bone Marrow (BM)-derived Dendritic Cells

Atefeh Yaftiyan¹, Maryan Skandaryan¹, Amir Hossein Jahangiri¹, Seyyed Mohammad Moazzeni¹

1. Department of Immunology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

Background: Dendritic cells (DCs) are professional antigen presenting cells which their activation and function have a major role in establishment of specific immunity responses. Different biological factors such as cytokines are involved in the development of immunogenic or tolerogenic functions of DCs. Leukemia Inhibitory factor (LIF) is a pleotropic cytokine from interleukin-6 family that play various roles in cell proliferation, differentiation and survival. Previous studies demonstrate that LIF is linked to several tolerogenic events and also to tolerogenic phenotype and function of immune cells; yet the exact effect of this cytokine on dendritic cells is not particularly identified. In current study, we investigate the probable effect of LIF on DC maturation.

Methods: In current study, immature dendritic cells were differentiated from mouse bone-marrow (BM) in a GM-CSF and IL-4 containing medium. To determine the role of LIF in maturation, the immature differentiated DCs were treated in four groups: with TNF- α (maturation inducer), TNF- α and LIF, LIF only, or without cytokine. The harvested cells were assessed for the expression of phenotypic surface markers, antigen uptake function and allogenic T-cell proliferation capacity.

Results: LIF treatment of dendritic cells while stimulating by TNF- α caused a significant decrement in the expression of MHCII, CD40 or CD86 molecules in comparison with TNF- α only stimulated cells; meanwhile it was accompanied with an increase in the antigen uptake function (an indication for a reduced functional maturation). The ability for induction of allogenic T-cell proliferation proved no statistically significant difference.

Conclusion: Based on these findings, LIF is efficiently effective in modulating the levels of phenotypic and functional maturation. Such overall state can lead to the development of semi-mature and tolerogenic DCs. This inquiry can lead to a more precise understanding of the mechanisms of dendritic cell-dependent regulation of immune responses.

Keywords: Leukemia Inhibitory Factor (LIF), Dendritic cell, Tolerogenic DC, Maturation

ICIA2018\Immune Cell Therapy\Poster Discussion\8331
8331

Engineering Recombinant Lentivirus Expressing MBP Targeted Chimeric Auto Antibody Receptor (CAR) for Treatment of Multiple Sclerosis

Maryam sahlolbei¹, Jafar kiani², Seyed Mahmoud Hashemi³

1. Department of Molecular Medicine Faculty of Advance Technologies in Medicine, Iran University of Medical Sciences, Tehran, Iran.

2. Department of Molecular Medicine Faculty of Advance Technologies in Medicine, Iran University of Medical Sciences, Tehran, Iran.

3. Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Cancer treatment has been the subject of many researches for many years. Using and reinforcement your own immune system is a new way of treating the disease, which has made significant advances in the field today. In CAR T cell therapy, T cells are engineered and injected against specific antigens. For treating different types of cancer, the results are fairly good and many clinical trials are underway.

Recently, the treatment of autoimmune diseases has been considered by researchers in the field of biology. To treat autoimmune diseases, pathogenic autoimmune cells should be eliminated.

Chimeric auto-antibody receptor (CAR) is a strategy for the removal of autoreactive B-cell in autoimmune diseases. CAR-T cells may be an effective and global strategy for targeting auto-reactive B cells in antibody-mediated autoimmune diseases.

One of the reasons that causes multiple sclerosis, MS, as an autoimmune disease, is the auto-reactive B cells that make antibodies against the MBP protein. Our goal is to engineer T cells, which express CAR at their surface to target the autoreactive B cell expressing the anti-MBP antibody.

Methods: The CAR structure was designed and synthesized via bioinformatics, transmitted to the HEK293T cell via a third generation lentivirus.

Results: The CAR-MBP design was expressed at the cell surface of HEK293T and its expression was confirmed by flowcytometry.

Conclusion: Finding new ways to treat MS is a new hope for patients, and maybe MS can be converted into a treatable condition by increasing studies in this field and simultaneous use of different immunotherapy techniques.

Keywords: MS, chimeric antigen receptor, CAR T cell, autoimmune,

ICIA2018\Immune Cell Therapy\Poster Discussion\8338
8338

Study of the Effect of Mesenchymal Stem Cell Derived Exosome- treated Dendritic Cells on T Cell Proliferation

Mehri Shahir¹, Esmail Mortaz¹, Seyed Mahmoud Hashemi¹

1. Department of Medical Immunology ,School of Medicine ,Shahid Beheshti University of Medical Sciences, Tehran ,Iran

Background: DCs are professional antigen-presenting cells with functional plasticity. Regulatory DCs play an important role in the maintenance of immunological tolerance via the induction of T cell unresponsiveness or apoptosis, and generation of regulatory T cells. Mesenchymal stromal cells that are adult multi-potent cells have also been isolated from virtually all postnatal organs. Exosomes play an important role as a messenger in communications of cells. They are releasing from cells and activates/suppress important signaling. In this study we aim to investigate the effects of isolated exosomes from MSCs on the proliferation of T lymphocytes.

Methods: Culture of C57BL/6 mouse AD-MSCs: Murine adipose-derived MSCs were isolated from C57BL/6 Mice. Extracellular matrix was digested with type I collagenase, centrifuged and the pellet was then cultured in DMEM.

Isolation and purification of exosomes: Exosome was isolated by Exo-spin kit and evaluated by SEM. The protein concentration was detected by BCA kit.

Preparation bone marrow derived DCs: BM mononuclear cells were prepared from C57BL/6 mouse femur and tibia and then cultured in 6-well plates containing RPMI-1640 medium supplemented with 20 ng/mL GM-CSF and 20 ng/mL IL-4.

Lymphocyte proliferation assay: The BALB/C mouse derived splenocytes were labelled by carboxy fluorescein diacetate succinimidyl diester (CFSE) and then suspended in RPMI 1640 with DC groups (immature DC (iDC), iDC+LPS(1µg/ml), iDC+exosome(100µg/ml), iDC+exosome(100µg/ml) +LPS(1µg/mL)) and cocultured at ratio of 1:3, 1:10 and 1:30 for 3 days in 96-well U-bottom plates. Finally 10000 cells were analyzed by Flowcytometry.

Results: The results showed that the T cell proliferation was decreased in the presence of MSC derived exosomes treated- DCs in comparison with control cells.

Conclusion: Our data suggests that the exosomes of MSC could effect on the proliferation of T cells in presence of DCs and modulates immunologic consequences.

Key words: Mesenchymal Stem Cell, Exosomes, Tolerogenic Dendritic Cell, In Vitro

ICIA2018\Immune Cell Therapy\Poster Discussion\8422
8422

The Effects of Caffeine and Nicotine on Mesenchymal Stem Cell Functions

Zahra Zeinaly¹, Seyyed Meysam¹, Abtahi Froushani¹

Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Background: Mesenchymal stem cells (MSCs) have especially used in the cell therapy plans for autoimmune and inflammatory diseases. MSCs functions are controlled by various factors in the microenvironment that surround them, like nicotinic receptors and adenosine receptors. Here, we investigated the effect of caffeine (as adenosine antagonist) or nicotine on some functions of MSCs.

Methods: After the isolation, MSCs were incubated with different concentrations of caffeine (0, 0.1, 0.5 and 1 mM) and or with different concentrations of nicotine (0, 0.1, 0.5 and 1 μ M) for 48 hours. The survivability of cells was determined by MTT and neutral red uptake method over a 7-day period. After 90% confluence, the cultured plates were scratched. MSCs were visualized using a time-lapse microscope and the cells were maintained at 37°C throughout the imaging period. Images were taken every 15 min over a 2-h period to estimate the regenerative potential of MSCs.

Results: Attained results indicated that caffeine at low to moderate doses cannot change the rate of vitality, proliferation and regenerative potential of MSCs. Caffeine at 1 mM and nicotine at any doses significantly reduced the vitality, proliferation and regenerative potential of MSCs.

Conclusion: Nicotine at any doses and caffeine at the logarithmic dose have shown to adversely affect the vitality, proliferation and regenerative potential of MSCs.

Keywords: Mesenchymal Stem Cell, Nicotine, Caffeine.

ICIA2018\Immune Cell Therapy\Poster Discussion\9839
9839

The effect of Phagocytosis of Apoptotic Mesenchymal Stem Cells by Mouse Peritoneal Macrophage on iNOS and Arginase Activation

Maryam Ghahremani Piraghaj¹, Seyed Mahmoud Hashemi^{*1}, Hosein Ghanbarian², Sara Soudi³

1. Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Department of Biotechnology, School of Advanced Technologies in Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

3. Faculty of Medical Sciences, Department of Immunology, Tarbiat Modares University, Tehran, Iran

Background: Efferocytosis or phagocytosis of apoptotic cells by macrophages is very important to protect tissues inflammation because of releasing pro inflammatory intracellular ingredients. Mesenchymal stem cells have substantial immunosuppressive properties and many of studies have focused on the immunomodulatory properties of MSCs and its therapeutic implication for autoimmune diseases.

Methods: MSCs were isolated from abdomen adipose tissue of C57/BL6 mice. After optimal confluency of MSCs, macrophages were obtained from peritoneal cavity of C57/BL6 mice four days after thioglycollate injection. MSCs were exposed to UV irradiation at 254 nm for 20 min and followed by incubation in 37 with 5% CO₂. After that apoptotic MSCs were added to macrophages and after phagocytosis of apoptotic cells by them (nearly 1.5 h), macrophages were washed with PBS and were incubate for additional 48 h for analysis of activation of iNOS and Arginase enzymes.

Results: Our study showed that phagocytosis of apoptotic MSCs limit activation of iNOS and production of Nitric Oxide and promotes activation of Arginase enzymes and production of Urea.

Conclusion: The phagocytosis of apoptotic mesenchymal stem cells, induce non-inflammatory phenotype in macrophages. So, injected Mesenchymal Stem Cells maintain their immunomodulatory properties even if they apoptosis in the body.

Key words: Macrophage, Mesenchymal stem cell, Efferocytosis, Phagocytosis



Immunodeficiency

Poster Discussion

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Immunodeficiency\Poster Discussion\5178

5178

Role of Treg Cells (CD4 + CD25 + FOXP3 +) in Allergic CVID Patients

Nahid Eskandari¹, Roya sherkat², Zahra Adnani Sadati¹, Vajiheh Ostadi¹

1. Department of Immunology, Faculty of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran.
neskandari@med.mui.ac.ir

2. Acquired Immunodeficiency Research Center, Isfahan University of Medical Sciences, Isfahan, Iran

Background: Common Variable Immunodeficiency (CVID) is a primary immunodeficiency which is characterized by the failure of antibody responses such as reduction in antibody levels and an increased sensitivity to bacterial infections. The most important immunologic deficiency is the lack of immunoglobulin production by B cells. In addition, the defect and disturbance of the T cells, including the regulatory T cells (Treg), have also been proven to play a role in some clinical manifestations such as autoimmunity and splenomegaly. Protein P-STAT5b plays a key role in the function, survival and differentiation of Treg cells. The purpose of this study is to evaluate the role of Treg cells in CVID patients with allergic clinical symptoms.

Methods: Peripheral blood mononuclear cells were isolated from blood of 10 healthy volunteers and 13 patients with CVID before treatment with intravenous immunoglobulin and 10 allergic patients using *Ficoll density* gradient. The percentage of Treg cells in these individuals was measured using flowcytometry. Then Treg cells were isolated by immunomagnetic method and after culture and stimulation, percentage of expression of P-STAT5b protein in Treg cells was measured using flowcytometry.

Results: The percentage of Treg cells in CVID patients was significantly lower than control groups. The percentage of P-STAT5b in the CVID patients with allergic symptoms was significantly lower than the control group.

Conclusion: Some of the clinical manifestations of CVID diseases are not only due to the defect in the production of antibody, but also due to the defect in Treg cells, which can play a role in the pathogenesis of the disease.

Keywords: Common Variable Immunodeficiency, Regulatory T cells, P-STAT5b protein, Allergy

ICIA2018\Immunodeficiency\Poster Discussion\5300

5300

Mannose-Binding Lectin Protein Deficiency among Patients with Primary Immunodeficiency Disease Receiving IVIG Therapy

Sina Abdolrahim Poor Heravi¹, Gholamreza Azizi^{1,2}, Asghar Aghamohammadi^{1,2}

1. Research Center for Immunodeficiency, Children's Medical Center, Tehran University of Medical Sciences, Tehran, Iran

2. Primary Immunodeficiency Diseases Network (PIDNet), Universal Scientific Education and Research Network (USERN), Tehran, Iran

Background: Primary immunodeficiencies (PIDs) are inherited disorders in which one or several components of the immune system are defective. Immunoglobulin replacement therapy is the mainstay of treatment for patients with impaired antibody production. However, recurrent infections would continue to occur in some patients due to the other high frequent concomitant defects, such as mannose-binding lectin (MBL) deficiency.

Methods: A total of 51 PID patients participated in this cross-sectional study. A detailed questionnaire was completed by interviewing patients in order to record demographic, clinical and laboratory data. The levels of MBL were determined in the serums of patients by a sandwich enzyme-linked immunosorbent assay (ELISA) technique.

Results: MBL deficiency was found in 29.4% of cases; 11.8% patients had mild, 3.9% patients had moderate and 13.7% patients had severe MBL deficiency. In patients with MBL deficiency, the rate of meningitis, sepsis, pneumonia, and otitis were higher than patients with normal MBL levels. Immunoglobulin replacement therapy reduced the rate of infectious complications in PID patients; however, these reductions were more apparent in patients with normal MBL levels than patients with MBL deficiency.

Conclusion: Antibody deficient patients with a concomitant immune defects in MBL production have higher rates of recurrent infections despite receiving Immunoglobulin replacement therapy.

Keywords: Primary immunodeficiency, mannose-binding lectin, infection.

ICIA2018\Immunodeficiency\Poster Discussion\7836

7836

Increased Expression of B lymphocyte Induced Maturation Protein 1 (BLIMP1) in Patients with Common Variable Immunodeficiency (CVID)

Faezeh Abbasi Rad¹, Nafiseh Esmaeil¹, Mazdak Ganjalikhani Hakemi¹, Roya Sherkat²

1. Department of Immunology, school of Medicine, Isfahan Isfahan University of Medical Sciences, Isfahan, Iran

2. Acquired Immunodeficiency Research Center, Isfahan University of Medical Sciences, Isfahan, Iran

Background: Common variable immunodeficiency (CVID) is a primary immune deficiency disorder characterized by hypogammaglobulinemia and defect in response to vaccines. B cells maturation and differentiation are defective in this disorder, and CVID patients demonstrate commonly reduced memory B cells and antibody-secreting plasma cells numbers. Since the BCL6 and BLIMP1 molecules are two important transcription factors in the maturation of B cells to plasma cells, we evaluated the expression levels of BCL6 and BLIMP1 in B lymphocytes from peripheral blood in patients with CVID.

Method: Blood samples were collected from 14 patients with CVID and 14 healthy donors. Peripheral blood mononuclear cells (PBMCs) were isolated using Ficoll density gradient separation. Subsequently, CD19 + B cells were purified using MACS. The expression of protein and transcriptional levels of BCL6 and BLIMP1 were assessed using flowcytometry and Real-time PCR, respectively.

Results: Our results showed that the percentage of plasma cells in patients was significantly lower than control subjects. Also the expression levels of BLIMP1 were significantly higher in patients compared to control subjects. In addition, we found that although the expression of BCL6 was slightly higher in patients compared with, this difference was not significant.

Conclusion: Our findings suggest that increased expression levels of BLIMP1 and BCL6 could be involved in defective maturation of B cells in CVID patients and provide mechanistic insights into the pathogenesis of this disorders.

Keywords: CVID, Plasma cell, BCL6, BLIMP1

ICIA2018\Immunodeficiency\Poster Discussion\8319

8319

Toll like Receptors Pathway in Common Variable Immune Deficiency (CVID) and X-linked Agammaglobulinemia (XLA)

Parsova Tavasolian¹, Asghar Aghamohammadi^{2,3}, Farshid Noorbakhsh¹, Rouzbeh Sanaei⁴, Mahsima Shabani^{2,5}, Nima Rezaei^{1,2,6*}

1. Department of Immunology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

2. Research Center for Immunodeficiency, Pediatrics Center of Excellence, Children's Medical Center, Tehran University of Medical Sciences, Tehran, Iran

3. Primary Immunodeficiency Diseases Network (PIDNet), Universal Scientific Education and Research Network (USERN), Tehran, Iran

4. Department of Immunology, School of Medicine, Iran University of Medical Sciences, Tehran, Iran

5. International Hematology/Oncology of Pediatrics Experts (IHOPE), Universal Scientific Education and Research Network (USERN), Tehran, Iran

6. Network of Immunity in Infection, Malignancy and Autoimmunity (NIIMA), Universal Scientific Education and Research Network (USERN), Tehran, Iran

Background: Common variable immunodeficiency (CVID) and X-linked agammaglobulinemia (XLA) are two major humoral immunodeficiencies, causing a high burden of morbidity and early age mortality in children. Although many studies have investigated the involved etiologic factors, the exact impaired pathway is yet to be elucidated. The activation of Toll-like receptors (TLRs) is considered to be essential in B cell activation, co-stimulatory molecule up-regulation, class-switch recombination, and antibody and cytokine production. Current studies focus on the role of TLRs and demonstrate the defects in different TLR pathways in immune cells of CVID and XLA patients.

Methods: Herein, we measured TLR-4 and TLR-9 RNA levels in peripheral blood mononuclear cells (PBMCs) of patients with CVID and XLA. Moreover, for understanding the functional defects in associated TLR pathways we evaluated the level of TNF- α and IFN- α production in PBMCs of patients with CVID and XLA. **Results:** We have found that in CVID and XLA patients, TLR-9 expression is not increased significantly after stimulation with CpG, the TLR-9 ligand, whereas lipopolysaccharide (LPS) induced TLR-4 expression is increased similar to PBMCs derived from healthy control group. IFN- α production was increased significantly in patients and healthy individuals after CpG stimulation but LPS stimulated TNF- α production was only increased in CVID patients and healthy controls.

Conclusion: Our data suggests that defects in TLR-9 activated pathways may be a result of decreased TLR-9 expression but defects in TLR-4 activated pathways are due to defects in TLR-4 downstream molecules activity. Although LPS induced TNF- α production is defected in XLA patients, the preserved IFN- α production in CVID and XLA patients despite decreased TLR-9 expression suggests that TLR-9 is not the key modulator in INF- α production in these patients.

Keywords: TLR-4 (Toll-Like Receptor-4), TLR-9 (Toll-Like Receptor-9), CVID (Common Variable Immunodeficiency), XLA (X-linked Agammaglobulinemia),

ICIA2018\Immunodeficiency\Poster Discussion\9954

9954

Differentiation Classification of CVID Patients Based on B cell Subpopulation

Reza Yazdani^{1,2*}, Mazdak Ganjalikhani Hakemi², Zahra Hamidi Esfahani³, Nima Rezaei¹,
Asghar Aghamohammadi¹

1. Research Center for Immunodeficiency, Pediatrics Center of Excellence, Children's Medical Center, Tehran University of Medical Science, Tehran, Iran

2. Department of Immunology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

3. Department of Immunology, School of Medicine, Tarbiat Modares University, Tehran, Iran

Background: Common variable immunodeficiency (CVID) is a heterogeneous disease, characterized by hypoaglobulinaemia leading to recurrent infections and various complications. The aim of this study was to classify CVID patients based on four known classifications (Paris, Freiburg, EUROclass, and B-cell patterns) by measurement of B-cell subsets and to assess the relation of each classification with clinical manifestations.

Methods: We measured all B-cell subsets as both absolute count and percentage in 30 CVID patients and 30 healthy individuals using four-color flow cytometry. Moreover, we evaluated antibody responses to pneumococcal vaccine in patients.

Results: A significant reduction in percentage of terminal B-cell subsets (total, marginal zonelike, switched memory, IgM-only memory, total memory B-cells and plasmablast) and absolute count of all B-cell subsets along with a strong increase in CD21^{low} B-cells has been observed in patients. Patients with splenomegaly and hepatomegaly were clustered in group Ia, smB + 21 low and group 1 based on known classifications, and significantly tended to have a decreased transitional and marginal zone-like B-cells count, as well as an increase in CD21^{low} B-cell counts. Patients with lymphadenopathy, bronchiectasis and allergy had a significant decrease in absolute count of total memory, switched memory and total B-cells, respectively.

Conclusions: Classification of patients could provide useful information to guide clinicians in long-term follow-up of CVID patients. Our data demonstrate that it may be more accurate to use absolute counts of B-cell subpopulations in CVID patients because absolute counts of B-cell subsets are more associated with clinical manifestations compared with their percentages and also four known classifications

Key words: B-cell subsets, Common variable immunodeficiency, Classifications

ICIA2018\Immunodeficiency\Poster Discussion\9977

9977

Heterogeneous features of Lipopolysaccharide-responsive Beige-like Anchor (LRBA) Deficiency in Siblings

Fereshte Salami ¹, Gholamreza Azizi ², Hassan Abolhassani ^{3,4}, Reza Yazdani ⁴, Asghar Aghamohammadi ⁴

1. University of medical sciences, Shahid Sadoughi yazd

2. Non-Communicable Diseases Research Center, Alborz University of Medical Sciences, Karaj, Iran; and Department of Laboratory Medicine, Imam Hassan Mojtaba Hospital, Alborz University of Medical Sciences, Karaj, Iran

3. Department of Laboratory Medicine, Karolinska Institute at Karolinska University Hospital Huddinge, Stockholm, Sweden; Research Center for Immunodeficiencies, Children's Medical Center, Tehran University of Medical Sciences, Tehran, Iran

4. Research Center for Immunodeficiencies, Children's Medical Center, Tehran University of Medical Sciences, Tehran, Iran

Background: LPS-responsive beige-like anchor protein (LRBA) deficiency is a novel primary immunodeficiency disorder (PID) caused by biallelic mutations in the LRBA gene that abolish its protein expression. Immune dysregulation as a consequence of LRBA deficiency leads to recurrent infections, autoimmunity, and enteropathy. Affected individuals mainly show reduced levels of at least two Immunoglobulin (Ig) isotypes. We reported on siblings from two different families with LRBA deficiency who were carrying the same mutation but had discordant serum Ig levels.

Methods: In this case report Whole Exome Sequencing (WES) has facilitated identification of genetic defects in these PID patients with the same mutation and variable clinical phenotypes.

Results: Mutations of LRBA result in diminished expression and a spectrum of clinical phenotypes that includes hypogammaglobulinemia, enteropathy, autoimmune disorders, respiratory infections, and combinations of these phenotypes. Patients with the same mutation in LRBA may present different clinical phenotypes or even be asymptomatic. We reported two immunologic phenotypes of LRBA deficiency in two female siblings, including hypogammaglobulinemia and normal Ig levels with autoimmunity. In summary, we reported two male siblings with LRBA deficiency with the same mutation (c.C4814G [p.S1605X, exon 30]) and discordant clinical and immunological presentations. The index patient had severe clinical complications, including chronic diarrhea, organomegaly, respiratory tract infection, and hypogammaglobulinemia, whereas his brother had no clinical complications and normal serum Ig and specific antibody levels.

Conclusion: Our findings are consistent with those of a report confirming that the same genetic mutation in the LRBA gene can manifest with a broad phenotypic spectrum and no genotype–phenotype correlation. Further studies are required to define the exact role of genetic and nongenetic parameters in the penetrance of LRBA deficiency.

Keywords: LRBA, Immunodeficiency, Hypogammaglobulinemia, Immunoglobulin

ICIA2018\Immunodeficiency\Poster Discussion\9979

9979

Association of Mycobacterium Infections in Patients with Mendelian Susceptibility to Mycobacterial Disease with Venous Thromboembolism

Maryam Alinejad Dizaj ¹, Esmail Mortaz ²

1. Chronic Respiratory Diseases Research Center, National Research Institute of Tuberculosis and Lung Diseases (NRITLD), Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Clinical Tuberculosis and Epidemiology Research Center, National Research Institute of Tuberculosis and Lung Diseases (NRITLD), Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: An association between a hypercoagulable state and Mendelian susceptibility to mycobacterial disease (MSMD) has been established in a few studies; resultant thrombosis is considered rare.

Methods: In a case-control study, the prevalence of factor V Leiden, prothrombin G20210A and methylenetetrahydrofolate reductase (MTHFR) C677T, A1298C mutations were investigated in mycobacterium-infected patients. The study comprised 30 patients with mycobacterial infections (invasive, disseminated and/or recurrent infections with Bacille Calmette–Guerin or non-tuberculosis mycobacteria and Mycobacterium Tuberculosis with positive results for acid-fast bacilli and tuberculin skin tests) and 30 normal healthy controls. Forty female (66.7%) and 20 male subjects (33.3%) aged from 3 to 70 years were recruited into this study. Genotyping of targeted genes was performed by RT-PCR and cytokine TNF- α concentrations were quantified using a commercially available ELISA kit.

Results: Significant associations between mycobacterial infection and TNF- α production after stimulating peripheral blood mononuclear cells with LPS alone and with IFN- γ plus LPS were identified. Moreover, genotyping analysis in the studied population revealed a significant association between MTHFR c.677C>T (OR, 3.28; 95% CI, 1.35–7.92; $P < 0.05$), MTHFR c.1298A>C (OR, 2.33; 95% CI, 1.10–4.93; $P < 0.05$) and mycobacterial infection in affected patients, indicating susceptibility to venous thromboembolism according to previous studies.

Conclusion: Mycobacterium-infected patients had a significantly greater prevalence of MTHFR C677T and A1298C mutations than controls.

Key words: MSMD, MTHFR A1298C, MTHFR C677T, TNF- α .

ICIA2018\Immunodeficiency\Poster Discussion\9982
9982

Susceptibility to Mycobacterial Disease Due to Mutations in IL-12R β 1 in Three Iranian patients

Maryam Alinejad Dizaj^{1,2}, Esmail Mortaz³

1. Department of Biotechnology, School of Advanced Technologies in Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Chronic Respiratory Diseases Research Center, National Research Institute of Tuberculosis and Lung Diseases (NRITLD), Shahid Beheshti University of Medical Sciences, Tehran, Iran

3. Clinical Tuberculosis and Epidemiology Research Center, National Research Institute of Tuberculosis and Lung Diseases (NRITLD), Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: In the last decade, autosomal recessive interleukin-12 receptor β 1 (IL 12R β 1) deficiency, the most common cause of Mendelian susceptibility to mycobacterial disease (MSMD), has been diagnosed in a few children and adults with severe tuberculosis in Iran.

Methods: Here, we report three cases referred to the Immunology, Asthma and Allergy ward at the National Research Institute of Tuberculosis and Lung Diseases (NRITLD) at Masih Daneshvari Hospital from 2012 to 2017 with Mycobacterium tuberculosis and nontuberculous mycobacteria infections due to defects in IL12R β 1 but with different clinical manifestations.

Results: All three were homozygous for either an IL-12R β 1 missense or nonsense mutation that caused the IL-12R β 1 protein not to be expressed on the cell membrane and completely abolished the cellular response to recombinant IL-12.

Conclusion: Our findings suggest that the presence of IL-12R β 1 deficiency should be determined in children with mycobacterial infections at least in countries with a high prevalence of parental consanguinity and in areas endemic for TB like Iran.

Keywords: IFN- γ , IL12RB1 , IL-12R β 1 , IMD30 , MSMD , PID



14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

Immunogenetics

Poster Discussion

ICIA2018\Immunogenetics\Poster Discussion\7687

7687

Susceptibility Pattern of HLA-DRB1 and DQB1 Alleles and Haplotypes for Pemphigus Vulgaris in an Iranian Population

Ghasem Solgi¹, Fatemeh Ahmadi Motamayel², Mahsa Mahdian², Hamzeh Rezaei¹, Zohreh Zarei Ghane¹, Mehrdad Hajilooi¹, Hamidreza Abdolsamadi², Ali Mahdavinezhad³

1. Department of Immunology, Medical School, Hamadan University of Medical Sciences, Hamadan, Iran.

2. Dental Research Center, Department of Oral Medicine, Hamadan University of Medical Sciences, Hamadan, Iran.

3. Department of Genetic and Molecular Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

Background: Genetic factors, particularly *HLA* class II genes have been implicated in pathogenesis of pemphigus vulgaris (PV). This study aimed to evaluate any correlation between HLA-DRB1 and DQB1 alleles and haplotypes frequencies and PV disease presentation in an Iranian population.

Methods: In this case-control study, 39 PV patients that diagnosed based on clinical and pathological criteria as well as 98 ethnic-matched healthy controls were included. HLA-DRB1 and HLA DQB1 specific alleles were determined by PCR-SSP methods. The alleles and haplotype frequencies were compared between both groups of the study as well as subgroups of the PV patients.

Results: Among *HLA*-DRB1 alleles, DRB1*04:02 ($P=2.3 \times 10^{-7}$), DRB1*04:03 ($P=0.02$) and DRB1*14 ($P=6.2 \times 10^{-5}$) were positively associated with PV and DRB1*15:01 ($P=0.01$) and DRB1*11:01 ($P=0.007$) showed negative association with PV disease. Also, HLA-DQB1*03:02 ($P=3.07 \times 10^{-11}$) and DQB1*05:03 ($P=7.1 \times 10^{-6}$) showed positive and DQB1*02:01, DQB1*03:01 and DQB1*05:01 alleles showed negative association with disease incidence. Two frequent haplotypes, HLA-DRB1*04:02-DQB1*03:02 ($P=2.3 \times 10^{-7}$) and HLA-DRB1*14-DQB1*05:03 ($P=9.3 \times 10^{-5}$) conferred susceptibility to PV whereas, DRB1*11:01-DQB1*03:01 and DRB1*13:01-DQB1*06:03 showed protective role for PV.

Conclusion: Our finding corroborates somewhat a different pattern for susceptible/protective HLA class II alleles and haplotypes among PV patients in Iranians.

Keywords: HLA-DRB1, HLA-DQB1, Alleles, Pemphigus Vulgaris

ICIA2018\Immunogenetics\Poster Discussion\8286

8286

rs352140 & rs187084 Polymorphisms in TLR9 Gene and Risk of Multiple Sclerosis

Saiedeh Erfanian¹, Maliheh Moradzadeh²

1. Department of Advanced Medical Sciences and Technologies, School of Medicine, Jahrom University of Medical Sciences (JUMS), Jahrom, Iran

2. Golestan Rheumatology Research Center, Golestan University of Medical Sciences, Gorgan, Iran

Background: Multiple sclerosis (MS) is a chronic inflammatory disease that degenerate myelin cells of the central nervous system (CNS), particularly in advanced societies

Methods: This study was conducted on 100 MS patients and 100 healthy people that were selected randomly by molecular study. After explaining the project and obtaining informed consent from patients and healthy subjects, 5 to 10 mL blood sample was taken from the cases. DNA samples were extracted using DNA extraction kit. To identify the gene region containing the polymorphism rs1044243, PCR method was used and then to enzymatic digestion, Afl II restriction enzyme for rs187084 polymorphisms and Bst UI restriction enzyme for rs352140 polymorphisms, was added to make a cutgel electrophoresis technique was used. The data collected were processed using the software spss 15.

Result: In genotype distribution, frequency of rs187084 polymorphism in healthy and patient groups, respectively was CC 10(10%) and 11(11.0%) (P=0.609), CT 65(65.0%) and 68(68.0%) (P=0.522) and TT 25(25.0%) and 21(21.0%) (P=reference) and no significant difference was observed between the two groups in any of the states. But in terms of the c allele frequency in rs187084 polymorphism among healthy subjects and patients with MS is 85(%42.5) and 110(%55) that two groups showed significant difference (P=0.016). In genotype distribution frequency of rs352140 polymorphism in healthy and patient groups, respectively was CC 25(%25) and 19(%19) (P=Reference), CT 53(%53) and 56(%56) (P=0.360) and TT 22(%22) and 25(%25) (P=0.340) and no significant difference was observed between the two groups in any of the states. The T allele frequency in rs352140 polymorphism among healthy subjects and patients with MS is 97(48.5%) and 94(47%) that the two groups showed no significant difference (P=0.841).

Conclusion: Despite the proof of association of rs352140 and rs187084 polymorphisms on TLR9 gene with MS in some study of out of Iran, our findings did not show any association in population of Fars province. However, we must consider this issue that besides the difference in terms of selection of the patients and controlling different study, some factors like: nationality, strips, nutrition, environmental factors and etc, must be analyzed and more studies need to be done to prove the role of this polymorphisms as an independent risk factor of MS.

Key words: Polymorphism rs352140, Polymorphism rs187084, TLR9 gene, MS disease

ICIA2018\Immunogenetics\Poster Discussion\8373

8373

Bioinformatics Analysis of Vitiligo Relationship with AllergiesElmira shiuokhi^{1, 2*}, Seyed saadi hosseini^{2, 3}*1. Department Science and Research .University of Islamic Azad. Central Tehran**2. Lab. of Regenerative Medicine and Biomedical Innovations, Pasteur institute of Iran**3. Department of Biotechnology Faculty of Bioscience University of Shahid Beheshti*

Background: Vitiligo is an autoimmune skin disease with depigmentation. Many of vitiligo genes identified. These genes encode proteins that involved in immune regulation, several others play roles in cellular apoptosis and others involved in regulating functions of melanocytes. vitiligo involving a complex relationship between programming and function of the immune system. The purpose of this study is to investigate the bioinformatics of vitiligo disease and its relation with allergic diseases.

Methods: nucleotide and proteins sequences of 35 genes that were candidates for vitiligo formation and 31 genes involved in allergic diseases that were likely to be associated with vitiligo were retrieved from the GenBank and UCSC. Then bioinformatics analyzes were performed with different software such as; Bioedit2007, Mega7, string-db, PROTPARAM, PROTSKALE, UNIPROT, InterPro.

Results: Phylogenetic analyzes showed that most of these genes are susceptible to single nucleotide mutations. Protein analyzes and genetic network showed that FOXP3, PTGS2, IL10, TNF and TSLP genes have the highest interactions and a key role in the network. Also, the analysis shows that TNF, FOXP3 and IL-10 genes has a direct effect on the interactions between genes. Analysis of the vitiligo gene network with the genes involved in the allergy disease showed that these genes, including the FASLG, STAT6, FAS, IRF4, CD44, CD80, CTLA4, STAT4, PTPRC, HLA-A, KiTLG and IL17 which may indicate that Vitiligo is associated with allergies, proves the need comprehensive bioinformatics and in vitro studies. Furthermore, the results of the genetic network showed that the FOXD3, C12orf10, TXNDC2, UVRAG, FBXO11 genes did not interact with other genes, which could indicate that these genes to Vitiligo disease had not meaning.

Conclusion: The results of this study showed that a number of genes had a significant effect on the vitiligo mechanism, and these genes have significant interactions with a number of allergenic agent genes which indicates the association of these diseases.

Keywords: Vitiligo, autoimmune, allergy. Bioinformatics

ICIA2018\Immunogenetics\Poster Discussion\8376

8376

Appropriate Housekeeping Gene in C57/B6 Bone Marrow Cells in Serum Starvation Condition for Real-time PCR

Gilda Parsamanesh¹, Amrolah Mostafazade¹, Davar Amani², Kaveh Baghaei³, Samaneh Tokhanbigli³,
Hamid Asadzadeh Aghdayi³

1. Department of immunology, School of medicine, babol University of medical scienc, Babol, Iran

2. Department of immunology, School of medicine, shahid beheshti University of medical sciences, Tehran, Iran

3. Basic and Molecular Epidemiology of Gastrointestinal Disorders Research Center, Research institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Real-time PCR (qPCR) assay is a conventional method for determining RNA transcription and gene expression. In this regard, housekeeping gene in Real-Time PCR is considered as a gene which its expression does not undergo too many changes in experimental conditions. Moreover, the reliability of a Real-Time PCR depends on a decent reference gene for normalization of the results. In some study, MHC class I, HPRT (Hypoxanthine Phosphoribosyl transferase) and GAPDH (Glyceraldehyde 3-phosphate dehydrogenase) are used as a reference gene. In current study, we aim to show that which of mentioned genes is a suitable candidate for serum starvation condition.

Methods: Bone marrow cells from C57/B6 were isolated and cultivated in complete media as a control group and without FBS as a starved group for 24, 48 and 72 hours. Consequently, after each time point RNA extraction and DNA synthesis were performed and given data were analyzed.

Results: HPRT and MHC class I expression after 48 and 72 hours starvation were decreased, whereas, GAPDH in 24, 48 and 72 hours had no significant difference in expression.

Conclusion: According to Real-Time PCR analysis, HPRT and MHC class I had shown substantial changes in their expression in serum starvation in comparison with the non-starved condition which suggests that these two genes could not be reliable housekeeping genes in Real-Time PCR in serum starvation circumstances.

Keywords: housekeeping gene/ MHC class I/ HPRT/ GAPDH/ Serum starvation/ Real-Time PCR

ICIA2018\Immunogenetics\Poster Discussion\8581

8581

V617F-independent Upregulation of JAK2 Gene Expression in Patients with Inflammatory Bowel Diseases

Khadijeh Koushki¹, Pedram Azimzadeh¹, Mohammad Rostami Nejad², Kazem Mashayekhi³, Davar Amani⁴, Hamid Asadzadeh Aghdaee¹, Mohammad Reza Zali².

1. *Basic and Molecular Epidemiology of Gastrointestinal Disorders Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran*

2. *Gastroenterology and Liver Diseases Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran*

3. *Immunology Department, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran*

4. *Immunology Department, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran*

Background: Inflammatory bowel disease (IBD) is one of the most important immune mediated disorders that affecting the GI tract. However, IBD can also be associated with numerous extraintestinal complications such as VTE or have an increased risk of vascular complications that leads to significant morbidity and mortality. The JAK2 (Janus kinase 2) V617F mutation is a well-known point mutation that involved in VTE and one of the most important gene involved in IBD pathogenesis. Therefore, the aim of this study was to evaluate the JAK2 V617F mutation detection and JAK2 gene expression in Iranian IBD patients.

Methods: A total of 452 individuals including 246 IBD patients (209 UC and 37 CD) and 206 healthy controls were enrolled in the study. The genomic DNA and total RNA were extracted from PBMCs. JAK2 V617F mutation detection was performed using Restriction Fragment Length Polymorphism (RFLP) method and JAK2 mRNA expression was analyzed by a Real-time PCR based on relative quantification method. The data were analyzed using SPSS software.

Results: Of 246 IBD (mean age $36 \pm SD=2.11$, 128 Male and 118 Female) and 206 controls (mean age $32 \pm SD=3.42$, 97 male and 109 female), our results showed no distinguishable relationship of JAK2 V617F mutation in IBD patient with thrombosis compared with IBD patients without thrombosis and healthy control. However, the mRNA expression was significantly different between IBD patients and controls enrolled in the study ($P=0.0001$).

Conclusions: Our results suggest a JAK2 V61F -independent upregulation of JAK2 mRNA expression in IBD patients. The absence of JAK2 V617F mutation in the thrombotic IBD patients suggests that other mechanisms such as transcriptional level play an important role in the pathogenesis of thrombosis in IBD. Our data confirms that thromboembolism in IBD is associated with elevated JAK2 mRNA level.

Keywords: Inflammatory Bowel Diseases (IBD), Janus Kinase 2 (JAK2), V617F, Thromboembolism

ICIA2018\Immunogenetics\Poster Discussion\9838

9838

Evaluation of Inflammasome Receptors Genes Expression of AIM2 and NLRP1 in Peripheral Blood of SM Injured Patients by Real-Time PCR

Mitra Sabetghadam Moghadam ¹, Jafar Salimian ², Jamileh Norouzi ³

1. Department of biology, Islamic Azad University, North Tehran Branch

2. Department of Chemical Injury Research, Baqiyatallah University of Medical Sciences, Tehran, Iran

3. Department of Microbiology, Islamic Azad University, North Tehran Branch

Background: Sulfur mustard (SM) exposed patients suffer from chronic lung injuries after 30 years exposure to mustard gas. It seems that the imbalance in the immune system is one of the most important molecular mechanisms involved in pathophysiology of late complications in mustard lung patients. Inflammasome is a protein complex that promotes inflammation and secretion inflammatory cytokines IL-18 and IL-1 β . NLRP1 and AIM2 molecules are innate immune cells receptors that can contribute to the formation of the inflammasome complex. In this study, the genes expression levels of NLRP1 and AIM2 receptors were evaluated in peripheral blood sample of SM patients and compared with chronic obstructive pulmonary disease patients (COPD) and healthy controls.

Methods: In this study, 15 SM patients as a case group, 15 COPD patients as control that referred to the lung department of Baqiyatallah hospital were selected, as well as 15 subjects as healthy controls. After extracting of RNA from their blood samples, cDNA was synthesized. The genes expression levels of NLRP1 and AIM2 were assayed using Real-Time PCR technique.

Results: The NLRP1 gene expression level was higher in SM patients (7 \pm), and COPD patients (14 \pm), compare to HC group ($P < 0.05$). There was no significant difference between the AIM2 gene expression level in SM and COPD patients group compared to the control group.

Conclusion: Increased gene expression level of NLRP1 in SM and COPD patients, proposed the inflammasome complex to be active in these patients. This indicates the role of NLRP1 in the inflammation process in SM and COPD patients. Due to the lack of significant difference in expression of AIM2 gene, it seems that the AIM2 is not active in these patients.

Keywords: SM, Inflammasome, NLRP1, AIM2

ICIA2018\Immunogenetics\Poster Discussion\9859

9859

Analysis of the Frequencies of HLA-A, B, and DRB1 Alleles in the Three Ethnic Groups, in East Azerbaijan, IranAva Nasrolahi^{1,2}, Roya Dolatkah², Hossein Nikkhah², Sima Shahmohammadi Farid²*1. Department of Molecular Medicine, Faculty of Advanced Medical Sciences, Tabriz University of Medical Sciences, Tabriz, Iran**2. Hematology and Oncology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran*

Background: Human Leukocyte Antigen (HLA) refers to the major histocompatibility complex (MHC), are one of the highly polymorphic loci of human genome. The HLA allele frequency and linkage disequilibrium patterns are different among human populations. The donors-recipients HLA matching for HLA-A, -B and -DRB1 loci have an important impact in the outcome prediction of allogeneic bone marrow transplantation. Aim of this study was calculated allele frequencies of the HLA-A, -B and -DRB1 genes and their relationship with ethnicity in 293 healthy individuals in East Azerbaijan province of Iran.

Methods: In this cross-sectional analytic study, HLA typing was performed using PCR-SSP method. Allele frequencies were estimated for the HLA-A, -B and -DRB1 classes for of 293 cases. For assessment of any relationship between different HLA class and mentioned variables Chi-square-test was performed.

Results: from 293 cases 132 (45.1%) were female and 161 (54.9%) were male, with the mean age of 30.30 ($\pm 8.70\%$) years. A total of 16 HLA-A alleles, 26 HLA-B alleles and 13 HLA-DRB1 alleles were detected. Among them, the most common were: HLA class I alleles A*02 with 97(16.6%), B*35 with 98(16.7%), and HLA class II alleles DRB1*11 with 96(16.4%) frequencies. there was not any significant association between ethnicity of the cases and HLA-A ($P = 0.88$), HLA-B ($P = 0.31$) and HLA-DRB1 ($P = 0.85$).

Conclusion: This was the first comprehensive study of frequency of different HLA types in East Azerbaijan, but any relationship between different HLA classes and ethnic groups was observed.

Keywords: Human Leukocyte Antigen, HLA typing, Transplantation

ICIA2018\Immunogenetics\Poster Discussion\9937
9937

Lack of association between Vitamin D binding protein (VDBP) gene polymorphisms and endometriosis

Mohammad Jafari^{1, 2}, Soheila Arefi¹, Sahel Heydari², Hadi Mohagheghian Yaghoubi¹, Ali Akbar Delbandi^{1, 2}, Nesa Rashidi^{1, 2}

1. Immunology Research Center, Immunology and Infectious Disease Institute, IRAN University of Medical Sciences

2. Immunology Department, School of Medicine, IRAN University of Medical Sciences

Background: Vitamin D binding protein (VDBP) is a multifunctional, polymorphic protein that is a carrier of vitamin D and its metabolites in serum. In addition, this protein enhances the neutrophil chemotactic activity of complement component 5a. In the previous studies observed variants of the VDBP gene seem to be connected to levels of the main circulating vitamin D metabolite [25(OH) D]. On the other hand, low levels of 25(OH)D in women associated with endometriosis. Then, polymorphisms in VDBP gene may be affected anti-inflammatory effects of vitamin D. The percent study aimed to investigate the distribution of variants of VDBP gene in Iranian endometriotic women.

Methods: We conducted an association study and included 118 Iranian endometriotic patients and 116 matched healthy controls. We examined the associations between the common variants of the VDBP (rs4588 and rs7041) gene using Restriction Fragment Length Polymorphism (RFLP) method after DNA isolation.

Results: There was no significant difference in genotypes and allele frequency of the investigated VDBP polymorphisms between patients and healthy women controls.

Conclusion: Our finding revealed that rs4588 and rs7041 polymorphisms of VDBP are not risk factor for susceptibility to endometriosis.

Keywords: Endometriosis, Vitamin D, Vitamin D binding protein, Genetic polymorphism

ICIA2018\Immunogenetics\Poster Discussion\10017
10017

Functional Promoter Polymorphism of Matrix Metalloproteinase (MMP)-3 5A/6A and Its Interaction with MMP-7 A-181G Polymorphism in Multiple Sclerosis

Ardeshir abbasi¹, zohreh rahimi^{2,3}, ziba rahimi³

1. Department of Immunology, Tarbiat Modares University of Medical Sciences, Tehran, Iran

2. Department of Clinical Biochemistry, Medical School, Kermanshah University of Medical Sciences, Kermanshah, Iran

3. Medical Biology Research Center, Kermanshah University of Medical Sciences, Kermanshah, Iran

Background: Multiple sclerosis (MS) is a progressive autoimmune disease of the central nervous system (CNS) in which axonal inflammation, demyelination and damage occurs. The aim of present study was to investigate the influence of matrix metalloproteinase (MMP)-3 5A/6A and its interaction with MMP-7 A-181G polymorphism on the risk and the clinical course of MS.

Methods: We studied 121 patients and 106 healthy individuals without family history of MS or any other autoimmune diseases from Kermanshah province. The MMP-3 and MMP-7 genotypes were detected using polymerase chain reaction (PCR)-restriction fragment length polymorphism (RFLP).

Results: The MMP-3 6A allele was more prevalent in patients (70.2%) than that in the controls (67%, $p=0.45$). The frequency of MMP-3 6A/6A genotype was higher in secondary progressive-MS (SP-MS) patients (57.9%) than that of relapsing remitting-MS (RR-MS) patients (39.2%) but it did not reach to a statistical significance. The concomitant presence of both MMP-3 6A and MMP-7 -181 G alleles compared to the combined presence of MMP-3 6A and MMP-7 -181A alleles increased the risk of MS by 1.64-fold ($p=0.05$).

Conclusion: finally, our study for the first time among a homogenous ethnic group of Iranians (Kurds) indicated the absence of an influence of MMP-3 5A/6A polymorphism on the risk of MS or its clinical course. However, we detected a gene-gene interaction between MMP-3 and MMP-7 that increased the risk of MS in the presence of MMP-3 6A and MMP-7 -181 G alleles.

Keywords: multiple sclerosis, MMP-3 5A/6A, MMP-7 A-181G, polymorphism, clinical course.



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Immunohematology

Poster Discussion

ICIA2018\Immunohematology\Poster Discussion\7478
7478

The Effect of *Tribulus terrestris* Aqueous Extract on Production and Secretion/Activity of Platelets Related Factors in Immune Thrombocytopenic Purpura Patient

Zahra Koolivand¹, Ali mostafaie², Mehrdad Payandeh³

1. Abadan School of Medical Sciences, Abadan, Iran.

2. Medical Biology Research Center, Kermanshah University of medical sciences, Kermanshah, Iran

3. Department of Hematology and Medical Oncology, Kermanshah university of medical sciences, Kermanshah, Iran

Background: Idiopathic thrombocytopenic purpura (ITP) is an autoimmune disease with decreased platelet levels. First line therapy consists of corticosteroids such as prednisone and intra venous immunoglobulin (IVIG) and anti-Rho (D) and in the case of unresponsiveness, the second line includes splenectomy and rituximab and thrombopoietin (TPO) receptors agonist. Both therapy plans are associated with side effects which could be avoided by using herbals such as *Tribulus terrestris*.

Methods: Aqueous extract of *Tribulus terrestris* was prepared, dried and final powder was distributed between twenty newly diagnosed ITP patients (40% of which were *Helicobacter pylori* positive) to take as pills three times a week for fourteen consecutive days. Blood samples were taken from patients on days 1, 14 and 28, after stop taking the pills to analyze plasma PF4, serotonin and VWF.

Results: In all of patients, PF4 was decreased and VWF showed significant increased on all three days. The serotonin was exhibited the increase on days 1 and 14.

Conclusion: Altogether, *Tribulus Terrestris* seems to be remarkable safe choice for treatment of ITP patients.

Keywords: *Tribulus terrestris*, ITP, serotonin, VWF, PF4

ICIA2018\Immunohematology\Poster Discussion\8150

8150

Percentage of Anti-A and Anti-B Hemagglutinins in “Dangerous O Blood Group Donors”

Hajar Shali1², Vahid Baghdadi², Mohammad Hossein Baradaran²

1. *Department of Immunology, School of medicine, Tabriz University of medicine Science, Tabriz, Iran*

2. *Qazvin Blood Transfusion Organization, Qazvin, Iran*

Background: Dangerous O blood group donors have high titers of anti-A, anti-B and/or anti-AB hemagglutinins in their plasma. These antibodies are mostly in class of IgM and IgG that causes agglutination of the erythrocytes of non-O recipients. Transfusion of blood products from these groups can cause severe complication such as an acute hemolytic transfusion reaction (HTR). To prevent transfusion reactions, titration of these antibodies is recommended.

Methods: For titration of anti-A and anti-B hemagglutinins (IgM class), serial dilutions were prepared with donor plasma collected in (EDTA) in saline solution (from 1:1 until 1:1024). To destroy IgM class and then titration of anti-A and anti-B hemagglutinins (IgG class), the donor plasma was treated with Mercaptoethanol (2ME). When the titers of anti-A or anti-B hemagglutinins were ≥ 128 and ≥ 256 related to IgM and IgG, respectively, donors were considered to be in the dangerous O blood group.

Results: Considerably, the percentage of cases with titer ≥ 256 for anti-A (IgG) and anti-B (IgG) are 4% and 3%, respectively. Percentage of anti-A (IgM) only was 2%. No one detected with anti-B (IgM).

Conclusion: To prevent the occurrence of dangerous transfusion reactions, O blood group products such as fresh-frozen plasma and cryoprecipitate, especially platelet concentrates obtained by apheresis should only be given to group O recipients.

Keywords: Dangerous O blood group, hemolytic transfusion reaction, hemagglutinins.



Immunology of Chemical Victims and Environmental Pollution

Poster Discussion

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Immunology of Chemical Victims and Environmental Pollution\Poster Discussion\7569
7569

Extremely low frequency electromagnetic fields (ELF-EMF) decreases serum levels of IL-17, TGF- β and down regulates Foxp3 expression in the spleen

Leila Mahdavinejad,¹ Mahdi Alahgholi Hajibehzad,^{1,2} Mohammad Mahdi Eftekharian,^{1,2} Zohre Zaerieghane¹, Iraj Salehi³, Mehrdad Hajilooi,^{1,2} Hanie Mahaki,^{1,2} Alireza Zamani^{1,2,*}

1. Department of Immunology, School of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

2. Molecular Immunology Research Group, Research Center for Molecular Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

3. Department of Physiology, School of Paramedical Sciences, Hamadan University of Medical Sciences, Hamadan, Iran

Background: The aim of this study was to determine the effects of extremely low frequency electromagnetic fields (ELF-EMF) on the serum levels of interleukin-17(IL-17) and transforming growth factor- β (TGF- β) that are secreted by Th17 and Treg cells respectively. Also to assess expression of transcription factors forkhead box P3 (Foxp3) and retinoid-related orphan receptor γ T(ROR γ T) that are lineage defining of Th17 and Treg cells in spleen and thymus of rats.

Methods: Eighty wistar male rats were randomly divided into four equal ELF-EMF exposed and a sham-exposed control group. The ELF-EMF groups were exposed to intensities of 1, 100,500 and 2000 μ T at frequency of 50 Hz, 2h/day and controls were placed in the solenoid without ELF-EMF. After one month of the exposures all rats were immunized by human serum albumin and the experiment was continued in the same manner for a month more.

Results: The results demonstrated that there were no significant change between the body weight and weight of spleen, whereas the weight of thymuses was significantly declined at intensities of 500 and 2000 μ T. At the pre-immunization phase, the serum levels of IL-17, TGF- β and expression of Foxp3 at intensities of 1 and 100 μ T in the spleen were significantly decreased but at the post-immunization phase, there were no difference among them. Whereas, total serum level of TGF- β in the post-immunization phase was increased comparing with the pre-immunization phase.

Conclusion: low intensities of ELF-EMF could reduce the serum levels of IL- 17 and TGF- β and down regulate the expression of Foxp3 in spleen.

Keywords: Interleukin-17, Transforming growth factor- β , Extremely low frequency of electromagnetic fields

ICIA2018\Immunology of Chemical Victims and Environmental Pollution\Poster Discussion\8573
8573

Evaluation of Serum Level of IL-8 in Chemical Victims and Its Relationship with Severity of Pulmonary Complications

Maryam Khabareh¹, Tooba Ghazanfari²

1. *Shahed University Tehran*

2. *Department of Research Center of Regulation the Immunology Response*

Background: Sulfur mustard (SM) is a toxic and chemical agent. One of the most important long-term effects of sulfur mustard exposure is pulmonary complication that chemical victims are suffering for years. Interleukin 8(IL-8) is one of the important cytokines contributes in chronic pulmonary inflammatory diseases. In this study, the serum level of IL-8 and its correlation with severity of pulmonary complications was evaluated in SM exposed individuals about three cascades after exposure.

Methods: In this work, 73 SM exposed cases and 86 unexposed as control were studied. Pulmonary function test was performed using spirometry. Chemical victims were classified into three groups of normal, moderate and severe pulmonary damage, according to classification of the medical committee of the foundation of martyr and veterans Affair. The serum level of IL-8 was measured by ELISA method.

Result: There was significant decrease in the serum level of IL-8 in the exposed group compared to control group also there was significant decrease in the serum level of IL-8 in all three exposed groups compared to control group. But no significant difference observed between the moderate and severe pulmonary damage groups with normal ones.

Conclusion: The serum level of IL-8 in chemical victims is affected by exposure to SM. But the serum level of IL-8 cannot play an important role in the severity of pulmonary Complications as an inflammatory factor.

Keywords: IL-8, Pulmonary Complications, sulfur mustard.

ICIA2018\Immunology of Chemical Victims and Environmental Pollution\Poster Discussion\8715
8715

Evaluation of Serum Level of IL-18 Binding Protein in Chemical Victims and Its Relationship with Severity of Pulmonary Complications

Somaye Mami¹, Tooba Ghazanfari²

1. Shahed University Tehran

2. Department of Research Center of Regulation the Immunology Response

Background: Exposure to mustard gas leads to acute and chronic toxic effects on the pulmonary. Results from several studies have been suggested the role of immune mediators including IL 18BP in the pathogenesis of pulmonary diseases. IL 18BP by binding to IL18 prevents its binding to the receptor and acts as a natural inhibitor for this cytokine.

Methods: In this study, the serum level of IL 18bp and its correlation with severity of pulmonary complications was evaluated in SM exposed individuals about three cascades after exposure. In this work, 93 SM exposed cases and 86 unexposed as control were studied. Pulmonary function test was performed using spirometry. Chemical victims were classified into three groups of normal, moderate and severe pulmonary damage, according to classification of the medical committee of the foundation of martyr and veterans Affair. The serum level of IL-18bp was measured by ELISA method.

Result: There was significant decrease in the serum level of IL 18bp in the exposed group compared to control group also. There was significant decrease in the serum level of IL-8bp in all three exposed groups compared to control group. But no significant different observed between the moderate and severe pulmonary damage groups with normal ones.

Conclusion: The serum level of IL-8bp in chemical victims is affected by exposure to SM, but the serum level of IL-8bp can 'not play an important role in the severity of pulmonary Complications as an inflammatory factor.

Keywords: IL-18, Pulmonary Complications, mustard gas.



Immunology of Exercise, Aging, and Nutrition

Poster Discussion

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Immunology of Exercise, Aging, and Nutrition\Poster Discussion\8071
8071

Effect of Inhibition of D2 Dopamine Receptors on Several Functions of Monocytes of Peripheral Blood in Rat under Food Restriction

Khaterreh Noormohammadi¹, Farrin Babaei Balderlou¹, Seyyed Meysam Abtahi Foroushani²

1. Department of Biology, Faculty of Sciences, Urmia University, Urmia, Iran

2. Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Background: In previous studies, the effects of food restriction on the changes in immune responses and brain dopamine content have been reported. On the other hand, it has been shown that immune cells, in addition to dopamine production, also have dopamine receptors. The purpose of this study was to evaluate the effect of inhibition of D2 dopamine receptors on several functions of monocytes of peripheral blood in rat under food restriction.

Methods: In this experimental study, 36 male Wistar rats (weighing 200-250 gr) were allocated into six groups (n=6), including control groups, food restriction (25%), food restriction (50%), food restriction (75%), food restriction 75% and Sulpiride and rats treated with Sulpiride. Sulpiride was injected Intracerebroventricular at a concentration of 50 µg / rat on day 21 after the study initiation. At the end, the mice were bled and peripheral blood mononuclear cells were isolated by ficoll gradient method.

Results: Food restriction caused a significant decrease in the activity of monocyte cells like neutral red uptake test and respiratory burst (NBT reduction test) simultaneously with decreasing lymphocytes proliferation after stimulation with phytohemagglutinin. Administration of Sulpiride with a 75% Food Restriction resulted in the improvement of these functions of monocyte cells as well as lymphocyte proliferation.

Conclusion: Intracerebroventricular administration of dopamine D2 receptor antagonists (sulpiride) effectively inhibited the effects of a severe dietary restriction on the suppression of immunity system.

Key words: Dopamine, Monocyte, Food Restriction Sulpiride, Rat.

ICIA2018\Immunology of Exercise, Aging, and Nutrition\Poster Discussion\8135
8135

An in vitro Evaluation of Anti-aging Effect of G2013 (G2013) Based on Enzymatic Oxidative Stress Gene Expression Using Healthy Individuals PBMCs

Mahsa Taebe, Seyed Shahabeddin Mortazavi Jahromib, Abdollah Jafarzadeha, Mohammad Reza Mirzaeic, Abbas Mirshafieyd

1. Department of Immunology, School of Medicine, Rafsanjan University of Medical Sciences, Rafsanjan, Iran
2. Department of Cellular and Molecular Biology, Kish International Campus, University of Tehran, Kish, Iran
3. Molecular Medicine Research Center, Rafsanjan University of Medical Sciences, Box: 771664-8798, Rafsanjan, Iran
4. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Box: 14155-6446, Tehran, Iran

Background: Aging is usually associated with increased levels of oxidants, and may result in damages caused by oxidative stress. There is a direct relationship between aging and increased incidence of inflammatory diseases. The present research intended to study the anti-aging and anti-inflammatory effects of the drug G2013 (guluronic acid) at low and high doses on the genes expression of a number of enzymes involved in oxidative stress (including SOD2, GPX1, CAT, GST, iNOS, and MPO) in peripheral blood mononuclear cells (PBMCs) of healthy individuals under in vitro conditions.

Methods: Venous blood samples were taken from 20 healthy individuals, the PBMCs were isolated and their RNAs extracted and their cDNAs were synthesized, and the genes expression levels were measured using the qRT-PCR technique.

Results: Our results indicated that this drug could, at both low and high doses, significantly reduce the expression of the genes for SOD2, GPX1, CAT, and GST compared to the LPS group ($p < 0.0001$). Moreover, it was noticed that the drug is able to significantly reduce gene expression levels at the high dose and at both doses (low and high), for iNOS and MPO compared to the LPS group ($p < 0.0001$), respectively.

Conclusions: The present research showed that G2013, as a novel NSAID drug with immunomodulatory properties, could modulate the expression levels of the genes for SOD2, GPX1, CAT, GST, iNOS, and MPO, to the level of healthy gene expression, and possibly it might reduce the pathological process of aging and age-related inflammatory diseases.

Keywords: G2013 Guluronic acid Anti-aging Oxidative stress NSAID Aging

ICIA2018\Immunology of Exercise, Aging, and Nutrition\Poster Discussion\8271
8271

Effect of Saturated and Unsaturated Fatty Acids on T Helper Differentiation in Vitro

Ayssouda Hossein zade¹, Maryam Moogooei¹, Shayan Mostafaei², Hossein Rahavi¹, Reza Mansouri^{1*}, Sepideh Soukhtezari³, Mojgan Ghaedi⁴

1. Department of Immunology, School of Medicine, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

2. Department of Biostatistics, School of Medical Sciences, Tarbiat Modares University, Tehran, Iran

3. Department of Medical Biotechnology, School of Medical Sciences, Tarbiat Modares University, Tehran, Iran

4. Department of Immunology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

Background: Autoimmunity shows a concerning growth recently. It seems that role of diet is important for development of autoimmunity. Nutritional elements can affect immune system functions. Oversupply or deficiency of specific metabolites may change performance of immune cells, especially that of T helper lymphocytes. For example, glucose, omega 3 and poly unsaturated fatty acids can induce specific Th subsets which play a critical role in autoimmune diseases. T lymphocytes, mostly T CD4⁺ (Th cells) not only play a critical role in orchestrating immune responses, but also they have major role in pathogenesis of some autoimmune disease.

Methods: We designed a research to find out effects of oleic and palmitic acids on differentiation of Th cells. We added oleic acid, palmitic acids and combination of them on peripheral blood mononuclear cells (PBMC) culture and cells were harvested for RT-PCR and flow cytometry analysis after seven days.

Results: Our results showed that palmitic and oleic acids induce Th1 and Th17 subsets (p-value<0.05).

Conclusion: All data showed that adding only 1mM of oleic and palmitic acids change Th cells metabolism and induce Th1 and Th17 induction. In this study saturated fatty acid, unsaturated fatty acid, and combination of both of them result in inflammatory Th subsets induction.

Keywords: T Helper Cells, Differentiation, Metabolism, Fatty Acids, mTOR



Immunology of Infectious Diseases

Bacteria and Fungi

Poster Discussion

14th International Congress of Immunology and Allergy (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Immunology of Infectious Diseases Bacteria and Fungi\Poster Discussion\7943
7943

OmpA Gene Targeted Real-time PCR Compared with the Conventional Culture Method for Detection of *Acinetobacter baumannii* in Pneumonic and *Zataria multiflora*-treated Balb/c Mice

Niloofer Hassannejad¹, Abbas Bahador², Kazem Parivar³, Hadi Ghazanfari⁴

1. Department of Cellular and Molecular Biology, Faculty of Sciences, Science and Research Branch of Islamic Azad University, Tehran, Iran.
2. Department of Microbiology, Faculty of Medicine, Tehran University of Medical Sciences, Tehran, Iran
3. Department of Biology, Faculty of Sciences, Research and Science Branch, Islamic Azad University, Tehran, Iran.
4. Department of Immunology, Faculty of Medicine, Tehran University of Medical Sciences, Tehran, Iran

Background: *Acinetobacter baumannii* is an important pathogen in health care-acquired infections and has free existence of multidrug-resistance responsible for severe nosocomial and community-acquired pneumonia. Currently, mouse model for *A. baumannii* pneumonia is essential for designing novel therapeutic agents.

Methods: In this report, for the first time we described a mouse model of *A. baumannii* using clinical and 19606R standard strains for developing a quantitative real time PCR. The qRT-PCR was used for rapid identification of *A. baumannii* infection from lung tissues of Balb/c mice with pneumonia and also for evaluation of antibacterial effects of *Zataria multiflora* extract on pneumonic Balb/c mice.

Results: Three doses of bacterial strains (0.5×10^8 , 1×10^8 , 1.5×10^8 cfu/ml) were used intranasally for three days to infect the mice. To treat the pneumonic mice, one day after infection with 1×10^8 cfu/ml bacterial strains, mice were treated with 60 mg/kg of *Z. multiflora* ethanolic extract I.P. for 3 days. Lung tissues of pneumonic and treated mice were cultured conventionally and the results were compared with the evaluation of *A. baumannii* *ompA* gene expression using qRT-PCR. In all experiments, clinical isolate had better positive results at day three with highest dose than 19606R strain either in culture (4 versus 3) or in qRT-PCR assay (5 versus 4). However, qRT-PCR detection was 100%, the specificity was 70%, and the positive predictive value was 27%. Our data also shown that *Z. multiflora* extract significantly decreased the number of positive samples in both clinical (from 4 to one) and in standard isolates (from 3 to zero).

Conclusion: The qRT-PCR detection of *A. baumannii* in lung tissue of Balb/c mice model had a high sensitivity as compared to the culture-based method. Also, *Z. multiflora* extract could be a new therapeutic agent for *A. baumannii* pneumonic infection.

Keywords: *Acinetobacter baumannii*, OmpA, Quantitative Real time PCR, *Zataria multiflora* Boiss, Pneumonia, Balb/c mice

ICIA2018\Immunology of Infectious Diseases Bacteria and Fungi\Poster Discussion\8084
 8084

Stimulation of PBMCs of Patients with Abdominal Aortic Aneurysm with *Helicobacter pylori* Antigens Increases IL-21 Production

Hamid Aria^{1,2}, Mehdi Kalani³, Hossein Hodjati⁴, Mehrnoosh Doroudchi¹

1. Department of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

2. Bioinformatics and Computational Biology Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

3. Prof. Alborzi Clinical microbiology research center, Nemazee hospital, Shiraz University of Medical Sciences, Shiraz, Iran

4. Department of Vascular Surgery, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Abdominal Aortic Aneurysm (AAA) is the local dilatation of abdominal aorta and an inflammatory condition in which cytokines may play a pathogenic role. AAA shares several pathogenesis criteria of atherosclerosis, therefore, we asked if stimulation with CagA+ and CagA- *Helicobacter pylori* has different effects on the level of cytokines produced by AAA patients' PBMCs compared to controls.

Methods: PBMCs were isolated from 5 men with diagnosis of AAA and 5 men with normal/insignificant angiography, CT-Scan and Ultrasonography results. Then, PBMCs were cultured in separate plates with bacterial extract of CagA+ and CagA- *Helicobacter pylori*. The supernatants were then removed to measure IL-2, IL-4, IL-5, IL-6, IL-9, IL-10, IL-13, IL-17A, IL-17F, IL-21, IL-22, IFN- γ and TNF- α using a commercial fluorescent-labeled bead assay.

Results: After stimulation with CagA+ *Helicobacter pylori*, patients' PBMCs (18.43 ± 12.91 Pg/ml) produced higher IL-21 compared to untreated PBMCs (10.84 ± 4.92 Pg/ml) ($P=0.06$). While after stimulation with CagA+ *Helicobacter pylori*, controls' PBMCs (9.90 ± 5.95 Pg/ml) produced higher levels of IL-13 compared to untreated PBMCs (4.59 ± 2.26 Pg/ml) ($P=0.06$). A remarkable observation was the huge production of IL-9 by the controls' PBMCs but not controls after stimulation by both CagA+ and CagA- *Helicobacter pylori* ($P=0.05$).

Conclusion: *Helicobacter pylori* is suggested to be one of the bacterial causes of endothelial lesion at the beginning of atherosclerotic process. Because Th2 cytokines are associated with the progression of AAA, the possible role of IL-13 in the onset of disease is noteworthy. Also it can be suggested that the lack of IL-9 may have an effect on AAA disease.

Keywords: Abdominal aortic aneurysm, cytokines, *Helicobacter pylori*

ICIA2018\Immunology of Infectious Diseases Bacteria and Fungi\Poster Discussion\8196
8196

Correlation between Recurrent Vulvovaginal Candidiasis and Dectin-1 Y238X Gene Polymorphism

Hamid Badali¹, Fardin Ahmadkhani^{2*}

1. Department of Medical Mycology and Parasitology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

2. Student Research Committee, Mazandaran University of Medical Sciences, Sari, Iran

Background: Vulvovaginal candidiasis is a frequent disease affecting approximately more than 75 % of all childbearing women at least once in their lifetime by overgrowth of opportunistic *Candida* species. Recurrent vulvovaginal candidiasis (RVVC) is common in otherwise healthy individuals. Several risk factors were reported to contribute to RVVC susceptibility. A polymorphism in Dectin-1 (Y238X, rs16910526) was identified in patients with RVVC and hypothesized that genetic factors play an important role in susceptibility to RVVC. Herein, we aimed to survey the polymorphisms in the Dectin-1 gene, linked to susceptibility to RVVC.

Methods: In the current study, blood samples were obtained from 25 patients who had frequent vulvovaginal candidiasis relapses and were diagnosed as RVVC. In addition, blood cultures were obtained from control group comprising of healthy individuals (n=25) with no history of RVVC, vaginal discharge, or itching on the day of examination. Dectin-1 Y238X gene polymorphism was investigated using DNA sequencing and bidirectional polymerase chain reaction (PCR) amplification of specific alleles (Bi-PASA), as previously described by Carvalho et al.

Results: The analysis revealed that all of the patients were wild-type homozygous for Dectin-1 Y238X polymorphisms. None of the individuals showed heterozygous or mutant homozygous Dectin-1 polymorphism.

Conclusion: No significant correlations were observed between the susceptibility to RVVC and Dectin-1 Y238X polymorphism in the Iranian population, which was not previously studied.

Keywords: *Candida* species, Dectin-1 Y238X gene polymorphism, RVVC.

ICIA2018\Immunology of Infectious Diseases Bacteria and Fungi\Poster Discussion\8260
8260

The Effects of Opium Addiction on the Immune System Function in Patients with Fungal Infection

Alireza Izadi¹, Seyyed Amin Ayatollahi Mousavi², Gholamreza Asadikaram³, Nouzar Nakhaee⁴

1. Department of Medical Parasitology and Mycology, School of Health, Tehran University of Medical Sciences, Tehran, Iran

2. Department of Medical Mycology and Parasitology, School of Medicine, Kerman University of Medical Sciences, Kerman, Iran

3. Endocrinology and Metabolism Research Center, Institute of Basic and Clinical Physiology Sciences, and Dept. of Biochemistry Kerman University of Medical Sciences, Kerman, Iran

4. Neuroscience Research Center, Kerman University of Medical Sciences, Kerman, Iran

Background: The use of narcotics such as opium exposes addicts as susceptible targets of different diseases so that they might easily be exposed to different diseases such as fungal infections. The present study aimed to investigate the effects of addiction to opium and fungal infection on plasma levels of certain cytokines including IL-4, IL-6, IL-17, IFN- γ and TGF- β .

Methods: Present study includes 72 individuals who were divided into 4 groups: 1) opium addicted with fungal infection; 2) opium-addicted without fungal infection; 3) non-opium-addicted with fungal infection; and 4) normal individuals (non-opium-addicted and non-fungal infection). The fungal samples, after being detected and confirmed by a physician, were prepared based on clinical symptoms and then analyzed by direct smear and culture method. The measurement of the plasma level of cytokines was done by ELISA method.

Results: The comparison of the mean of the plasma level of cytokines shows that addiction to opium and fungal infection had respectively the significant effects on the plasma levels of IL-17, IFN- γ , TGF- β cytokines in all studied groups. The interaction of addiction to opium and fungal infection was only significant in the case of plasma level of IL-6.

Conclusion: Addiction to opium and fungal infection, either separately or simultaneously with each other, pose significant effects on the immune system and cause disorders in the cytokine network and the immune system and also provides a suitable environment for fungal infection.

Keywords: Opium; Addiction; Fungal infection; Cytokine

ICIA2018\Immunology of Infectious Diseases Bacteria and Fungi\Poster Discussion\8569
8569

Investigating the Association between *Helicobacter pylori* Infection and IgE Increase in Children Under the Age of 15 Years

Nima Rostami Koshki¹, Mehdi Ebrahimiyan², Mohammad Salehi³

1. Yong Reserch and Elit Club, Marand Branch, Islamic Azad University, Marand, Iran

2. Department of Biology, Naein Branch, Islamic Azad University, Naein, Iran

3. Department of Biology, Urmia Branch, Islamic Azad University, Urmia, Iran

Background: Almost half of the world population is infected with *Helicobacter pylori*, a major risk factor for chronic gastritis, peptic ulcer and gastric cancer. The most effective way to improve the eradication of *H. pylori* peptic ulcers is recommended. Recent studies indicate that there is a relationship between infection with *Helicobacter pylori* and an increase in the level of IgE in the blood of patients. The purpose of this study was to determine the association between total IgE and *Helicobacter pylori* infection as a reliable diagnostic method.

Methods: From 45 patients referred to Children Hospital Tabriz, suspected Gastric biopsies were obtained. Biopsies were transported to the laboratory in Stuart transport medium. First, endoscopy and then biopsy specimens were prepared and further confirmed using urease test. Subsequently, the samples were confirmed by the PcR test. Anti-H. Pylori IgG anti-H. Pylori-CagA IgG, and total plasma IgE levels were measured using ELISA and then statistically analyzed.

Results: 19 out of 45 patients had H. pylori infection, out of which 16 patients were CagA positive.

Conclusion: The findings of the present study demonstrated that there is a significant relationship between *Helicobacter pylori* infection and elevated levels of IgE in the blood of patients.

Keywords: *Helicobacter pylori*, IgE, ELISA, Endoscopy

ICIA2018\Immunology of Infectious Diseases Bacteria and Fungi\Poster Discussion\9768
9768

Immediate Hypersensitivity and Serum IgE Antibody Responses in Patients with Dermatophytosis

Pro.AR.Khosravi¹, Hadis khamevar²

1. Mycology Research Center, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran

2. Student of Mycology, Faculty of Veterinary, University of Tehran, Tehran, Iran

Background: The association of dermatophytes with atopic patients and improvement in allergic signs with antifungal treatment suggest a possible link between chronic infection and atopy. The purpose of this study was to determine skin reactivity and serum IgE antibody responses in patients with chronic and acute dermatophytosis.

Methods: One hundred and sixty-three patients with chronic dermatophytosis, 35 patients with acute dermatophytosis, 41 atopic patients and 49 healthy subjects were enrolled in this study. Sensitization to *Trichophyton mentagrophytes* (*T. mentagrophytes*), *Candida albicans* and *Aspergillus fumigatus* antigens has been evaluated in patients by skin prick test (SPT) and by the presence of specific IgE antibody in enzyme-linked immuno-sorbent assay (ELISA).

Results: Positive immediate hypersensitivity (IH) reactions were obtained in 95.1% of the atopic patients with chronic infection for *T. mentagrophytes*, representing a significant difference from other patient groups ($P < 0.05$). Specific anti-*T. mentagrophytes* IgE antibodies were detected in atopic patients with chronic (65.9%) and acute (50%) dermatophytosis, while none of the atopic subjects had positive IgE reactions to *T. mentagrophytes*.

Conclusion: The results showed significant higher positive IH and specific anti-*T. mentagrophytes* IgE responses in atopic patients with chronic dermatophytosis than the other groups.

Keywords: IgE, hypersensitivity, dermatophytosis, *T. mentagrophytes*

ICIA2018\Immunology of Infectious Diseases Bacteria and Fungi\Poster Discussion\9796
9796

Probiotic *Lactobacillus* Strains Stimulate the Innate Immune Response and Modulate the TLR Expression against *Salmonella* Enteritidis Infection *in vitro*

Mona Moshiri¹ (Presenter author), Mohammad Mehdi Soltan Dallal^{1,2}, Farhad Rezaei³, Masoumeh Douraghi¹, Laleh Sharifi⁴, Zahra Noroozbabaei³, Mehrdad Gholami⁵, Abbas Mirshafiey⁶

1. Division of Microbiology, Department of Pathobiology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

2. Food Microbiology Research Center, Tehran University of Medical Sciences, Tehran, Iran

3. Department of Virology, Tehran University of Medical Sciences, Tehran, Iran

4. Uro-Oncology Research Center, Tehran University of Medical Sciences, Tehran, Iran

5. Department of Microbiology, Faculty of Medicine, Iran University of Medical Sciences, Tehran, Iran

6. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

Background: Lactic acid bacteria (LAB) have been observed to modify the immune responses and could have therapeutic effects in inflammatory disorders. The aim of the present work was to evaluate the effects of *Lactobacillus acidophilus* (*L. acidophilus*) and *Lactobacillus casei* (*L. casei*) strains on toll-like receptor (TLR2 and TLR4) expressions in HT29 cell line infected with *Salmonella enterica* serovar Enteritidis (*SesE*).

Methods: HT29 cells were cultured in Roswell Park Memorial Institute medium supplemented with 10% fetal bovine serum and 1% penicillin/streptomycin. The cells were treated with the *Lactobacillus* strains after or before challenge with *SesE*. At 2 and 4 hours post-infection, we measured changes in the expression levels of TLR2 and TLR4 via real-time polymerase chain reaction method.

Results: Probiotic strains increased the proportion of TLRs expressions and stimulate the innate immune response, when uninfected HT29 cells were treated with lactobacilli. While, the administration of the probiotic strains reduced the TLRs expression previous and also after the challenge with *SesE*.

Conclusion: Our findings show that both of the *L. acidophilus* and *L. casei* boosted innate immune responses, as following the stimulation of TLR signalling and the downregulation of a broad array of pro-inflammatory cytokines after infectin with *SesE*. Although, *L. acidophilus* display greater anti-inflammatory activity than *L. casei* in this work. Further *in vivo* and *in vitro* studies are required to further elucidate the mechanisms underlying this anti-inflammatory effect.

Keywords: Probiotics, *Lactobacillus*, Toll-like receptors, Innate immunity, *Salmonella* Enteritidis

ICIA2018\Immunology of Infectious Diseases Bacteria and Fungi\Poster Discussion\9801
9801

Gastrointestinal Inflammation: Microbial and non-Microbial Factors

Shila Jalalpour¹, Vahid Mirzaee², Abdollah Jafarzadeh^{3,4,5} Afshin Tahmasebi²

1. Molecular Medicine Research Center, Rafsanjan University of Medical Sciences, Rafsanjan, Iran, shilla.
2. Rafsanjan University of Medical Sciences, Rafsanjan, Iran
3. Immunology of Infectious Diseases Research Center, Rafsanjan University of Medical Sciences, Rafsanjan, Iran
4. Department of Immunology, School of Medicine, Rafsanjan University of Medical Sciences, Rafsanjan, Iran
5. Department of Immunology, School of Medicine, Kerman University of Medical Sciences, Kerman, Iran

Background: Gastrointestinal inflammation can be caused by microbial and non-microbial factors, such as life style: alcohol use, stress, or the use of certain medications such as aspirin or other anti-inflammatory drugs and *Helicobacter pylori*. *H. pylori* is common, infecting around 50% of the global human population. The aim of present study was investigate and compare prevalence of microbial and non-microbial gastrointestinal inflammation in visited patients in endoscopy of unit of Aliebn-Abitaleb Hospital in Rafsanjan city. This descriptive cross-sectional study was conducted in the second half of 1396 in patients refer to gastroenterologist. Gastrointestinal inflammation diagnosis by a specialist physician with endoscopy method.

Results: According to the result, of the 87 patients with inflammation, 51 (58.62%) patients had *H. pylori* infection and 36 (41.37%) patients were negative *H. pylori* infection.

Conclusion: The results of this study indicate a higher prevalence of inflammation in people who are infected with *H. pylori* colonization. Due to the importance and role of *H. pylori* in gastrointestinal inflammation and the fact that the persistence of gastrointestinal inflammation can lead to peptic ulcers and cancer, it is important *H. pylori* treatment and elimination in patients.

Keywords: Gastrointestinal inflammation, Gastric, *Helicobacter pylori*, Endoscopy, Colonization

ICIA2018\Immunology of Infectious Diseases Bacteria and Fungi\Poster Discussion\9891
9891

Evaluation of Antibacterial and Cytotoxic Effects of K4 Synthetic Peptide

Mohammad Mehdi ADIBZADEH SERESHGI¹, Kazem Ahmadi¹, Mahdi Fasihi-Ramandi^{*,1}

1. Molecular Biology Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran

Background: With increasing antimicrobial resistance to common antibiotics, development of alternative therapeutic strategies is necessary. Also, antibacterial peptide with numerous behavior and different properties such as net charge, hydrophobicity, length, etc. could act against pathogenic microorganism. In recent years, novel peptides with activity against wide range of bacteria have been introduced.

Methods: In this study, seventeen pathogenic bacteria were chosen to study the antibacterial effect of K4 peptide using MIC and MBC assays. The therapeutic index (TI) of this peptide experimentally calculated by the ratio of HC50 to MIC as a parameter to represent the specificity of AMP. *In silico* analysis was performed to predict the physico-chemical properties, structures, and behavior of this peptide. *In vitro* cytotoxic effect of peptide was evaluated on the Hela cell line using MTT assay, and the amount of macrophage nitric oxide production was measured by Griess method on the J774 cell line.

Results: Peptide concentrations of 25-400 µg/ml was seen as the MIC value results for different bacteria. MBC assay showed such a result with concentration of more than 25-400 µg/ml. The result of hemolysis assay was 24 percent at 1 mg/ml concentration. The amount of nitric oxide production of macrophage cell line J774 was 25.9873 µM at 6.3 µg/ml peptide concentration.

Conclusion: K4 peptide had strong antibacterial effect on some bacteria such as *B. melitensis*. This peptide may have a role in immunity with nitric oxide production. Additionally, it enhances bacterial killing mechanism of macrophage, making this peptide a potential agent against pathogens.

Keywords: Antimicrobial peptide; Cationic peptide; MIC; MBC; Nitric oxide.

ICIA2018\Immunology of Infectious Diseases Bacteria and Fungi\Poster Discussion\9913
9913

Egg Yolk Immunoglobulin Against *ETEC*, Prevention and Treatment of Infections

Razieh Dawodi¹, Seyed. Latif Mousavi Gargari², Shahram Nazarian³

1. Student Master, Department of Cell Biology, Faculty of Basic Sciences, Shahed University, Tehran, Iran

2. Professor, Department of Cell Biology, Faculty of Basic Sciences, Shahed University, Tehran, Iran

3. Assistant Professor, Biology Research Center, Imam Hossein University, Tehran, Iran

Background: Diarrheal illnesses are a severe public health problem and a major cause of morbidity and mortality in diarrhea in developing countries and travelers to these countries. 30-40% of cases of diarrhea in the world due to infection with *E. coli* strains such as *ETEC* occurred infection with *ETEC* is the leading cause of travelers' diarrhea and a major cause of diarrheal disease in lower-income countries, especially among children. Irrational use of antibiotics for the treatment of diarrhea is a global problem.

Due to the increasing prevalence of antibiotic resistance around the world and spreading to humans, concern about the presence of antibiotic residues in the environment, the protection of intestinal flora against the side effects of antibiotic and considering that vaccines have not been effective in treating the infection so far, the use of antibodies as an agent for passive immunity to prevent and treat infections.

The properties of IgY such as easy and inexpensive production compared to mammalian serum antibodies, non-invasive and animal rights protection, non-activating the complement system and non-response to rheumatoid factors, have paved the way for its utilization as an anti-infectious agent within the mammalian gut.

Methodes: Chickens were immunized intramuscularly with *ETEC* formalin killed emulsified with an equal volume of Freund's adjuvant to obtain *ETEC*-specific IgY loaded eggs. After collecting eggs the water-soluble fraction was isolated from egg yolk to utilize for the analysis of IgY properties. the antibody titer was measured by ELISA technique.

Results: In examining ELISA results, increased antibody titers showed a significant increase compared to control samples. The produced bird's immunoglobulin has the ability to detect and attach to the antigen and, by means of the neutralization mechanism, prevents the binding, colonization and replication of the animal's intestines and causes the disease.

Keywords: Formalin killed, *ETEC*, Egg yolk immunoglobulin



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Immunology of Infectious Diseases Viruses

Poster Discussion

ICIA2018\Immunology of Infectious Diseases Viruses\Poster Discussion\6316
 6316

Study of the Effects of Combined Therapy with Sodium Valproate, Peg-interferon and Corticosteroid on Proviral Load of HTLV-1 and Expression of Rel-A, Creb and IL-1 Genes in Patients with HTLV-1 Associated Spastic Paraparesis (HAM/TSP)

Davoud Ghaffarian¹, Reza Farid Hosseini², Hadi Atabati¹, Reza Boostani³, Abdolrahim Rezaei^{4*}, Houshang Rafat Panah⁵

1. Department of Immunology, Mashhad University of Medical Sciences, Mashhad, Iran

2. Department of Immunology, Mashhad University of Medical Sciences, Mashhad, Iran

3. Department of Neurology, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

4. Inflammation and Inflammatory Diseases Research Center, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

5. Department of Immunology, Mashhad University of Medical Sciences, Mashhad, Iran

Background: Human T-Cell Lymphotropic Virus Type I (HTLV-I) is endemic in northeast of Iran. Although chronic inflammation of spinal cord seems to play a major role in the disease pathogenesis, immunomodulatory treatments have not shown significant improvement in disease symptoms. This study was conducted to evaluate the effect of sodium valproate in combination with Peg-Interferon (Peg-IFN) and prednisolone on proviral load and inflammatory factors of patients with HTLV-I associated myelopathy/tropical spastic paraparesis (HAM/TSP).

Methods: In this pilot clinical trial, 10 HAM/TSP patients, referred to HTLV- 1 clinic of Qaem Hospital of Mashhad, Iran in 2012, were treated with sodium valproate, Peg-IFN and prednisolone for 6 months. Peripheral blood mononuclear cells (PBMCs) were isolated using Ficoll Hypaque and the RNA was extracted from PBMCs. Complementary DNA was synthesized using TaqMan Gold RT-PCR Kit. A real-time PCR TaqMan method was designed and optimized for evaluation of Rel-A, Creb and IL-1 genes expression.

Results: The analysis in this study indicated that proviral load and IL-1 mRNA expression in patients were significantly lower after treatment. Despite this decline, Creb and IL-1 mRNA expression was not significant.

Conclusion: The combination of sodium valproate, Peg-IFN and prednisolone seems an effective treatment for proviral load and relatively efficient for damaging inflammatory factors of HAM/TSP; however, further studies with sufficient sample sizes are necessary.

Keywords: Sodium valporate, Peg-Interferon, prednisolone, HAM/TSP

ICIA2018/Immunology of Infectious Diseases Viruses/Poster Discussion/7877
7877

The Effects of Curcumin on Apoptosis and Cytotoxicity-Related Molecules in HTLV-1-Associated Myelopathy/Tropical Spastic Paraparesis (HAM/TSP) Patients

Houshang Rafatpanah¹, Asadollah Mohammadi¹, Reza Boostani², Zohre Poursina¹, Shadi Zamanian¹

1. Immunology Research center, Division of Inflammation and Inflammatory Diseases, Mashhad University of Medical sciences, Mashhad, Iran

2. Department of Neurology, Mashhad University of Medical sciences, Mashhad, Iran

Background: Apoptosis is a universal cellular defense mechanism against viral infection. Curcumin, an anti-inflammatory phytochemical, induces apoptosis through mitochondrial and receptor-mediated pathways, as well as activation of caspase cascades. Here, we investigated the impact of supplementation with curcumin on the expression of a panel of apoptosis- and cytotoxicity-related genes in patients suffering from human T lymphotropic virus Type 1 (HTLV-1)- associated myelopathy/tropical spastic paraparesis (HAM/TSP), a progressive demyelinating neuroinflammatory disease caused by HTLV-1 infection.

Methods: We enrolled 21 HAM/TSP patients in this study. Curcumin nanomicelles (80 mg/day, orally) were administered once a day for 12 weeks. The mRNA levels of total Fas (tFas), membrane-bound Fas (mFas), Fas-Ligand (FasL), TNF-related apoptosis-inducing ligand (TRAIL), perforin, granzyme A, granzyme B and granulysin were analyzed before and after treatment in peripheral blood lymphocytes. Protein levels of Fas, FasL, TRAIL and granulysin were also measured in serum using ELISA.

Results: Curcumin supplementation inhibited FasL mRNA production and up-regulated the expression of pro-apoptotic molecules granzyme A (at the mRNA level) and granulysin (at the protein level), suggesting degranulation of granulysin-bearing cells following curcumin supplementation. Conversely, Curcumin did not affect Fas, TRAIL, perforin, granzyme B at the mRNA level, and anti-apoptotic molecules sFas, sFasL and sTRAIL at the protein level.

Conclusions: The present results suggest that curcumin supplementation increases cytotoxicity-related molecules granzyme A and granulysin in patients with HAM/TSP.

Keywords: HTLV-1; Curcumin; Apoptosis; HAM/TSP

ICIA2018\Immunology of Infectious Diseases Viruses\Poster Discussion\7892

7892

Sequence and Phylogenetic Analysis of LTR region of Human T-Cell Lymphotropic Virus Type 1 from Mashhad, Iran

SeyedAliMirhosseini¹, MehranMoharreri¹, SeyedRohollahArab¹, BarataliMashkani², SeyedAbdolrahimRezaee³, RezaFaridHosseini⁴, HoushangRafatpanah¹

1. Inflammation and Inflammatory Diseases Research Center, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

2. Biochemistry and Nutrition Research Center, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran^[1]_{SEP}

3. Immunology Research Center, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran^[1]_{SEP}

4. Allergy Research Center, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran^[1]_{SEP}

Background: Human T-cell lymphotropic virus type 1 (HTLV-1) is widespread around the globe, with endemic areas in Southern part of Japan, Africa, the Caribbean, South America and Northeast of Iran. The virus is responsible for causing adult T-cell leukemia (ATL) and the neurological disorder HTLV-1 associated myelopathy or tropical spastic paraparesis (HAM/TSP) in less than 5% of the infected individuals. Variations in the provirus sequence of HTLV-1 have been used as a tool for molecular epidemiology investigation of HTLV-1 infection.

Methods: In current study, the amplified LTR region was purified, cloned and sequenced from 15 infected individuals living in Mashhad, Iran. The sequences were compared with the sequences from Neyshabur, Torbat-e Heydarieh and Sabzevar. It was also compared with a reference sequence obtained from GenBank (Accession No. J02029) from Japan.

Results: The LTR sequence in the samples belonged to the transcontinental subgroup A. There was no mismatch in the LTR sequences in 13 out of 15 samples collected from Mashhad with those from Neyshabur, Torbat-e Heydarieh, and Sabzevar. However, these LTR sequences were difference in 2.6% of the nucleotides from the J02029.

Conclusion: These results indicate that HTLV-1 infection in the Northeast of Iran had a common origin. However, further investigation is required to identify the source of small differences were observed in those two samples.

Keywords: HTLV-1, HAM/TSP, ATL, LTR, DNA sequencing

ICIA2018/Immunology of Infectious Diseases Viruses/Poster Discussion/8015

8015

Evaluation of Gene and Protein Levels of IL-33 and Soluble Receptor (ST2-Receptor) Patients with HAM/TSP and HTLV-1 Carriers

Saeedeh Mehraban Moghadam¹, Reza Boostani², SeyedAbdolrahim Rezaee¹, Bahareh Fazeli¹, Narges Valizadeh¹, Houshang Rafatpanah¹

1. Immunology Research Center, Division of Inflammation and Inflammatory Diseases, Mashhad University of Medical Sciences, Mashhad, Iran

2. Department of Neurology, Mashhad University of Medical Sciences, Mashhad, Iran

Background: HAM/TSP is a chronic, progressive and neurologic disease in CNS which is caused by HTLV-1. Secretion of pro and inflammatory cytokines such as: IFN- γ and TNF- α have been shown to be associated with HAM/TSP. IL-33 is an inflammatory cytokine released from necrotic tissue as an alarmin because of its high expression in endothelial and epithelial cells exposed to tissue damage or pathogen. In the present study we examined the role of IL-33 and its soluble receptor named sST2 in hTLV-1 infection.

Methods: The population study was included HAM/TSP patients, asymptomatic HTLV-1 carriers (ACs) and healthy controls (HCs). Peripheral blood mononuclear cells were collected and RNA was extracted and cDNA was synthesized. The mRNA expression of IL-33 was quantified by real time PCR. IL-33 and soluble ST2 serum levels was measured by ELISA.

Results: The IL-33 expression was higher in AC group compared with HAM/TSP patients ($p=0.001$) and HCs ($p=0.012$). There was no significant difference in IL-33 expression between HAM/TSP patients and HCs ($p>0.05$). IL-33 serum level was higher in HAM/TSP patients compared with AC group ($p=0.009$). There was no significant difference in IL-33 serum level between HAM/TSP patient and HCs ($p=0.345$). No significant difference in IL-33 serum levels was observed between ACs and HCs groups ($p=0.158$). Finally, there was no significant difference in sST2 serum levels was detected among there groups ($p=0.385$).

Conclusion: The serum levels of IL-33 was markedly elevated in HAM/TSP patients compared to HTLV-1 carriers which might suggest it might be involved in the pathogeneses of HTLV-1 infection.

Keywords: HTLV-1, HAM/TSP, IL-33, sST2

ICIA2018\Immunology of Infectious Diseases Viruses\Poster Discussion\8117
8117

Predictive Factors of Treatment Failure in Patients with Chronic Hepatitis C

Maryam Khosravi¹, Hamid Salimi², Amir Mohammad Alborzi³, Pedram Nazari³, Farzaneh Chahardahcherik³

1. Department of Immunology, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

2. Department of Internal Medicine, Washington University School of Medicine, St. Louis, MO, USA

3. Cancer Research Center Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

Background: Drug resistance and relapse of disease are the main challenges encountered when treating patients with chronic Hepatitis C Virus (HCV) infection. Production of interferon gamma (IFN- γ), IL-29 and other inflammatory cytokines appears to play a critical role in limiting virus replication and thus impacting on patients' response to antivirals therapies. The aim of this study was to assess effects of the above mentioned factors on the lack of response to treatment among a group of chronically infected HCV patients.

Methods: This study included 57 HCV patients who were admitted to *Shariati* Hospital for treatment. Blood samples were collected from each subject before the onset of treatment with ribavirin and peg-interferon. Levels of IL-29, IFN- γ , liver enzymes as well as HCV titer were then measured in each serum sample. To determine HCV titer, RNA was extracted from peripheral blood mononuclear cells (PBMCs) of patients using High Pure Viral RNA Kit. RT-nested PCR was performed on the RNA extracts using primers designed for the Core region of HCV genome. Patients were followed for 24 months post-treatment. Significance level was considered as $P < 0.05$.

Results: While all non-responder (NR) patients carried HCV in their PBMCs, 41.1% and 58.7% of those in sustained virologic response (SVR) group and relapse group were positive for the virus genome, respectively. The level of liver enzymes was significantly higher in patients who contained HCV genome in their PBMCs compared to those with no detectable viral load. The highest level of IL-29 and IFN- γ was detected in SVR group, followed by the relapse and NR groups, respectively

Conclusion: The results from this study revealed a direct correlation between levels of IL-29 and IFN- γ in patients' sera and their response to antiviral therapy. Induction of these immune responses in HCV patients may, therefore, be beneficial in controlling virus replication and thereby preventing disease relapse/treatment failure.

Keyword: Hepatitis C Virus, treatment response, predictive factors

ICIA2018/Immunology of Infectious Diseases Viruses/Poster Discussion/8392
8392

Effect of Aromatherapy Massage on Salivary Cortisol and Psychological Assessments in a Depressed Patient with Hepatitis B: A Reversal-to-Baseline Design

Bijan Pirnia¹, Dr. Kambiz Pirnia², Hamid Reza Taheri Nakhost³, Dr. Ali Reza Moradi⁴, Soheila Rahmani⁵, Dr. Rasool Roshan⁶

1. *Department of Psychology, Faculty of Humanities, University of Science and Culture, Tehran, Iran*

2. *Technical Assistant in Bijan Center for Substance Abuse Treatment, Tehran, Iran*

3. *Iranian National Center for Addiction Studies and Tehran University of Medical Sciences, Tehran, Iran*

4. *Department of Psychology, Faculty of Humanities, School of Psychology, Kharazmi University*

5. *Islamic Azad University, Karaj, Iran*

6. *Department of Psychology, Faculty of Humanities, Shahed University, Tehran, Iran*

Background: Studies show that 33-78% of patients with chronic Hepatitis B (CHB) suffer from depression in Iranian society. Mental disorders such as depression may affect the immune response in these patients, hence it is clinically important. This was conducted aimed to evaluate the efficacy of aromatherapy as a complementary therapy on the level of cortisol and mood index in an opioid dependent patient.

Methods: The patient was a 41 years old man with hepatitis B with a history of pseudoaneurysm and a long history of heroin using who was under methadone treatment. In an ABAB reversed design with a multiple baseline, methadone treatment was evaluated over a 24-week period and aromatherapy massage was presented at two periods last for forty minutes each. The level of salivary cortisol was considered as primary outcomes and the changes in anxiety, stress and depression indices were considered as secondary outcomes. The results were analyzed by linear mixed model (LMM) and Kolmogorov-Smirnov (K-S) tests.

Results: The primary outcomes showed that inhalation aromatherapy led to a decrease in salivary cortisol levels. Secondary outcomes showed that aromatherapy has a significant effect on the improvement of anxiety, stress and depression. Also, there was a significant correlation between the levels of cortisol and triple indices (all<0.01).

Conclusion: Aromatherapy as an alternative and complementary therapy can be used in the management of psychological indices in drug dependent individuals.

Keywords: aromatherapy, hepatitis B, cortisol, pseudoaneurysm, depression, anxiety

ICIA2018\Immunology of Infectious Diseases Viruses\Poster Discussion\8633
8633

Detecting the Effects of Anti-HCV Drug “Daclatasvir” on Apoptosis’ Factors

Somaieh Nasereslami ^{1*}, Taravat Bamdad ¹, Asghar Abdoli ²

1. Department of Virology, School of Medical Sciences, University of Tarbiat Modares, Tehran, Iran

2. Department of Virology, Pasture Institute, Tehran, Iran

Background: Direct-acting antiviral agents (DAAs) have revolutionized the treatment of hepatitis C virus (HCV) infection over the last 5 years. As a result of our better understanding of the HCV life cycle, specific DAAs have been developed for HCV that are able to target the viral proteins implicated in replication of the virus, i.e., the NS3/4A protease, NS5B polymerase, and multifunctional NS5A replication complex. The first-generation protease inhibitors significantly improved the sustained virologic response (SVR) in genotype 1-infected patients, but at the cost of increased side effects, a complex pattern of drug–drug interactions, and viral resistance. In addition, the first-generation drugs still required the use of PEGylated interferon (PEG-IFN) for 24–48 weeks. Oral IFN-free combinations containing at least two DAAs enabled less complex dosing, tolerable side effects, and fewer drug–drug interactions

Methods: In this study MTT dose of Daclatasvir that has minimum cytotoxicity for cell. Flow cytometry test in order to demonstrate. LC3 protein accumulation was carried out on Huh7 cells. After the culture of cells and incubation with 60 pM daclatasvir, transfected in pcDNA-Beclin1, NS5A, P53, Caspase, Bax and Bcl2. After cDNAs synthesis Real-time PCR was carried out on their expression (with HPRT), were T assay was applied to indicate evaluated. The result had been analyzed by Prism software.

Results: To examine the potential mechanism underlying the effects of daclatasvir on cell cycle and apoptosis, we analyzed the expression of genes involved in the cell cycle regulation and apoptosis. We evaluated P53, BCL-2, BAX, and CASP3 gene expression, 48 hours after the transfection with these factors. P53, a known target of daclatasvir, showed a 2.53 fold increase in the infected cells in comparison with the control cells. The expressions of CASP3 and BAX were upregulated in the infected cells in comparison with the control cells (3-fold and 2-fold increase, respectively) ($p < 0.05$). The BCL-2, an anti-apoptotic gene, showed no significant change in response to the increased apoptotic factors expression level.

Conclusion: These findings suggest that DAA-based therapy (Daclatasvir) could be effective in advanced patients and could improve the current therapeutic strategies for treatment hepatocarcinoma.

Keywords: Daclatasvir, HCV, apoptosis, NS5A.

ICIA2018/Immunology of Infectious Diseases Viruses/Poster Discussion/8704
8704

Accessing of TCD8+ Cells Response Against Adipophilin, Epstein - Barr virus, Cytomegalovirus, Influenza Derived Poly Peptide in Patients with Atherosclerosis Compared to Healthy Individuals

Solmaz khansalar¹, Zahra farzaneh¹, Hossein Golmoghdam¹, Mohamad Ali asadi¹,
Saeed Malek Hosseini¹, Mehrnoosh Doroudchi¹, Kurosh kalantar¹

1. Department of Immunology, School of Medicine, Shiraz University of Medical sciences, Shiraz, Iran

Background: Atherosclerosis, a chronic inflammatory process that affects the walls of large and medium-sized arteries, is the leading causes of morbidity and premature death worldwide. Many cell types, including endothelial cells, monocytes, dendritic cells (DCs), lymphocytes, eosinophils, mast cells and smooth muscle cells, contribute to the formation of atherosclerotic plaques. Inflammatory responses at the atherosclerotic plaques can damages self-tissue and subsequently release self-peptides that are engulfed by APCs and will be presented to T cells. CD8+ T lymphocytes may play an important role in atherogenesis. As adipose differentiation-related protein (ADRP, adipophilin) is a prominent component of the cytosolic lipid droplets (CLD) present in most of the cells of the atherosclerotic plaque, the possible use of HLA class I restricted peptide derived from this protein was analyzed as a T-cell epitope presented by APCs. In addition, several reports showed that the role of some pathogens in the immunopathology of atherosclerosis. So, Peptide from Epstein-Barr virus, Cytomegalovirus, influenza also was selected to explore this possibility.

Methods: 28 patients with atherosclerosis and 40 age and sex matched healthy donors were entered the study. PBMCs were isolated and HLA typing was done using flow cytometry. HLA-A2 positive PMCs were cultured and stimulated using specific peptides (Adipophilin 129-137, Flu 58-66, CMV 495-503, EBV 280-288). Cell culture supernatant was used to measure IFN- γ using human ELISA kit.

Results: 11 (39.3%) out of 28 patients with atherosclerosis and 16 (40%) out of 40 healthy individuals were positive for HLA-A02. PBMCs stimulation revealed no statistically significant differences in the levels of IFN- γ between HLA-A2 positive patients or healthy controls against selected peptides.

Conclusion: Adipophilin is not a self Ag presented by APCs at the site of atherosclerotic plaque, but further experiments are needed to completely rule out its importance in the etiopathogenesis of arthrosclerosis.

Keywords: Atherosclerosis, CD8+ T lymphocytes, adipose differentiation-related protein (ADRP), adipophilin, cytosolic lipid droplets (CLD).

ICIA2018\Immunology of Infectious Diseases Viruses\Poster Discussion\9810

9810

HTLV-1 Infection Dampens NK cell Function

Asadollah Mohammadi¹, Reza Boostani², Houshang Rafatpanah³

1. Cellular & Molecular Research Center, Faculty of Medicine, Kurdistan University of Medical Sciences, Sanandaj, Iran

2. Department of Neurology, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

3. Inflammation and Inflammatory Diseases Research Center, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

Background: HTLV-1 is the etiological agent of HTLV-1-associated myelopathy/tropical spastic paraparesis (HAM/TSP). Natural killer (NK) cells are necessary for first defense against virally-infected and stressed cells. They kill target cells by receptor-mediated and granule exocytosis pathways.

Methods: A total of 21 asymptomatic carriers (ACs), 21 HAM/TSP patients and 8 healthy controls (HCs) were recruited. NK cell numbers, phenotype, function, and effector molecules expression were analyzed by flow cytometry. The cytotoxic activity of NK cells was assessed by CD107a degranulation. Also, the effects of NK cell anti-viral functions on HTLV-1 proviral load, Tax and HBZ mRNA expression were analyzed.

Results: Total NK cell numbers showed a decrease in HAM/TSP patients. NK cells from HAM/TSP patients and ACs showed substantial decrease in their function and had lower cytotoxic-related molecules such as Fas, FasL and TRAIL than HCs, but the levels of perforin, granzyme B, and granulysin were similar. No significant correlations between NK cell number, phenotype and function with proviral load, Tax and HBZ mRNA expression were found.

Conclusion: Lower numbers of NK cells and specific defects in their capacity to degranulate may play a role in HTLV-1 persistence. Also, lower numbers of NK cells and activity in HAM/TSP patients could be an effect of a high proviral load of HTLV-1, antigen abundance and/or persistence, rather than a cause. Restoration of NK cell capacity, as achieved by viral load reduction, could therefore contribute to viral control and subsequently HAM/TSP disease susceptibility.

Keywords: HTLV-1; NK cell; Neuroinflammation; HAM/TSP.

ICIA2018/Immunology of Infectious Diseases Viruses/Poster Discussion/9933
9933

Comparison of IL-12, IL-18 and IFN- γ Gene Expression in HTLV-1-Associated Myelopathy / Tropical Spastic Paraparesis (HAM/TSP) Patients, HTLV-1 Carriers and HTLV-1 Negative Individuals

Hamid Reza Bidkhori¹, Houshang Rafatpanah^{2*}, Seyyed Abdolrahim Rezaee², Narges Valizadeh²,
Mohsen Tadayyon, Sanaz Ahmadi Gezeldasht¹, Rahele Miri¹, Arezo Gohari Shabgah²,
Mohammad Reza Hedayati Moghaddam^{1*}

1. Blood-borne Research Center, Academic Center for Education, Culture and Research (ACECR)-Khorasan Razavi, Mashhad, Iran

2. Immunology Research Center, Inflammation and Inflammatory Diseases Division, Mashhad University of Medical Sciences, Mashhad, Iran

Background: Chronic inflammation is among the parameters that may increase the risk of developing HAM/TSP in the HTLV-1 carriers. The imbalance between Th1 and Th2 cells is highly determinative due to variations in the level of cytokines. Interleukins 12 and 18, along with interferon gamma, induce Th1 response. The present study aimed to determine the gene expression and serum levels of IL-12, IL-18 and IFN- γ in patients with HAM/TSP and asymptomatic HTLV-1 carriers compared to healthy subjects.

Methods: In this study, 21 asymptomatic HTLV-1 carriers, 20 HAM/TSP patients and 21 HTLV-1-negative healthy volunteers were enrolled. Real time PCR technique was used to measure the expression level of IL-12, IL-18, IFN- γ genes and proviral load. The plasma levels of cytokines were obtained using ELISA assay.

Results: IL-12 gene was expressed in all healthy controls, nine asymptomatic HTLV-1 carriers and five HAM/TSP patients. The gene expression and the serum levels of IL-18 were lower in the HTLV-1-positive group than in the control group ($p = 0.001$ and $p = 0.012$, respectively). However, there was no significant difference between HTLV-1 carriers and HAM/TSP patients ($p = 0.335$ and $p = 0.695$, respectively). The mean viral load in HAM/TSP patients (3693.2 ± 2444) was higher than non-symptomatic carriers (1388.4 ± 1365.2) ($p = 0.002$). The correlation coefficient between serum level of IFN- γ and viral load in HAM/TSP patients was not statistically significant ($r = -0.142$, $p = 0.55$) but there was a positive correlation between serum level of IL-18 and viral load in this group ($r = 0.654$, $p = 0.002$).

Conclusion: In this study, increased HTLV-1 proviral load and high serum IFN- γ levels in the HAM/TSP group were determined in comparison to the asymptomatic carriers. Moreover, the expression level of IL-18 in the HTLV-1-positive group and the healthy control group was significantly different. Our findings highlight the fact that the proviral load and the inflammatory cytokine profile during HTLV-1 infection may be independent events leading to the occurrence of HAM/TSP.

Keywords: HTLV-1 infection, Non-symptomatic carriers, HAM/TSP, IL-12, IL-18 and IFN- γ .



Immunology of Mycobacterium Tuberculosis

Poster Discussion

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018 Immunology of Mycobacterium Tuberculosis\Poster Discussion\7961
7961

Assessment of B cell Lymphoma 6 (BCL6) Expression Level in Peripheral Blood CD38+B Lymphocytes of Common Variable Immunodeficiency (CVID) Patients

Faezeh Abbasi Rad¹, Fahimeh Khadem¹, Nafiseh Esmail¹, Mazdak Ganjalikhani Hakemi¹, Shokrolah Farrokhi², Roya Sherkat

1. Department of Immunology, school of Medicine, Isfahan Isfahan University of Medical Sciences, Isfahan, Iran
2. Department of Immunology and Allergy, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Research Institute, Bushehr University of Medical Sciences, Bushehr, Iran
3. Aquired Immunodeficiency Research Center, Isfahan University of Medical Sciences, Isfahan, Iran

Background: Common variable immunodeficiency (CVID), the most common symptomatic primary immunodeficiency, is a heterogeneous set of immunological abnormalities including decreased serum levels of antibodies, impaired antibody response to infections or vaccination. The syndrome includes impaired B-cell maturation, impaired somatic hypermutation, reduced numbers of circulating memory B cells, and absent or reduced plasma cells. BCL6 (B cell Lymphoma 6) is a transcription factor which is important for the evolution and proliferation of B cells. This study aimed to investigate the expression of BCL6 in peripheral blood CD38+B lymphocytes of the patients with CVID.

Methods: Blood samples were collected from 14 CVID patients with substitutive immunoglobulin (Ig) therapy before immunoglobulin infusion and 14 normal controls. Then peripheral blood mononuclear cells were isolated by Ficoll-Hypaque density centrifugation. CD19+ B lymphocytes were purified from PBMCs by positive selection using B-cell isolation kit. Flow cytometry method was employed to determine the expression of the BCL6 in CD38+ B cells.

Results: In this study, the expression of BCL6 in CD38 +B lymphocytes in patients was 1.51% and in healthy subjects was 0.58%, respectively. According to the statistical analysis, the difference was not significant ($P>0.05$).

Conclusion: The results showed that there is no significant difference in the mean expression of BCL6 of the CD38 + B cells in the patients with CVID, compared with control group. However, the average BCL6 expression of CD38+ B lymphocytes in patients was more than in control group.

Keywords: B lymphocyte; Plasma cell; PBMC; BCL6; CD38

ICIA2018\Immunology of Mycobacterium Tuberculosis\Poster Discussion\7994
7994

IL-10 -592 A/C Gene Polymorphism and Cytokine Plasma Levels Are Associated with Susceptibility to Drug Resistance Tuberculosis

Atefeh Sadeghi Shermeh¹, Mehdi Shekarabi^{*1, 2}, Ali Akbar Delbandi^{1, 2}, Payam Tabarsi³, Esmail Mortaz³, Mohammad Reza Bolouri¹, Majid Khoshmirsafa¹, Elham Alipour favez¹

1. Department of Immunology, School of Medicine, Iran University of Medical Sciences, Tehran, Iran

2. Immunology Research Center, Institute of Immunology and Infectious Diseases Iran University of Medical Sciences, Tehran, Iran

3. Clinical Tuberculosis and Epidemiology Research Center, National Research Institute of Tuberculosis and Lung Diseases (NRITLD), Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Tuberculosis (TB) as a global public health problem, still remains a major cause of mortality worldwide. In recent decades, Mycobacterium tuberculosis (Mtb) strains resistant to anti-tubercular agents have been emerging as critical issues in tuberculosis treatment. Immune responses to all pathogens are cytokine dependent, Mtb is no exception and variation in cytokine genes, especially single nucleotide polymorphisms (SNPs) may lead to abnormal or ineffective immune response.

Methods: So we investigated SNPs in *IFN-γ* (+874 T/A) and *IL-10* (-592 A/C) genes in relation to these cytokines level and pulmonary tuberculosis treatment responsiveness. A total of 87 patients and 100 healthy individuals were included in this study. According to drug sensitivity test using polymerase chain reaction (PCR), patients divided into two groups, including 67 drug-sensitive (DS-TB) and 20 drug-resistant (DR-TB). Genomic DNA was genotyped for two SNPs by PCR-based methods. Serum cytokine levels were measured using ELISA.

Results: In both patients and controls, the *IL-10* -592A allele and AC genotype were presented in a greater proportion, but these differences were only significant between DR-TB and controls ($p < 0.05$). TB patients showed higher IL-10 serum levels in comparison to healthy individuals ($p < 0.05$). Although, no significant differences were observed for allele and genotype frequencies in *IFN-γ* +874, the serum levels of IFN- γ was higher in DS-TB patient compared with DR-TB and healthy individuals.

Conclusion: Thus, our findings suggest that association of the *IL-10* (-592 A/C) polymorphism with susceptibility to DR-TB in the studied population.

Keywords: Tuberculosis, Polymorphism, IFN- γ , IL-10, Drug resistance tuberculosis

ICIA2018\Immunology of Mycobacterium Tuberculosis\Poster Discussion\8070

8070

Association of P2X7 Gene Common Polymorphisms with Pulmonary Tuberculosis in Lur Population of Iran

Farhad Shahsavari¹, Toomaj Sabooteh^{2*}, Seyyed Amir Yasin Ahmadi³, Ali Amiri⁴, Alireza Azargoon⁴

1. Department of Immunology, Lorestan University of Medical Sciences, Khorramabad, Iran

2. Faculty of Medicine, Lorestan University of Medical Sciences, Khorramabad, Iran

3. Student Research Committee, Lorestan University of Medical Sciences, Khorramabad, Iran

4. Department of Internal Medicine, Lorestan University of Medical Sciences, Khorramabad, Iran

Background: Different genetic and environmental factors are associated with susceptibility to pulmonary tuberculosis (TB) in different individuals of different populations. Based on previous studies role of P2X7 gene common polymorphisms in susceptibility to pulmonary TB was associated with ethnicities. Aim: We intend to perform this study on genetic reservoir (gene pool) of Lur population of western Iran.

Methods: For the present case-control study, 100 unrelated pulmonary TB patients and 100 unrelated controls were enrolled through convenient sampling. TB confirmation was through smear and culture of sputum. Polymerase chain reaction with restriction fragment length polymorphism (PCR-RFLP) was used for molecular assay. This study has been approved in the ethic committee of Lorestan University of Medical Sciences with registration number LUMS.REC.1396.253.

Results: Among the genotypes of polymorphism 1513A/C, AA genotype was associated with susceptibility to pulmonary TB ($P=0.0001$; $OR=4.750$) whereas AC genotype was a protecting factor ($P=0.0001$; $OR=0.192$). Higher genetic reservoir of A allele was associated with more susceptibility to pulmonary TB ($P=0.0001$; $OR=2.879$) whereas C allele was a protecting factor ($P=0.0001$; $OR=0.347$). No significant result was found for -762T/C polymorphism.

Conclusion: In Lur population of Iran, 1513A/C polymorphism of P2X7 is associated with susceptibility to pulmonary TB. It is suggested that bio-information banks should be established and developed in countries.

Keywords: P2X7, Pulmonary Tuberculosis, Lur population, Iran.

ICIA2018\Immunology of Mycobacterium Tuberculosis\Poster Discussion\8261
8261

Cloning, Expression and Immunological Assessment of EspB Protein from *Mycobacterium tuberculosis*

Zahra Hashemi¹, Soheila Ajdary², Fariborz Bahrami², Alireza Rafiee¹, Farhad Riazi Rad²

1. Department of Immunology, Faculty of Medicine, Mazandaran University of Medical Sciences
2. Department of Immunology, Pasteur Institute of Iran

Background: Tuberculosis (TB) is the second leading cause of mortality in the world. One third of the world's population are infected with *Mycobacterium tuberculosis* (MTB), the causative agent of the disease. The most widely used vaccine against TB is BCG; however it provides inadequate protection. Therefore, a more effective vaccine is needed to control TB and for this purpose, identifications of the virulent factors which are both immunogenic and specific for the infection, are highly in demand. The genes residing in "region of difference 1" (RD1) are involved in pathogenesis of MTB. EspB is a newly identified protein, encoded by a gene located in this region, appears to be essential for the virulence of the bacterium. The objective of the present study was to produce recombinant EspB and evaluate its immunoreactivity in humans.

Methods: The encoding gene for MTB was commercially synthesized which was then used as a template to produce an amplicon, flanked by appropriate restriction enzymes by PCR. The PCR product was cloned in-frame into a pET21a expression vector and transformed into *Escherichia coli* BL21 (DE3) for a histidine-tagged recombinant protein production by IPTG induction. The expression and purity of the protein (rEspB) was confirmed by SDS-PAGE and Western blotting analyses using anti-His antibody. PBMCs of patients with confirmed TB and healthy donors were isolated and stimulated with rEspB. Lymphocyte proliferation and IFN- γ production were assessed by thymidine incorporation assay and ELISA.

Result: EspB gene was successfully cloned and expressed in *E. coli*. PBMCs from TB patients recognized rEspB and proliferated in response to this protein. IFN- γ was produced by PBMCs from the patients.

Conclusion: Being a virulent factor of MTB, EspB is a target for anti-mycobacterial immune responses in humans and can be considered as a vaccine candidate against TB.

Keywords: *Mycobacterium tuberculosis*, EspB, PBMC, Immune response.

ICIA2018\Immunology of Mycobacterium Tuberculosis\Poster Discussion\8334
8334

Interferon- γ Gene Polymorphisms in Pulmonary Tuberculosis Patients

Ali Pourmohammad¹, Mohammad Javad Nasiri¹

1. Department of Microbiology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Upon infection with *Mycobacterium tuberculosis*, the IFN- γ plays an essential role for the elimination of pathogen. Subtle variations in IFN- γ gene may influence susceptibility to *M. tuberculosis*. This study investigates the relationship of the single base change polymorphic variants in interferon- γ and tuberculosis (TB) susceptibility.

Methods: We studied a population of 100 patients with culture-proven pulmonary TB and 90 healthy tuberculin-negative control subjects. Polymorphic variants in IFN- γ gene were assessed using PCR-RFLP.

Results: Statistical analyses revealed that IFN- γ 2109 A/G polymorphism was marginally associated with the risk of TB ($P=0.04$).

Conclusions: The present results revealed that certain genetic variants in IFN- γ gene may be associated with TB development, which may be useful preliminary data for future investigation.

Keywords: Susceptibility, Tuberculosis, Interferon-gamma, Polymorphisms

ICIA2018\Immunology of Mycobacterium Tuberculosis\Poster Discussion\8386
8386

Evaluation and Comparison of *IFN- γ* and *IL-2* Gene Expression in Peripheral Blood of Patients with Positive and Negative Quantiferon-TB Test

Mahsa Jafari¹, Dr. Reza mansouri²

1. *Shahid sadoughi university of medical sciences*

2. *Reza Mansouri MD-PhD Assistant Professor Department of Immunology Faculty of Medicine Shahid Sadoughi University of Medical Sciences Safaieye, Bo-Ali street Yazd/Iran*

Background: Tuberculosis (TB) is a leading cause of mortality and morbidity worldwide. This infection is a major public health problem. TB control is largely dependent to early diagnosis and managements of suspected cases. Today several bacteriologic and immunologic methods such as microscopic study of Smear microscopy and sputum Culture. And tuberculin skin test is used for laboratory diagnosis of TB infection, but they were not as successful in decreasing TB as expected. In order to modify mentioned shortcomings, in recent years, IGRA, an immunologic test, is recommended for TB patients as a diagnostic test, and meanwhile releasing *IFN- γ* from lymphocytes is studied. Immunologic methods applied for diagnosing TB are IGRA and TST. These measuring are tentative. They also cannot make distinction between recent TB infection and the current one. *IFN- γ* has a key role in removing mycobacterium TB from the body. Considering this matter that 10% of patients with latent TB are subject to active TB and studding the amount of solution *IFN- γ* alone cannot be helpful in diagnosing latent TB and determining different stages of disease. Thus it is possible that other cytokines such as *IL-2* can help in diagnosing latent TB. Because of the importance of screening and controlling latent TB infection in restricting the disease, besides *IFN- γ* studding gene-expression of relevant biomarkers such as *IL-2* may have an important role in diagnosing latent TB and preventing disease progression. In this study expression of two genes, *IFN- γ* and *IL-2*, in patients with positive or negative Quantiferon-TB test are done. **Methods:** In this research 25 subjects with positive result in Quantiferon-TB test (13 females and 12 males with average age of 58.56 ± 19.296) As the target group and 25 subjects with negative result in Quantiferon-TB test (13 females and 12 males with average age of 58.56 ± 19.296) as witness group were chosen. In this study expression of two genes, *IFN- γ* and *IL-2*, in patients with positive or negative Quantiferon-TB test are done. After RNA extraction from the present blood in Quantiferon tube for mentioned two groups, cDNA syntheses was done and eventually gene-expression by means of real-time PSR technic was studied. Then amount of gene expression of these genes, their relations together and with *IFN- γ* release in Quantiferon-TB test were analyzed statistically. **Results:** In this study with two target and witness groups, findings indicated that *IFN- γ* gene-expression rate has significant correlation with becoming positive in Quantiferon-TB test. But between *IL-2* gene-expression rate and becoming positive in Quantiferon-TB test is not significant correlation. There is no significant correlation between *IFN- γ* and *IL-2* gene-expression rate with *IFN- γ* release in Quantiferon-TB test. **Conclusion:** By knowing the results of current study, *IFN- γ* gene-expression in patient's blood with positive Quantiferon-TB test can only considered as one of Para clinical indexes in diagnosing mycobacterium TBinfection.

Keywords: mycobacterium tuberculosis, Tuberculin Skin test, Interferon Gamma Release Assay, Interleukin 2, Interferon Gamma

ICIA2018 Immunology of Mycobacterium Tuberculosis\Poster Discussion\9870
9870

Study of The Exosomal miRNAs in Patient with Active Pulmonary Tuberculosis

Shamila D. Alipoor^{1,2,3}, Esmaeil Mortaz¹, Payam Tabarsi¹, Johan Garssen^{4,5}, Mohammad Varahram⁶, Ian M. Adcock^{7, 8}, Ali Akbar Velayati⁶

1. Clinical Tuberculosis and Epidemiology Research Center, National Research Institute of Tuberculosis and Lung Diseases (NRITLD), Shahid Beheshti University of Medical Sciences, Tehran, Iran
2. Department of Biotechnology, School of Advanced Technologies in Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran
3. Institute of Medical Biotechnology, Molecular Medicine Department, National Institute of Genetic Engineering and Biotechnology (NIGEB), Tehran, Iran
4. Division of Pharmacology, Utrecht Institute for Pharmaceutical Sciences, Faculty of Science, Utrecht University; Utrecht, Netherlands
5. Nutricia Research Centre for Specialized Nutrition, Utrecht, Netherlands
6. Mycobacteriology Research Centre, National Research Institute of Tuberculosis and Lung Disease (NRITLD), Shahid Beheshti University of Medical Sciences, Tehran, Iran
7. Priority Research Centre for Healthy Lungs, Hunter Medical Research Institute, The University of Newcastle, Newcastle, NSW, Australia

Background: Tuberculosis (TB) remains a major threat to human health. Due to the limited accuracy of the current TB diagnostic tests it is critical to determine novel biomarkers for this disease. Circulating exosomes have been used as diagnostic biomarkers in various human diseases.

Methods: serum-derived exosomes were isolated from TB patients and matched control subjects. The expression of miR-1224, -1293, -484, -425 and -96 were examined by RT-qPCR. Receiver operating characteristic (ROC) curve was performed to evaluate the diagnostic potency of each individual serum exosomal miRNA.

Results: only miR-484, -425 and -96 were significantly increased in the exosomes of TB patients as determined and significantly correlated with the TB infection level in patients. ROC curve analysis showed the diagnostic potency of each individual serum exosomal miRNA with an area under the curve (AUC) =0.72 for miR-484 ($p<0.05$), =0.66 for miR-425 ($p<0.05$) and =0.62 for miR-96 ($p<0.05$). The predictive value of the individual miRNAs was greatest in the patients with high bacterial burden particularly for miR-484 (AUC=0.82) and miR-425 (AUC=0.868) in 3+ patients. ROC curves using combined miRNA expression levels gave greater AUC values with the combination of all 3 being more predictive of TB per se (AUC=0.78).

Conclusion: These results demonstrate that exosomal miRNAs have diagnostic potential in active tuberculosis. The diagnostic power may be improved when combined with conventional diagnostic markers.

Key words: Exosomes, Tuberculosis, miRNA

ICIA2018\Immunology of Mycobacterium Tuberculosis\Poster Discussion\9872

9872

A Computational Study on the Roles of Serum Exosomal miRNAs in Patient with Active Tuberculosis.

Shamila D. Alipoor¹, Esmail Mortaz², Payam Tabarsi², Johan Garssen³, Ian M. Adcock⁴, Mohammad Varahram⁵, Neda Dezfouli², Ali Akbar Velayati⁵

1. Institute of Medical Biotechnology, Molecular Medicine Department, National Institute of Genetic Engineering and Biotechnology (NIGEB), Tehran, Iran

2. Clinical Tuberculosis and Epidemiology Research Center, National Research Institute of Tuberculosis and Lung Diseases (NRITLD), Shahid Beheshti University of Medical Sciences, Tehran, Iran

3. Division of Pharmacology, Utrecht Institute for Pharmaceutical Sciences, Faculty of Science, Utrecht University; Utrecht, Netherlands

4. Airways Disease Section, National Heart & Lung Institute, Imperial College London, London, UK

5. Mycobacteriology Research Center (MRC) National Research Institute of Tuberculosis and lung diseases (NRITLD), Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Tuberculosis (TB) remains a major threat to human health. Due to the limited accuracy of the current TB diagnostic tests it is critical to determine novel biomarkers for this disease. Circulating exosomes have been used as diagnostic biomarkers in various human diseases. In this study a miRNAs network analysis was performed on the changed miRNAs after BCG infection.

Method: MicroRNA target genes were determined using miRTarBase and microT-CDs algorithms. Pathway analysis was performed using EnrichR and the KEGG pathway database. Protein-protein interaction (PPI) networks were constructed and analyzed by STRING online server with an annotation score ≥ 0.5 as the cut-off and visualized and analyzed by cytoscape v.3.4.0. The highly interconnected regions (clusters) were extracted using MCODE plugin in cytoscape. Topological analysis of this network was performed to determine properties of nodes including score and connectivity degree (k) parameters.

Result: The extended network consisted of 295 nodes connected via 2552 edges. A total of three clusters were detected by MODE plugin implemented in cytoscape. The result showed the highest score belonged to cluster A with score 20.83. The number of nodes in clusters A, B, C was 56, 16, 9 with scores: 20.83; 9.6 and 8.75 for clusters A, B and C respectively. Pathway analysis showed that all the clusters contain nodes that involved in the immunological and metabolism pathways.

Conclusion: Our results confirmed that infection with mycobacterium affected exosomal miRNAs which may reflect the intracellular status and pathology of the cells. These data illustrated the potential role of exosomal miRNAs as a biomarker for TB diagnosis.

Keywords: Exosome, Tuberculosis, Computational study



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Immunology of Rheumatic Diseases

Poster Discussion

ICIA2018\Immunology of Rheumatic Diseases\Poster Discussion\7506
7506

Nanocurcumin Impacts on Aberrant T cells Related MicroRNAs Profile Expression in Ankylosing Spondylitis Patients

Yousef Daneshmand pour^{1,2}, Sanam Dolati^{1,3}, Mehdi Yousefi^{1,3}

1. Stem Cell and Regenerative Medicine Institute, Tabriz University of Medical Sciences, Tabriz, Iran

2. Department of medical Genetics , Faculty of Medicine, Tabriz University of Medical sciences

3. Department of Immunology, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Ankylosing spondylitis (AS) is a chronic, progressive inflammatory rheumatic disease that sacroiliac joints and the axial skeleton are involved and back pain and progressive stiffness of the spine are main clinical features. Oligoarthritis of the hips and shoulders, enthesopathy, and anterior uveitis are common but, involvement of the heart and lungs is rare. The aim of current study was to evaluate the effect of Nanocurcumin on expression level of significant miRNAs associated with Th17 and Treg function.

Methods: 24 AS patients were divided in two groups. One group received 80 mg nanocurcumin every day for 16 weeks and 12 other patients received placebo as control group. The peripheral blood mononuclear cells (PBMCs) were collected from blood samples of patients and then, expression level of significant miRNAs was evaluated by Real time PCR (mir-141, mir-155 and mir-200, mir-27, mir-17 and mir-146a).

Results: Our study showed in case of Th17 related miRNAs, expression level of mir-141, mir-155 and mir-200 (P value=0.04, 0.02 and <0.0001, respectively) were significantly decreased in case group. In Treg related miRNAs, mir-17 and mir-27 were significantly decreased while mir-146a was significantly increased (p value=0.0026 and p value=0.0003, p value=0.02 respectively).

Conclusion: Our data indicates significance of Th17 and Treg function in AS and well therapeutic effect of Nanocurcumin on these patients. Also, our results shows designing therapeutic strategies based on changing miRNA expression in Treg and Th17 can be possible treatment for AS patients in future.

Keywords: Ankylosing spondylitis, Nanocurcumin, miRNA, T Cells

ICIA2018\Immunology of Rheumatic Diseases\Poster Discussion\7563

7563

The Potent Suppressive Effect of β -D-mannuronic Acid (M2000) on Molecular Expression of the TLR/NF-kB Signaling Pathway in Ankylosing Spondylitis Patients

Maryam Roozbehkia¹, Mahdi Mahmoudi^{2*}, Somaye Aletaha¹, Nima Rezaei³, Abbas Mirshafiey¹

1. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran)

2. Rheumatology Research Center, Shariati Hospital, Tehran University of Medical Science, Tehran, Iran)

3. Network of Immunity in Infection, Malignancy and Autoimmunity (NIIMA), Universal Scientific Education and Research Network (USERN), Sheffield, UK

Background: Ankylosing spondylitis (AS) is a chronic inflammatory rheumatic disease characterized by the inflammation of sacroiliac joints and axial skeleton. A combination of genetic, environmental and immunological factors are involved in AS's pathogenesis. TLRs are type I transmembrane glycoproteins that play a crucial role in the innate immune responses against invading pathogens. Observational studies have demonstrated a possible association between TLR dysregulation and AS. The β -D-mannuronic acid (M2000), as a novel NSAID with immunosuppressive property, has shown an inhibitory effect on Toll-like receptor (TLR) 2, 4 signaling in HEK293 cells.

Methods: In the present study, we investigated the gene expression of Myd88, IKB-alpha, NF-kB and MAPK14 (genes of the TLR/NF-kB Signaling Pathway) by Real time PCR in AS patients in comparison to healthy subjects and also the effect of β -D-mannuronic acid on disease activity and mRNA expression of these molecules in affected patients.

Results: We showed for the first time that the gene expression level of Myd88, IKB-alpha, NF-kB and MAPK14 was higher in AS patients in comparison to healthy subjects. Moreover we confirmed that the β -D-mannuronic acid not just reduced significantly the disease activity of AS individuals compared to placebo, but also it could significantly decrease the expression level of genes associated with TLR/NF-kB Signaling Pathway in treated patients with M2000.

Conclusion: These results may provide a new therapeutic approach to attenuate inflammatory responses in AS patients,(Identified; IRCT2013062213739N1).

Key words: Ankylosing spondylitis, M2000, TLRs, Mannuronic acid, Myd88, IKB-alpha, NF-kB, MAPK14

ICIA2018\Immunology of Rheumatic Diseases\Poster Discussion\7572.doc
7572

Analysis of PD-1 and Tim-3 Expression on CD4⁺ T Cells of Patients with Rheumatoid Arthritis; Association to Disease Severity

Zohreh Koohini¹, Hadi Hossein Nataj¹, Maryam Mobini², Aref Hosseinian Amiri², Alireza Rafiei¹, Hossein Asgarian Omran^{1,3}

1. Department of Immunology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

2. Department of Rheumatology, Imam Khomeini Hospital, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran.

3. Immunogenetics Research Center, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

Background: T-cell immunoglobulin and mucin-domain containing-3 (Tim-3) and Programmed cell death-1 (PD-1) are inhibitory receptors which involved in various immunological functions like regulating Th1 and Th17 activity, induction of tolerance and reducing production of inflammatory cytokines. In this study, the expression of Tim-3 and PD-1 were evaluated on CD4⁺ T cells of patients with Rheumatoid Arthritis (RA) and their association with disease severity was also addressed.

Methods: A total of 37 RA patients and 31 sex and age-matched healthy controls were included in this study. Disease severity of RA patients was determined by Disease Activity Score of 28 joints scoring system (DAS28). A three-color flow cytometry method was applied to determine the frequency of Tim-3⁺/PD-1⁺/CD4⁺ T cells. To measure the cytokine production, PBMCs were stimulated with PMA/ionomycin. Concentrations of IL-17, IL-10, IFN- γ , and TNF- α were measured in culture supernatants by ELISA.

Results: The frequency of PD-1⁺/CD4⁺ and Tim-3⁺/PD-1⁺/CD4⁺ T cells were significantly higher in patients with RA compared to controls ($p=0.0013$ and $p=0.050$, respectively). The percentage of Tim-3⁺/CD4⁺ T cells was similar in patients and controls ($p=0.4498$). The RA patients have produced significant higher levels of TNF- α , IL-17, and IFN- γ than those of healthy controls ($p=0.0121$, $p=0.0417$ and $p=0.0478$, respectively). Interestingly, an inverse correlation was found between the frequency of Tim-3⁺/CD4⁺ cells and disease severity of RA patients ($r=-0.4696$, $p=0.0493$). Similarly, the percentage of Tim-3⁺/PD-1⁺/CD4⁺ T cells was also revealed an inverse correlation with DAS28 score ($r=-0.5268$, $p=0.0493$). Moreover, significant positive correlations were detected between the concentrations of TNF- α ($r=0.6418$, $p=0.0023$) and IL-17 ($r=0.4683$, $p=0.0373$) with disease activity of RA patients.

Conclusions: Our results indicate that Tim-3 and PD-1 are involved in immune dys-regulation mechanisms of rheumatoid arthritis and could be considered as useful biomarkers for determination of disease severity and progression.

Keywords: Rheumatoid Arthritis, Tim-3, PD-1, Disease Activity Score

ICIA2018\Immunology of Rheumatic Diseases\Poster Discussion\7783

7783

***Lactobacillus delbrueckii* and *Lactobacillus rhamnosus* encompass a Tolerogenic Effect on Pristane Induced Lupus Murine Model**

Sahar Khorasani^{1,2}, Maryam Rastin¹, Mahmoud Mahmoudi¹, Fahime Lavi Arab¹, Fatemeh Mardani¹, Nafiseh Tabasi¹, Alireza Esmaceli¹, Mahdiah Khazaei¹

1. Immunology Research Center, BuAli Research Institute, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

2. Faculty of Ferdows Paramdical Sciences, Birjand University of Medical Sciences, Birjand, Iran

Background: It was shown that some *Lactobacillus* probiotics may induce regulatory T cells and so have beneficial role in SLE remission. The aim of this study was to evaluate the tolerogenic effect of *Lactobacillus delbrueckii* and *Lactobacillus rhamnosus* on pristane induced lupus murine model in both prophylactic and therapeutic manner.

Methods: Fifty-four female BALB/c mice were randomly divided into 9 groups. After SLE induction by pristane, four groups were treated from day 0 and 4 groups treated 2 months later. The treatments were as follows: *Lactobacillus rhamnosus*, *Lactobacillus delbrueckii*, mix of both probiotics and prednisolone. In addition, one group was also considered as an induction control group without any treatment. Finally we evaluated the presence of anti-nuclear antibody (indirect immunofluorescence), anti-double strand DNA and ribonucleoprotein (ELISA), renal function (proteinuria, serum level of creatinine and urea and also histological staining of kidney mice), the expression of FoxP3, IL-10, TGF- β and IL-6 (real time) and Treg number (flow cytometry).

Results: Our study demonstrated the probiotics therapy lead to decrease ANA, anti-dsDNA, anti-RNP, kidney dysfunction, IL-6 and increase IL-10, TGF- β , FoxP3 and also Tregs compared to induce lupus control group. In addition, the prednisolone receiving groups show decrease IL-6 and increase TGF- β more prominent than the probiotic ones.

Conclusion: The results indicate the role of *Lactobacillus rhamnosus* and *Lactobacillus delbrueckii* in the enhancement of Tregs number and decrease of inflammatory cytokines and disease severity. Also it seems the tolerogenic effect of probiotics (by increase Tregs, FoxP3 and IL-10) was more prominent than prednisolone receiving once, whereas the prednisolone has more anti-inflammatory effect (by decrease IL-6 and maybe Th17) compared to the *Lactobacillus*. Concerning the results, this study suggests managing SLE patients with a lower dose of prednisolone with the *Lactobacillus* supplementation; further studies will confirm the results.

Keywords: *Lactobacillus rhamnosus*, *Lactobacillus delbrueckii*, Lupus erythematosus, Pristane, Regulatory T cell

ICIA2018\Immunology of Rheumatic Diseases\Poster Discussion\7842

7842

Foxp3 Gene Polymorphism in Rheumatoid Arthritis Patients Associated with Tregs Frequency in Iranian Population

Amir Salek Farrokhi¹, Vida Hashemi², Rana Ezzeddini³, Mohammad Taher Tahoori⁴

1. *Department of Immunology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran*

2. *Department of basic science, Faculty of Medicine, Maragheh University of Medical Sciences, Maragheh, Iran*

3. *Department of Clinical biochemistry, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran*

4. *Department of Immunology, Faculty of Medicine, Shahid Sadoughi University of Medical Sciences, Yazd, Iran*

Background: Rheumatoid arthritis (RA) is characterized by infiltration of T lymphocytes. Regulatory T cells (Tregs) and Forkhead box P3 (Foxp3) have a crucial role in preventing autoimmunity and undesirable T cell responses. The aim of this study was to analyze the association of Foxp3 gene polymorphism and Tregs frequency in Iranian RA patients.

Methods: RA patients (n=240) and control subjects (n=240) were selected from Iran in order to carry out the study. Genomic DNA was genotyped for -3279 C/A Foxp3 gene SNP using the PCR-RFLP. The frequency of Tregs and serum level of IL-10, TGF- β , anti-CCP and RF were determined by flow cytometry and ELISA methods, respectively.

Results: The results showed a significant association of Foxp3 -3279 A allele with augmented risk of RA in Iranian patients compared to wild-type allele. The frequency of CA and AA genotypes were significantly higher in patients. RA patients with AA genotype have a significant reduction in frequency of Tregs compared to patients with CC and CA genotypes. TGF- β and IL-10 significantly diminished in patients with AA genotype compared to patients with CA and CC genotypes.

Conclusion: Our findings indicated that the Foxp3 polymorphism was associated with Tregs frequency and susceptibility to RA in the Iranian populations.

Keywords: Rheumatoid Arthritis, Foxp3, Regulatory T Cells, Polymorphism

ICIA2018\Immunology of Rheumatic Diseases\Poster Discussion\8246

8246

Increased Autophagy in Peripheral Blood Mononuclear Cells of Rheumatoid Arthritis Patients

Bahareh Kardideh¹, Mahdi Taghadosi¹, Zahra Samimi², Hamid Reza Mohammadimotlagh², Mona Sadeghalvad³

1. Department of immunology, School of medicine, Kermanshah University of Medical sciences, Kermanshah, Iran

2. Student research committee, School of medicine, Kermanshah University of Medical sciences, Kermanshah, Iran

3. Student research committee, School of medicine, Tehran University of Medical sciences, Tehran, Iran

Background: Rheumatoid arthritis (RA) is a chronic inflammatory autoimmune disease that primarily affects the joints. During RA, T cells and other immune cells recruit to the synovial tissue and promote RA. Autophagy is a process in which intracellular organelles and compounds are degraded and their products are reused by cells. Autophagy as a regulator of cell homeostasis can affect immune cells activation and contribute in RA pathogenesis. The aim of this study was to evaluate the autophagy-related genes (Atgs) expression in two groups of RA patients and healthy persons.

Methods: Peripheral Blood was obtained from three groups of donors including 20 patients with early RA, 20 under treatment RA patients (with Methotrexate, Hydroxychloroquine and Prednisolone therapy) and 20 age- and sex-matched healthy persons. The expression of three autophagy related genes including Lc3b (microtubule associated Light chain 3b), Beclin-1 and Atg5 was investigated by Real time PCR technique.

Results: The expression of Atgs was significantly increased in patients with early RA compared to healthy persons. The expression of these genes in under treatment patients was significantly lower than early RA patients.

Conclusion: This study showed that in early RA patients, the increased expression of Atgs can promote RA pathogenesis. In the other Words our findings suggest that the decreased autophagy can reduce RA severity.

Keywords: Autophagy, Rheumatoid Arthritis, Autoimmune Disease, Autophagy related genes

ICIA2018\Immunology of Rheumatic Diseases\Poster Discussion\8281
8281

The Effect of Disease Modifying Anti-rheumatic Drugs (DMARDs) in Combination with Prednisolone on TNF- α Plasma Levels and Disease Activity in Rheumatoid Arthritis Patients

Zahra Samimi¹, Bahareh Kardideh², Cyrus Jalili², Mahdi Taghadosi²

1. Student research committee, School of medicine, Kermanshah University of Medical sciences, Kermanshah, Iran
2. Department of immunology, School of medicine, Kermanshah University of Medical sciences, Kermanshah, Iran

Background: Rheumatoid Arthritis (RA) is an inflammatory autoimmune disease that leads to destruction of joints and adjacent tissues. Several inflammatory cytokines such as tumor necrosis factor- α (TNF- α) have critical roles in RA pathogenesis. Disease Modifying Anti-Rheumatic Drugs (DMARDs) are the mainstay of therapy in the treatment of RA. The aim of this study was to assess the plasma levels of TNF- α between three groups untreated, under treatment patients and healthy subjects and also evaluate the effect of DMARDs in combination with Prednisolone on the plasma levels of TNF- α and severity of RA disease.

Methods: Peripheral Blood was obtained from three groups including, 30 untreated, 30 under treatment patients and 30 age- and sex-matched healthy subjects. Under treatment patients were received combinational DMARDs therapy, including Methotrexate, Hydroxychloroquine plus Prednisolone. The plasma levels of TNF- α in three groups were measured by ELISA technique. The disease severity was assessed by disease activity score (DAS-28) and patients based on DAS-28 were categorized into four different groups, including remission, low, moderate and high disease activity.

Results: The plasma levels of TNF- α had no significant difference between three groups ($p>0.05$) but the DAS-28 was significantly higher in the untreated group in comparison with under treatment patients ($p<0.001$). DAS-28 was significantly correlated with TNF- α level in untreated patients ($p=0.001$, $r_s=0.594$) but we could not find a significant correlation between DAS-28 and TNF- α in under treatment group ($p=0.12$, $r_s=-0.29$).

Conclusion: It can be concluded that the plasma levels of TNF- α could be a good predictor of disease activity in newly diagnosed untreated RA patients, but quantification of its levels in plasma is not a good biomarker for diagnosis of RA. Also we found that DMARDs combination and prednisolone therapy do not have effect on the plasma levels of TNF- α .

Keywords: Rheumatoid Arthritis, DMARD, Disease activity score, TNF- α

ICIA2018\Immunology of Rheumatic Diseases\Poster Discussion\8306.doc

8306

Synbiotic Supplementation and the Effects on Clinical and Metabolic Responses in Patients with Rheumatoid Arthritis: a Randomized, Double-blind, Placebo-controlled Trial

Batol Zamani¹, Shima Farshbaf², Hamid R. Golkar¹, Fereshteh Bahmani², Zatollah Asemi²

1. Department of Gastroenterology, Kashan University of Medical Sciences, Kashan, Iran

2. Research Center for Biochemistry and Nutrition in Metabolic Diseases, Kashan University of Medical Sciences, Kashan, Iran

Background: Synbiotic intake may be associated with reduced inflammation in patients with rheumatoid arthritis (RA) due to optimized inflammatory markers, oxidative stress and insulin resistance. This research was conducted to assess the effects of synbiotic supplementation on the clinical and metabolic parameters of patients with RA.

Methods : A total of fifty-four patients with RA were allocated into two groups to receive either a synbiotic capsule (n 27) or a placebo (n 27) for 8 weeks in this randomised, double-blind, placebo-controlled trial. Fasting blood samples were taken at baseline and week 8 of the study to quantify related markers.

Result: After the 8-week intervention, compared with the placebo, synbiotic supplementation resulted in a significant reduction in serum high-sensitivity C-reactive protein (hs-CRP) levels (-1427.8 (SD 3267.2) v. $+2833.4$ (SD 5639.7) ng/ml, $P=0.001$). In addition, compared with the placebo, synbiotic supplementation improved disease activity score-28 joints (DAS-28) (-1.6 (SD 0.8) v. -0.3 (SD 0.5), $P<0.001$) and visual analogue scales (VAS) pain (-30.4 (SD 18.7) v. -11.5 (SD 15.9), $P<0.001$). In addition, a significant elevation in plasma nitric oxide (NO) ($+0.8$ (SD 4.4) v. -2.6 (SD 4.5) $\mu\text{mol/l}$, $P=0.008$), and significant reductions in insulin values (-13.8 (SD 26.4) v. $+4.2$ (SD 28.2) pmol/l, $P=0.01$), homoeostasis model of assessment-estimated insulin resistance (HOMA-IR) (-0.5 (SD 1.0) v. $+0.1$ (SD 1.1), $P=0.03$) and homoeostatic model assessment- β -cell function (HOMA-B) (-9.4 (SD 17.9) v. $+3.3$ (SD 18.9), $P=0.01$) following supplementation with the synbiotic compared with the placebo. Compared with the placebo, synbiotic supplementation also resulted in a significant increase in plasma GSH ($+36.6$ (SD 63.5) v. -58.5 (SD 154.4) $\mu\text{mol/l}$, $P=0.005$).

Conclusion: Overall, our study demonstrated that synbiotic supplementation for 8 weeks among patients with RA had beneficial effects on hs-CRP, DAS-28, VAS, NO, insulin levels, HOMA-IR, HOMA-B and GSH levels.

Keywords: Synbiotics, Supplementation, Rheumatoid arthritis, Metabolic profiles

ICIA2018\Immunology of Rheumatic Diseases\Poster Discussion\8507
8507

The Effect of Ginger Powder on IL2, TNF α and IL1 β Cytokines Gene Expression Levels in Patients with Active Rheumatoid Arthritis

Aryaeian Naheed^{1*}, Mahmoudi Mahdi², Shahram Farhad², Poursani Shiva², Jamshidi Fatemeh², Tavakoli Hajar³

1. Department of Nutrition, School of Public Health, Iran University of Medical Sciences, Tehran, Iran.

2. Rheumatology Research Center, Tehran University of Medical Sciences, Tehran, Iran.

3. Departments of Cellular and Molecular Nutrition, School of Nutritional Sciences and Dietetics, Tehran University of Medical Sciences, Tehran, Iran.

Background: Rheumatoid arthritis (RA) is a chronic inflammatory disease that affects the joints and consequently leads to the destruction of cartilage and bone lesions. Ginger is a native plant to India, which traditionally has been used in treatment of osteoarthritis, joint and muscle pain, neurological diseases, and inflammation of gums, tooth pain, asthma, stroke, diabetes, and constipation. This study aimed to determine the effect of ginger on some immunological and inflammatory factors in patients with rheumatoid arthritis.

Methods: In this study, 66 patients with active rheumatoid arthritis were enrolled. Patients were randomly divided into two groups that were treated using ginger and placebo (wheat flour) respectively. To determine the effect of confounding factors on the findings of the study, questionnaires for nutrient intake, physical activity, medication, and determination of body mass index were filled for each participant, at the beginning and end of the study. Serum CRP and mRNA levels of IL-1 β , IL-2, and TNF- α were measured by ELISA and Quantitative Real-Time PCR respectively.

Results: Results of the study showed ginger supplementation caused a significant decrease in CRP ($P=0.05$), and IL-1 β ($P=0.02$). TNF α mRNA level in ginger group compared to placebo group decreased, although the difference was not significant between the two groups ($P=0.09$). Ginger had no effects on IL2 expression.

Conclusions: The study showed that ginger reduces inflammatory markers, CRP, and IL-1 β in patients with active RA and it seems that ginger can improve the inflammation in the patients.

Keywords: Ginger, Active rheumatoid arthritis, Inflammatory markers.

ICIA2018\Immunology of Rheumatic Diseases\Poster Discussion\8617
8617

Molecular analysis of ERAP1 Allelic Variations in Patients with Ankylosing Spondylitis

Farhad Babaie^{1,2}, Hamed Mohammadi^{1, 2}, Sorayya Salimi^{1, 2}, Alireza Rezaeiemanesh³, Behzad Baradaran^{1, 2}, Zohreh Babaloo^{1, 2}

1. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

2. Department of Immunology, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

3. Department of Immunology, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

Background: Ankylosing spondylitis (AS) is a group chronic inflammatory arthritis termed seronegative spondyloarthropathies. It typically affects the joints of the spinal and axial skeleton and exhibits typical clinical features and genetic factors such as HLA-B27 and ERAP1. Among the non-HLA loci, the strongest association was observed for the ERAP1 of single nucleotide polymorphisms (SNPs).

Methods: In our study, we have determined the frequencies of ERAP1 allelic variants and genotypes for three non-synonymous SNPs in AS patients and healthy individuals from the Iranian population. Both the AS patients and healthy group consist of HLA-B27 positive and HLA-B27 Negative individuals. We implemented the SSP-PCR system for genotyping of 160 AS patients and 160 healthy controls from the Iranian population.

Results: Considerable differences in allele's frequencies within patients vs control cohort were shown for 3 SNPs including rs30187, rs2287987, and rs10050860 under investigation. 3 SNPs were associated with the risk of AS [odds ratio (OR) 0.778, 95% CI 0.516–0.968, P =0.037 for rs30187, OR 1.57, 95% CI 1.07–2.33, P =0.025 for 10050860 and OR 1.60, 95% CI 1.10–2.32, P =0.015 for rs2287987].

Conclusion: The ERAP1 gene polymorphism might be a risk factor in the pathogenesis of AS. In contrast, ERAP1 gene polymorphisms may serve a protective role in AS.

Keywords: Ankylosing spondylitis, ERAP1, single nucleotide polymorphism



Immunomodulation and Immunoregulation

Poster Discussion

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018/Immunomodulation and Immunoregulation/Poster Discussion/8220

8220

Mesenchymal Stem Cell-Conditioned Medium Ameliorate inflammation in Mouse Model of Acute Colitis

Sedigheh Pouya¹, Maryam Heidari¹, Kaveh Baghaei², Hamid Asadzadeh Aghdaei², Afshin Moradi³, Saeed Namaki¹, Mohammad Reza Zali³, Seyed Mahmoud Hashemi¹

1. Department of Immunology, School of medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Basic and Molecular Epidemiology of Gastrointestinal Disorder Research Center, Research Institute for Gastroenterology and Liver Disease, Shahid Beheshti University of Medical Science, Tehran, Iran

3. Gastroenterology and Liver Disease Research Center, Research Institute for Gastroenterology and Liver Disease, Shahid Beheshti University of Medical Science, Tehran, Iran

Background: Inflammatory bowel disease (IBD) is a multifactorial disease that the imbalanced immune responses have an important role in development and progression of the disease. On the other hand, studies on mesenchymal stem cells (MSCs) and its immunomodulatory properties have shown that these cells can be effective in immune/inflammatory response regulation through their secretions. Therefore, this study was designed to investigate the effects of the mesenchymal stem cell-conditioned medium (MSC-CM) injection in mouse model of acute colitis.

Methods: Acute colitis was induced in C57BL/6 mice with 2.5 % dextran sulfate sodium (DSS) dissolved in drinking water. The disease was induced in 1 cycle (4 days use of water containing DSS, followed by 6 days of water). The MSC-CM injection was performed 3 times (500 µl/mouse). During the study, changes in body weight, bleeding, stool consistency, disease activity index (DAI), and mortality rate were recorded. After euthanizing the mice, weight, and length of the colon, the percentage of Treg cells, the levels of TGF-β, IL-17, and IL-10 were measured. The pathology examination of the colon was also performed.

Results: According to the results, MSC-CM injection inhibited body weight loss, bleeding, DAI, and mortality rates. The stool consistency improved. The percentage of Treg cells and the TGF-β production was increased while the level of IL-17 decreased. In pathological observations, it was found that after injection of MSC-CM, infiltration of inflammatory cells and epithelial destruction decreased.

Conclusion: The results showed that MSC-CM has the ability to modulate the immune response and decrease inflammation. Then it can be used as a cell-free therapy in the treatment of inflammatory bowel disease.

Keywords Inflammatory bowel disease, Mesenchymal stem cells, Conditioned Medium, Dextran Sulfate Sodium

ICIA2018\Immunomodulation and Immunoregulation\Poster Discussion\8278
8278

MicroRNA-150 Targets PU.1 and Regulates Macrophage Differentiation and Function in Experimental Autoimmune Encephalomyelitis

Leila Shakerian^{1,2}, Samira Ghorbani¹, Farideh Talebi¹, Farshid Noorbakhsh^{1*}

1. Department of Immunology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

2. Shefa Neuroscience Research Center, Khatam Al-Anbia Hospital, Tehran, Iran

Background: MicroRNAs are regulatory RNA species which are involved in various physiological and pathological processes, including central nervous system (CNS) diseases. Multiple sclerosis (MS) is an inflammatory demyelinating disease which is characterized by leukocyte infiltration and monocytoid cell activation, followed by demyelination. PU.1 is a key transcription factor which is involved in differentiation and activation of monocytoid cells. Herein, we investigated the expression of PU.1 transcription factor and its potential regulation by miRNAs in the CNS of mice affected with experimental autoimmune encephalitis (EAE) as well as in vitro cultured primary macrophages. Physical interactions between PU.1 and miRNAs as well as the effect of miRNAs on macrophage inflammatory response and polarization were also investigated.

Methods: Three miRNAs, i.e. miR-18a, miR-150 and miR-155, which could potentially target PU.1, were selected based on bioinformatic criteria. Gene and microRNAs expression were first analyzed in the CNS of EAE mice at different time points after disease induction as well as in primary macrophages using real-time RT-PCR. MiRNA species which showed a negative correlation with the target gene both in vivo and in vitro were selected for further analyses. as Transfection experiments in macrophages as well as luciferase assays using vectors expressing luciferase coding sequence with the 3'UTR of PU1 were performed to investigate the physical interaction of miRNA sequences with PU.1 transcripts. Following the confirmation of interaction, the effects of miRNA on macrophage response and polarization were evaluated.

Results: Our findings showed a significant increase in PU.1, miR-155 and miR-18a at both acute and chronic phases of EAE disease. While showing an increase at the acute phase, miRNA-150 expression was significantly reduced at the chronic phase of disease. Expression analyses on activated macrophages revealed a substantial induction of PU.1 and miR-155 following LPS stimulation. On the contrary, miR-150 showed a remarkable decrease following treatment with LPS. Luciferase assays as well as transfection experiments confirmed that PU.1 transcripts were targeted by miR-150. Overexpression of miRNA-150 modulated macrophage inflammatory response and shifted polarization toward M2 phenotype.

Conclusion: These findings point to the potential role of miRNA-150 in regulating inflammatory responses in CNS through targeting PU.1. Our results raise the possibility that miR-150-PU1 interaction might be a likely target for therapeutic interventions in autoimmune neuroinflammation.

Keywords: microRNA, miR-150, PU1, macrophage, Experimental autoimmune encephalomyelitis

ICIA2018/Immunomodulation and Immunoregulation/Poster Discussion/8279

8279

miR-101a Modulates Lymphocyte Activation in Experimental Autoimmune Encephalomyelitis through targeting c-fos

Samira Ghorbani¹, Neda Heidari¹, Farideh Talebi¹, Farshid Noorbakhsh^{1*}

1. Department of Immunology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

Background: Multiple sclerosis (MS) is a chronic autoimmune disease of the central nervous system. MicroRNAs are small, noncoding RNA molecules which function via base-pairing with complementary sequences within mRNA molecules, usually resulting in gene silencing. Herein, we analyzed the expression of Fos family T cell transcription factors and the miRNAs which are known to target them in the spinal cord of mice affected by EAE, an animal model of MS.

Methods: The expression of Fos family members(c-fos, Fosb, Fosl1, Fosl2) and their potentially targeting miRNAs were measured in the spinal cords of EAE as well as activated lymphocytes by real time RT-PCR. To examine the direct interaction between miRNA and target gene, luciferase reporter assay system was used.

Results: Our data showed a significant increase of c-fos and Fosl2 levels in EAE mice at the acute and chronic phases of disease. Although miRNAs-targeting c-fos (miR-7b, miR-101a, miR-101b) were down-regulated in chronic phase of EAE, only down regulation of miR-101a was statistically significant. Our data also revealed a rapid upregulation of c-fos in lymphocytes at 5 minutes post activation, whereas increased levels of other members of Fos family were not observed. Activated T cells showed early reduced levels of miR-7b, miR-101a and miR-101b 5 minutes after stimulation. Cells transfected with a c-fos-3'UTR containing plasmid showed a significant reduction in the luciferase activity following transfection with miR-101a, indicating the interaction between miRNA and the 3'UTR region.

Conclusions: Our findings indicate that miR-101a might be involved in regulating c-fos expression level during EAE and lymphocyte activation, hence, leading to development of new therapeutic approaches for T cell mediated autoimmune disorders.

Keywords: MicroRNA, EAE, c-fos

ICIA2018\Immunomodulation and Immunoregulation\Poster Discussion\8310.doc

8310

The Role of P21 in Different Molecules Ubiquitination n TCR Signaling PathwayRahman Shokri^{1,2}, Dimitrios Balomenos¹1. *Department of Immunology and Oncology, University of Autonoma de Madrid*2. *Current address: Pasteur Institute of Iran-Karaj*

Background: p21 can act as a cell cycle inhibitor, cellular senescence mediator or control DNA damage repair. Our latest data with p21^{-/-} T cells revealed a novel p21 function as a negative regulator of the activation machinery in repeatedly stimulated T cells. Protein ubiquitination is an important post-translational process that can affect protein targets in different ways. In this study, we checked the role of p21 on different protein ubiquitination like p50 (data not shown) and TRAF6 in T cell receptor (TCR) signaling pathway.

Methods: T cell cultures were incubated with proteasome inhibitor MG132 before harvest, washed in PBS/N-ethylmaleimide (N-EM), lysed in SDS buffer by boiling (5 min), and diluted (1/10) in RIPA buffer (without SDS) supplemented with 20 μ M N-EM and protease/phosphatase inhibitor cocktail. Equal amounts of protein were incubated with appropriate antibody. After SDS-PAGE, proteins were transferred to a nitrocellulose membrane and immunoblotted with an anti-Ub antibody (P4D1; Santa Cruz Biotechnology) to detect ubiquitination.

Results: TRAF6 ubiquitination was reduced in p21-deficient compared with wild type CD4⁺ T cells at 30 and 60 minutes after second stimulation, which shows p21 induces TRAF6 ubiquitination.

Conclusion: TRAF6 is recruited to the T cell immunological synapse and interacted with LAT to activate T cells. Also MALT1 promotes TRAF6 auto-ubiquitination, which then recruits the TAK1/TAB2 adaptor protein complex to activate IKK. Therefore, TRAF6 is an important molecule in TCR signaling pathway and it's ubiquitination by p21 leads to T cell activation and regulation. Overall, p21 is an important factor that negatively regulates CD4⁺ T cells activation after second stimulation by different process like ubiquitination.

Keywords: TCR, T CD4⁺, Activation, Ubiquitination, TRAF6, p21

ICIA2018/Immunomodulation and Immunoregulation/Poster Discussion/8459

8459

Evaluation of the Immunoregulatory and Proliferative Capacities of Adipose Derived Mesenchymal Stem Cells (Ad-MSCs) following pre-treatment with Human Menopausal Gonadotropin (hMG)

Shadi Kesmati¹, Moein Farshchian², Sepideh Sadat Hosseini¹, Halimeh Hassanzadeh², Majid Momeni Moghaddam¹, Fatemeh Sadeghifar^{1*}, Maryam Moghaddam Matin^{2,3}, Hamid Reza Bidkhori^{2*}

1. Department of Biology, Hakim Sabzevari University, Sabzevar, Iran

2. Stem Cells and Regenerative Medicine Research Department, Academic Center for Education, Culture and Research (ACECR)-Khorasan Razavi, Mashhad, Iran

3. Department of Biology, Faculty of Sciences, Ferdowsi University of Mashhad, Mashhad, Iran

Background: The hypothesis behind the clinical application of Mesenchymal Stem/Stromal Cells (MSCs) is supported on two characteristics of them: Their proliferative capability based on their multipotential state and MSCs' trophic activity via their anti-inflammatory and immunomodulatory effect. Prolonged culture of MSCs *in vitro* reduces their proliferative ability and changes them to into mature phenotypes. Pre-treatment of the cells with some chemical agents during *in vitro* expansion before transplantation is one strategy to overcome this limitation by enhancing both aforementioned capacities. Human menopausal gonadotropin (hMG) is a natural medication used to induce ovulation in women with infertility problems. hMG consists of both LH and FSH, hormones needed to help trigger ovulation. The aim of this study is to determine whether pre-treatment of Ad-MSCs with hMG, will enhance the expression of some genes involved in proliferation and immunomodulation (c-Myc, Oas2 and TGF- β).

Methods: MSCs were harvested from human processed lipoaspirate. Adipose-derived MSCs were treated at passage 2 for targeted gene expression profiling. MSCs were pre-treated with 2, 4 and 8 IU hMG for 24h, 48h and 72h. QPCR experiments were performed by SYBR Green method in CFX-96 Bio-Rad Real-Time PCR. c-Myc, Oas2 and TGF- β were compared at mRNA level in both groups.

Results: Results show hMG significantly up-regulated the c-Myc and Oas2 gene expression compare with the control group while it decreased TGF- β expression significantly.

Conclusion: Pre-treatment of Ad-MSCs with hMG may allow the *in vitro* cultured cells to regain their proliferation and immunoregulatory capacities.

Keywords: Adipose-derived mesenchymal stem cells, pretreatment, hMG, c-Myc, Oas2, TGF- β , Immunomodulation.

ICIA2018\Immunomodulation and Immunoregulation\Poster Discussion\8717

8717

The Role of Regulatory T cells in the Failure of Kidney Function in Patients with Autoimmune and Non-autoimmune Disorders

Farshid Fathi¹, Hossien Motedayyen², Nilofar Keyhanmehr¹, Nahid Ekandari¹, Noushin Lotfi¹, Shohreh Mohammadi³, Hossien Hafezi⁴, Mehdi Raie⁵

1. Department of Immunology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

2. Department of Laboratory Medicine, Kashan University of Medical Sciences, Kashan, Iran

3. Khorshid Hospital, Isfahan University of Medical Sciences, Isfahan, Iran

4. Department of Dermatology, Isfahan University of Medical Sciences, Isfahan, Iran

5. Health Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran

Background: Autoimmune response is one of the causes of the failure in the renal function in patients with end-stage renal disease. Regulatory T cells (Tregs) are a subset of CD4⁺T cells that play an indispensable role in regulating immune responses and preventing the development of autoimmune diseases. Therefore, the aim of this study was to investigate the percentage of Tregs in kidney patients who need to renal transplantation due to autoimmune responses and other causes.

Methods: Heparinized whole blood (5 ml) was obtained from patients with end-stage renal diseases and healthy subjects. Peripheral blood mononuclear cells (PBMCs) were isolated using Ficoll density centrifugation. PBMCs were stained with FITC anti-human CD4, PE anti-human CD25, and PE-cy5 anti-human Foxp3 antibodies. The percentage of the stained cells was measured by a FACS Calibur flow cytometer and analyzed using the CellQuest software.

Results: Our results indicated that the percentage of Tregs in kidney patients with autoimmune disorders was significantly decreased compared to healthy individuals and patients with renal failures owing to non-autoimmune disorders ($p < 0.01-0.05$). However, there was no difference in Tregs percentage between kidney patients without autoimmune disorders and healthy individuals.

Conclusion: These findings suggest that the reduction of Tregs number in kidney patients can contribute to the risk of developing autoimmune responses that lead to the loss of kidney function.

Key words: Autoimmune disorders, Regulatory T cells, Kidney patients.



Immunoparasitology

Poster Discussion

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Immunoparasitology\Poster Discussion\7935

7935

Lack of IL-17a May Be Involved in Healing of the Lesions in Patients Afflicted by Cutaneous LeishmaniasisMohammad Moafi¹, Hossien Rezvan², Roya Sherkat³, Roya Taleban⁴1. *Acquired Immunodeficiency Research Center, Isfahan University of Medical Sciences, Isfahan, Iran*2. *Department of Pathobiology, Faculty of Veterinary Science, Bu-Ali Sina University, Hamedan, Iran*3. *Acquired Immunodeficiency Research Center, Isfahan University of Medical Sciences, Isfahan, Iran*4. *Acquired Immunodeficiency Research Center, Isfahan University of Medical Sciences, Isfahan, Iran*

Background: Cutaneous Leishmaniasis (CL) is assumed as neglected tropical disease, which spontaneously heals less than one year. However, the lesions of CL rarely persist more than one year. The persistency of the lesions could be attributed to different factors, including pro-inflammatory cytokines. In this context, the role of IL-4 and IFN-gamma in duration of lesion healing has been evaluated in some of the previous studies; however the role of IL-17a in duration of healing lesion is not completely understood. The current study aims to assess the role of IL-17a in non-healing form of CL (NHCL).

Methods: This was a cross-sectional study implemented from 2015-1015 in immunodeficiency research center, Isfahan University of Medical Sciences (IUMS). This study included 10 cases afflicted by NHCL and 33 cases suffering from healing form of CL. The peripheral blood mononuclear cells (PBMCs) of the patients were isolated by Ficoll and were harvested in three different medium. The stimulators were as follow: purified protein derivative (PPD), Phytohemagglutinin (PHA), and Soluble *Leishmania* Antigen (SLA). The supernatant was isolated and the levels of IL-17a were determined using ELISA.

Results: The levels of IL-17a produced in PPD stimulated wells was significantly lower, compared with PHA and SLA stimulated wells ($P<0.05$). Furthermore, the level of IL-17a produced in PHA stimulated wells was significantly higher, compared with PHA stimulated wells ($P<0.05$). The level of IL-17a produced in NHCL cases was significantly lower than healing cases ($P<0.05$).

Conclusion: Our findings support the notion that IL-17a plays a crucial role in the duration of lesion healing in those afflicted by CL.

Keywords: Cutaneous Leishmaniasis, IL-17a

ICIA2018/Immunoparasitology/Poster Discussion/8011

8011

In vitro Efficacy of Rapamycin on Old World *Leishmania* Strains

Fatemeh khadir^{1,2}, Ahmad Oryan^{2*}, Tahereh Taheri¹, Maryam Heidari-kharaji^{1*}, Sima Rafati^{1*}

1. Immunotherapy and *Leishmania* Vaccine Research Dept., Pasteur Institute of Iran, Tehran, Iran

2. Department of Pathology, School of Veterinary Medicine, Shiraz University, Shiraz, Iran

Background: Leishmaniasis is a serious health problem in different countries. There is no human vaccine and the disease is generally controlled by chemotherapy, but most of the available drugs are highly toxic or result in drug resistance. Here we investigated Rapamycin as an anti-leishmanial drug due to its mTOR (mammalian target of Rapamycin) blockage activity that affects the parasite propagation. Introducing Food and Drug Administration (FDA)-approved Rapamycin for anti-leishmanial activity may open up a new approach in the treatment of this important disease.

Methods: In the present study, in vitro efficacy of Rapamycin was investigated against two species of *Leishmania* including *L. major* and *L. tropica*. In vitro potency was evaluated using MTT (3-(4,5-Dimethylthiazol-2-Yl)-2,5-Diphenyltetrazolium Bromide) method on promastigote and amastigote form of *Leishmania* parasites, scanning electron microscopy (SEM) and transmission electron microscopy (TEM) were done for observing the effect of the drug on the shape and function of the host cells and parasites.

Results: Rapamycin is significantly effective to inhibit the growth and survival of both *L. major* and *L. tropica*, throughout different analyses using scanning and transmission electron microscopy. There are some changes in the appearance and size of the promastigote form of both species. In addition, our result in in vitro showed the interaction of host cell and amastigote form of both species were changed dramatically in the presence of Rapamycin.

Conclusion: Based on in vitro analysis, Rapamycin seems to be a potent anti-leishmanial agent. Rapamycin as FDA approved drug may open a new window in order to control cutaneous leishmaniasis as sole or combinational chemotherapy approach against *L. tropica* and *L. major* infection.

Keywords: *Leishmania major*, *Leishmania tropica*, Rapamycin, TEM, SEM.

ICIA2018\Immunoparasitology\Poster Discussion\8172

8172

A Simple Method for Evaluation of *Leishmania* Parasite Burden among Experimental ModelsMahdieh Azizi¹, Sepideh Haghdooost¹, Mostafa Haji Molla Hoseini¹, Farshid Yeganeh¹*1. Department of Medical Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran*

Background: The model of *Leishmania* infection in mice has been widely used in various studies. A limitation dilution assay (LDA) is commonly used method for evaluating the efficacy of therapeutic methods and/or immunization effectiveness. This method suffers from the risk of contamination, loss of samples, and time consuming. Furthermore, utilizing from it requires having familiarity and access to special results analysis software. The purpose of present study was to introduce a simple and alternative method for measuring the burden of *Leishmania* among the experimental model of this infection.

Methods: At the first *Leishmania Major* that express Green Fluorescent Protein was produced (GFP- *Leishmania*). Then the Stationary phase of promastigotes (The standard L. major strain of MRHO/IR/75/ER) were used for infection of ten female BALB/c mice. Finally, mice with different wounds diameter were selected to compare the parasite burden using LDA method (described elsewhere) and our fluorescence microscopic technique. Briefly, in our technique, we count florescent cell in the same area of the microscopic field following by adjusting cell count.

Results: At least 3.14 (log of parasite/Lymph node) and the maximum of 7.85 (log of parasite/Lymph node) estimated by LDA method. In fluorescence microscope method, we calculate 0.03-33.3 infected cells per lymph node. Correlation analysis was performed between two methods and the correlation coefficient of 0.96 was obtained between the two methods ($p < 0.0001$).

Conclusion: Our finding showed GFP- *Leishmania* is an appropriate tool for parasite burden estimation using our fluorescent microscopic technique. In addition, expressing GFP makes it possible to continuously evaluate infected mice by in-vivo imaging method during the study.

Keywords: GFP-*Leishmania*, Parasite burden, Fluorescent microscopy

ICIA2018\Immunoparasitology\Poster Discussion\8249

8249

***Leishmania tropica*: Infectious Dose Affects Pathogenicity and Immunogenicity in a Mouse Model**

Mosayeb Rostamian¹, Maryam Abolghazi², Davood Jafari³, Hadiseh Farahani², Fatemeh Doroudian¹,
Hamid M. Niknam^{1*}

1. Immunology Department, Pasteur Institute of Iran, Tehran 13164, Iran

2. Department of Microbiology, Qom Branch of Islamic Azad University, Qom, Iran

3. Islamic Azad University of Borujerd, Science and Research Unit, Borujerd, Lorestan, Iran

Background: *Leishmania (L.) tropica* is a causative agent of cutaneous leishmaniasis in humans. An experimental model is needed for this species. The effect of parasite dose on *L. tropica* infection was studied in the current paper. High and low doses of *L. tropica* were used for ear infection of naïve BALB/c mice and lesion development, parasite load, and cytokine response were assessed. *L. major* infection was used for comparison.

Methods: Mice were challenged in footpad by a fixed high dose of *L. tropica*. Infection with high dose *L. tropica* in contrast to low dose resulted in dissemination of parasite to spleen.

Results: Primary infection with high dose *L. tropica* in comparison to low dose resulted in lower lesion diameter and lower parasite load of lymph nodes after secondary *L. tropica* challenge. *L. tropica* in comparison to *L. major* results in lower lesion diameter, more growth in lymph nodes at early phases of infection, dissemination to spleen, and lower levels of interleukin-10.

Conclusion: Our findings show only the high dose of *L. tropica* results in visceralization of parasite and protection at the site of infection after secondary challenge of *L. tropica*. Therefore dose of infection is an important factor in experimental model of *L. tropica*.

Keywords: Leishmaniasis, Infectious dose, *Leishmania tropica*, *Leishmania major*, Spleen, Cytokine, BALB/c mice.

ICIA2018\Immunoparasitology\Poster Discussion\8256

8256

Haematological and Serum Biochemical Parameters in Horses with Natural Babesiosis in Urmia, Northwest of IranNeda Ahmadi afshar¹, Farnaz Malekifard¹, Mousa Tavassoli¹

1. Department of Pathobiology, Faculty of Veterinary Medicine, Urmia University, Urmia, West Azarbaijan

Background: Equine piroplasmosis (EP) is a tick-borne disease caused by hemoparasites *Theileria equi* and *Babesia caballi*. This disease affects all members of the Equus genus (horses, donkeys and zebras. The aim of this study was to determine the effect of babesiosis on some hematological and biochemical parameters in horses infected with *T.equi* and *B.caballi* in Urmia suburb, West Azerbaijan province, Iran.

Methods: A total of 240 blood samples collected randomly from horses of 25 villages. The specimens were transferred to the laboratory and the blood smears stained with Geimsa. The parasitological diagnosis was confirmed using multiplex PCR analysis. As a control group, 20 clinically healthy horses reared under the same management and environmental conditions were also sampled.

Results: The results of the PCR assays showed 26(10.83%), 14(5.83%) and 4 (1.66%) were distinguished as *T.equi*, *B.caballi* and mixed infection, respectively. Compared to controls, BUN, creatinine, cholesterol, triglyceride, HDL and LDL level showed a significant increase ($P < 0.05$). Significant elevation ($P < 0.05$) of total leukocyte count, number of lymphocyte, neutrophil, monocyte and eosinophil were found in infected horses. While, hemoglobin concentration (Hb), packed cell volume (PCV), red blood cells (RBCs), mean corpuscular volume (MCV) and mean corpuscular hemoglobin concentration (MCHC) significantly decreased ($P < 0.05$) in infected horses.

Conclusion: Hematological and biochemical parameters changes suggesting that these indicators would indirectly promote the presence of these parasites in the horses.

Keywords: *Theileria equi*, *Babesia caballi*, hematological and biochemical parameters, Horse, Iran

ICIA2018\Immunoparasitology\Poster Discussion\8257

8257

Serologic and Microscopic Studies on Piroplasmosis in Donkeys (*Equus asinus*) in the Northwest of Iran

Neda Ahmadi afshar¹, Farnaz Malekifard¹, Mousa Tavassoli¹

1. Department of Pathobiology, Faculty of Veterinary Medicine, Urmia University, Urmia, West Azarbaijan

Background: Equine piroplasmosis is a tick-borne protozoal disease of horses, mules, donkeys and zebra; it causes economic losses to the equine industry. The aim of this study was to determine the seroprevalence of *T. equi* and *B. caballi* infection in donkeys of West Azerbaijan province in North West of Iran.

Methods: From April to September 2014, a total of 120 blood samples collected randomly from apparently healthy donkeys and examined by microscopy and indirect immunofluorescent antibody test.

Results: Microscopic observation on 120 blood smears determined 3(2.5%) and 1 (0.83%) samples were infected by *T. equi* and *B. caballi*, respectively. Antibodies against *T. equi* and *B. caballi* were found in 13(10.83%), and 5 (4.16%) serum samples, respectively.

Conclusion: We couldn't detect mix infection with both parasites. No significant difference was observed between the seroprevalence of piroplasm infection with risk factors such as age and gender in donkeys. This is the first report of detection of *T. equi* and *B. caballi* infection using IFAT in Iran. Our findings indicated that *T. equi* and *B. caballi* were prevalent among donkey population in Iran.

Keywords: *Babesia caballi*, donkey, Iran, Indirect Immunofluorescent antibody test, *Theileria equi*

ICIA2018\Immunoparasitology\Poster Discussion\8538
8538

Evaluation of NOHA, an Arginase Inhibitor, on *Leishmania tropica* on BALB/c Mice Using Reporter Genes

Shima Nahidi¹, Elham Gholami¹, Yasaman Taslimi¹, Negar Seyed¹, Sima Habibzadeh¹, Elahe Davarpanah^{1,2}, Behrooz Vaziri³, Sima Rafati¹, Tahereh Taheri^{*1}

1. Department of Immunotherapy and Leishmania Vaccine Research, Pasteur Institute of Iran, Tehran, Iran

2. Department of Genetics Tehran Medical Sciences Branch, Islamic Azad University, Tehran, Iran

3. Biotechnology Research Center, Pasteur Institute of Iran, Tehran, Iran

Background: Leishmaniasis is one of the five most important parasitic diseases worldwide. Among the *Leishmania* (*L.*) species, causative agent of this disease, *L. tropica* is responsible for both cutaneous and visceral leishmaniasis transmitted from human to human by sand fly. Arginase (ARG) is essential for cell growth and parasitic pathogens is where converts L-arginine into ornithine and urea. We examined the inhibitory impact of ARG using N-hydroxy-nor-arginine (nor-NOHA) as a powerful ARG inhibitor on *L. tropica* infectivity in mice model.

Methods: Different groups of BALB/c mice were infected subcutaneously (in footpad) or intradermally (in ear) with *L. tropica*^{EGFP-LUC}. Two infected groups received nor-NOHA intraperitoneally two times a week within 6 months. Two infected groups remained untreated. One group was treated/uninfected as a control. All groups were compared in respect to parasite load, *in vivo* bioluminescence imaging, footpad swelling and ARG/nitric acid activity.

Results: Infection rate in all groups was monitored during 6 months. In infected ear, milder swelling and inflammation was observed which subsided gradually but, in infected footpad swelling was increased up to the end of experiment. In both injected sites, parasite load correlated with swelling size or redness. Also, *in vivo* bioluminescence monitoring of infected footpad mice was clearly observable increasing of parasite burden in infected footpad with *L. tropica*^{EGFP-LUC}. The comparison between different groups (untreated and treated) did not show any significant difference in footpad size. Also, there was observable not significant decreasing of parasite burden and ARG activity in treated and untreated groups).

Conclusion: Our results first demonstrated that *L. tropica*^{EGFP-LUC} could be used to follow parasitemia progression in infected footpads of mice. Second arginase inhibition partially controls *L. tropica* infection reflected in parasite burden and ARG activity decrement. Furthermore *in vivo* imaging is practically more applicable in the study of metabolic pathways in *L. tropica*.

Keywords: *Leishmania tropica*, Arginase, nor-NOHA, nitric oxide, *in vivo* imaging.

ICIA2018/Immunoparasitology/Poster Discussion/9748
9748

Achievement Amastigotes of *Leishmania infantum* and Investigation of Pathological Changes in the Tissues of Infected Golden Hamsters

Sajad Rashidi¹, Kurosh Kalantar², Gholamreza Hatam³

1. Department of Parasitology and Mycology, Shiraz University of Medical Sciences, Shiraz, Iran

2. Department of Immunology, Shiraz University of Medical Sciences, Shiraz, Iran

3. Basic Sciences in Infectious Diseases Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

Background: *Leishmania (L) infantum* is identified as a causative agent of visceral leishmaniasis (VL). Amastigote form is used for investigations on vaccines, treatment, and diagnosis. This study aims to achieve the *L. infantum* amastigotes in the tissue of infected golden hamsters and J774 macrophages and investigate the pathological changes which occur in the liver and spleen of the hamsters with VL.

Methods: Four male golden hamsters were infected with promastigote (intra-peritoneal (10^8) and intra-cordial (2×10^7)) of *L. infantum*. After 5 months, the hamsters were euthanized and touch and pathology smears were prepared from the livers and spleens. The tissues were homogenized and centrifuged at $100 \times g$. Supernatants were collected and centrifuged at $2000 \times g$ and the pellet was collected. Also, J774 macrophages were infected with *L. infantum* promastigotes (1 macrophage/10 promastigotes) and the infected macrophages were ruptured. Then, centrifuge stages were done the same as the previous part.

Results: The amastigotes were observed in the touch and pathology smears of both livers and spleens. The amastigote load in the livers was more than the spleens in the touch and pathology smears. Although the structure of the livers underwent the pathological changes, the spleens were unchanged. The macrophage infectivity ratio was 95%.

Conclusion: This study presents a simple and accessible way to achieve lots of pure and real *L. infantum* amastigotes. In addition, it seems that the pathological changes in the liver of the hamsters with VL are different from the pathological changes in their spleen which might be due to the genetic and immune process. Also, experimental surveys in vivo cannot be an appropriate way for the investigation into VL pathological changes because the results in experimental and natural infected animals vary. It is recommended that we should rely on the results of human experiments in this regard.

Keywords: *L. infantum*, Amastigote, J774 macrophage, Golden hamster, Pathological change

ICIA2018\Immunoparasitology\Poster Discussion\9962
9962

***Toxocara canis* Infection in Children with Hypereosinophilia (2–15 Years Old) Referred to Health Centers of Lorestan Province, Iran**

Pardis Bayat¹, Maryam Moslehi Baharanchi¹, Nadereh Taei², Hossein Mahmoudvand^{3,*}

1. Student Research Committee, Lorestan University of Medical Sciences, Khorramabad, Iran.

2. Department of Pediatric, Lorestan University of Medical Sciences, Khorramabad, Iran

3. Department of Medical Laboratory Sciences, School of Allied Medicine, Lorestan University of Medical Sciences, Khorramabad, Iran

Background: The present study aims to assess the seroprevalence of *Toxocara canis* infection in children with hypereosinophilia (2–15 years old) referred to health centers of Lorestan province, Iran.

Methods: This cross-sectional study was performed from August 2016 to March 2017 on 73 children with hypereosinophilia (eosinophilia>10%) (2–15 years old) referred to health centers of Lorestan province, Iran. All serum samples were tested using the commercially available anti IgG-*Toxocara* kit (IBL, Germany) according to manufacturer's instructions. The variables used to evaluate association between risk factors and status of anti-*T. canis* IgG antibodies were age, gender, area of residence, contact with dogs, and consumption of raw or unwashed vegetables and fruits.

Results: Of the 73 children with hypereosiniphilia, 3 (4.1%) tested seropositive for anti-*T. canis* IgG antibody. Risk factors that were significantly related to *T. canis* seropositivity included living in rural regions ($p<0.05$) and contact with dogs ($p<0.05$). However, other demographic and risk factors did not demonstrate any association with *T. canis* seropositivity.

Conclusion: We found that *T. canis* infection is prevalent among children with hypereosinophilia referred to health centers of Lorestan province, Iran with an overall seroprevalence rate of 4.1%. These findings suggested that *T. canis* infection can be considered as one of the reasons for hypereosinophilia in children from Lorestan Province Iran.

Keywords: Toxocariasis; ELISA; eosinophilic; children

ICIA2018\Immunoparasitology\Poster Discussion\9967

9967

Relationship between Helminthic Infections and Hyper eosinophilia in Children Referred to Health Centers of Lorestan Province, Iran

Maryam Moslehi Baharanchi¹, Pardis Bayat¹, Nadereh Taei², Hossein Mahmoudvand^{3,*}

1. Student Research Committee, Lorestan University of Medical Sciences, Khorramabad, Iran.

2. Department of Pediatric, Lorestan University of Medical Sciences, Khorramabad, Iran

3. Department of Medical Laboratory Sciences, School of Allied Medicine, Lorestan University of Medical Sciences, Khorramabad, Iran

Background: Intestinal helminthic infections are well-known as one of the most important socioeconomic and health problems around the world. The present study aims to evaluate the prevalence of the helminthic infections in children with hyper eosinophilia referred to Health Centers of Lorestan Province, Iran.

Methods: This cross-sectional study was performed from August 2016 to March 2017 on 73 children with hyper eosinophilia (eosinophilia>10%) referred to Health Centers of Lorestan Province, Iran. The microscopic analysis was accomplished on 73 stools by means of the direct smear, and formol-ether methods.

Results: Out of the 73 children with hypereosiniphilia, 5 (6.8%) children were infected with at least one or more intestinal parasites including *Enterobius vermicularis*, *Hymenolepis nana*, *Trichostrongylus* spp, and *Ascaris lumbricoides*. Statistical analysis showed that some risk factors were significantly associated to the prevalence intestinal helminthic parasites included gender ($p<0.02$), living in rural regions ($p<0.001$), hands washing habit ($p<0.001$) and consumed raw or unwashed vegetables and fruits ($p<0.001$).

Conclusion: These findings suggested that intestinal helminthic parasites can be considered as one of the reasons for hyper eosinophilia in children from Lorestan Province Iran.

Keywords: Intestinal parasites, stool, eosinophilia, children



Immunopharmacology and Medicinal Plants

Poster Discussion

14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

ICIA2018/Immunopharmacology and Medicinal Plants/Poster Discussion/7387

7387

The Effect of Mesenchymal Stem Cells Treated with Caffeine in Alleviating the Rheumatoid Arthritis Induced in Wistar rat

Elham Moases Ghafari¹, Seyyed Meysam Abtahi Froushani¹

1. Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Background: Previous documents indicated that Caffeine treated mesenchymal stem cells (MSCs) augment the instruction of anti-inflammatory macrophages. This study was done to investigate the effect of Caffeine treatment on the potential of MSC in alleviating the animal model of Rheumatoid Arthritis (RA).

Methods: Bone marrow-derived MSCs were co-cultured with 0.5 mM of Caffeine for 24 h. RA was induced by injection of complete Freund's adjuvant into the footpad of Wistar rats. One week after immunization, rats were divided into 3 groups: treated with MSCs alone, treated with MSCs co-cultured with Caffeine and untreated group. The change in the diameter of wrists and ankles of each rat was recorded every 5 days until 33 days after induction. At the end, nitric oxide production by Griess method, the ability of yeast opsonized phagocytosis and intensity of respiratory burst (NBT Test) in the splenocytes population were analyzed.

Results: The edema and swelling of ankles of RA rats treated with Caffeine pulsed MSCs were significantly regressed compared to RA rats received MSCs alone. Nitric oxide, respiratory burst, and phagocytosis rate of neutral red in RA rats received MSCs exhibited significant decrease compared to RA rats without treatment. Albeit, these reductions were more prominent in RA rats treated with MSCs pulsed with Caffeine compared to RA rats received MSCs alone.

Conclusion: Caffeine pulsed MSCs caused more favorable outcomes in RA model compared to un-treated MSCs.

Keywords: Caffeine, Mesenchymal stem cell, Rheumatoid arthritis, Immunity Responses

ICIA2018\Immunopharmacology and Medicinal Plants\Poster Discussion\7414
7414

Role of 13-Cis Retinoic Acid in Mycobacterium tuberculosis Infection Control by Human U937 Macrophage

Bahareh Abd Nikfarjam^{1,2}, Marjan Nassiri Asl^{1,3}, Mehri Hajiaghayi^{1,2},
Taghi Naserpour Farivar^{1,4}

1. Cellular and Molecular Research Center, Qazvin University of Medical Sciences, Qazvin, Iran

2. Department of Immunology, School of Medicine, Qazvin University of Medical Sciences, Qazvin, Iran

3. Department of Pharmacology, School of Medicine, Qazvin University of Medical Sciences, Qazvin, Iran

4. Department of Microbiology, School of Medicine, Qazvin University of Medical Sciences, Qazvin, Iran

Background: Mycobacterium tuberculosis survives and proliferates within the main cells of the innate immune system, macrophages. The goal of our study was to investigate the immunostimulatory effects of 13-Cis retinoic acid in PMA-induced human U937 macrophages against Mtb R37Ra infection by evaluating its potential role in the cell surface expression of HLA-DR, CD14 molecules and NO production as well as prevention of the Mtb growth within macrophages.

Methods: In this study, we investigated the immunostimulatory effects of 13-Cis retinoic acid on Mycobacterium tuberculosis infected macrophages using Flowcytometry and Griess methods.

Results: 13-Cis retinoic acid enhanced the cell surface expression of HLA-DR and CD14 molecules on U937 macrophages and prevented the growth of Mtb within macrophages suggesting an activated phenotype for 13-Cis retinoic acid treated U937 cells. Also, 13-Cis RA, has significantly increased NO generation compared to untreated control macrophages ($P < 0.001$). 13-Cis retinoic acid has a significant inhibitory effect on Mycobacterium growth ($P < 0.05$). Moreover, 13-Cis RA is potent in stimulation of surface markers expression on infected macrophages ($P < 0.05$).

Conclusion: The results of our study showed that infected U937 macrophages treated with 13-Cis retinoic acid represented significant increases in NO production, CD14 and HLA-DR expression and also prevents intracellular survival of Mtb. Therefore, 13-Cis retinoic acid may have a significant therapeutic approach in the control of Mtb infection.

Keywords: Mycobacterium, Tuberculosis, 13-Cis retinoic acid, U937 Macrophages

ICIA2018 Immunopharmacology and Medicinal Plants\poster discussion\7417

7417

***In vitro* Assessment of G2013 (guluronic acid) on microRNA-155 in TLR4 Signaling Pathway**

Seyed Shahabeddin Mortazavi Jahromi^{1, 2}, Mehdi Malek Jamshidi^{1, 2}, Mehdi Yousefi³, Farzaneh Tofighi Zavareh⁴, Abbas Mirshafiey⁴

1. Department of Cellular and Molecular Biology, Kish International Campus, University of Tehran, Kish, Iran

2. School of Biology, University College of Science, University of Tehran, Tehran, Iran

3. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

4. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

Background: microRNAs are considered one of the main epigenetic mechanisms in regulating gene expression, and it has been shown that impairment in their expression causes autoimmune diseases including rheumatoid arthritis and multiple sclerosis. The present research intended to investigate the effects of G2013 (guluronic acid) on the gene expression level of microRNA-155 in the HEK-Blue TLR4 cell line.

Methods: Total RNA was extracted, and cDNA was synthesized for microRNAs and gene expression level of microRNA-155 was evaluated using the Real-Time PCR technique.

Results: It was found that this medication could not significantly reduce the gene expression level of microRNA-155 at low and high doses in HEK-Blue hTLR4 compared to the control group ($p > 0.05$). It was also observed that when used lipopolysaccharide (LPS) to HEK-Blue hTLR4 cells led to a significant increase in gene expression level of microRNA-155 while the addition of LPS four hours after treating the cells with guluronic acid could not increase the gene expression level of microRNA-155 ($p < 0.05$).

Conclusion: This study revealed that guluronic acid, as a novel non-steroidal anti-inflammatory drug, is able to reduce gene expression level of microRNA-155 after stimulation by LPS.

Keywords: G2013, Guluronic acid, microRNA-155, TLR4

ICIA2018\Immunopharmacology and Medicinal Plants\Poster Discussion\7418

7418

The Evaluation of the Effect of β -D-mannuronic Acid (M2000) on MicroRNA-146a and Related Molecules at the Level of Gene Expression in HEK Blue hTLR2 Cell Line

Seyed Shahabeddin Mortazavi Jahromi^{1,2}, Mehdi Malek Jamshidi^{1,2}, Ali Farazmand^{1,2}, Mehdi Yousefi³, Zahra Aghazadeh⁴, Abbas Mirshafiey⁴

1. Department of Cellular and Molecular Biology, Kish International Campus, University of Tehran, Kish, Iran

2. School of Biology, University College of Science, University of Tehran, Tehran, Iran

3. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

4. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

Background: microRNA-146a is a post-transcriptional negative regulator of immune cell signaling through TLRs (TLR2, TLR4). Changes in the gene expression of microRNA-146a were reported in the pathogenesis of many autoimmune diseases including rheumatoid arthritis, systemic lupus erythematosus, and psoriasis. This research aimed to study the effects of β -D-mannuronic acid (M2000) on the expression of microRNA-146a and target molecules (IRAK1, TRAF6) in the HEK-BluehTLR2 cell line.

Methods: M2000 cytotoxicity was assessed by the MTT assay. The expression levels of the mentioned genes were detected by quantitative real-time PCR method.

Results: The MTT findings showed that M2000 ($< 500 \mu\text{g/ml}$) has no cytotoxic effect on HEK-Blue hTLR2 cells. According to our findings, low and high doses of M2000 could significantly reduce gene expression level of microRNA-146a compared to control group ($p < 0.05$). It was also revealed that two doses of this medicine could significantly decrease the gene expression level of IRAK1 and TRAF6 compare to control group ($p < 0.05$).

Conclusion: This research showed that M2000, as novel NSAID with immunosuppressive properties, was able to moderate TLR2 signaling and minimize the probability of inflammatory reactions by inhibiting microRNA-146a, IRAK1, and TRAF6.

Keyword: β -D-mannuronic acid, M2000, microRNA-146a, Target molecules (IRAK1 and TRAF6)

ICIA2018/Immunopharmacology and Medicinal Plants/Poster Discussion/7732
7732

Thymol enhances the *in vivo* Generation of Foxp3⁺ regulatory T cells in Mice

Haideh Namdari¹, Maryam Izad², Zahra Amirghofran^{1, 3*}

1. Department of Immunology, Medical School, Shiraz University of Medical Sciences, Shiraz

2. Department of Immunology, Faculty of Medicine, Tehran University of Medical Sciences, Tehran

3. Autoimmune Diseases Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Regulatory T cells (Tregs) play a critical role to maintain tolerance and prevent autoimmunity, thus selective *in vivo* increase of Tregs has wide therapeutic implications for autoimmune and inflammatory diseases. We investigated the effects of thymol, a main component of thyme on Tregs generation in mice.

Methods: Splenocytes from non-immunized and ovalbumin (Ova)-immunized mice treated and untreated with thymol were isolated for enumeration of Treg cells by flow cytometry. Isolated cells were cultured in the presence of Ova to measure *ex vivo* expression of Tregs and their suppression activity using CFSE labeling.

Results: We demonstrated that *in vivo* treatment of Ova-immunized mice with thymol increased the number of Tregs to more than 8% compared to the untreated mice ($P < 0.01$). *Ex vivo* Ova challenge of splenocytes from Ova-immunized mice treated with thymol showed reduced proliferation of stimulated naïve T cells to $< 30\%$ of control ($p < 0.01$). The percentage of CD4⁺Foxp3⁺T cells in *ex vivo* culture of splenocytes from thymol-treated Ova-immunized mice increased to 11% compared to those from untreated Ova-immunized mice ($p < 0.01$). Intracellular staining of cells for TGF- β demonstrated an increased level of this cytokine in CD4⁺Foxp3⁺ Treg cells ($48.1 \pm 5.9\%$, $p < 0.05$).

Conclusion: These data showed the ability of thymol to improve Tregs generation and their suppressor activity by increasing TGF- β production and decreasing stimulated naïve T cells proliferation. These findings suggest thymol as a helpful natural immunosuppressive compound for treating autoimmune and inflammatory diseases.

Keywords: Thymol; immunomodulation; CD4⁺ T cells; regulatory T cells; ovalbumin

ICIA2018\Immunopharmacology and Medicinal Plants\Poster Discussion\8283

8283

The Cytotoxic Effect of Kombucha, a Fermented Microbial Product, on the 4T1, Mouse Breast Cancer Cell Line

Somayeh Ahmadiashar¹, Mahsa Alem¹, Norouz Delirezhi¹

1. Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Background: Kombucha has long been used in traditional medicine to treat a variety of ailments. It has also been seen that Kombucha ceases to grow cancerous cells. It can renew the chemical and ionic balance in the cell membrane so prevent many disorders. This study was done to the evaluation of the cytotoxic and apoptotic effects of the Kombucha on 4T1 as mouse breast cancer cell line.

Methods: For this purpose, after cell line culture, cells were divided into different groups such as negative control, treatment with different doses of Kombucha such as 50, 100, 150 and 200 µl/ml, for 24 h. The cell morphology of all groups was studied via invert microscope. In order to evaluation of cell vitality and viability tetrazolium dye (MTT) assay and natural red (NR) respectively were tested. Finally, propidium iodide (PI) and acridine orange (AO) staining was done to assess apoptosis and necrosis situation of the cancerous cells.

Results: The NR assay test showed that administration of Kombucha could statistically significant induces changes in the cell membrane viability. Moreover, MTT reduction assay test showed that Kombucha could decrease vitality and biological activity of Kombucha treated mouse breast cancer cell line compared to control group. PI and AO staining manifested that Kombucha induced apoptosis and necrosis in treated 4T1 cell line, but not in control cells.

Conclusion: The administration of Kombucha could improve the cytotoxic effects on breast cancer and could alter the phenotype of these cells.

Keywords: Kombucha, 4T1, Mouse breast cancer cell line.

ICIA2018/Immunopharmacology and Medicinal Plants/Poster Discussion/8621

8621

Evaluation of the immunomodulatory effects of *Mentha longifolia* main compound on proliferation and cytokine levels in human lymphocytes and Jurkat cell line

Maryam Bayat¹, Zahra Amirghofran^{1,2}

1. Department of Immunology, Shiraz University of Medical Sciences, Shiraz, Iran

2. Autoimmune Disease Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

Background: *Mentha longifolia* is a medicinal plant used in traditional medicine for various purposes. The present study assessed the immunomodulatory effects of *M. longifolia* main compound (MLMC), on peripheral blood mononuclear cells (PBMCs) and Jurkat T cell line.

Methods: PBMCs and Jurkat cell line were treated by different concentrations of MLMC and the proliferation inhibition was measured by BrdU and MTT colorimetric assays. Analysis of cell viability was performed by propidium iodide (PI) staining using flow cytometry. Cytokine levels were measured by enzyme-linked immunosorbent assay (ELISA).

Results: Our results showed that MLMC decreased the proliferation of PBMCs and Jurkat cell line dose and time-dependently; at the highest concentration (800 µg/ml), 43.9 ± 17% and 53.5 ± 10.8% of Jurkat cells were inhibited after 24 and 48 h of treatment, respectively. Corresponding rate of proliferation inhibition in PBMCs in the presence of the same concentration of MLMC was 96.37±0.27.

Analysis of cell viability using PI staining showed no significant cytotoxic effects at all concentrations used in both Jurkat and PBMCs. Measuring cytokine levels by ELISA indicated a decreased dose dependent IFN-γ secretion in treated PBMCs; The highest concentration (800µg/ml) of the compound reduced this cytokine level to 18.3 ± 2.4 pg/ml compared to the control (141 ± 20 pg/ml). The compound had no effects on IL-4 production by PBMCs.

Conclusion: These data showed the inhibitory effects of MLMC from *Mentha longifolia* on the proliferation and cytokine levels in PBMCs and Jurkat cells which imply the immunomodulatory effects of this compound.

Keywords: Immunomodulatory effect, *Mentha longifolia*, PBMCs, Jurkat cell line

ICIA2018\Immunopharmacology and Medicinal Plants\Poster Discussion\8629
8629

Influences of the Ginger Extract on the Expression of Chemokines CCL20 and CCL22 and Chemokine Receptors CCR6 and CCR4 in Experimental Autoimmune Encephalomyelitis

Abdollah Jafarzadeh^{*1}, Zahra Arabi², Rayhaneh Ahangar Parvin², Marziyeh Mohammadi Kordkhaiyali¹, Zahra Taghpour³, Maryam Nemati⁴

1. Department of Immunology, Medical School, Kerman University of Medical Sciences, Kerman, Iran
2. Department of Immunology, Medical School, Rafsanjan University of Medical Sciences, Rafsanjan, Iran
3. Department of Histology, Medical School, Rafsanjan University of Medical Sciences, Rafsanjan, Iran
4. Department of Laboratory Sciences, Para-Medicine School, Kerman University of Medical Sciences, Kerman, Iran

Background: Chemokines play a central role in the leukocytes infiltration into the central nervous system (CNS) which is an essential step in the pathogenesis of multiple sclerosis. Ginger possesses a wide range of immunoregulatory and anti-inflammatory properties. This study aimed to evaluate the effects of ginger extract on the expression of CCL20 and CCL22 and CCR6 and CCR4 in experimental autoimmune encephalomyelitis (EAE).

Methods: EAE induction was induced in female C57BL/6 mice by immunization with myelin oligodendroglial glycoprotein. Then, the EAE mice were treated with PBS or ginger extract, from day +3 to +30. At day 31, mice were scarified and the expression of aforementioned markers measured in the spinal cord measured using real time-PCR.

Results: The gene expression of CCL20, CCL22 and CCR4 in the spinal cord of PBS-administrated EAE mice was significantly increased as compared with healthy control group ($P<0.04$, $P<0.05$ and $P<0.02$, respectively). Treatment of EAE mice with 200- and 300 mg/kg ginger extracts reduced the expression of CCL20, CCL22 and CCR4 in comparison with PBS-administrated EAE group ($P<0.04$, $P<0.01$ and $P<0.002$ for 200 mg/kg ginger and $P<0.01$, $P<0.005$ and $P<0.004$ for 300 mg/kg ginger, respectively). The CCR6 expression in EAE mice treated with 200- or 300 mg/kg ginger extracts was lower than PBS-administrated EAE mice ($P<0.01$ and $P=0.07$, respectively).

Conclusion: Ginger extract down-regulates the expression of CCL20 and CCL22 and their receptors in EAE mice. The possible therapeutic potential of ginger for treatment of MS can need to be considered in more investigations.

Keywords: Experimental autoimmune encephalomyelitis, ginger extract, CCL20, CCL22, CCR4, CCR6

ICIA2018\Immunopharmacology and Medicinal Plants\Poster Discussion\9787

9787

Immunomodulatory Effect of APS on the Interaction of Peripheral Blood Mononuclear Cells with Cervical Cancer CellsElham Shokati¹, Elahe Safari¹, Kobra Entezami¹, Samaeh Khorami¹*1. Immunology Department, School of Medicine, Iran University of Medical Sciences*

Background: Immunomodulator components from traditional medicinal plants are explored as a beneficial therapeutic approach to improve the effectiveness of common therapy because of the severe side effects of common cancer therapies. Astragalus membranaceus is one of the important medicinal herbs. Polysaccharides are the main constituents of Astragalus membranaceus. Astragalus polysaccharides (APS) have been shown different immunostimulatory effects like anti-inflammatory, anti-oxidative and anti-cancer properties. In this study, the potential effect of APS on the proliferation of PBMC and induction of MDSCs in co-culture with cervical cancer cell line was evaluated.

Methods: The proliferation characteristics of PBMCs in co-culture with cell lines were performed with CFSE dilution assay. Cell viability of PBMCs and HeLa cell line was assessed with MTT test. The frequencies of MDSCs in the PBMCs of healthy individuals after co-culture between PBMCs and HeLa cell line was examined by using FACS analysis.

Results: The result showed that APS could significantly promote the viability of treated PBMCs co-cultured with HeLa cancer cell line ($p < 0.05$). APS could improve proliferation of treated PBMCs co-cultured with HeLa cancer cell line ($p < 0.01$). Moreover, no significant inhibition was observed on MDSCs expansion in treated PBMCs co-cultured with HeLa cell line.

Conclusion: Our results indicate that APS can improve immune response in vitro by increasing PBMCs proliferation. However, no effect was observed on rate of MDSC. But, it seems that APS as an immune modulator can enhance the anti-tumor immune response.

Keywords: APS, Immunomodulation, Cancer

ICIA2018\Immunopharmacology and Medicinal Plants\Poster Discussion\9886.doc
9886

Effect of *Ganoderma. Lucidum* (*G.Lucidum*) on Specific Immunoglobulin Isotype against Tetanus Vaccine

Sajjad Ahmadi Renani¹, Kazem Ahmadi²

1. Tehran University of Medical Science, Tehran, Iran

2. Department of Immunology, Faculty of medicine, Baqiyatallah University of Medical Science, Tehran, Iran

Background: The immuno-modulating effects of *Ganoderma. Lucidum* (*G.Lucidum*) polysaccharides have been widely studied, including promoting the function of antigen-presenting cells, humoral and cellular immunity. Previous studies have also shown its effect on cytokine production by peritoneal macrophages and enhancement of CD40/CD86 marker on human peripheral blood mononuclear cells. The aim of this study was evaluation of its effect on mice antibody isotype.

Methods: Twenty four mice were divided in 4 groups each of six. The first six mice were kept as control. The second group was orally treated with 40 µg/day/mouse *G.Lucidum* in drinking water for 30 days. The third group was subjected to three intra dermal administration of tetanus toxoid as vaccine without *G.Lucidum*. The last six mice in addition to *G.Lucidum* in drinking water, subjected to three intra dermal injection of tetanus toxoid as vaccination. All mice were anaesthetized and blood samples were collected at 45 days of first vaccination. Sera were separated and specific antibody isotype against tetanus toxoid were measured by ELISA.

Results: The results showed that *G.Lucidum* enhanced specific antibody level comparing with group having vaccine without *G.Lucidum* ($p < 0.001$). It has also been shown that the amount of IgG antibody against tetanus toxoid is sharply higher in group having both vaccine and *G.Lucidum* ($p < 0.001$).

Conclusion: The data presented are in agreement with previous research indicating the higher expression of CD40/CD86 marker on human blood mononuclear cells and also higher cytokine release in response to *G.Lucidum* treatment. Taken together, it seems that *G.Lucidum* could be a good immune modulator both in humoral and cellular immunity.

Keyword: *G.Lucidum*, tetanus vaccine, antibody isotype.



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Innate Immunity and Inflammation

Poster Discussion

ICIA2018\Innate Immunity and Inflammation\Poster Discussion\7348
7348

Decrease in Activated Tregs is accompanied by Their IL-17 Expression in the Peripheral Blood of Patients with Atherosclerosis

Mohammad Reza Yazdani¹, Shahdad Khosropanah², Mehrnoosh Doroudchi¹

1. Department of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

2. Department of Cardiology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Human atherosclerotic plaques contain 1-5% Treg cells compared to other chronically inflamed tissues (25%). A protective role is suggested for CD4+CD25+Foxp3+ natural Tregs (nTreg), while the role of Th17 cells in the atherosclerosis is controversial. In this study we enumerated nTreg subsets as well as IL-17 producing Tregs in the peripheral blood of patients with atherosclerosis.

Methods: Fifteen ml heparinized blood was obtained from each of the 13 non-diabetic patients and 13 control individuals with normal/insignificant coronary artery disease confirmed by angiography. Peripheral Blood Mononuclear Cells were separated by Ficoll. Flowcytometry was performed in both group without stimulation, for detection of CD4/CD25/CD45RO/Foxp3 markers, and after stimulation (PMA/ION: 50/250 ng/ml) for detection of CD4/ CD45RO/ IL-17/FoxP3. The data were analyzed with the FlowJo software and analyzed using Mann-Whitney test by SPSS.

Results: Higher frequencies of resting nTregs (CD45RO-CD25+Foxp3^{lo}) and activated nTregs (CD45RO+CD25^{hi}Foxp3^{hi}) were found in controls. The frequency of effector/memory T-cells (CD4+CD25+CD45RO+Foxp3-) was higher in patients. In CD4+CD25+ and CD4+CD25+Foxp3+ populations, Foxp3-MFI was higher in controls. CD4+IL-17+Foxp3+ and CD4+IL-17+Foxp3- subsets showed higher frequencies in patients before and after stimulation. In CD4+ population, the MFI of IL-17 was increased in controls compared with patients both before and after stimulation. After stimulation, CD4+CD45RO+Foxp3+IL-17- T-cells (aTregs) increased in controls, while Th17 (CD4+CD45RO-IL-17+Foxp3-) and CD4+CD45RO+IL-17+Foxp3- T-cells were more frequent in patients. The frequency of effector/memory T-cells remained higher in patients while the percentage of nTregs was higher in controls after stimulation. The frequency of intermediate cells (CD4+CD45RO+IL-17+Foxp3+) with a high IL-17-MFI was higher in patients.

Conclusion: We showed that in patients with atherosclerosis, the frequencies of aTregs and nTregs decrease and the frequencies of Th17 and effector T-cells increase compared to controls. IL-17 production by an intermediate population with an aTreg phenotype suggests a population shift from FoxP3+ to FoxP3+IL-17+ and FoxP3- cells in atherosclerosis.

Keywords: IL-17, Treg, Atherosclerosis, FoxP3

ICIA2018\Innate Immunity and Inflammation\Poster Discussion\7484

7484

Effects of Mannuronic Acid, as an Anti-inflammatory Drug, by targeting SHIP Signaling Pathway in Inflammation Process

Fatemeh Pourgholi ^{1,2}, Mahsa Hajivalili ^{1,2}, Zohreh Hekmati ², Mehdi Yousefi ^{1,2}

1. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

2. Department of Immunology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

Background: B-D-mannuronic acid (M2000) is a newly designed nonsteroidal anti-inflammatory drug (NSAID) that also exhibits immunosuppressive and antioxidant properties in various experimental models. Src Homology-2 domain containing inositol-5'-phosphatase (SHIP) is an adaptor protein in downstream TLR2 signaling pathway. SHIP has a vital role in controlling inflammation functions via its ability to down regulate PI3K signaling pathway. The aim of this research was evaluate M2000 effects on expression levels of SHIP in Lipopolysaccharide (LPS) induced inflammation in HEK-TLR2 cell line.

Methods: Optimum dose of M2000 was estimated by MTT assay and HEK-TLR2 cell line was treated by different concentrations of M2000 and LPS (1µg/ml). After 24 hours incubation, total RNA was extracted using TRIZOL in all of conditions. Then cDNA was synthesized according to instructions and then expression level of SHIP was assessed by quantitative Real time PCR method.

Results: Our MTT results shown that optimum doses of M2000 was up to 25µg/ml ($p < 0.05$). According to MTT results we considered 5 µg/ml and 25 µg/ml as low and high doses.

Our findings demonstrated that expression levels of SHIP in 5 and 25 µg/ml doses of M2000 were significantly up regulated to 1.68 (1.68 ± 0.1207 , $p = 0.01$) and 1.89 (1.89 ± 0.04 , $p = 0.004$) fold, respectively, in comparison to control group. SHIP expression level after treatment by LPS (1µg/ml), 2.33 fold decreased (2.33 ± 0.1635 , $p = 0.03$). After co treatment of LPS and M2000 in low and high doses, expression levels of SHIP 1.53 (1.53 ± 0.0346 , $p = 0.04$) and 1.85 fold increased (1.85 ± 0.0206 , $p = 0.04$) according to control group.

Conclusion: M2000 can be used as a safe NSAID in LPS induced inflammation and decrease inflammatory cytokine productions by targeting SHIP in different inflammatory diseases. Also M2000 can be a good candidate in treatment of autoimmunity and inflammatory disorders.

Keywords: M2000, SHIP, inflammation, NSAID, HEK-TLR2

ICIA2018\Innate Immunity and Inflammation\Poster Discussion\7582

7582

Evaluation of Gene Expression of NLRP3 and NLRC4 as Inflammasome Receptors in Peripheral Blood of SM Injured Patient by Real-Time PCRSholeh Etehad asnaf¹,*1. Department of Biology School of Science University of Islamic Azad North Tehran Branch*

Background: Mustard gas is one of the most widely used chemical warfare agents used during the Iraqi wars against Iranian warriors. Some mechanisms of chronic pulmonary damage caused by mustard gas such as oxidative stress, protease-anti protease imbalance and inflammation have been studied. In the context of inflammation, the role of inflammasome and the genes involved in it, especially the receptors of this inflammatory complex, is important. Therefore, the present study evaluated the expression of NLRP3 and NLRC4 genes as inflammatory receptors in peripheral blood of veterans of pulmonary chemo using Real-Time PCR.

Methods: The present study was conducted as a cross-sectional analytical study on 15 chemical warfare victims exposed to mustard gas, 15 COPD patients and 15 healthy individuals as controls that were referred to the lung clinic of Baqiyatallah Hospital (AS). After extracting RNA from the blood sample and synthesizing cDNA, expression of the two NLRP3 and NLRC4 genes was evaluated using Real Time PCR. Finally, the data were analyzed by SPSS version 20.

Results: The two NLRP3 and NLRC4 genes did not change significantly in veterans and people with COPD compared to healthy subjects. Although the expression of these two genes was higher in COPD patients than veterans (0.17 to 0.05 and 0.15 to 0.04, respectively), there was no statistically significant relationship.

Conclusion: According to the results of the present study, it has been found that two NLRP3 and NLRC4 genes have the potential to be involved in inflammation and chronic obstructive pulmonary embolism. As a result, it is hoped that with the full knowledge of the process of chronic damage to the lungs caused by mustard gas, one can use it to create a pattern of diagnosis and prevent its further complications.

Keywords: Mustard gas, chemical veterans, inflammasome, NLRP3 gene expression, NLRC4 gene expression

ICIA2018\Innate Immunity and Inflammation\Poster Discussion\7719
7719

Effect of Aqueous Extract and Saponin Fraction of *Tribulus terrestris* L. on the Expression of ICAM1 and VCAM1 in the Human Endothelial Cell Lines in Vitro

Elahe Amirinezhad Fard¹, Zahra Fereydouni¹, Ali Mostafaie¹

1. Department of immunology, Kermanshah University of Medical sciences, Kermanshah

Background: Atherosclerosis is a common chronic inflammation which during of it low density cholesterol and lipid deposition on the wall of medium- and large-sized arteries are formed, known as fibrotic-lipid plaques (atheroma). In this study we aimed at investigating the effect of aqueous extract and saponin fraction of *Tribulus terrestris* L. on the expression of Intracellular Adhesion Molecule-1 (ICAM1) and Vascular Cell Adhesion Molecule-1 (VCAM1) in the Human Umbilical Vein Endothelial Cell (HUVEC) and Human Bone Marrow Endothelial Cell (HBMEC) lines in vitro during normal and LPS-induced conditions.

Methods: HUVEC and HBMEC lines were cultured and induced with LPS. Afterwards, the effect of aqueous extract and saponin fraction of *T. terrestris* on the expression pattern of endothelial proteins especially VCAM-1 and ICAM-1 in the HUVEC and HBMEC lines in vitro during normal and LPS-induced conditions was characterized by SDS-PAGE and blotting.

Results: Our findings showed that the aqueous extract and saponin fraction of *T. terrestris* decreased the expression of adhesion molecules (ICAM-1 and VCAM-1) in both LPS-induced HUVEC and HBMEC lines.

Conclusion: Aqueous extract and saponin fraction of *T. terrestris* clearly affected protein pattern of HUVEC and HBMEC lines, so that the expression of proteins with molecular weight of more than 90 and 65 KDa decreased. In addition, the extract and saponin fraction of *T. terrestris* down regulate the expression of ICAM-1 and VCAM-1. Anti-inflammatory activity of the aqueous extract is more than the effect of saponin fraction of *T. terrestris*.

Keywords: Atherosclerosis, ICAM1, VCAM1, *Tribulus terrestris* L.

ICIA2018\Innate Immunity and Inflammation\Poster Discussion\7937
7937

The Inhibitory Role of M2000 (Beta-D-Mannuronic acid) on the Expression of Toll-like Receptor 2 and 4 in HT29 cells

Laleh Sharifi¹, Mona Moshiri², Mohammad Mehdi Soltan Dallal², Saied Bokaie³, Zahra Norouzababaei⁴, Abbas Mirshafiey⁵

1. Uro-Oncology Research Center, Tehran University of Medical Sciences, Tehran, Iran

2. Department of Microbiology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

3. Department of Epidemiology, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran

4. Department of Virology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

5. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

Background: Anti-inflammatory agents play main roles in controlling inflammatory diseases such as inflammatory bowel disease (IBD); but due to the vast side effects their use are restricted. M2000 (Beta-D-Mannuronic acid) is a new immunomodulatory drug. According to the key role of TLRs in inflammatory responses, in this study we aimed to evaluate the role of M2000 on the expression of TLR2 and TLR4 in HT29 cell line as colonic epithelial cells.

Methods: MTT assay was carried out to determining the viability rate of cells after M2000 treatment. HT29 cells were cultured and treated with low (5µg/ml) and high (25µg/ml) doses of M2000. Total RNA was extracted and cDNA synthesized, quantitative real-time PCR was done to quantify the TLR2 and TLR4 mRNA synthesis.

Results: We found that M2000 at concentration of ≤ 1000 µg/ml had no obvious cytotoxicity outcome. The results of quantitative real-time PCR showed that the expression of TLR2 mRNA in HT29 cells in the treatment group of low-dose M2000 were decreased significantly compared with untreated control group, 1.00 ± 0.16 vs 0.32 ± 0.03 ($p=0.02$). Also we showed that high dose of M2000 decrease the expression of TLR2 significantly 1.00 ± 0.16 vs 0.05 ± 0.03 ($p=0.01$) Treating the cells with low-dose M2000 led to significant reduction in TLR4 compared to control cells, 1.00 ± 0.09 vs 0.52 ± 0.08 ($p=0.01$); similarly high dose of M2000 decreased gene expression of TLR4, 1.00 ± 0.09 vs 0.16 ± 0.03 ($p=0.008$).

Conclusion: The results of present study in a model of colonic epithelial cells indicate that M2000 can be consider as a new treatment of inflammation in IBD. However more studies are required to recognize the comprehensive molecular mechanism of M2000 and evaluate its therapeutic effects.

Keywords: M2000, Beta-D-Mannuronic acid, IBD, TLR2, TLR4

ICIA2018\Innate Immunity and Inflammation\Poster Discussion\7979

7979

Amelioration of High-Glucose Induced Inflammation by Quercetin Through Macrophage Polarization in RAW624.7 Cell Line

Reyhaneh Babaei Khorzoughi¹, Maryam Shabani^{*1}

1. Department of Clinical Biochemistry, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

Background: Macrophages are divided into two phenotypes (M1 and M2) based on their surface markers. M1 markers include CD11, iNOS, etc. and M2 markers include CD206, CD163, etc. These markers activations are affected by several mechanisms such as ER stress, hypoxia, lipotoxicity, ROS production and NF- κ B activation. High glucose concentrations give rise to lipotoxicity and ROS products that result in M2 to M1 shift and lead to inflammation. Pro-inflammatory cytokines are released by M1 macrophages and M2 macrophages release anti-inflammatory cytokines. Furthermore, the balance between M1/M2 has an important role in inflammation mechanisms. Many studies have been shown polyphenols have anti-inflammatory potential and safety effects. Quercetin is a flavonol, one of the six subclasses of flavonoid compounds. This polyphenol possesses strong anti-inflammatory capacities; through an increase of anti-oxidative activities, reduction of lipogenesis and macrophage polarization regulation.

Methods: This study has surveyed the glucose (53mM) modulation of RAW 264.7 macrophages activation and effects of Quercetin (25 μ M) on high glucose-induced lipotoxicity and macrophage polarization. We have measured high glucose-induced lipogenesis by oil red O staining. For investigating macrophage polarization, we assess M2 marker, CD206, as an anti-inflammatory factor and M1 marker, CD11c, as an inflammatory factor via flow cytometry.

Results: Our results suggest that high glucose induce RAW264.7 lipogenesis and modify RAW264.7 morphology. Flow cytometry analyses showed that high glucose increase M1 marker, CD11c, significantly (about 80%) in vitro. RAW264.7 were treated with Quercetin strongly reduce lipid droplet in oil red O staining and decrease M1 marker, CD11c to approximately 20%. Our data have shown quercetin Caused a very slight increase in M2 marker, CD206 (4%) that isn't signed.

Conclusion: These results show that Quercetin produces a potential anti-inflammatory effect by modulating macrophage polarization and attenuate high glucose-induced lipogenesis in vitro. Decreasing M1 phenotype involved in the anti-inflammatory property of Quercetin.

Keyword: Inflammation, Quercetin, High Glucose, Macrophage polarization

ICIA2018\Innate Immunity and Inflammation\Poster Discussion\8043
8043

Increased Interleukin-1 β and Caspase 1 Production through Activation of Inflammasome Contributes to the Neuroinflammation and Memory Impairment in a STZ-Induced Model of Alzheimer Disease

Mahdiye Saadi ^{1,2}, Ahmad Karkhah^{1,2}, Fereshteh Pourabdolhossein ², Hamid Reza Nouri ²

1. Student Research Committee, Babol University of Medical Sciences, Babol, I.R. Iran

2. Cellular and Molecular Biology Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, I.R. Iran

Background: Alzheimer's disease (AD) is an inflammatory neurodegenerative disease considered by neuronal cell death and progressive dementia. Recent evidence demonstrated that IL-1 β plays a critical role in the pathogenesis of AD. Thus, the present study was conducted to investigate which inflammasome pathway contribute in IL-1 β production and memory impairment in Alzheimer disease.

Methods: In this study, Alzheimer disease was induced in male Wistar rats by two bilaterally injection of Streptozotocin 3 mg/kg into the lateral ventricles (each side 3 μ l) with 48 h interval. Morris Water Maze was used to evaluate learning and memory. In addition, Hematoxylin and Eosin staining was used to assess inflammatory parameters in hippocampus region. Expression level of several genes involved in inflammasome pathway including NLRP1, AIM2, NLRP3, NLRC4, ASC, IL18, IL-1 β and Caspase 1 in the hippocampal tissue was measured using Real-time PCR.

Results: Behavioral study revealed that STZ injection in the lateral ventricles impaired learning and memory function ($p < 0.05$). Histological study using H & E staining confirmed the inflammatory response in the hippocampus region of STZ treated animals. Gene expression studies demonstrated a significant ($p < 0.05$) increase in NLRC4 mRNA expression in STZ treated group when compared to control group. The expression level of ASC, IL-18 and Caspase 1 in the hippocampus of STZ treated group was more than control, but there was no statistically significant difference between the two groups. In addition, mRNA expression level of NLRP1, AIM2 decreased in STZ treated group as compared with control group, however this decrease was not statistically significant.

Conclusion: We concluded that increased interleukin-1 β and caspase 1 production through activation of NLRC4 inflammasome may be involved in neuroinflammation and memory impairment in Alzheimer disease which can introduce promising targets for treatment of Alzheimer disease.

Keywords: Alzheimer, interleukin-1 β , Caspase 1, NLRC4, Neuroinflammation, Memory impairment.

ICIA2018\Innate Immunity and Inflammation\Poster Discussion\8064
8064

Development of Experimental Fibrotic Liver Diseases Animal Model by Carbon Tetrachloride

Atoosa Gitiara¹, Samaneh Tokhanbigli¹, Sogol Mazhari¹, Kaveh Baghaei^{1*}, Behzad Hatami², Seyed Mahmoud Hashemi³, Ali Asadi Rad³, Afshin Moradi², Nakisa Zarrabi Ahrabi⁴, Hamid Asadzadeh Aghdaei¹, Mohammad Reza Zali²

1. Basic and Molecular Epidemiology of Gastrointestinal Disorders Research Center, Research institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran.
2. Gastroenterology and Liver Diseases Research center, Research institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran.
3. Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran
4. Department of Biology Central Tehran Branch, Islamic Azad University, Tehran, Iran

Background: Liver fibrosis is a result of inflammation and liver injury caused by wound healing responses which ultimately lead to liver failure. Consequently, after liver fibrosis, the progression will be continued to liver cirrhosis and at the end stage hepatocellular carcinoma (HCC). Many studies have demonstrated that one of the most important causes of liver fibrosis is Non-alcoholic steatohepatitis (NASH). Fibrotic Liver is affected by an excessive accumulation of extracellular matrix (ECM) proteins like collagen and α -SMA.

Methods: in two different experiment, male Vistar, and Sprague Dawley Rat models ranging from 200 \pm 60, corresponding to an age of approximately 10 weeks were utilized in order to induce CCL4 treated liver fibrosis.

Results: After 6 weeks of CCL4 injection, different tests have been carried out to verify the liver fibrosis including serum markers such as Aspartate aminotransferase (AST) and Alanine aminotransferase (ALT), molecular tests containing, laminin and α -SMA and also pathological observation by Hematoxylin and eosin staining in both fibrosis and control group.

Conclusion: The results of Pathology and Real-time PCR showed that fibrosis was induced much more effective in Sprague Dawley rat model in compare with Wistar rats.

Keywords: Liver Fibrosis, CCL4, Animal Model.

ICIA2018\Innate Immunity and Inflammation\Poster Discussion\8169

8169

The Effects of Marhame-Mafasel in IL-1 β ,IL-6 ,TNF α Expression in Animal Induced Arthritis

Mohammad Majidi^{1, 2}, Roya Yaraee^{1*}, Shahin Bonakdar², Mohsen Naseri¹, Mohammadreza Jalali³, Fatemeh Heidarnejad², Arsalan Jalili A⁴ Maryam Salimi⁵

1. Department of Immunology, Shahed University, Tehran, Islamic Republic of Iran

2. National Cell Bank Department, Pasteur Institute of Iran, Tehran, Iran

3. Department of Pathology, School of Medicine, Shahed University, Tehran, Iran

4. Department of Stem Cells and Developmental Biology at Cell Science Research Center, Royan Institute for Stem Cell Biology and Technology, ACECR, Tehran, Iran.

5. Department of Biology and Anatomical Sciences, Faculty of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Rheumatoid arthritis (RA) is a systemic chronic disease with synovial membrane, tendon and auricular tissue inflammation. Two fundamental criteria of RA include chronic inflammation and bone absorption. Current treatments of RA have many side effects and are so expensive. Currently, new treatments procedures and inexpensive herbal drugs are developed. Marhame-Mafasel made of 2 traditional herbs (Arnebiaeuchroma and Martricariachamomilla) and produced by pharmacology department of Shahed university of Iran. Marhame-Mafasel has been used for RA patient's treatment for many years. In this study, the effects of Marhame-Mafasel on a part of immune system were evaluated by an animal model of RA.

Methods: 15 male wistar rats were used in three group of Marhame-Mafasel, Paraffin (negative control) and piroxicam (positive control). Volume of hind paw was measured every other day from 0 to 19 using water changed volume approach. Gene expression of TNF α ,IL-1 β and IL-6 were evaluated using Real-Time PCR and inflammation in joint was evaluated using histopathology assay.

Results: Hind paw swelling of Marhame-Mafasel and piroxicam groups at day 10 and 19 that are most important days in animal model of RA(10: acute inflammation, 19: chronic inflammation), was reduced. This note should be taken, reduction between each groups weren't statistically significant. Histopathology and Real-Time PCR evaluation between three groups were not statistically significant.

Conclusion: Although the hind paw swelling of Marhame-Mafasel and piroxicam groups was reduced, Because of the small number of animals and the lack of precise measurement instrument, results have large variance then more future studies must be carried out.

Keywords: Marhame-Mafasel, IL-1 β ,IL-6 ,TNF α , Animal induced arthritis

ICIA2018\Innate Immunity and Inflammation\Poster Discussion\9840

9840

Phagocytosis of Apoptotic Mesenchymal Stem Cells by Mouse - Peritoneal Macrophages Effects on Cytokine Production and Phagocytosis Ability

Maryam Ghahremani Piraghaj¹, Seyed Mahmoud Hashemi^{1*}, Hosein Ghanbarian², Sara Soudi³

1. Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Department of Biotechnology, School of Advanced Technologies in Medicine, Shahid Beheshti of Medical Sciences, Tehran, Iran University

3. Faculty of Medical Sciences, Department of Immunology, Tarbiat Modares University, Tehran, Iran

Background: Macrophages are one of the most important immune cells. Macrophages can be divided into two main subgroups of classical (M1) and alternative or (M2), due to different stimuli. One of the factors that cause the macrophage to orient towards M2 is the phagocytosis of apoptotic cells (efferocytosis). The phagocytosis of mesenchymal stem cells, can be very important in cell therapy due to immunomodulatory properties and their ability to modulate macrophage function

Methods: Adipose-derived mesenchymal stem cells were isolated from CB57bl/6 Mice and characterized by flow cytometry as well as differentiation to *osteoblasts and adipocytes*. MSCs in the passage 2 exposed to UV light for induction of apoptosis for thirty minutes and followed by incubation for two hours. The cells were then isolated and added to macrophage in a ratio of 4 to 1. After that cells were incubated for 48 hours, after incubation time the production of TNF α and IL10 cytokines was measured by ELISA. Macrophages phagocytosis ability was also measured by yeast and apoptotic thymocytes phagocytosis in different groups.

Results: The phagocytosis of apoptotic mesenchymal stem cells by macrophages reduces the production of inflammatory cytokine TNF α and increases the production of inhibitory cytokine IL-10. Also after phagocytosis of mesenchymal stem cells, the ability of yeast phagocytosis in these macrophages is reduced and phagocytosis of apoptotic thymocyte was increased

Conclusion: The phagocytosis of apoptotic mesenchymal stem cells, induce non-inflammatory phenotype in macrophages. So, injected Mesenchymal Stem Cells maintain their immunomodulatory properties even if they apoptosis in the body.

Keywords: macrophage, Mesenchymal stem cell, Efferocytosis, Phagocytosis



Monoclonal Antibody Diagnostic and Therapeutic

Poster Discussion

14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Monoclonal Antibody Diagnostic and Therapeutic\Poster Discussion\5240

5240

The Effect of All-trans Retinoic Acid Alone and Combination with Osahexaenoic Acid on Monoclonal Antibody Production in Hybridoma Cells: an *in vitro* and *in vivo* Study

Mohsen Rokni¹, Ali Akbar Saboor Yaraghi¹, Ali Reza Razavi¹, Fazel Shokri¹, Tayebbeh Sohrabi², Maryam Kargar³, Masoome Keykha³, Masoud haghshenas⁴

1. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

2. Department of Hematology and blood bank, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

3. Department of Immunology, Faculty of Medicine, Zahedan University of Medical Sciences, Zahedan, Iran

4. Blood Transfusion Research Center, High Institute for Research and Education in Transfusion Medicine, Tehran, Iran

Background: Hybridoma cells can produce monoclonal antibody (MAb) and the productions of these antibodies can be affected by different factors including stimulants, inhibitors, and supplements. Several types of research have been done to improve MAb production in cell culture and in animal model. However, the impact of micronutrients on the production of MAb in mouse hybridoma cells and in the peritoneum of animals is not fully understood.

Methods: The hybridoma cells, M₃C₅, were cultured and treated with different concentrations of all trans-retinoic acid (ATRA) and osahexaenoic acid (DHA), alone and in combinations and at different time of exposure, then the changes in production of Mab in culture medium were evaluated using ELISA and compared with vehicle-treated controls. The hybridoma cells, after single and combined treatment with ATRA, DHA, and vehicles were IP injected to Balb/c mice and the changes in production of Mab in ascites of mice, were determined by ELISA. The effect of these metabolites on the cloning efficiency of hybridoma cells was also explored.

Results: The results showed that single and combined treatment of ATRA and DHA significantly elevated the productions of MAb by M₃C₅ hybridoma cells, in both *in vivo* and *in vitro* conditions. The production of MAb following *in vitro* single treatment with 1μM of ATRA ($p<0.003$) and 10μM of DHA ($p<0.007$) for 2 days was significantly increased. The *in vitro* effects of ATRA on up-regulation of MAb production was obtained more than DHA however, at *in vivo* conditions, DHA was more effective in increasing the MAb production($p<0.001$).

Conclusion: The results of this study, for the first time, showed that ATRA and DHA are effective supplements that can increase production of MAb in mouse hybridoma cells under both *in vitro* and *in vivo* conditions.

Keywords: Hybridoma cells, Monoclonal antibody, osahexaenoic acid, All-trans retinoic acid, Cloning efficiencies, Ascites.

ICIA2018\Monoclonal Antibody Diagnostic and Therapeutic\Poster Discussion\7720
7720

Characterization of Monoclonal and Polyclonal Antibodies Recognizing Prostate Specific Antigen: Implication for Design of a Sandwich ELISA

Sahar Raoofi Mohseni¹, Forough Golsaz Shirazi¹, Mostafa Hosseini², Jalal Khoshnoodi¹, Tannaz Bahadori³, Mohammad Ali Judaki¹, Mahmoud Jeddi Tehrani³, Fazel Shokri^{1,3}

1. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

2. Department of Epidemiology and Biostatistics, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

3. Monoclonal Antibody Research Center, Avicenna Research Institute, ACERCR, Tehran, Iran

Background: Prostate cancer is the second common cancer in men. Prostate-specific antigen (PSA) is a tumor-associated glycoprotein with enzymatic activity which is secreted by the prostate gland. Following entry to the blood, 70-90 % of PSA forms complexes with protease inhibitors and its enzymatic activity is inhibited. The serum level of PSA is increased and the rate of free PSA (fPSA) to total PSA is decreased in prostate cancer patients. Therefore, measurement of PSA and fPSA in serum is very valuable for diagnosis and prognosis of prostate cancer.

Methods: In the present study, five anti PSA monoclonal antibodies (mAb) were characterized by enzyme-linked immunosorbent assay (ELISA) and immunoblotting. For design of a sandwich ELISA, epitope specificity of these antibodies was studied by a competition ELISA. Free PSA was purified by electroelution technique from seminal plasma and used to produce polyclonal anti-fPSA antibody in rabbit. Purified Polyclonal antibody (pAb) and mAbs were conjugated with HRP enzyme and Biotin (Bio) to set up the sandwich ELISA.

Results: Three of the mAbs were found to recognize PSA, similarly. One of these mAbs (2G3) was paired with anti-fPSA pAb to detect fPSA in serum. Eventually serum fPSA concentration of 356 subjects was measured and compared by our designed ELISA and a commercial ELISA kit. Our results demonstrated a significant correlation ($r=0.68$; $P<0.001$) between the two assays. Sensitivity and specificity of our designed ELISA is 72.4% and 82.8%, respectively

Conclusion: These results imply suitability of our designed ELISA for detection of fPSA in patients with prostate cancer.

Keywords: Free PSA, prostate cancer, sandwich ELISA test, monoclonal antibodies

ICIA2018\Monoclonal Antibody Diagnostic and Therapeutic\Poster Discussion\8164
8164

Evaluation of the Binding Activity of a Specific VH Antibody Fragment for *Neisseria Meningitidis* Factor H Binding Protein (fHbp) via ELISA

Hediyeh Rafighdoust¹, Fatemeh Yarian^{2,3}, Fattah Sotoodehnejad Nematalahi¹, Mona Salahshoor dahr⁴

1. Department of Biology, School of Basic Sciences, Science and Research Branch, Islamic Azad University, Tehran, Iran

2. Department of Biotechnology, School of Advanced Technologies in Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

3. Cellular & Molecular Biology Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

4. Department of Biology, School of Basic Sciences, Islamshahr Azad University, Tehran, Iran

Background: Although widespread use of whole antibodies in medicine and pharmacy is undeniable, they also have many limitations for most diagnostic and therapeutic applications. Antibody fragments have shown high potential detection of infectious agents, and studies in this area are potentially ongoing. Previously, we obtained a specific scFv antibody for *N. meningitidis* factor H binding protein (fHbp) and it's binding to the fHbp was characterized by bioinformatics tool. The bioinformatics data showed that the VH fragment of the scFv has more binding to the antigen. The main study was evaluation of the binding activity of a specific VH fragment to the recombinant fHbp antigen.

Methods: The T.Vector contained VH antibody fragment gene and pET28a vector were cut with NcoI, NotI enzymes. It was successfully cloned into pET28a vector, transformed into *E.coli* strain BL21 and expressed in optimal condition. The expressed protein was approved by SDS-PAGE and Western blotting. The recombinant His-tagged VH antibody was purified via affinity chromatography using Ni²⁺-NTA agarose resin. Binding activity of the VH fragment antibody for recombinant factor H binding protein (fHbp) was evaluated via ELISA.

Results: The cloned VH fragment antibody into the pET28a (+) vector was expressed. The expressed protein (about 12 kDa) was investigated by SDS-PAGE and approved in western blotting by anti-his tag antibody. We used the ELISA technique to determine the functions of the purified VH fragment antibody.

Conclusion: cloning, expression and purification of the VH fragment antibody was successfully performed. ELISA was performed to determine the functions of the purified VH antibody. The results indicate that despite the small size of the antibody fragment, it was exhibited more binding activity to recombinant fHbp. The VH fragment antibody may provide the basis for the development of a diagnostic kit.

Key words: VH fragment antibody, Cloning, *Neisseria meningitides*, factor H binding protein

ICIA2018\Monoclonal Antibody Diagnostic and Therapeutic\Poster Discussion\8192
8192

Design and Expression of a Specific Diabody for *Neisseria Meningitidis* Factor H Binding Protein (fHbp)

Mona Salahshoor Dahr¹, Fatemeh Yarian^{2,3}, Maryam Ghane¹, Mojgan bandehpour^{2,4}, Hediye Rafighdoust⁵

1. Department of Biology, School of Sciences, Islamic Azad University Islamshahr Branch, Islamshahr, Iran

2. Cellular & Molecular Biology Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

3. Department of Biotechnology, School of Advanced Technologies in Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

4. Department of Biotechnology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

5. Department of Biology, School of Basic Sciences, Science and Research Branch, Islamic Azad University, Tehran, Iran

Background: The single-chain fragment variable antibody, scFv, is an antibody fragment that has shown a unique capability for binding to many antigens, such as proteins and even whole pathogens. By engineering the monovalent antibody fragments (such as Fab and scFv) into multivalent molecules as diabodies, triabodies or larger aggregates, it can be significantly increased their functional affinity and specificity to the target antigens. Previously, we obtained two specific scfvs against *N.meningitidis* factor H binding protein (fHbp). The functionality of these novel scFvs were characterized by ELISA. The main purpose of this study is design and expression of a specific diabody for this protein.

Methods: In this study, the nucleotide sequence of the diabody was designed and synthesized. The pGH vector contained the diabody gene and pET28a vector were cut with Nco I, Not I enzymes. The cloned diabody into the pET28a (+) vector were transformed into *E.coli* strain BL21 and expressed in optimal condition. The expressed protein was approved by SDS-PAGE and Western blotting.

Results: The desired gene is 1430 nucleotides. The expressed protein (about 51 kDa) was investigated by SDS-PAGE and was approved in western blotting by anti-his tag antibody.

Conclusion: The design antibody was successfully cloned into pET28a vector and expressed. It is expected that the designed diabody will be efficient for diagnosing of all *N. meningitis* serogroups in CSF in the future.

Keywords: Diabody, Cloning, factor H binding protein, *Neisseria Meningitidis*

ICIA2018\Monoclonal Antibody Diagnostic and Therapeutic\Poster Discussion\8202
8202

Isolation of MKN-45 Gastric Cancer cell Recognizing Single Chain Variable Fragment (scFv) Antibodies Using Phage Display Technology

Tayebeh Mehdipour^{1,2}, Mohammad Reza Tohidkia², Ali Akbar Rahim Rahimi^{2,3}, Mohammad Nouri^{2,4}

1. Department of Medical Biotechnology, School of Advanced Medical Science, Tabriz University of Medical Science, Tabriz, Iran

2. Research Center for Pharmaceutical Nanotechnology, Tabriz University of Medical Sciences, Tabriz, Iran

3. Department of Microbiology, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

4. Department of Biochemistry, School of Medicine, Tabriz University of Medical Sciences, Iran

Background: Aim of this study was to isolate recombinant antibody fragments recognizing poorly differentiated metastatic gastric cancer (GC) cell line, MKN-45, to direct discovery of potential tumor markers related to diffuse type adenocarcinoma.

Methods: A semi-synthetic human scFv library (Tomlinson I) was used for identification of novel antibody fragments recognizing MKN-45 GC cell line. Diverse antibody phage-scFv libraries were prepared from *E.coli* transformants infected with two different helper phages, KM13 and hyperphage, in parallel. Overall four rounds of subtractive selection were carried out, each consisting of three pre-absorptions using NIH 3T3 and AGS cell lines prior to positive selection on MKN-45 cells. Characterization and specificity analysis of the selected scFv antibodies were determined through whole cell ELISA and flow cytometry.

Results: ELISA-based screening of 192 phage-displayed scFv clones indicated that 2 and 3 scFv candidates rescued by KM13 and hyperphage respectively, produced strong binding activity against MKN-45 and low reactivity to negative AGS and NIH 3T3 control cells. Further analysis by flow cytometry illustrated that B12 clone from KM13 and G1clone from hyperphage had the most binding activity and specificity for MKN-45 GC cell line.

Conclusion: The selected scFvs may be potential immunereagents in diagnosis and even novel therapeutic agents of metastatic GC. Additionally, efficient and sit-specific delivery of the drugs and nanoparticles can be done with the aim of these antibody fragments.

Keywords: Phage display, Whole cell panning, Gastric cancer, Diffuse adenocarcinoma

ICIA2018\Monoclonal Antibody Diagnostic and Therapeutic\Poster Discussion\8332
8332

Optimization of Refolding Procedures for Novel Humanized Immunotoxin Produced in the *E. coli*

Bahman akbari¹, Safar farajnia²

1. Department of Medical Biotechnology, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

2. Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

Background: High-level expression of many recombinant proteins in *E. coli* leads to formation of inclusion bodies, highly aggregated proteins; they can also form in mammalian, insect, and yeast cells. Over expression of a humanized immunotoxin (IT) against epidermal growth factor receptor (EGFR) led to the production of inclusion bodies. The aim of this study was optimization of refolding process for recombinant humanized anti-EGFR IT produced in the *E. coli*.

Methods: The BL21 (Plys S) cells containing a pET-22b- humanized anti-EGFR IT construct -constructed and characterized in previous work- were induced by 0.25mmol/l IPTG at 25 C° for 4 h and the amount of expression was checked by SDS-PAGE, even though the majority of expressed proteins appeared as inclusion bodies. These inclusion bodies were individually solubilized in 8 M urea and 6 M guanidine hydrochloride and then purified by Ni-NTA affinity chromatography which was seen as a single band in SDS-PAGE analysis.

Results: The proper refolding humanized IT achieved by stepwise dialysis method. Reactivity assessment of humanized IT obtained from the urea approach with A431 carcinoma cells by relative ELISA test revealed that the refolded humanized IT had a high reactivity, indicating suitable folding of purified antibody. The 50% binding activity of humanized IT achieved from urea and guanidine hydrochloride approaches were 0.5 and 1.75 µg/ml concentrations, respectively.

Conclusion: The results of this study revealed that the urea approach was very effective in solubilizing and refolding of IT that expressed in bacteria cells as inclusion bodies.

Keywords: Guanidine hydrochloride, Inclusion body, Refolding, Urea.

ICIA2018\Monoclonal Antibody Diagnostic and Therapeutic\Poster Discussion\8335

8335

An Investigation on the Structure and Function of Cetuximab Monoclonal Antibody in Quasi-physiological Conditions

Fatemeh Hasheminia¹, Razieh Farazmand ^{2*}, Nazanin Gholampour Faraji², Mona Abbasi kalabeh¹, Aliakbar Haddad Mashadrizeh³, Farzaneh Rostami⁴

1. Pharmaceutical Research Group, Khayyam Bioeconomy Institute, Mashhad, Iran

2. Structural biology and Bioinformatics Research Group, Khayyam Bioeconomy Institute, Mashhad, Iran

3. Cell and Molecular Biotechnology Research Group, Institute of Biotechnology, Ferdowsi University of Mashhad, Mashhad, Iran.

4. Department of Biology, Faculty of Science, Ferdowsi University of Mashhad, Mashhad, Iran.

Background: The Cetuximab with brand Erbitux is one of the monoclonal antibodies that target epithelial growth factor-associated cancers (EGFR) and approved by FDA. So, today is using for a range of cancers such as colorectal, head and neck, lung and pancreatic carcinoma. So, achievement to the sequence and its modeling in quasi-physiology situation could be confirm the presented sequence point of view of structure and function, as well as opportunity for its optimization and development which are considered in this study.

Methods: In this study, the Protein Atlas database was used to determine the level and tissues expressing-EGFR. The NCBI, and RCSB provided the sequence and protein structure of the drug and the antigen. Antigen modeling and assembling the fragment sequences of drug were performed by using Modeller software by homology modeling method. ERRAT, Verify 3D and RAMPAGE programs were used to determine the quality of the 3D structures based on amino acid location. Assessment the stability of drug in quasi-physiological condition was performed with Gromacs at 37 ° C. Moreover, the functionality of this drug was evaluated via its affinity to corresponding antigen in comparison to other antigens as control. The PYMOL software was used to visualize structures and interactions.

Results: Assessment the expression of EGFR has led to reveal the high expression of this antigen on the surface of cancer cells of Head and neck, glioma, kidneys, skin and liver. Surveying the sequence of the Cetuximab reveal the presence of the variable light and heavy chain of antibodies with disulfide bonds, which are derived from mice and humans. Structural modeling of this drug has led to reveal the suitable quality of it based on the location of amino acid as well as its stability at 37 ° C during 20 nanosecond after assembling. Moreover, affinity of this drug to EGFR showed significantly difference in comparison to other antigens.

Conclusion: In conclusion, the obtained results of this study confirm the structure and function of the model of the Cetuximab sequence which is available in database. Moreover these results provide opportunity for Cetuximab optimization as well as immunotoxin development based on fusion to cytotoxic domains which is agenda of our group.

Keywords: Cancer, monoclonal antibody, Cetuximab, EGFR, *in-silico* biology

ICIA2018\Monoclonal Antibody Diagnostic and Therapeutic\Poster Discussion\8565
8565

Expression and Optimization of Anti-HER2 scFv in *E.coli*

Farzaneh Farshdari¹, Elham Mohit², Maryam Ahmadzadeh²

1. Department of Biotechnology, Faculty of Advanced Sciences & Technology, Pharmaceutical Sciences Branch, Islamic azad University, Tehran- Iran (IAUPS)

2. Department of Pharmaceutical Biotechnology, School of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: The human epidermal growth factor receptor (HER) family plays an important role in many different kinds of cancer. Relationship between expression of human epidermal growth factor 2 (HER2) and malignity of breast tumors has been resulted in the production of monoclonal antibodies targeting HER2⁺ tumors. Single chain variable fragment (scFv), the smallest functional region for antigen binding, has improved ability to penetrate into tumors while retaining specific affinity and having low immunogenicity. Particularly, anti-HER2 scFv can efficiently target the therapeutic proteins, drugs, and nanoparticles toward HER2⁺ cancer cells. The main aim of this study is to express anti-HER2 scFv protein in prokaryotic system, to determine protein solubility and to evaluate the effect of host type, induction temperature, and inducing conditions on the amount and solubility of the anti- HER2 scFv protein.

Methods: The synthesis of scFv against HER2 gene was cloned in expression vector pET22 b (+) and its expression was investigated in the BL21 (DE3) and Origami. Furthermore, different concentrations of IPTG, induction period and induction temperature were examined in the BL21 (DE3). The effect of temperature on the solubility of this protein was also considered. Finally, the expression of anti HER2-scFv protein was evaluated using Western blot analysis.

Results: Western blot analysis confirmed the correct expression of anti HER2-scFv protein. Strain BL21, was used for optimizing anti HER2-scFv protein. Totally, the optimum condition for the expression of anti HER2-scFv protein in the BL21 (DE3) was 37°C, 0.5 mM IPTG and 24 hours of induction

Conclusion: In this study anti HER2-scFv of Trastuzumab was expressed and optimized for the first time in BL21 (DE3) successfully.

Keywords: Breast cancer, scFv, Human Epidermal Growth receptor 2 (HER2)

ICIA2018\Monoclonal Antibody Diagnostic and Therapeutic\Poster Discussion\8713
8713

In-silico Assay of the Structure, Function and Modeling of ONTAK in Real-like Condition

Mona Abbasi kalabe¹, Nazanin Gholampour Farogi ^{*2}, Behrouz Jalali Ghassam¹, Aliakbar Haddad Mashadrizesh³, Ali Eshaghi⁴

1. Pharmaceutical Research Group, Khayyam Bioeconomy Institute, Mashhad, Iran

2. Structural biology and Bioinformatics Research Group, Khayyam Bioeconomy Institute, Mashhad, Iran

3. Cell and Molecular Biotechnology Research Group, Institute of Biotechnology, Ferdowsi University of Mashhad, Mashhad, Iran

4. Department of Biology, Islamic Azad University, Khorasan Razavi, Mashhad, Iran

Background: Today, the prevalence and fatality of cancer have led to develop smart drugs including immunotoxins. In 1999, ONTAK was introduced to the world market as the first immunotoxin which approved by the FDA, and presently it is utilized to treat the lymphoma disease. Hence, recognizing the properties of this drug in a quasi-physiological circumstance allows enhancing the function of it, which is on the agenda.

Methods: The sequences of drug were retrieved from the pharmaceutical database as well as the relevant patent. Then, its structural features were examined by computational methods. Subsequently, modeling and assembling of fragments were achieved by using Modeller software based on homology modeling method. In addition, evaluating the structure quality of the models were determined using ERRAT and RAMPAGE tests. Moreover, the structural simulation of the drug model was performed to confirm its stability in pseudo-physiological conditions. AutoDock and IEDB software were used for analyzing of affinity and immunogenicity of model, respectively.

Results: The obtained results from molecular investigation of desired drug demonstrated the presence of a short length form of diphtheria toxin and human IL-2 as fatal part and fusion domain of this immunotoxin, respectively. Moreover, it also revealed the flexible glycine-rich linker in its structure. In addition, assembling of these fragments was resulted in providing an optimal quality model from the point of view of the amino acids location in the proper position as well as structural stability and immunogenicity at real-like condition at 37 ° C for 20 k ps. Furthermore, evaluation of drug affinity to IL-2R as target antigen showed a high binding affinity related to negative control.

Conclusion: Generally, the results of this study revealed a model of ONTAK with an optimal structure and function, which could be an appropriate choice for synthesis of this drug as well as opportunity for its optimization.

Keywords: Cancer, Smart drugs, Immunotoxin, ONTAK

ICIA2018\Monoclonal Antibody Diagnostic and Therapeutic\Poster Discussion\9743
9743

Monoclonal Antibody Production Against Human CD105 (endoglin)

Esmail Jarrahi¹, Mohammad Reza Gharaati², Mehdi Alikhani³, Masoud Nafari⁴

1. Department of Molecular and cellular Biology Faculty of Basic Sciences and Advanced Technologies in biology University of Science and Culture, Tehran, Iran

2. Department of Stem Cells at Royan Institute

3. Department of Stem Cells at Royan Institute

4. Department of Molecular and cellular Biology Faculty of Basic Sciences and Advanced Technologies in biology University of Science and Culture, Tehran, Iran

Background: CD105 is one of the most important angiogenesis factors. Pervious investigations revealed that CD105 protein had considerable expression during metastasis tumors angiogenesis in comparison with before. Also, CD105 is considered as the main marker in mesenchymal stem cells. Mesenchymal stem cells are one of the most basic cells and have differentiation capability to other cells. Therefore, detecting this protein plays an important role in research and treatment. Producing and applying monoclonal antibodies is significant in detecting and treating.

Methods and Results: In order to produce monoclonal antibody against human CD105, firstly the sequences being capable of using as epitope were extracted by bioinformatics, then entered expressive system as a gene construct. Expressed protein was injected to Balb-c mouse as timetable. Following, the spleen of mouse was extracted. Fusion process was carried out by mouse myeloma cell hybridation and extracted cells from mouse spleen using polyethylene glycol. For creating monoclonal characteristics, limiting dilution was carried out make the developed cells have equal lineage. Finally, the validity of this valuable molecule was proved by conducting molecule confirmation tests.

Keywords: Monoclonal antibodies, CD105, Mesenchymal stem cell



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Mucosal and Gastrointestinal Immunology

Poster Discussion

ICIA2018\Mucosal and Gastrointestinal Immunology\Poster Discussion\5273

5273

IRF4-dependent Upregulation of Interleukin 9 Protein is Associated with Ulcerative Colitis PathogenesisMojtaba Shohan¹, Hedayatollah Shirzad²

1. Department of Microbiology and Immunology, Faculty of Medicine, Shahrekord University of Medical Sciences, Shahrekord, Iran

2. Cellular and Molecular Research Center, Basic Health Sciences Institute, Shahrekord University of Medical Sciences, Shahrekord, Iran

Background: Ulcerative colitis is a type of inflammatory bowel diseases histologically characterized by chronic inflammation in the large intestine. Disease occurrence is dependent on the combination of environmental factors and genetical predisposition. Secretion of inflammatory mediators and immune system hyperactivity leads to sustained inflammation in the colon and it can be life threatening if remains untreated. Our aim was to measure the mRNA expression of IRF4 and protein level of IL-9 by means of real time PCR and ELISA in patients with UC, respectively.

Methods: 20 patients with ulcerative colitis and 21 healthy subjects were enrolled in the study. mRNA expression of IRF4 gene was determined by sybr green master mix detection method and real-time polymerase chain reaction (RT-PCR). Also protein contents of tissues were extracted and interleukin 9 protein level was measured by enzyme linked immunosorbent assay.

Results: Quantitative measurement of IL-9 protein and mRNA expression of IRF4 showed notable increases in patient's samples in comparison to healthy subjects.

Conclusion: Interleukin 9 is a multifunctional protein, which has been recently reported to be involved in the pathogenesis of autoimmune diseases and also experimental models of UC. Interferon regulatory factor 4 is a key regulator transcription factor of IL-9 production, which also targeted recently in various autoimmunities. As our data demonstrates, interleukin 9 protein is significantly upregulated in the inflamed lesions of UC patients so it may be a remarkable point of research in the novel therapeutic approaches for IBD.

Keywords: Ulcerative colitis, IL-9, IRF4

ICIA2018\Mucosal and Gastrointestinal Immunology\Poster Discussion\8252

8252

The Association Study of TNF α Gene Polymorphism -1031 C>T, -376 G>A with Susceptibility to Celiac Disease in Iranian population

Zohreh Nasiri¹, Abdolrahim Nikzamir¹, Majid Sirati Sabet¹, Vahid Chaleshi³, Elham Aghamohammadi², Mohammad Reza Zali³, Mohammad Rostami Nejad³

1. Biochemistry Department, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Basic and Molecular Epidemiology of Gastrointestinal Disorders Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

3. Gastroenterology and Liver Diseases Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: TNF α is an important cytokine in inflammatory responses and plays an important role in the pathogenesis of autoimmune disorders such as celiac disease (CD). Regarding the position of TNF α gene in the original set of tissue constructs and in view of the role of TNF α in the pathogenesis of celiac disease, our goal in this study was to determine the relationship between -1031 C / T and -376 G / A TNF α polymorphisms with susceptibility to celiac disease in correlation with its serum level in this population compare with healthy people.

Methods: 104 patients with celiac disease whom their disorder confirmed by serological and pathological test and 102 control people were included in this study. DNA of both groups were extracted and gene polymorphisms was determined using PCR-RFLP technique. Also serum concentration of TNF α was evaluated ELISA method was used to determine the.

Results: The-1031 C / T and -376 G / A polymorphisms TNF α gene were not significantly correlated with BMI, age and sex. No significant difference was observed between CD patients and healthy controls for -1031 C / T (p=0.0566 and p=0.608 respectively) and -376 G / A gene polymorphisms (p= 0.8256 and p=0.806 respectively). Also no significant difference in serum concentration of TNF α was detected between celiac patients and healthy subjects (p= 0.189).

Conclusion: 1031C/T and -376G/A were not associated with CD nset in the Iranian population. This study reinforces the importance of other variants of TNF α in unraveling the pathogenic mechanisms of CD, therefore further genetic studies with larger number of patients are needed.

Keywords: Celiac disease, PCR-RFLP, TNF α gene polymorphism, autoimmune and inflammatory reaction

ICIA2018\Mucosal and Gastrointestinal Immunology\Poster Discussion\8272

8272

The Correlation between Serum Level of IL-15 and Abnormal Histopathology Degrees in Iranian Patients with Celiac Disease

Elham Aghamohamadi¹, Parviz Kokhaei¹, Fateme pak¹, Kamran Rostami², Afshin Moradi³, Mohamad Amin Pourhoseingholi³, Vahid Chaleshi³, Mohammad Reza Zali⁴, Mohammad Rostami Nejad³

1. Immunology Department, School of Medicine, Semnan University of Medical Sciences, Semnan-Iran
Milton University Hospital, UK

2. Basic and Molecular Epidemiology of Gastrointestinal Disorders Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

3. Gastroenterology and Liver Diseases Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Celiac disease (CD) is a intestinal *autoimmune disorder* caused by the ingestion of wheat, barley and rye (gluten) in *people* who are *genetically predisposed*. Different studies showed that IL-15 is a key regulator in celiac disease immunopathology. Therefore, in this study we investigated the correlation between serum level of IL-15 and abnormal histopathology degrees in Iranian patients with celiac disease.

Methods: Serum samples were collected from 51 confirmed CD patients including 12 patients with Marsh I, 14 with Marsh II and 25 with Marsh III during 2014. Serum concentrations of IL-15 determined by enzyme-linked immunosorbent assay (eBioscience, Affymetrix, Inc). Demographic and clinical data were collected by questionnaire and the result were analyzed using SPSS version 21.

Results: The mean age of subjects in this study was 38.16 years. The mean serum concentration of interleukin 15 in patients with Marsh I was 102.738, March II 482.526 and in March III 344.288. The serum concentration of IL-15 showed the greatest increase in March II but were not significantly different between three groups. The serum concentration of IL-15 showed the greatest increase in March II but were not significantly different between three groups.

Conclusion: Our results indicate that there are no significant correlate between serum concentration of IL-15 and abnormal histopathology degrees. These findings provide new insights into the IL-15 signaling pathway and the pathogenesis of celiac disease.

Keywords: Celiac disease, IL-15, Marsh classification

ICIA2018\Mucosal and Gastrointestinal Immunology\Poster Discussion\8352

8352

Comparison of Dextran Sodium Sulfate Different Doses effects in the Induction of Chronic Colitis in C57BL /6 Mice

Maryam Heidari ¹, Seyed Mahmoud Hashemi^{1,4}, Sedigheh Pouya¹, Kaveh Baghaei², Mohammad Reza Zali³

1. Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Basic and Molecular Epidemiology of Gastrointestinal Disorder Research Center, Research Institute for Gastroenterology and Liver Disease, Shahid Beheshti University of Medical Science, Tehran, Iran

3. Gastroenterology and Liver Disease Research Center, Research Institute for Gastroenterology and Liver Disease, Shahid Beheshti University of Medical Science, Tehran, Iran

4. Department of Applied Cell Sciences, School of Advanced Technologies in Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: inflammatory bowel disease (IBD) is a multifactorial disease in two general title of Crohn's disease and ulcerative colitis. Animal models have provided extensive tools for identifying the etiology of the disease. Several methods, including the use of chemical compounds, have been introduced to induce experimental colitis. In this study, in order to find an optimal dose of dextran sodium sulfate (DSS), C57BL /6 mice strain were used to induce colitis and the severity of clinical sign and survival, caused by different doses of 1.5, 2 and 3% DSS, were evaluated.

Methods: In order to induce chronic colitis, the DSS solution administered in drinking water of mice for 4 days and continued with 7 days of normal water intake. This process repeated in 3 cycles. Clinical symptoms of colitis in mice included, body weight loss, bleeding, diarrhea and stool consistency were daily monitored. The mice sacrificed on day 34 and the colon changes were analyzed.

Results: In 3% DSS receiving-group, a high incidence of mortality was observed with severe clinical symptoms, but no significant changes were recorded in the group which received 1.5% DSS in comparison to healthy control. Using 2%, DSS resulted in induction of colitis with moderate symptoms and no mortality in mice.

Conclusion: According to the results, using 2% DSS, can be suggested as an optimal dose for induction of colitis in mice.

Keywords: Inflammatory bowel disease, Animal model, Experimental colitis, Dextran sodium sulfate.

ICIA2018\Mucosal and Gastrointestinal Immunology\Poster Discussion\8363

8363

Association of TNF α and IL1 β Serum Levels, the Presence of Helicobacter and Peptic Ulcer in the Population of Northern Iran

Mehdi Tourani, Maryam Habibzadeh, Javad Shokri Shirvani, Hamidreza Nouri

1. Cellular and Molecular Biology Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, Iran

2. Department of Internal Medicine, Babol University of Medical Sciences, Babol, Iran

3. Department of Immunology, School of Medicine, Babol University of Medical Sciences, Babol, Iran

Background: Peptic ulcer is a major cause of death all around the globe. Inflammatory cytokine such as TNF α and IL1 β is one of the most important immune response to gastric infection. Regarding the significance of these cytokines in peptic ulcer and high prevalence of the disease in northern Iran, the relationship of TNF- α and IL-1 β with peptic ulcer and presence of H. pylori was conducted by measuring the serum level of said cytokines.

Methods: Subjects were divided into two groups of 50 people with ulcers and without ulcers. Blood and biopsy samples were taken from each person. Each group were measured primarily in terms of the presence of H.Pylori by measuring the serum level and the presence in tissue. Then both groups were compared in terms of serum levels of IL1 β and TNF α using standard ELISA kit.

Results: 56% of the subjects were infected with H. pylori. The infection was significantly higher in PUDs in comparison to NPUDs. The mean serum level of TNF- α in PUDs and H.Pylori+ subjects was higher than that of NPUDs and H.Pylori-, respectively. However, no significant difference of IL1 β serum level was observed between PUDs and NPUDs as well as between H.Pylori positive and negative.

Conclusion: the findings of this research indicate that increase of the TNF- α cytokine is an important risk factor in intensification of peptic ulcer.

Keywords: TNF-a, IL1 β , Helicobacter Pylori

ICIA2018\Mucosal and Gastrointestinal Immunology\Poster Discussion\8505

8505

Evaluation of the Expression of CXCR2 and CXCR6 in Animal Models of Ulcerative Colitis Induced by Acetic Acid

Poorya foroutan pajoohian^{1*}, Parinaz mahdavi¹, Mohammad Amin Boshagh¹, Ali Jalili², Shohreh Fakhari²

1. Student Research Committee, Kurdistan University of Medical Sciences, Sanandaj, Iran

2. Department of Immunology, Faculty of medicine, Kurdistan University of Medical Sciences, Sanandaj, Iran

Background: The IBD (Inflammatory Bowel Disease) is classified into two types, which include Ulcerative Colitis (UC) and Crohn's Disease (CD). The pathology of these diseases suggests the prevalence of neutrophil infiltration in ulcerative colitis and the dominant infiltration of mononuclear cells in Crohn's disease. CXCRs are a series of chemokine receptors which are expressed on the surface of leukocytes. Various types of these receptors have been identified, which are categorized nearly from CXCR1 to CXCR7, Each is attached to a specific chemokine and goes to the inflammation site in a specific tissue. Therefore, the aim of this study was to evaluate the expression of CXCR2 and CXCR6 in animal models of ulcerative colitis induced by acetic acid.

Methods: In this experimental study, 6 male Wistar rats were randomly divided into two groups: control and treated groups. In the treated group, colitis was induced by rectal injection of acetic acid. After 48hours, the animal was anesthetized and then the mucosa samples of its Colon were separated in order to consider the histopathological indices and Polymerase Chain Reaction.

Results: based on PCR results the expression of these chemokine receptors in the treated group has changed in comparison with the control group. CXCR2 up regulated but CXCR6 down regulated in treated group in comparison with the control group.

Conclusion: According to the information obtained from this study, the likely expression of these receptors in the animal model of the ulcerative colitis has changed. Given the fact that this study is part of my thesis, the final confirmation, and publication of exact information, require more samples, as well as quantitative techniques such as Real-Time PCR, as well as verification methods such as Immunohistochemistry. Finally, in the current study, we intend to evaluate the expression of the whole CXCRs in this model.

Keywords: Inflammatory bowel disease, Ulcerative colitis, Chemokine receptors, Acetic acid

ICIA2018\Mucosal and Gastrointestinal Immunology\Poster Discussion\9925
9925

Evaluation of miRNA-146a and miRNA-193a Expression in Gastric Cancer cell lines

Amir Baghbanzadeh^{1,2}, Elham Baghbani², Saeed noorolyai^{1,2}, Ahad Mokhtarzadeh^{2,1}, Behzad Baradaran^{*2}

1. Department of Biochemistry, Higher Education Institute of Rab-Rashid, Tabriz, Iran

2. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Gastric cancer is the second most popular reason of malignancy dependent death through the world. MicroRNA is short non-coding RNA with 18 to 22 bp. Many studies evidenced that miRNAs such as miRNA-146a & miRNA-193a, have an important role in gastric cancer development. The miRNA-146a and miRNA-193a act as tumor suppressor in gastric cells and the evidence demonstrated that these miRNAs act as oncomir in gastric cancer cells. In this study, we evaluated the expression of miRNA-146a & miRNA-193a in three cell lines: KATO III, MKN45 and AGS.

Methods: in this study, KATO III, MKN45 and AGS cell lines were cultured in RPMI 1640 with 10% FBS and 5% of CO₂ condition. Total RNA was extracted by Trizol kit and cDNA was synthesized. Real-time PCR was used to measure changes in the expression of miRNA-146a & miRNA-193a genes. We used U6-miRNA as an internal control.

Results: Real-time PCR data indicated that the expression level of both miRNA-146a & 193a in KATO III is less than two other cell lines. The expression of miRNA-146a in AGS cell line is less than MKN45. The expression of miRNA-193a in MKN45 cell line was shown low level compared to AGS.

Conclusion: according to our finding, the expression of following miRNAs in KATO III is less than other cell lines.

Keywords: gastric cancer, microRNA, migration, replacement therapy

ICIA2018\Mucosal and Gastrointestinal Immunology\Poster Discussion\9930
9930

Evaluation of TLR 8 mRNA Expression in Peripheral Blood Mononuclear Cells and Association with Clinical Outcome of IBD Patients

Vahid Chaleshi¹, Mahyar Nourian¹, Hamid Asadzadeh Aghdaei¹, Shabnam Shahrokh¹, Mohammad Reza Zali².

1. *Basic and Molecular Epidemiology of Gastrointestinal Disorders Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran*

2. *Gastroenterology and Liver Diseases Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran*

Background: Ulcerative colitis (UC) and Crohn's disease (CD) are two major type of inflammatory bowel disease (IBD). Toll-like receptors (TLRs) play a critical role in the innate immune response to invading pathogens by sensing microorganism. Toll like receptors family are a member of immune system, which could identify a wide range of microorganism. The aim of present study was to evaluate expression of TLR8 between both groups of patients in different phases called remission and flare up and comparison of these by control group.

Methods: The present study 42 patients with UC, 6 patients with CD and 20 controls were included. RNA extraction from PBMC and reverse transcriptase PCR used. To evaluate the expression of target gene, qRT-PCR method was applied. Statistical analysis was performed using One-way ANOVA, Graph the chi-square test.

Results: The assessment of expression of TLR8 in CD and UC patients and control groups exposed no difference ($P=0.4936$). In addition, comparison of flare-up and remission groups with controls illustrate no significant change ($P=0.5419$). Likewise, no correlation exists in expression of TLR8 and five different groups of drugs which involved in the present study ($P=0.3703$).

Conclusion: Expression of TLR8 gene in IBD has not any significant relationship with control, so we cannot consider this as a biomarker for prognosis of IBD.

Keywords: Inflammatory Bowel Diseases, Crohn Disease, Colitis, Ulcerative, Gene Expression, Toll-Like Receptors.

ICIA2018\Mucosal and Gastrointestinal Immunology\Poster Discussion\9942
9942

Evaluation of KCNQ1 Level in CRC Tumors and Its Association with Clinical Features

Elaheh azizi¹, Sedigheh Arbabian², Ehsan Nazemolhosseini Mojarad³

1. Department of cellular and molecular, North Tehran Branch, Islamic Azad University, Tehran, Iran

2. Department of cellular and molecular, North Tehran Branch, Islamic Azad University, Tehran, Iran

3. Gastroenterology and Liver Diseases Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Colorectal cancer (CRC) is the third known cancer worldwide that has been mostly diagnosed at advanced stages. Therefore, finding new biomarkers with marked sensitivity and specificity is matter of urgent. Studies show that alteration of KCNQ1 expression pattern is effective in the initiation and progression of tumorigenesis. Therefore, we evaluated the level of this lncRNA within the CRC patients.

Methods: 52 tumor tissue samples and 10 normal controls were collected along with from the cases that had been diagnosed and approved by the Taleghani Hospital, Tehran, Iran. Samples were subjected to RNA isolation and then reverse transcript. The obtained cDNAs were amplified by Real time-PCR method. Relative expression abundances of KCNQ1 were measured by normalizing to Ribosomal 18S RNA using the $2^{-\Delta\Delta CT}$ method.

Results: The data showed a in the expression pattern of KCNQ1 within the tumor samples compared to the controls. These observations were statistically associated with patient TNM stage, tumor location and differentiation.

Conclusion: Taking together, increasing of the KCNQ1 provokes tumorigenesis in CRC patients. So, this lncRNA could be applied as diagnosis biomarker for cancer detection.

Keywords: KCNQ1, Colorectal cancer, Diagnosis biomarker, Tumorigenesis

ICIA2018\Mucosal and Gastrointestinal Immunology\Poster Discussion\9958
9958

The Relationship between the Expression of PTENP1 LncRNA in People with Colorectal Cancer and its Clinical Relevance

Niloofer Sadat Ebrahimi¹, Ehsan Nazemalhosseini Mojarad²

1. *Tehran shomal University, Tehran, Iran*

2. *Department of Pathology, Shohada Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran*

3. *Gastroenterology and Liver Disease Research center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran*

4. *Tehran shomal university, Tehran, Iran*

Background: Colorectal Cancer (CRC) is the third most frequent neoplasm worldwide, that being more frequent in developed countries than in developing countries. Long non-coding RNAs (lncRNAs) are defined as transcripts longer than 200 nucleotides that are not translated into protein. Among them, PTENP1 is considered as a marked tumor suppressor, which decrease cancer progression by increasing PTEN stability and PI3K/Akt signaling pathway depression.

Methods: 52 fresh tumor tissues plus 10 normal controls were obtained from the Taleghani Hospital, Tehran, Iran. The RNA were isolated/purified and reverse transcribed. The yielded cDNAs were assessed using qPCR technology. Relative expression abundances of targets were determined by normalizing to Ribosomal 18S RNA using the $2^{-\Delta\Delta CT}$ method.

Results: The data showed that PTENP1 was markedly downregulated in tumors comparing to normal samples. This decrement was statistically associated with patient TNM stage, tumor location and differentiation.

Conclusion: The alteration in the expression pattern of PTENP1 is assessable during CRC development, and therefore could be used as diagnosis biomarker for cancer detection.

Keywords: Colorectal cancer, Long none coding RNAs, PTENP1, Tumor suppressor

ICIA2018\Mucosal and Gastrointestinal Immunology\Poster Discussion\9996
9996

Evaluation of *lncRNA IFNG-AS1* Expression Level with Clinical Outcome of IBD Patients

Fariba Rashid¹, Vahid Chaleshi², Gholamreza Javadi¹, Hamid Asadzadeh Aghdaei²

1. Department of Biology, Islamic Azad University, Science and Research Branch, Tehran, Iran

2. Basic and Molecular Epidemiology of Gastrointestinal Disorders Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Inflammatory bowel disease (IBD) causes chronic and relapsing gastrointestinal inflammation that classified into Crohn's disease and ulcerative colitis. lncRNAs that can affect the expression of gene products. LncRNAs are a family of non-coding RNAs that dysregulation in their expression involves the pathogenesis of IBD. In this study, the expression of *lncRNA IFNG-AS1* was examined.

Methods: In this case-control study, 15 Crohn's patients, 20 ulcerative colitis and 20 controls who referred to the Liver and Gastro intestinal Research Institute RNA isolated from tissue samples using by AllPrep DNA/RNA/miRNA Universal kit (QIAGEN). cDNA was synthesized using by Thermo Scientific RevertAid RT Kit. Quantitative Real Time PCR (qRT-PCR) was used to evaluate the expression of the *lncRNA IFNG-AS1* gene (BioFact™ 2X Real-Time PCR Smart mix Sybergreen-Rotator gene Q MDX QIAGEN).

Results: Based on the findings, there was a significant difference between the CD, UC and control groups (P value = 0.0017). Also, significant difference between *IFNG-AS1* gene expression and phases of disease (Remission, Flare up) and control groups was absorbed (p value = 0.0028). However, expression level of *IFNG-AS1* in male and female groups have been assessed and there were no significant differences (P>0.05).

Conclusion: According to our result level of *IFNG-AS1* gene expression may be considered as a prognostic factor in the susceptibility to inflammatory bowel disease and also may be as a biomarker to distinguish between the CD and UC group.

Keywords :Inflammatory bowel disease (IBD), *IFNG-AS1*, Crohn's disease, ulcerative colitis, LncRNA



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Nanoimmunology

Poster Discussion

ICIA2018\Nanoimmunology\Poster Discussion\6321

6321

Synthesis and Characterization of Novel Nickel-Salen Supported Paramagnetic Nanoparticles for 6×His-tag Recombinant Protein Affinity PurificationZahra Rashid¹, Ramin Ghahremanzadeh¹, Amir Hassan Zarnani^{2, 3}*1. Nanobiotechnology Research Center, Avicenna Research Institute, ACECR, Tehran**2. Iran.Reproductive Immunology Research Center, Avicenna Research Institute, ACECR, Tehran, Iran**3. Immunology Research Center, Iran University of Medical Sciences, Tehran, Iran*

Background: Affinity separations based on affinity tags that exploit the unique property of extremely specific biological interactions to achieve separation and purification is theoretically capable of giving robust purification, even from complex mixtures in a single process. In this research, a simple, efficient, inexpensive, rapid and high yield method for the purification of 6×histidine-tagged recombinant protein was developed.

Methods: Manganese ferrite magnetic nanoparticles (MNPs) were synthesized through a co-precipitation method and then they were conveniently surface modified with tetraethyl orthosilicate (TEOS) in order to prevent oxidation and form high density of hydroxyl groups. The salen ligand was then prepared from condensation reaction of salicylaldehyde and 3-aminopropyl (trimethoxy) silane (APTMS) in 1:1 molar ratio; followed by complexation with Ni (OAc) 2.4H₂O. Finally, the prepared Ni (II)-salen complex conjugated to silica-coated MNPs and MnFe₂O₄@SiO₂@Ni-Salen complex nanoparticles were obtained. The prepared MNPs were used purification of 6×His-tag recombinant proteins.

Results: The functionalized nanoparticles were spherical with an average diameter around 70 nm. The obtained MNPs had a saturation magnetization about 54 emu/g and had super paramagnetic character. These MNPs were used efficiently to enrich recombinant histidine-tagged (His-tagged) protein-A from bacterial cell lysate. In about 45 minutes, highly pure His-tagged recombinant protein was obtained, as judged by SDS-PAGE analysis and silver staining. The amount of target protein in flow through and washing fractions was minimal denoting the high efficiency of purification process. The average capacity of the matrix was found to be high and about 180±15 mg. g⁻¹ (protein/ MnFe₂O₄@SiO₂@Ni-Salen complex).

Conclusion: Purification process with MnFe₂O₄@SiO₂@Ni-Salen complex nanoparticles is rapid, efficient, selective and whole purification can be carried out in only a single tube without the need for expensive systems.

Keywords: Magnetic nanoparticles, Protein purification, Immobilized metal affinity chromatography (IMAC), Histidine-tagged protein

ICIA2018\Nanoimmunology\Poster Discussion\6323

6321

Fabrication of Anti-CD4 Monoclonal Antibody-conjugated Magnetic Nanoparticles for Positive Selection of Peripheral Blood T CD4+ Lymphocytes

Zahra Rashid¹, Fazel Shokri¹, Mahmood Jeddi Tehrani², Amir Hassan Zarnani^{3,4}

1. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

2. Monoclonal Antibody Research Center, Avicenna Research Institute, ACECR, Tehran, Iran

3. Reproductive Immunology Research Center, Avicenna Research Institute, ACECR, Tehran, Iran

4. Immunology Research Center, Iran University of Medical Sciences, Tehran, Iran

Background: Magnetic-activated cell sorting (MACS) is a gentle and fast method for separation of various viable and functionally active cells by magnetic labeling. The present study aimed at synthesis of a novel MACS-based system for isolation of peripheral blood T CD+ lymphocytes.

Methods: Magnetic nanoparticles were first synthesized through a co-precipitation method from ferrous and ferric iron solutions and then SiO₂ shell was coated on the magnetic core with tetraethyl orthosilicate (TEOS) through a silanization reaction to prevent oxidation, agglomeration and expansion of the density of OH groups on the surface of Fe₃O₄. Subsequently, Fe₃O₄@SiO₂@PMIDA magnetic nanoparticles were obtained as a result of the reaction between N-(phosphonomethyl) iminodiacetic acid (PMIDA) and Fe₃O₄@SiO₂. Anti-CD4 monoclonal antibody was conjugated to the surface of Fe₃O₄@SiO₂@PMIDA magnetic nanoparticles through covalent binding between carboxylic group activated with 1-Ethyl-3-(3-dimethylaminopropyl)-carbodiimide)/ N-hydroxysuccinimide (EDC/NHS) on the particle surface and primary amine groups of the antibodies. The fabricated immunomagnetic beads were then utilized for isolation of CD4+ lymphocytes from peripheral blood.

Results: The generated immunomagnetic particles efficiently separated CD4+ lymphocytes from whole blood with purity of about 93±3 %, comparable to that of commercial MACS system.

Conclusion: The presented approach for fabrication of anti-CD4 magnetic nanoparticles is inexpensive, easy and efficient for isolation of T CD4+ lymphocytes. Based on the consistency of results, this platform might be applicable for separation of various cell population.

Keywords: MACS, Purification, Positive selection, Magnetic Nanoparticles, CD4

ICIA2018\Nanoimmunology\Poster Discussion\7813

7813

A Nanobody-based chimeric antigen Receptor Can Function as an Anti-angiogenesis by Targeting a VEGFR2 in Solid Tumor

Fatemeh Hajari Taheri¹, Mahmoud Hassani^{1, 2}, Arash Arashkia³, Zahra Sharifzadeh¹, Mehdi Behdani⁴, Mohsen Abolhassani¹

1. Department of Immunology, Hybridoma lab, Pasteur Institute of Iran, Tehran, Iran

2. Department of Molecular Medicine, Tehran University of Medical Sciences (TUMS), Tehran, Iran

3. Department of Virology, Pasteur Institute of Iran, Tehran, Iran

4. Department of Molecular Medicine, Pasteur Institute of Iran, Tehran, Iran

Background: Solid tumors account for more than 85% of cancer mortality. Since cancer cells able to form new blood vessels (angiogenesis), many attempts were made to prevent angiogenesis. VEGFR2 is a protein over expressed on the surface of metastatic cancer cells and is a target for immunotherapy of solid tumors. Chimeric antigen receptor (CAR)-T cells has shown promising results on solid tumors. In this study, we have developed a 3rd generation nanobody based CAR T cell targeting tumor cells.

Methods: The CAR-T cell was developed using VHH against VEGFR2 that is linked to IgG1 Fc domain as spacer, CD28 and CD3 ζ signaling domains, and co-stimulatory domain of OX-40. T cells were isolated from healthy donor blood and activated with CD3/CD28 beads in the presence of rIL-2 (20U/ml). Activated T cells were electroporated with the vector encoding the VEGFR2-CAR and then biological activity were evaluated *in vitro*.

Results: The surface expression of activation markers, CD69 and CD25 were measured upon coculturing with VEGFR2 expressing 293-KDR tumor cells (50% and 40% respectively), as compared with non-expressing HEK-293 as negative control. IL-2 and IFN- γ productions in response to VEGFR2-expressing target cells were 1200 and 900 pg/ml respectively. Cell surface expression of degranulation marker, CD107a, was evaluated on engineered T cells upon coculturing with VEGFR2 expressing 293-KDR cells (32% versus 6% of negative cells). Although the CD107a assay does not directly measured target cell lysis, it does provide an indication of the cytotoxic potential of the responding CD8⁺ T cells.

Conclusion: For the first time we constructed a nanobody-based anti-VEGFR2-CAR T cells. VHH is preferred to scFv because of its small size, low immunogenicity, enhanced stability, and penetration into the dense solid tumors. Therefore, this engineered VEGFR2-CAR T cells may be developed for the adoptive T-cell immunotherapy of solid tumors.

Keywords: CAR T cells, VEGFR2, Angiogenesis, Nanobody, VHH, Adoptive transfer

ICIA2018\Nanoimmunology\Poster Discussion\8516

8516

Encapsulation of Tumor cell Lysate in PLGA Nanoparticles and Evaluation of Their Proliferative Activity in Mice

Sanaz Sheikhzadeh¹, Nowruz Delirezhi¹, Rahim Hobbenaghi²

1. Department of Microbiology, Faculty of Veterinary Medicine, Urmia University

2. Department of Pathology, Faculty of Veterinary Medicine, Urmia University

Background: Breast cancer is the second leading cause of cancer death in women following lung cancer. Nanoparticles-based delivery systems can be used to facilitate the cancer therapy targeting immune system by enhancing the humoral and cellular immune responses. Encapsulation of tumor antigens in biodegradable nanoparticles increases its immunostimulatory capacity than when tumor antigens are given in the soluble form. Poly (lactic-co-glycolic acid) (PLGA) is one of the most developed biodegradable polymers that use for preparation of nanocapsule drugs. In this study, we assessed Nanoparticles (NPs) size, polydispersity index and capacity of PLGA NP-based delivery system on cellular proliferation in mouse model.

Methods: Tumor cell lysate was obtained from Four cycles of freeze-thaw of murine 4T1 mammary tumor cells and then encapsulated in to PLGA copolymers (50% lactide: 50% glycolide) by double-emulsion solvent evaporation technique. Nanoparticles were characterized in terms of average particle size and size distribution using dynamic light scattering (DLS). The mice were challenged subcutaneously with 4T1 cells. After tumor growth, mice in the experimental groups were vaccinated with free or nano-encapsulated forms of breast tumor cell lysate and animals in the control groups received PBS. One week after the last immunotherapy, mice spleens were removed for MTT assay test.

Results: The tumor cell lysate loaded nanoparticles size was in the range of 200-300 nm with a polydispersity index of 0.3 (n=3). A significant increase in the numbers of immunocyte was found in mice that received PLGA nano-encapsulated form of tumor cell lysate in comparison with mice that received free tumor cell lysate.

Conclusion: Our results demonstrated that tumor cell lysate loaded PLGA nanoparticles promote proliferation of immune cells and could be used as a potential therapeutic approach for the breast cancer treatment.

Keywords: PLGA nanoparticles, Cellular proliferation, Tumor cell lysate, Breast cancer

ICIA2018\Nanoimmunology\Poster Discussion\8521

8521

Targeting of HMGA2 by siRNA-loaded Nanoliposome Induces Apoptosis and Metastasis in Gastrointestinal Cancer CellsAli Mohammadi^{1,2}, Behzad Mansoori^{1,2}, Behzad Baradaran¹*1. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran**2. Student Research Committee, Tabriz University of Medical Sciences, Tabriz, Iran*

Background: Considering the complex nature of gastrointestinal cancer, different methods including surgery, radiotherapy, and chemotherapy are considered for the treatment; however the explained methods couldn't effective in these patients the novel strategies including silencing of oncogenes by the safe delivery method could consider as a novel therapy in colorectal cancer. The aim of this study is to understand the silencing effect of nano-liposomes loaded HMGA2 siRNA on gastrointestinal cancers.

Methods: The small interfering RNA-lipoplexes prepared with DOTAP/Chol/DOPE and dissolved separately in butyl alcohol then the aqueous sucrose and siRNA solution prepared and added to butyl alcohol solution to form a monophasic solution. The size, polydispersity index (PDI), and zeta potential of nano-liposomes evaluated by Zetasizer analyzer. The silencing of the HMGA2 in cancer cells were evaluated by the qRT-PCR method. The effect of liposomes on cell cytotoxicity was evaluated by MTT assay. The migratory effect of the liposomes was evaluated by wound healing assay. To evaluate if this liposome could induce apoptosis the annexin/PI assay was performed.

Results: The result of size, polydispersity index (PDI), and zeta potential of nano-liposomes showed the liposome has less than 350nm size and 0.67 PDI and -79 zeta potential. The result of gene silencing showed the optimum condition of HMGA2 silencing was 80 pmolHMGA2 concentration and 48 hr after treatment in each cancer cell lines. The result of MTT assay showed silencing of HMGA2 in optimal condition could reduce the viability of the cancer cell in the three cell lines. The result of apoptosis assay showed more than 50% of the cell deaths are related to the apoptosis in all three cells. The gene expression evaluation confirms the apoptosis induced by inner pathway by inducing Caspase 3 and Caspase 9 expression. Also reduction in bcl2 expression confirms the activation apoptosis pathway in the treated cancer cells. The wound healing assay showed the suppression of cancer cell migration after treatment of this nano-liposome.

Conclusion: The result of this study showed the nano-liposomes loaded HMGA2 siRNA could have effects in the treatment of gastrointestinal cancers and it might have potential have more study in order to develop the new agent for the treatment of gastrointestinal patient base on target therapy.

Keywords: HMGA2, Nano-liposome, Gastrointestinal cancer

ICIA2018\Nanoimmunology\Poster Discussion\9871

9871

Preparation and Characterization of a Novel Nanobody against T-cell Immunoglobulin and Mucin-3 (TIM-3)

Vida Homayouni ¹, Mazdak Ganjalikhani-hakemi ², Abbas Rezaei ², Hossein Khanahmad ³, Mahdi Behdani ⁴, Fatemeh Kazemi Lomedasht ⁴

1. Acquired immunodeficiency research center, Isfahan University of Medical Sciences. Isfahan, Iran

2. Immunology Department, Faculty of Medicine, Isfahan University of Medical Sciences. Isfahan, Iran

3. Genetic Department, Faculty of Medicine, Isfahan University of Medical Sciences. Isfahan, Iran

4. Biotechnology Research Center, Biotechnology Department, Venom & Bio-therapeutics Molecules Lab, Pasteur Institute of Iran, Tehran, Iran

Background: As T-cell immunoglobulin and mucin domain 3 (TIM-3) is an immune regulatory molecule. Based on the unique features of Nanobodies, we aimed to construct an anti-TIM-3 Nanobody as an appropriate tool for manipulating immune responses for future therapeutic purposes.

Methods: We immunized a camel with TIM-3 antigen and then, synthesized a VHH phagemid library from its B cell's transcriptome using nested PCR. Library selection against TIM-3 antigen was performed in three rounds of panning. Using phage-ELISA, the most reactive colonies were selected for sub-cloning in soluble protein expression vectors. The Nanobody was purified and confirmed with a nickel-nitrilotriacetic acid (Ni-NTA) column, SDS-PAGE and Western blotting.

Results: Specific 15kD band representing for Nanobody was observed on the gel and confirmed with Western blotting. The Nanobody showed significant specific immune-reactivity against TIM-3 with a relatively high binding affinity.

Conclusion: Finally, we successfully prepared a functional anti-human TIM-3 specific Nanobody with a high affinity and an anti-proliferative activity on an AML cell line in vitro.

Keywords: Antibody, Nanobody, HcAbs, Phage Display, TIM-3

ICIA2018\Nanoimmunology\Poster Discussion\11071
11071

The effect of *Leishmania* cysteine peptidase A and B conjugated to PLGA on Nitric oxide production

H. Noormehr ¹, A. Zavarani Hosseini ^{1*}, S. Soudi ¹

1. Department of Immunology, School of Medical science, University of Tarbiat Modares

Background: Nanocarriers with optimized physicochemical and biological properties are taken up by cells more easily than larger molecules, so they can be successfully used as antigen delivery tool. Protein antigens have drawn a lot of attention from investigators working on leishmaniasis vaccines. *Leishmania major* contains many distinct cysteine peptidase genes which contain cathepsin L-like enzymes (CPA and CPB). Here, we examined the effect of CPA, CPB conjugated to PLGA nanoparticle on nitric oxide production by macrophages against *L.major* infection.

Methods: For this purpose *L.major* RNA were extracted and coding DNA of *cpa* and *cpb* genes were amplified by PCR method. Both *cpa* and *cpb* were sub-cloned in pET28a vector and expressed in *E.coli*. Produced rCPB and rCPA purified from inclusion bodies by Ni-NTA agarose affinity column. rCPA, rCPB were conjugated to PLGA nanoparticle. Morphology and Conjugation efficacy rCPA/ rCPB-PLGA were assessed by SEM and FTIR, respectively. 5 microgram of protein content of rCPA, rCPB, and their conjugated form were injected intraperitoneal to four different BALB/c mice groups respectively. After 24 hours, NO production of macrophages were measured in response to 10:1 *L.major* infection.

Results: No morphologically, difference were seen between CPB/ CPA – PLGA with non-conjugated form. CPA and CPB conjugation to PLGA were confirmed by FTIR method. Internalization of CPB/ CPA – PLGA were confirmed by FITC labeling of antigen. Both CPB/ CPA – PLGA treated groups showed a significant increase in the amount of NO production by macrophages in response to *L.major* compared with control groups.

Conclusion: Our results proposed that PLGA nanoparticles may be a promising adjuvant in the development of therapeutic leishmania vaccines. It seems that activated macrophages with nanoparticles lead to increase NO production resulting in inhibiting of leishmania major growth.

Keywords: *L.major*, cysteine peptidase A, B, PLGA, NO.



14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

Oral Immunology

Poster Discussion

ICIA2018\Oral Immunology\Poster Discussion\8178

8178

Evaluation of RANKL/OPG in Chronic Periodontitis Patients after Nonsurgical TherapyAbbas Azadmehr¹, Zahra Alizade Tabari², Niloufar Sadat Miri¹, Mohammad Amir Alizade Tabrizi²

1. Department of Immunology, Babol University of Medical Sciences, Babol, Iran

2. Department of Periodontics, Dental School, Qazvin University of Medical Sciences, Qazvin, Iran

Background: Periodontitis (PD) is an inflammatory disease that involves progressive loss of the alveolar bone around the teeth. The receptor activator of nuclear factor kappa B (RANK)/RANK ligand (RANKL)/osteoprotegerin (OPG) system signaling pathway is implicated in bone resorption through its key function in osteoclast differentiation and inflammatory response. The aim of the present study was to determine the alternation salivary concentration of sRANKL /OPG ratio in response to nonsurgical therapy.

Methods: Twenty-five patients with chronic periodontitis were selected based on clinical evaluation and twenty-five practically healthy volunteers selected as control group. Saliva samples were collected from patients and control group at first session and then the samples were recollected in periodontitis group four weeks after non-surgical periodontal therapy in order to measure the level of sRANKL and OPG by ELISA method.

Results: There was a significant difference between sRANKL and sRANKL/OPG concentration in patients before treatment and healthy group (Respectively, $P=0.004$, $P=0.001$). Also, the concentration of sRANKL and sRANKL/OPG in patients was significantly reduced as a result of treatment (Respectively, $P=0.012$, $P=0.011$). However, there was no significant difference between OPG concentration in healthy and patient groups ($P=0.455$).

Conclusion: As the ratio of sRANKL/OPG in saliva has a diagnostic value for chronic periodontitis, it seems to be clinically a good predictor of successful treatment outcomes. However, to validate these results, future studies with more samples are required.

Keywords: Saliva, Chronic Periodontitis, sRANKL , OPG

ICIA2018\Oral Immunology\Poster Discussion\8311

8311

Expression of NLRP3 in Aggressive Periodontitis

Shideh Mehrmofakham^{1*}, Mandana Sattari¹, Mohammadreza Talebi Ardakani¹, Sarira Shahnava¹,
Aram Mohammadi¹

1. Shahid Beheshti University of Medical Sciences, Tehran, IRAN.

Background: Periodontal diseases are poly microbial infections resulting in bone destruction and tooth loss. Host's response to these infections includes both mechanisms of innate immune responses and acquired immune responses. Bacteria and especially gram-negative bacteria play an important role in prevalence of periodontal diseases. Also, it is determined that some periodontopathic bacteria make inflammatory cells to express NLRP3 which is one of the most important parts of inflammasome. According to this, the current research tries to survey the expression of NLRP3 in generalized aggressive periodontitis.

Methods: Gingiva tissue samples were collected from 20 people having clinically healthy gingiva and 25 people having aggressive periodontitis. After RNA extraction and cDNA preparation from each sample the amount of expression of NLRP3 was surveyed by the Real-time PCR technique.

Results: Expression of NLRP3 in tissues with aggressive periodontitis was significantly higher than the expression of NLRP3 in healthy gingiva ($P \approx 0.034$), but we could not find any significant correlation between expression of NLRP3 and clinical parameters.

Conclusion: It is concluded that in aggressive periodontitis the increased expression and production of NLRP3 is observed which results in inducing release of inflammatory cytokines and aggravating inflammation in the periodontal tissues.

Keywords: NLRP3, Aggressive Periodontitis, Poly Microbial Infections, Healthy Gingiva

ICIA2018\Oral Immunology\Poster Discussion\8312

8312

Immunohistochemical Analysis of Visfatin Expression in Gingival Tissue of Periodontitis Patients

Niloufar Sadat Miri¹, Abbas Azadmehr¹, Zahra Alizadeh Tabari², Baharak Qa'edi², Forouz Keshani³, Marzieh Nejati Fard²

1. Department of Immunology, Babol University of Medical Sciences, Babol, Iran.

2. Periodontics Department, Dental School, Qazvin University of Medical Sciences, Qazvin, Iran.

3. Torabinejad Dental Research Center, Department of Oral and Maxillofacial Pathology, School of Dentistry, Isfahan University of Medical Sciences, Iran

Background: Visfatin is a pro-inflammatory cytokine that has been associated with several immunomodulation processes and there is little clinical information on the relationship between periodontopathogens and visfatin. However, studies on visfatin as a potential therapeutic target in periodontal diseases are scarce. In the present study, we analyzed the visfatin in gingival tissue of patients with generalized aggressive periodontitis.

Methods: 13 patients with aggressive periodontitis and 20 periodontal healthy individuals enrolled in this study based on the inclusion and exclusion criteria. Tissue samples were obtained in both groups during surgery. The degree of inflammation infiltration and visfatin expression were determined by H&E staining and Immunohistochemistry (IHC) methods respectively. The data was analyzed by SPSS software and $P < 0.05$ was considered statistically significant.

Results: Visfatin was detectable in all samples. Visfatin levels were higher in patients with aggressive periodontitis compared to those of healthy subjects ($P = 0.00$, $P = 0.037$). The relationship between inflammatory, cell infiltration and visfatin expression in patients group were statistically significant, positive and relatively strong ($P = 0.025$, $R = 0.617$).

Conclusion: Visfatin expression increased in gingival tissue of periodontitis patients group. These results demonstrated that the high expression level of visfatin in periodontitis tissues were correlated with the degree of tissue inflammation. However, more studies with larger sample sizes are necessary to validate these findings.

Keywords: Aggressive periodontitis, Inflammatory Cytokine, Visfatin, Immunohistochemistry

ICIA2018\Oral Immunology\Poster Discussion\8322

8322

Assessment of Non-surgical Periodontal Therapy on IL22 and S100A12 Concentration in Gingival Crevicular Fluid of Periodontal Patient

Ferial Taleghani¹, Mandana Sattari², Rokhsareh Sadeghi³, Razieh Nejadi⁴

1. Department of Periodontics, Dental School, Shahed University, Tehran, Iran.

2. Department of Immunology, Shahid Beheshti University of medical sciences, Tehran, Iran

3. Department of Periodontics, Dental School, Shahed University, Tehran, Iran.

4. Corresponding Author, Dental School, Shahed University, Tehran, Iran

Background: Periodontitis is a destructive and progressive inflammatory condition of tooth-supporting tissues, the pathogenesis of which is attributed to several risk factors, including bacteria, the host immune responses and genetics. The present study was undertaken to evaluate the effect of phase I periodontal treatment on the concentrations of IL-22 and S100A12 in patients with chronic periodontitis.

Methods: In this study, 22 patients (14 female and 8 male), with a mean age of 39 years, and moderate to severe periodontitis were sampled. phase I periodontal treatment was rendered. The subjects were recalled after 4 weeks for collecting samples. Wilcoxon's signed-rank test was used to analyze concentration of S100 and IL22 before and after treatment with CAL and PD.

Results: There was an inverse correlations between the mean PD and S100 before treatment ($P < 0.05$). In relation to CAL, although an inverse correlation was expected between CAL and S100 concentration, no significant correlation was found between them ($P \approx 0.079$).

Conclusion: The periodontal tissues are weak cellular sources for the secretion of IL-22 and S100. Given the inverse correlation between the concentration of S100 and the mean PD before treatment and also the inverse and significant correlation between the concentration of S100 and the decrease in CAL and PD after treatment, a possible protective role might be considered for S100 for the incidence of periodontitis and success of phase I treatment.

Keywords: Chronic Periodontitis – Periodontal Phase I Treatment - S100A12 – IL22

ICIA2018\Oral Immunology\Poster Discussion\8323

8323

Visfatin as an Inflammatory Marker in Periodontitis Patients

Maryam Mehri¹, Abbas Azadmehr¹, Niloufar Sadat Miri¹, Zahra Alizadeh Tabari², Baharak Qa'edi², Ali Nohekhan²

1. Department of Immunology, Babol University of Medical Sciences, Babol, Iran

2. Periodontics Department, Dental School, Qazvin University of Medical Sciences, Qazvin, Iran

Background: Visfatin is an adipocytokine that suggested as one of the periodontal inflammation regulators that increased in periodontitis and can affect patient's systemic condition. It is indicated that serum/plasma level of visfatin increase in a number of inflammatory disorders. The present study aim to comparison the salivary visfatin concentration in chronic periodontitis patients and healthy individuals.

Methods: saliva samples were collected at first session from 20 patients with chronic periodontitis (mean age: 38.45±9.98) and 20 healthy individuals (mean age 33.85±6.84) as control group which selected based on clinical evaluation. Then four weeks after non-surgical therapy, saliva sample were recollected from patients group in order to assay the visfatin by enzyme linked immunosorbent assay method (ELISA).

Results: our results showed, visfatin concentration mean in patients with chronic periodontitis before non-surgical therapy (33.43ng/ml±15.72ng/ml) were significantly higher than control group (23.38ng/ml±7.85ng/ml) (p=0.015) and decrease after therapy (23.00ng/ml±9.75ng/ml) (p=0.003). The level of visfatin in patient and control groups were same after non-surgical therapy.

Conclusions: Our findings indicated that salivary visfatin concentration decrease after non-surgical periodontal therapy. Also visfatin value may be consider as an "inflammatory marker" and can be detect in future as a potential therapeutic target in treatment of chronic periodontitis.

Keywords: Visfatin, Chronic Periodontitis, Non-surgical periodontal therapy, Biomarker, Saliva

ICIA2018\Oral Immunology\Poster Discussion\8360

8360

Association between gene expression of TRPV1 (Transient receptor potential vanilloid1) and IL8 (CXCL8) in pulp with superficial caries

Shaghayegh Pezeshki Naraghi¹, Mandana Sattari¹

1. Department of Immunology, School of medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: In view of the importance of pulpitis pathobiology and in order to assist innovation in its treatment methods, the present study focuses on investigating the role of superficial caries by expressing TRPV1(Transient Receptor Potential Vanilloid1) and IL-8 (CXCL8) genes.

Methods: The research is carried out by designing a cohort study on the pulp of 50 extracted premolar teeth divided into 2 groups as healthy (control) and surface caries. After extracting pulp and homogenizing the samples through quantitative real time PCR, the extent of gene expression for TRPV1 and IL-8 is separately calculated. Afterwards data gathered from each group is compared with case groups and with each other in pairs.

Results and Conclusion: Carrying out ANOVA test on 2 study groups in view of CXCL8 gene we obtain significant results ($P \approx 0.001$). Upon conducting Post Hoc Tukey test it becomes evident that there is a significant difference between healthy teeth and superficial caries groups ($P \approx 0.001$) and moderate caries & deeply complicated caries ($P \approx 0.012$) in such a way that in both cases the highest expression of CXCL8 gene is related to caries with involvement of the pulp.

Keywords: CXCL-8, TRPV-1, superficial caries

ICIA2018\Oral Immunology\Poster Discussion\8397

8397

RANKL Expression and GingivitisMandana Sattari¹, Niussha Amiri¹, Nikoo Hossein Khan Nazer^{1*} Mahdi Samani¹, Shideh Mehrmofakham¹*1. Department of Immunology, School of medicine, University of Shahid Beheshti*

Background: The tissue destruction found in periodontal disease results, for the most part, from actions of the immune system and related effector mechanisms. Although periodontal disease is initiated and sustained by microorganisms, especially bacteria in dental plaque, the progression and severity of tissue destruction are mostly caused by immune host defense mechanisms to infecting bacteria. Immune responses have a crucial role in the pathogenesis and tissue destruction associated with infectious disease. In spite of the existing evidence for the association of periodontitis with RANKL/OPG ratio, reports of the roles of RANKL in the pathogenesis of this disease are still controversial. So the present study was undertaken to assess the expression of RANKL in gingival tissue of patients with gingivitis.

Methods: Gingival biopsies were obtained from patients with gingivitis (n=17), as well as healthy subjects (n=15). RANKL expression were determined by using Real- time PCR (SYBR green Assay). GAPDH gene was used as housekeeping gene.

Results and Conclusion: We found that RANKL expression was significantly higher in the inflamed periodontal tissues (Gingivitis) compared to healthy controls. Furthermore, there was a positive correlation between RANKL and clinical attachment loss ($P<0.05$).

Keywords: Gingivitis, RANKL

ICIA2018\Oral Immunology\Poster Discussion\8555
8555

The Effectiveness of Acupuncture on Addiction Severity Index, a single case experimental design in a Case of Methamphetamine Abuser Patient with Trismus Syndrome

Bijan Pirnia¹, Dr Kambiz Pirnia², Zohreh Mortezaei³

1. Department of Psychology, Faculty of Humanities, University of Science and Culture, Tehran, Iran

2. Internal disease specialist, Technical Assistant in Bijan Center for Substance Abuse Treatment, Tehran, Iran

3. Payam Noor University of Dubai, UAE.

Background: Trismus is one of the side effects of methamphetamine using, which is associated with contractions of the jaw muscles. Acupuncture is a traditional and complementary treatment that is effective on pain and psychological indices. The purpose of this study was to investigate the effectiveness of the acupuncture in a case of trismus caused by methamphetamine abusing.

Methods: The patient was a 31-year-old man with a history of chronic methamphetamine use, reported to have dependence and severe jaw pain. In a single case study and in an ABAB design with multi baselines, we used acupuncture for three weeks to reduce pain and addiction severity index. Data were analyzed through generalized estimating equation.

Results: The results showed that there was no significant correlation between severity of addiction and pain ($all > 0.05$).

Conclusion: Due to damage in the process of production of tyrosine hydroxylase and mitogen-activated protein kinase and the role of these precursors in the production of dopamine as an effective factor in the acupuncture, the effectiveness of this treatment can be limited.

Keywords: Acupuncture, Stimulus, Trismus, Addiction

ICIA2018\Oral Immunology\Poster Discussion\8562

8562

Does TLR4 and Its Co-receptor Have Important Role in Pathogenesis of Aggressive Periodontitis?Hediyeh Ghalikhani¹, Sarira Shahnava¹, Aram Mohammadi¹, Mandana Sattari¹, Shideh Mehrmofakham¹*1. Department of Immunology School of medicine Shahid Beheshti University of medical sciences*

Background: Periodontal diseases are poly-microbial infections that cause bone destruction and tooth loss. Host responses to these infections involving both innate and adoptive immune responses. Moesin plays role in the attachment of cytoskeleton to the cell membrane. This protein has a variety of tasks, such as a new co-receptor for TLR4. It was also shown that in the case of moesin deficiency, the incidence of inflammatory responses to LPS fell three times. Regarding to the role of moesin in recognition of LPS by TLR4 and since there is not enough information about the role of moesin in periodontal disease, the aim of this study was to evaluate the relationship between moesin and TLR4 expression and aggressive periodontitis disease.

Methods: 25 samples for aggressive and 25 samples for healthy group 20 samples were collected (totally 50 gingival samples). After extracting the RNA, cDNA prepared from all of samples and for determining of the quality of the cDNA, PCR amplification was performed. Then Beta actin (Housekeeping gene) and Moesin and TLR4 gene expression was investigated by Real-time PCR technique.

Results: There was a significant lower expression of moesin in aggressive periodontitis compared to control group ($P \approx 0.025$). There was also a significant lower expression of TLR4 in aggressive periodontitis group compared to control group ($P \approx 0.013$).

Conclusion: It is concluded that recognition of periopathogens by TLR4 and moesin does not have an important role in aggressive periodontitis.

Keywords: Aggressive Periodontitis, TLR4, Moesin



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Psychoneuroimmunology and Immunoendocrinology

Poster Discussion

ICIA2018\Psychoneuroimmunology and Immunoendocrinology\Poster Discussion\7619
7619

Evaluation of Jak1, T-bet and Foxp3 Gene Expression in Major Depression Disorder

Marzie Askari ^{1,2}, Mohammad Haghighi ³, Iraj Salehi ⁴, Zohreh Zareighane ¹, Reza Shahbazi ³, Alireza Zamani ¹, Ghasem Solgi ¹, Mahdi Alahgholi Hajibehzad ^{1,*}

1. Department of Immunology, Hamadan University of Medical Sciences, Hamadan, Iran

2. Student Research Committee, Hamadan University of Medical Sciences, Hamadan, Iran

3. Research Center for Behavioral Disorders and Substance Abuse, Hamadan University of Medical Sciences, Hamadan, Iran

4. Neurophysiology Research Center, Hamadan University of Medical Sciences, Hamadan, Iran

Background: Major depression disorder (MDD) is a psychiatric illness that is associated with the immune system. The inflammation play a major role in development of MDD. The purpose of this study was the evaluation of Jak1, T-bet and Foxp3 gene expression in patients with MDD compare to healthy control subsets (HC).

Methods: Blood samples were collected from 40 MDD patients (mean age 38.5 ± 13.2) and 30 (40.5 \pm 12) HCs. Twenty patients were taken Selective Serotonin Reuptake Inhibitor (SSRI⁺) as a medication and 20 of MDD patients were new case (SSRI⁻). Total RNA was isolated from PBMC (Peripheral blood mononuclear cells) of both patients and HCs and cDNA was synthesized using reverse transcriptase kit. Expression of genes was performed with the SYBR green Real Time PCR method. The relative genes expression level for each sample were calculated by $\Delta\Delta C_t$ method.

Results: The relative level of Foxp3 was significantly increased in SSRI⁺ compared to SSRI⁻ group, whereas there was no significant difference in the Jak1 and T-bet expression. No significant difference was observed between HCs and SSRI⁺/SSRI⁻ groups separately in the expression level of T-bet, JAK1 and Foxp3. T-bet gene expression was slightly decreased in patients compared to HCs, whereas there were no significant changes in Foxp3 and Jak1 expression.

Conclusion: This result show a changed level of T-bet, JAK1 and Foxp3 expression in MDD, which could contribute to the pathogenesis of the disease.

Keywords: MDD, Jak1, T-bet, Foxp3

ICIA2018\Psychoneuroimmunology and Immunoendocrinology\Poster Discussion\7905
7905

Distribution of Dendritic Cells Subpopulations (Myeloid and Plasmacytoid) in Peripheral Blood of Hyperprolactinemic Women Compared to Normal Subjects

Fatemeh Raeesi Nejad¹, Mohammad Mahdi Mohammadi², Mojgan Sanjari³, Reza Malk poor⁴, Elham Jafari⁵, Mahbubeh Zolala⁶, Mohammad Javad Abasszadeh Afshar⁷

1. Department of Immunology, School of Medicine, Jiroft University of Medical Sciences
2. Department of Immunology, School of Medicine, Kerman University of Medical Sciences.
3. Department of Endocrinology, School of Medicine, Kerman University of Medical Sciences
4. Department of Pathology, School of Medicine, Kerman University of Medical Sciences
5. Department of Pathology, School of Medicine, Kerman University of Medical Sciences
6. Department of Immunology, School of Medicine, Kerman University of Medical Sciences.
7. Department of Parasitology, School of Public Health, Tehran University of Medical Sciences

Background: Interaction between immune and endocrine system is an inextricable two-way communication. In recent years, due to possible role of prolactin in the pathogenesis of autoimmune disorders, this hormone and its relationship with the immune system has attracted much attention. High prolactin levels in different autoimmune diseases as well as increased levels of autoantibodies in hyperprolactinemic patients are evidences for this claim. Dendritic cells subpopulations and their balance play an important role in the pathogenesis of many autoimmune diseases. The aim of the present study was to evaluate the effect of hyperprolactinemia on distribution of DCs subpopulations (myeloid and plasmacytoid) and helping to clarify the complex relationship between PRL and autoimmunity.

Methods: This case-control study was conducted on 70 women, including 35 hyperprolactinemic patients as case group and 35 matched healthy controls. The patients had been admitted into Shahid Bahonar hospital affiliated to Kerman University of Medical Sciences. After obtaining written informed consent from all individuals, blood samples for the cytometric analysis of peripheral blood myeloid and plasmacytoid DCs subtypes were collected. Statistical analyses were carried out using SPSS v18.

Results: The population of myeloid and plasmacytoid DCs in hyperprolactinemia patients were decreased compared to control group. This difference was statistically significant ($P_{pDCs}=0.002$, $P_{mDCs}=0.005$). Also, the study found that hyperprolactinemia does not have significant effect on mDC / pDC balance in peripheral blood ($P=0.928$).

Conclusion: According to the results of this study it can be stated that increased serum prolactin reduces the number of myeloid and plasmacytoid dendritic cells in peripheral blood but does not affect their proportion and balance.

Research on the effects of this hormone on the maturity and function of these cells, as well as its effect on autoimmune diseases, seems necessary.

Keywords: Prolactin, Hyperprolactinemia, Dendritic Cells, Myeloid, Plasmacytoid, Flow Cytometry

ICIA2018\Psychoneuroimmunology and Immunoendocrinology\Poster Discussion\9875
9875

Serum Levels of CCL5 in Type 2 Diabetic Patients with and without Nephropathy

Vahid Bagheri¹, Gholamhossein Hassanshahi¹, Hossein Khorramdelazad¹

1. *Molecular Medicine Research Center, Rafsanjan University of Medical Sciences.*

Background: Type 2 diabetes (T2D) is a chronic metabolic disorder and its prevalence is increasing globally. A number of genetic and environmental factors play a role in the development of T2D. T2D is associated with microvascular complications (e.g. nephropathy) and macrovascular complications. Diabetic nephropathy is one of the main causes of chronic kidney disease and affects about 40% of diabetic patients. CCL5 (RANTES) is a member of the CC chemokine subfamily and actively participates in recruitment of leukocytes to the site of inflammation. The aim of this study was to evaluate the serum level of CCL5 in type 2 diabetic patients with and without nephropathy.

Methods: Peripheral blood samples were collected from 100 patients with T2D, 100 T2D patients with nephropathy, and 100 healthy controls. Ethical approval was obtained from the Ethics Committee at Rafsanjan University of Medical Sciences. Exclusion criteria included presence of inflammatory diseases, infections, and allergic conditions. Demographic and clinical features of patients and controls were also collected. CCL5 serum levels were detected using ELISA (eBioscience) in both groups immediately after blood collection.

Results: Our results showed that serum levels of CCL5 were significantly elevated in patients with T2D (586.37 ± 19.7 pg/ml) compared with healthy controls (502.14 ± 15.35 pg/ml) ($P < 0.004$). However, there was no significant difference in CCL5 levels between nephropathic group (547.11 ± 21.34 pg/ml) and control group ($P = 0.228$).

Conclusion: The results suggest that CCL5 may be involved in the pathogenesis of T2D. Moreover, CCL5 does not seem to be associated with diabetic nephropathy in patients with T2D.

Keywords: CCL5, Type 2 diabetes, Chemokine, Nephropathy



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Reproductive Immunology

Poster Discussion

ICIA2018\Reproductive Immunology\Poster Discussion\6336
6336

Immunomodulatory Effects of Human Amniotic Epithelial Cells on Naïve CD₄⁺ T cells Differentiation from Women with Unexplained Recurrent Spontaneous Abortion

Hossein Motedayyen^{1*}, Nader Tajik², Amir Hassan Zarnani², Abbas Rezaei¹

1. Department of Immunology, Faculty of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

2. Immunology Research Center (IRC), Iran University of Medical Sciences, Tehran, Iran

Background: Unexplained Recurrent spontaneous abortion (URSA) is a common disorder in 1%–5% of women of reproductive age that is largely associated with the failure of feto-maternal immunologic tolerance. As human amniotic epithelial cells (hAECs) have immunomodulatory properties, we investigated immunomodulatory effects of hAECs on differentiation of naïve T cells from USRA patients.

Methods: Naïve CD₄⁺ T cells were isolated from 10 patients with URSA using immunomagnetic separation method. hAECs were isolated from the amnion delivered by healthy women with uncomplicated pregnancies during elective cesarean deliveries. hAECs were co-cultured at different ratios (2:1, 1:1, 1:2, 1:5, 1:10) with naïve CD₄⁺ T cells (4×10^5) in 24-well plates and stimulated with anti-CD3/CD28. After 6 days, the cells were collected and stained with PE anti-human CD25, PE/Cy5 anti-human CD4 and anti-human Foxp3 antibodies. Data was acquired using a FACS Calibur flow cytometer. In this experiment, naïve CD₄⁺ T cultured alone and stimulated with anti-CD3/CD28 were considered as control.

Results: Our data showed that in the presence of hAECs at all ratios, there was a significant differentiation toward Tregs after 3 and 6 days of co-culture ($p < 0.0001$). The effect of hAECs on production of Treg cells was in a dose-dependent manner.

Conclusion: hAECs have the ability to induce differentiation of naïve T cells from URSA patients into Tregs *in vitro*. Based on these findings, hAECs may be considered as potential candidate in immunotherapy of URSA women. However, future studies using animal model and more information are required to decide whether hAECs have clinical applications in URSA.

Keywords: Amniotic Epithelial Cells, Naïve T cells, Recurrent Spontaneous Abortion, Immunomodulatory effects.

ICIA2018\Reproductive Immunology\Poster Discussion\6338
6338

Human Amniotic Epithelial Cells Inhibit Activation and Pro-Inflammatory Cytokines Production of Naive CD4⁺ T cells from Women with Unexplained Recurrent spontaneous Abortion

Hossein Motedayyen^{1*}, Abbas Rezaei¹, Amir-Hassan Zarnani², Nader Tajik²

1. Department of Immunology, Faculty of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

2. Immunology Research Center (IRC), Iran University of Medical Sciences, Tehran, Iran

Background: Unexplained recurrent spontaneous abortion (URSA) has been assumed to be caused by a defect in maternal immunological tolerance to the fetus. Human amniotic epithelial cells (hAECs) have pregnancy-friendly features and the ability to modulate the innate and adoptive immune responses. This study aimed to investigate whether hAECs have immunomodulatory effects on naive CD4⁺ T cells from URSA patients.

Methods: hAECs were obtained from 15 healthy pregnant women and phenotypic profile of hAECs was determined by flow cytometry. Naive CD4⁺ T cells were isolated from 25 URSA patients using an immunomagnetic separation method. Naive T cells were stimulated with anti-CD3/anti-CD28 antibodies and co-cultured with different numbers of hAECs for 3 and 6 days. Immunomodulatory effect of hAECs on activation of stimulated T cell was assessed by flow cytometry and Enzyme-linked immunoabsorbent assay (ELISA). The hAECs effect on pro-inflammatory cytokines production of activated T cells was also measured by ELISA.

Results: Our results indicated that hAECs significantly inhibited the activation of naive T cells in a dose-dependent manner ($p < 0.05-0.0001$). They significantly reduced the production of transforming growth factor-beta1 (TGF- β 1) of stimulated CD4⁺ T cells ($p < 0.05-0.001$). Moreover, hAECs had potent immunomodulatory effects on the production of interferon-gamma (IFN- γ) and interleukin-17A (IL-17A) of activated T cells ($p < 0.01-0.0001$). **Conclusion:** These findings suggest that hAECs may be a suitable cell source to modulate abnormal immune responses in women with URSA.

Keywords: Immunomodulatory Effects, Human Amniotic Epithelial Cells, Naive T cells, Unexplained Recurrent Spontaneous Abortion

ICIA2018\Reproductive Immunology\Poster Discussion\7520
7520

The Immune Profile of Peripheral Blood of Recurrent Implantation Failure Patients

Mahnaz Ghaebi^{1,2}, Majid Ahmadi^{1,2}, Mehdi Yousefi^{1,2}

1. Department of Immunology, School of Medicine, Tabriz University of Medical Sciences

2. Stem Cell and Regenerative Medicine Institute, Tabriz University of Medical Sciences

Background: Recurrent implantation failure (RIF) is a major cause of failure of in vitro fertilization. It is defined when a woman experience failure of good quality embryos to implant following a number of IVF cycles. Immunological abnormalities may be one of the main causes of reproductive failure in RIF.

Methods: 20 women with RIF and 20 healthy women participated in the study. We evaluated the pre-conception level of regulatory T cells and Th17 in peripheral blood of women with RIF using flowcytometry. The expression level of genes involved in T cell responses including GATA-3, GTR, IRF-4 and T-bet was evaluated by quantitative real time PCR.

Results: The results showed that Treg cells were decreased in women with RIF compared with healthy women. As expected, a high level of Th17 cells was observed in patients. Expression of GATA-3 and GTR which regulate Th2 cell differentiation and are involved in Treg function, respectively, were significantly downregulated in RIFs compared with normal women. Expression of IRF-4 and T-bet were higher in women experiencing IVF failure than normal women.

Conclusion: The high frequency of Th17 and lower level of Tregs in peripheral blood of RIF patients compared with normal women show the critical role of Tregs in successful embryo implantation and the detrimental function of pro-inflammatory Th17 cells in this process. In addition, expression pattern of genes involved in Treg and Th2 cell responses compared with Th1 and Th17 related transcripts indicate that Th1 and Th17 cells play a detrimental role while Treg and Th2 cells are protective.

Keywords: Recurrent Implantation Failure, Treg, Th17.

ICIA2018\Reproductive Immunology\Poster Discussion\7527
7527

Evaluation of TH17 and Treg Frequency in Preeclampsia Patients

Shadi Eghbalfard ^{1,2}, Majid Ahmadi ^{1,2}, Mehdi Yousefi ^{1,2}

1. Department of Stem Cell and Regenerative Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

2. Department of Immunology, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Preeclampsia (PE) is an immune-mediated syndrome that occurs in 5-10% of all pregnancy during the second and third trimesters of pregnancy, and is generally characterized by hypertension (>140/90mmHg), proteinuria (>300mg/24 h) and maternal systemic inflammation. It has been proposed that inappropriate activation of the immune system, particularly the imbalance of Th17/Treg may be involved in the pathophysiology of PE.

Methods: A total of 25 pre-eclamptic patients and 25 healthy pregnant women, who were in the second or third trimesters of their pregnancies enrolled in this study. First, the peripheral blood mononuclear cells (PBMCs) were isolated from peripheral blood, and the frequency of Tregs and Th17 cells were analyzed by flowcytometry. Then, the expression level of FoxP3 and RORc genes were evaluated using real-time qPCR. In addition, ELISA was carried out to determine the levels of cytokines such as TGF- β and IL-17 in the peripheral blood of these patients.

Results: Results showed higher percentage of Th17 cells and lower percentage of Treg cells in patients with PE compared to healthy pregnant women ($P<0.02$ and $P<0.03$, respectively). Additionally, the expression of RORc was significantly elevated in PE patients compared with healthy group, whereas expression of FOXP3 was significantly attenuated ($P=0.02$ and $P=0.03$, respectively). As expected, a significant elevation in serum levels of IL-17 was observed in PE patients ($P<0.04$), while the opposite was the case for TGF- β ($P<0.02$).

Conclusion: The study indicates that a negative correlation between Treg and Th17 cells are present in women with PE, characterized by down regulation of Treg cells along with Treg cell-associated transcripts (FOXP3 and TGF- β) and up-regulation of Th17 cells and its associated factors (RORc and IL-17). It seems possible that imbalance of Th17/Treg cells and alteration in cytokines profile may break the maternal tolerance to the fetus and activate inflammatory response in PE patients.

Keywords: Preeclampsia, Regulatory T cell, Th17, FOXP3, Transforming growth factor-beta (TGF-beta), Interleukin-17

ICIA2018\Reproductive Immunology\Poster Discussion\7844
7844

MSC Therapy Protects Fetus from Rejection in Abortion-prone Mouse Model with Involvement of Co-stimulatory and Co-inhibitory Molecules

Amir Salek Farrokhi¹, Amir Hassan Zarnani², Seyed Mohammad Moazzeni¹

1. Department of Immunology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

2. Nanobiotechnology Research Center, Avicenna Research Institute, ACECR, Tehran, Iran

Background: Immunological tolerance has a crucial role in the successful pregnancy since normally the maternal immune system does not reject the semi-allogeneic conceptus. Co-stimulatory and co-inhibitory molecules have essential roles in T cell activation and immune responses. Mesenchymal stem cells (MSCs) have been shown to possess broad immunoregulatory capabilities and inhibit immune cells proliferation, maturation and suppress immune reactions.

Methods: Abdominal fat derived MSCs from female CBA/J mice were administered intraperitoneally (i.p) to the DBA/2 mated CBA/J female mice on day 4.5 of pregnancy. On day 13.5 of pregnancy, abortion rates were calculated and CD80, CD86, CD28 and CTLA-4 gene expression in deciduas and placentas were evaluated by Real-Time PCR.

Results: Our results demonstrated that MSCs administration could decrease the abortion rate significantly in MSCs treated group compared to untreated group. In addition, the findings indicated a significant reduction of CD80, CD86 and CD28 as well as up-regulation of CTLA-4 gene expression following MSCs therapy in both deciduas and placenta.

Conclusion: Our findings demonstrated that administration of MSCs improves pregnancy outcome in abortion prone mouse model through induction of immunological tolerance by reducing the co-stimulatory molecules and increasing co-inhibitory molecules at feto-maternal interface.

Keywords: Mesenchymal stem cells, recurrent spontaneous abortion, Co-stimulatory molecules, Co-inhibitory molecules, Pregnancy

ICIA2018\Reproductive Immunology\Poster Discussion\7849

7849

Mesenchymal Stem Cells Administration Improves Pregnancy Outcome and Induces Th2 Type Cytokines Profile in Abortion Prone Mouse Model

Amir Salek Farrokhi¹, Amir-Hassan Zarnani², Seyed Mohammad Moazzeni¹

1. Department of Immunology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

2. Nanobiotechnology Research Center, Avicenna Research Institute, ACECR, Tehran, Iran

Background: The imbalance of Th1/Th2 cytokines is well known in recurrent spontaneous abortion (RSA) mouse model. Mesenchymal stem cells (MSCs) possess potent immunoregulatory properties that could modulate the Th1 cytokine responses in benefit of Th2 types. In this study, we aimed to analyze the local and systemic balance of Th1/Th2 cytokines following MSCs therapy.

Methods: Syngeneic adipose derived MSCs were administered to abortion prone mice during the implantation window. The abortion rate was determined and IL-4, IL-6, IL-12, IL-2, IFN- γ and GM-CSF gene expression was evaluated by Real-Time-PCR in decidual and placental tissues of pregnant mice at day 13.5 of pregnancy. Splenocytes of pregnant mice were co-cultured with mitomycin C treated paternal splenocytes and IL-2, IL-4, IL-10 and IFN- γ cytokines were measured in co-cultures supernatants by ELISA method. Proliferation response of female splenocytes to paternal antigens was also evaluated using the CFSE method.

Results: Our results showed a significant reduction in abortion rate following MSCs administration in abortion prone mice. We also observed a significant down-regulation of IL-2 and IFN- γ as well as up-regulation of IL-4 and IL-10 production from pregnant mouse splenocytes following MSCs therapy along with a significant reduction of splenocytes proliferation against paternal antigens. Our findings revealed that MSCs therapy increased the IL-4, IL-6, IL-10 and GM-CSF and at the same time decreased the IL-12, IL-2 and IFN- γ gene expression at feto-maternal interface.

Conclusion: Here, we showed that MSCs therapy could modulate the systemic as well as local Th1/Th2 cytokines production along with protection of fetus from resorption in abortion prone mice. The fine balance of Th1/Th2 cytokine response could be considered as one of the possible mechanisms for fetal protection following MSCs therapy.

Keywords: Mesenchymal stem cells, recurrent spontaneous abortion, Th1/Th2 balance, Pregnancy

ICIA2018\Reproductive Immunology\Poster Discussion\7873
7873

A Novel Sperm Protein Target in Azoospermia Men

Mohammad sadegh Soltani Zangbar^{1,2}, Behrouz Ghareesi fard^{1,3}, Jaleh Zolghadr^{3,4}

1. Department of Immunology, Medical School, Shiraz University of Medical Sciences, Shiraz, Iran

2. Department of Immunology, Medical school, Tabriz University of Medical Sciences, Tabriz, Iran

3. Infertility Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

4. Department of Obstetrics and Gynecology, Medical School, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Infertility is one of the most important and worldwide increasing problems of couples who are in reproductive age. Male infertility has many reasons such as azoospermia, defined as complete absence of spermatozoa in the semen. Azoospermia occurs in two forms, including obstructive azoospermia (OA) and non-obstructive azoospermia. OA accounts for around 40 % of azoospermia cases. In obstructive azoospermia due to problems with the ductal system or issues with ejaculation, blood-testis barrier might breakdown and the immune system meets the sperm cells and results in producing the anti-sperm antibodies (ASA). Although the presence of the ASA is reported in 88 % of patients with OA, interestingly there is no data regarding ASA targets in OA individuals. The aim of this study was identification of sperm antibody targets in a group of obstructive azoospermic men. **Methods:** Two-dimensional gel electrophoresis (2-DE) technique was used for separation of the human sperm proteins. After separation, protein spots were transferred on PVDF membranes and blotted with pooled sera from OA patients and compared with membranes blotted with normal men sera. MALDI TOF/TOF mass spectrometry was used for identifying the desirable blotted spots. The results of mass were confirmed using RT-PCR.

Results: The result indicated that OA patients may produce antibody against ODF2 and other sperm proteins such as Tektin-2 and TPI1P1. Our result indicated that normal men don't produce antibody against this identified protein. The mass result of ODF2 was confirmed at RNA level by RT-PCR.

Conclusion: The data of the present study candidate ODF2 as a sperm protein target in OA men.

Keywords: Azoospermia, ODF2, 2DE, Western blot

ICIA2018\Reproductive Immunology\Poster Discussion\7973
7973

Difference in Protein Expression of the Peripheral Blood CD4⁺ T Lymphocytes between Polycystic Ovary Syndrome (PCOS) and Healthy Women

Fatemeh Nasri¹, Fatemeh Hesam Pour¹, Behrouz Ghareasi Fard^{1,2*}

1. Department of Immunology, Shiraz University of Medical Sciences, Shiraz, Iran

2. Infertility Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Polycystic ovary syndrome (PCOS) is the most common endocrinopathy in women of reproductive age. Although the etiopathogenesis of this syndrome is almost clear, proteome profile analysis of CD4⁺ T lymphocytes may represent the proteins involved in the pathogenesis of the disease. The present study aimed to compare the protein expression profiles of the peripheral blood of CD4⁺ T lymphocytes between PCOS patients and healthy women.

Methods: We used two-dimensional gel electrophoresis (2-DE) followed by mass spectrometry (MS) of selected protein spots. Moreover identified protein spots were confirmed by western blot technique.

Results: Despite the overall proteome similarities between patients and healthy women, the analysis of protein spots revealed that at least seven spots were differently expressed ($P < 0.05$). Protein identification was successfully achieved for 3 out of 7 spots by Mass technique and confirmed by western blot. All 3 identified proteins including Phosphatidyl ethanolamine-binding protein 1 (PEBP1), Proteasome activator complex subunit 1 (PSME1), and Triosephosphate isomerase 1 (TPI) showed over-expression in PCOS patients compared with the healthy subjects. Differentially expressed proteins might be involved in oxidative processes and cardiac pathology.

Conclusion: In summary, this evidence highlights T lymphocytes competence as a living biosensor system to record the alteration of metabolism and gene expression and would be a good substitution for tissue biopsies in neural disorders.

Keywords: CD4⁺ T lymphocytes, proteomics, PCOS, mass spectrometry

ICIA2018\Reproductive Immunology\Poster Discussion\8259

8259

The Effect of Mesenchymal Stem Cells Therapy on the Expression Pattern of Activating and Inhibitory Receptors on Uterine Natural killer cells in Abortion-prone Mouse ModelFatemeh Rezaei¹, Seyed Mohammad Moazzeni¹*1. Immunology Department, Faculty of Medical Sciences, Tarbiat Modares Universty, Tehran, Iran*

Background: Uterine natural killer cells (uNKs) are the major population of immune cells in the maternal – fetal interface and play an important role in establishment and maintenance of normal pregnancy. Recurrent spontaneous abortion is one of the most common complications of pregnancy which in many cases is related to the immune cells disorders. We have shown that mesenchymal stem cells (MSCs) therapy could reduce the abortion rate in abortion prone mice. In this study we aim to evaluate the effect of MSCs therapy on the expression pattern of activating and inhibitory receptors on uNK cells.

Methods: MSCs were injected (IP) at day 4 of gestation to female CBA/J mice following their mating with DBA/2 male. The same mating pairs received PBS as control group and CBA/J x BALB/c mating was also used as normal pregnancy control. On day 12.5 of pregnancy embryo resorption rate was determined and decidual cells were isolated by enzymatic digestion. The expression pattern of activating and inhibitory receptors by NK cells were examined through flow cytometric analysis.

Results: MSCs administration dramatically decreased embryo resorption rate compared with control group. MSCs could also affect the phenotype of NK cells in uterine and changed the pattern of activating and inhibitory receptor on cell surface to a more regulatory types. **Conclusion:** This finding indicate that administration of MSCs improved the pregnancy outcome and corrected the functions and phonotype of uNK cells in abortion prone mice. However, the changes in other properties of uNK cells and other aspects of immune systems are remained to be determined and is under investigation in our laboratory.

Keywords: Recurrent Spontaneous Abortion, Mesenchymal stem cell, NK cells, activating and inhibitory receptors.

ICIA2018\Reproductive Immunology\Poster Discussion\8365

8365

Protective Effect of Berberine against Lipopolysaccharide-Induced Abortion

Tahereh Jalilvand¹, Hasan Namdar Ahmadabad², Maryam Nezafat Firizi²

1. Department of Molecular Science, School of Medicine, North Khorasan University of Medical Sciences, Bojnord, Iran

2. Department of Pathobiology and Laboratory Sciences, North Khorasan University of Medical Sciences, Bojnord, Iran

Background: Berberine is an isoquinoline derivative alkaloid with anti-inflammatory activity. In present study, we investigated the protective effects of berberin in prevention of LPS-induced abortion.

Methods: On gestation day (GD) 9.5, the pregnant mice were injected with low, middle and high doses of berberine or with PBS. After 4 hours, berberine or PBS-pretreated mice were injected with LPS. On GD 11.5, blood samples and uterine tissues were collected from treated mice and percentage of abortion and serum levels of NO, TNF- α , IL-10 and IL12p70 determined by macroscopic examination and sandwich ELISA, respectively.

Results: Our findings showed that mice injected with berberine resistant to LPS-induced abortion. We also found that this treatment prevents the reduction of IL-10 and the enhancement of NO, TNF- α , and IL-12p70 in LPS-treated pregnant mice.

Conclusion: Taken together, our results suggest that berberine as an anti-inflammatory agent has protective effects on LPS-induced abortion by modulation of pro- and anti-inflammatory factors.

Keywords: Abortion, Berberine, Cytokine, Lipopolysaccharide, Nitric Oxide.

ICIA2018\Reproductive Immunology\Poster Discussion\9905
9905

Dysregulation of Fas in Endometrial, Peritoneal Fluid, and Blood Mononuclear Cells May Propose a Role for Resistance to Apoptosis in Patients with Endometriosis

Tahereh Mohammadi^{1,2}, Sepide Khodaverdi³, Nader Tajik^{1,2}, Zahra Moradi^{1,2}, Ali Akbar Delbandi^{1,2*}

1. Immunology Research Center, Immunology and Infectious Disease Institute, IRAN University of Medical Sciences

2. Immunology Department, School of Medicine, IRAN University of Medical Sciences

3. Endometriosis Research Center, IRAN University of Medical Sciences

Background: Endometriosis is characterized by the presence of endometrial tissues composed of stroma and glands with capacity to avoid apoptosis outside the uterine cavity. *This study aims at investigating the expression levels of Fas in ectopic (EESCs) and eutopic (EuESCs) endometrial stromal cells, as well as peritoneal fluid (PFMCs) and peripheral blood mononuclear cells (PBMCs) in patients with endometriosis in comparison to non-endometriotic controls.*

Methods: Ectopic and eutopic endometrial tissues were taken from 29 patients with endometriosis and 10 eutopic endometrial tissues taken from 10 individuals without endometriosis. We also collected peritoneal fluid and peripheral blood from 10 endometriosis patients, as well as 10 individuals without endometriosis. Endometrial tissues were digested by enzymatic method and the endometrial stromal cells (ESCs) were characterized by flowcytometry. Peritoneal fluid and blood mononuclear cells were isolated by ficol density separation method. Gene and protein expression levels of Fas in separated cells were evaluated by Real-Time PCR and Western blotting, respectively.

Results: We found that EESCs and EuESCs produced lower amounts of Fas in gene and protein levels compared to CESC (P<0.01). Interestingly, the expression level of this gene was higher in PFMCs of patients than in that of the control group (P<0.05). In addition, the expression levels of Fas was lower in PBMCs of patients in comparison with the control group (P<0.01).

Conclusion: It seems that lower expression of Fas in EuESCs and EESCs is one mechanisms that explain why these cells are resistant to apoptosis. Furthermore, probably the elevated expression of this gene in PFMCs of patients can describe the reasons of increasing apoptosis in the mentioned cells.

Keywords: Endometriosis, Stromal cell, peritoneal fluid mononuclear cells, Fas

ICIA2018\Reproductive Immunology\Poster Discussion\9951
9951

1, 25-Dihydroxy Vitamin D3 reduced HGF Production in Endometriosis

Sahel Heydari^{1,2}, Sepideh Khodaverdi³, Nader Tajik^{1,2}, Ali Akbar Delbandi^{1,2}

1. Immunology Research Center, Immunology and Infectious Disease Institute, IRAN University of Medical Sciences

2. Immunology Department, School of Medicine, IRAN University of Medical Sciences

3. Endometriosis Research Center, IRAN University of Medical Sciences

Background: Endometriosis is an inflammatory disease characterized by the presence and growth of endometrial tissue outside the uterine cavity. HGF plays an important role in the proliferation of ectopic endometrial tissues in endometriosis. In this study, we examined the production levels of HGF by peritoneal (PFMCs) and peripheral blood (PBMCs) mononuclear cells, as well as Ectopic (EESCs) and Eutopic (EuESCs) stromal cells from patients with endometriosis compared to nonendometriotic controls. Moreover, the effects of vitamin D3 on the production of HGF in mentioned cells were investigated.

Methods: EESCs, EuESCs, PBMCs, and PFMCs were isolated from 10 laparoscopically-confirmed endometriotic patients. Furthermore, 10 separated CECs, PBMCs, and PFMCs from 10 non-endometriotic women were enrolled as controls. The gene and protein expression levels of HGF were measured in all cell types by Real-time PCR and ELISA, respectively. For determining the effect of vitamin D3 on HGF expression levels, the cells were treated with the optimized concentration of 1, 25-Dihydroxy Vitamin D3. Then, the gene and protein expression levels of HGF were measured by Real-time PCR and ELISA, respectively.

Results: The gene and protein expression levels of HGF were higher in EESCs and PFMCs in endometriosis patients compared to nonendometriotic controls ($P<0.05$ and $P<0.01$, respectively). We found that vitamin D3 could decrease the HGF gene ($P<0.05$) and protein expression levels ($P<0.01$) in PFMCs, EuESCs and EESCs in endometriotic patients. However, this vitamin, has no effect on control group. Treatment of PBMCs with vitamin D3, caused a significant decreases in the HGF protein level in both patients and controls ($P<0.05$).

Conclusion: These results showed HGF is contributing to the pathogenesis of the endometriosis. In addition, these findings propose the unique therapeutic potential for 1, 25-Dihydroxy Vitamin D3 for endometriosis treatment.

Keywords: Endometriosis; Vitamin D3; HGF; Stromal cell

ICIA2018\Reproductive Immunology\Poster Discussion\9961

9961

1, 25-Dihydroxy Vitamin D3 Regulate Apoptosis in Endometrial Stromal and Peritoneal Fluid Cells in Endometriosis

Tahereh Mohammadi^{1,2}, Sepide Khodaverdi³, Nader Tajik^{1,2}, Zahra Moradi^{1,2}, Ali Akbar Delbandi^{1,2}

1. Immunology Research Center, Immunology and Infectious Disease Institute, IRAN University of Medical Sciences

2. Immunology Department, School of Medicine, IRAN University of Medical Sciences

3. Endometriosis Research Center, IRAN University of Medical Sciences

Background: Endometriosis is a chronic inflammatory disease characterized by the dysregulated apoptosis in ectopic endometrial cells. The objective of this study was to investigate the effects of 1, 25-Dihydroxy vitamin D3 on FasL expression in Ectopic (EESCs) and eutopic (EuESCs) endometrial stromal and peritoneal fluid mononuclear cells in patients with endometriosis compared to endometrial stromal cells (CESCs) and PFMCs from nonendometriotic controls.

Methods: Stromal cells were separated through enzymatic digestion of ectopic and eutopic endometrial tissues from 29 endometriotic patients and 10 nonendometriotic controls and characterized by flow cytometry. Moreover, we collected peritoneal fluid from 10 endometriosis patients, as well as 10 non-endometriotic controls. Following cell isolation, the gene and protein expression levels of FasL were evaluated by Real-time PCR and Western blot, respectively. Also, the cells were treated with the optimum concentration of 1, 25-Dihydroxy vitamin D3 in different time manner and then the effects if this vitamin were assessed in gene and protein expression of FasL in mentioned cells.

Results: EESCs exhibited a significantly higher FasL gene and protein expression compared to EuESCs or CESCs ($P<0.01$). Furthermore, significantly lower gene expression levels of FasL were found in PFMCs in endometriotic patients compared to cells from nonendometriotic controls ($P<0.01$). 1, 25-Dihydroxy vitamin D3 reduced the FasL protein expression of EuESCs ($P<0.01$) and EESCs ($P<0.05$). Interestingly, this vitamin increased the gene and protein expression of FasL in PFMCs from endometriotic patients ($P<0.05$).

Conclusion: These findings propose that 1, 25-Dihydroxy vitamin D3 have beneficial effects on the regulation of apoptosis in patients with endometriosis.

Keywords: Endometriosis; Stromal cells; FasL; 1, 25-Dihydroxy vitamin D3

ICIA2018\Reproductive Immunology\Poster Discussion\9969
9969

MSC Therapy Alters the Expression Levels of Crry and Adipsin at the Feto-maternal Interface of Abortion Prone Mice

Shahab Shahgaldi¹, Amir Salek Farrokhi¹, seyed Mohammad Moazzeni¹

1. Department of Immunology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

Background: Recurrent spontaneous abortion, with a prevalence of 2-5% among pregnant women, is one of the most common complications of pregnancy. Today, it's well established that a great proportion of recurrent spontaneous abortions (RSAs) is due to immunological factors. In this regard, complement activation and regulation plays a pivotal role.

In this study, we measured the impact of MSCs on the expression levels of Crry and Adipsin at the feto-maternal interface of abortion prone mice.

Methods: Adipose tissue-derived mesenchymal stem cells (AT-MSCs) were isolated from the abdominal fat of CBA/J mice. On the 4.5th day of gestation, the test group received an IP injection of 1×10^6 of AT-MSCs. On the 13.5th day of gestation, tissue samples (placentae and deciduae of pregnant mice) were analyzed to measure the expression levels of Crry and Adipsin using real-time PCR.

Results: Following MSC therapy, the average weight of fetuses was increased. Besides, the real time PCR analysis of the expression levels of Crry and Adipsin demonstrated that Adipsin expression levels was decreased significantly upon MSC administration while the expression levels of Crry was elevated at the feto-maternal interface of abortion prone mice.

Conclusion: In the present study, we showed for the first time that MSCs improved fetal developmental conditions in abortion-prone mice by adjusting the expression levels of complement regulatory molecules. Our results suggested that MSCs altered the expression levels of Adipsin and Crry, which ultimately contributed to better pregnancy outcomes.

Keywords: Mesenchymal stem cell, recurrent spontaneous abortion, Cell therapy, Complement regulatory molecules, Complement system, Pregnancy, Adipsin, Crry.

Stem Cells Based Immunotherapy

Poster Discussion

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Stem Cells Based Immunotherapy\Poster Discussion\5302

5302

The Effect of Mesenchymal Stem Cell Pulsed with 17- β Estradiol in Ameliorating Rat Model of Ulcerative Colitis

Siavash Mashhour¹, Seyyed Meysam Abtahi Froushani²

1. Graduate of Veterinary Medicine, Department of Microbiology, Faculty of Veterinary Medicine, University of Urmia

2. Department of Microbiology, Faculty of Veterinary Medicine, University of Urmia

Background: One of the natural factors which has a proliferative effect on Mesenchymal stem cells (MSCs), is 17- β estradiol (E2). However, there is no/or limited documents about the effects of treatment of ulcerative colitis (UC) with E2-primed MSCs. Therefore, this research was set out to assess the effects of E2-primed MSCs therapy in the rat model of UC.

Methods: In this experimental study, after isolation of MSCs, they were co-cultured with 0 or 100nM 17- β estradiol for 24 h. Colitis induced by acetic acid in four groups of male Wister rats; control group, MSCs treated group, co-culture MSCs by estrogen treated group and control group. MSCs (2×10^6 cell) were injected into the peritoneum in different group separately. After 10 days, the rats were euthanized and evaluated for level of malondialdehyde, myeloperoxidase, nitric oxide and total protein in the homogenate of gut tissues.

Results: The cell therapy in the UC rats with 17- β estradiol primed MSCs showed a more desirable outcome, causing the regression of the cumulative clinical score to be more favorable than the therapy with un-treated MSCs. The levels of Myeloperoxidase, Nitric oxide and malondialdehyde were significantly decreased and conversely the total protein levels were significantly increased in UC rats received 17- β estradiol primed MSCs more pronounced than UC rats received un-treated MSCs.

Conclusion: 17- β estradiol treated MSCs caused more favorable regression the signs and induce better outcome in rats with UC compared to un-treated MSCs.

Keywords: 17- β estradiol, Mesenchymal stem cell, Ulcerative Colitis.

ICIA2018\Stem Cells Based Immunotherapy\Poster Discussion\7465
7465

The Effect of LPS Treated Adipose Tissue-Derived Mesenchymal Stem Cells Conditioned Media on Nitric Oxide Production and Phagocytostic Activity of Peritoneal Macrophages

Zahra khosrowpour.¹, Seyed.Mahmoud Hashemi.², Sara Soudi.³

1. Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences

2. Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences

3. Department of Immunology, Faculty of Medical Sciences, Tarbiat Modares University

Background: Adipose tissue-derived mesenchymal stem cells (AD-MSCs) are capable of migration to inflamed site and exerts its immunomodulatory effects on tissue resident immune cells like macrophages. To investigate the effect of MSCs treated with LPS soluble factors on macrophage function, we assess nitric oxide and phagocytosis potential production of macrophages after treatment with MSCs treated with LPS conditioned media.

Methods: MSCs were isolated from adipose tissue of C57BL/6 mice. MSCs surface markers were analysed using flowcytometry method. MSCs were cultured and divided into two groups at passage two, first group treated with LPS for three days and the second one had no treatment. C57BL/6 peritoneal macrophages were harvested and treated for 72h with MSCs conditioned media. After treatment, macrophages were treated with yeast particle at 1:10 ratio and phagocytosis percent was measured by microscopic examination. Nitric oxide production was measured in the supernatants of treated macrophages by Griess method.

Results: According to obtained results we found that the level of nitric oxide production of macrophage was decreased after co-culture with MSCs treated with LPS CM compared to the control. Phagocytosis percent of yeast particles were reduced significantly in macrophages cultured with treated MSCs with LPS CM compared to the control.

Conclusion: In this research, we show that MSCs stimulation with LPS will change the immunomodulatory properties of macrophages by reduction of NO production, beside the decrease in phagocytosis percent.

Keywords: AD-MSCs, LPS, macrophage, C57BL/6

ICIA2018\Stem Cells Based Immunotherapy\Poster Discussion\7515

7515

Evaluation of T-regulation of Lymphocytes and Anti-inflammatory and Inflammatory Cytokines Profile of Spleen Cells in Diabetic Mouse after Injection of Exosomes Isolated from Adipose Tissue Mesenchymal Stem Cells

Shahrzad Nojehdehi^{1,2}, Seyed Mahmoud Hashemi^{3,4}, Ardeshir Hesampour⁵, Sara Soudi⁶, Masoud Soleimani⁷

1. Department of Biology, Islamic Azad University Central Tehran Branch, Tehran, Iran

2. Stem Cell Technology Research Center, 1997775555, Tehran, Iran

3. Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

4. Department of Applied Cell Sciences, School of Advanced Technologies in Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

5. Department of Biology, Islamic Azad University Central Tehran Branch, Tehran, Iran

6. Department of Immunology, Faculty of Medical Science, Tarbiat Modares University, Tehran, Iran.

7. Department of Hematology, School of Medical Sciences, Tarbiat Modares University, Tehran, Iran

Background: Type 1 diabetes is one of the most common autoimmune diseases that are due to the activation of T cells against beta cells, resulting in degradation of them, which may have various causes, such as genetic factors and environmental stimuli, including viral infection. Continuous high blood sugar and metabolic disturbances caused by diabetes cause secondary pathophysiologic changes in many organs in the body, which can lead to many problems and costs for people with diabetes and the community. Existing evidence showed that stem cells release exosomes and nano-vesicles known as exosomes that may be used as cell-cell-linkers and potentially transmit stem cell phenotype to living cells and maintain, Facilitate differentiation, self-renewal, and repair of stem cells. In this study, the therapeutic use of exosomes as a non-cellular therapeutic regimen has advantages that can be a good justification for its replacement in conventional cell therapies. The main innovation of this project is the use of isolated exosomes from the use of soup and suppository mesenchymal stem cell culture that does not have the dangers and problems of cell therapy and also has more efficacy than the supernatant cell culture. **Methods:** First, by disease induction using STZ and based on disease symptoms, mice were separated into three groups: healthy group, control group, and treatment group. The treatment group were treated by intraperitoneal injection (2 times per week) of exosomes extracted by ultracentrifuge subtraction of mesenchymal stem cells derived from adipose tissue, Then after 8 weeks the mice were dislocated and spleen cells were isolated .the spleen lymphocytes were examined in order to evaluate their alternation in cytokine profiles by ELISA and for evaluate Regulatory T (iTreg) cell T lymphocytes (CD4 + CD25 + Foxp3 +) using flow cytometry. **Results:** The data showed, the mice that were not treated had a shorter life span and also exosomes derived from mesenchymal stem cells were able to reduce the secretion of inflammatory cytokines and increasing anti-inflammatory. Then a significant increase in the population of Treg cells in diabetic mice treated with exosomes derived from AD-MSCs in comparison to the untreated diabetic group have been reported (P<0.05) **Conclusion:** In this study, the therapeutic use of exosomes as a non-cellular therapy has a number of benefits that could properly explain the reason for replacing it with the common methods of cellular therapy.

Keywords: Exosome, Cytokines, Regulatory T Cells

ICIA2018\Stem Cells Based Immunotherapy\Poster Discussion\7552
7552

Immunochemical Evaluation of Mesenchymal Stem Cells Derived from Wharton's jelly Encapsulated by Alginate, Which are Transplantation to the Peritoneal Cavity of the Rat

Saeed Azandeh¹, Masoud Moghimi¹, Negar Varae¹

1. cellular and molecular research center, Ahvaz jundishapur university of medical sciences, ahvaz, iran

Background: Since previous studies claimed that human umbilical cord WJ-MSCs are low immunogenic, with the assumption in this study, the evaluation of immune responses of WJ-MSCs in vitro with and without alginate was investigated.

Methods: Cellular bioavailability against B cell antibody toxicity evaluated by MTT assay and anti-MSCs antibodies were also measured by ELISA test. Also, in the evaluation of cellular immunity against MSCs, the level of IFN- γ secretion measured by ELISA method and proliferation of spleen and lymph cells against MSC was evaluated by CFSE staining and flow cytometry.

Results: In the evaluation of cellular and humoral immunity, one month after cell, scaffold with cell and without cell transplantation did not receive any immune response in the rats and control group that didn't received any graft.

Conclusion: Our result showed because of low immunogenicity of MSCs and major role of alginate with high G scaffold in immunization of component of host immunity from graft donor.

Keywords: alginate; cell encapsulation, immunogenicity, cellular and humoral immunity

ICIA2018\Stem Cells Based Immunotherapy\Poster Discussion\8337

8337

Flow cytometry Analysis of Maturation and Costimulatory Markers of Bone Marrow Dendritic Cells Cultured by Isolated Exosomes of Mesenchymal Stem Cells

Mehri Shahr¹, Esmail Mortaz¹, Seyed Mahmoud Hashemi¹

1. Department of Medical Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: DCs orchestrate innate inflammatory responses and adaptive immunity through activation of T cells. MHCII molecules present the first classical signal in the process of antigen presentation, and co-stimulatory molecules such as CD40 represent the second signal. Since DCs are so well equipped to initiate adaptive immune responses, they are considered prime targets for modulating immune responses. MSCs have anti-inflammatory effects. Exosomes are microparticles secreted from cells and conduct cell-cell communication. In this study we aimed to investigate the effects of isolated exosomes derived from MSCs on surface expression of maturation markers of myeloid DCs.

Methods: Culture of C57BL/6 mouse AD-MSCs: Murine adipose-derived MSCs were isolated from C57BL/6 Mice. Extracellular matrix was digested with type I collagenase, centrifuged and the pellet was then cultured in DMEM. Isolation and purification of exosomes: Exosome was isolated by Exo-spin kit and evaluated by SEM. The protein concentration was detected by BCA kit. Preparation bone marrow derived DCs: BM mononuclear cells were prepared from C57BL/6 mouse femur and tibia and then cultured in 6-well plates containing PRMI-1640 medium supplemented with 20 ng/mL GM-CSF and 20 ng/mL IL-4. Flow cytometry: After co-culture of exosomes (100µg/mL) with DCs (1 million/ml) the surface expression of CD11c, CD83, CD86, CD40 and MHCII were analyzed in following groups: (immature DC (iDC), iDC+LPS (1µg/ml), iDC+exosome (100µg/ml), iDC+exosome (100µg/ml)+LPS (1µg/ml)) was determined in 10000 cells by Flow cytometry.

Results: The results showed that MSC derived exosomes could decrease the expression of the studied markers in comparison to control cells.

Conclusion: The current data indicates that exosomes isolated from MSCs suppress the maturation of bone marrow derived DCs. Besides, our data could suggests that exosomes might be important modulator for the immune responses induces by DCs.

Key words: Mesenchymal Stem Cell, Exosomes, Tolerogenic Dendritic Cell, In Vitro

ICIA2018\Stem Cells Based Immunotherapy\Poster Discussion\8377

8377

The Effect of Serum Starvation on C57B6 Bone Marrow Cells

Gilda Parsamanesh¹, Amrolah Mostafazade¹, Davar Amani², Ali Asadirad², Kaveh Baghaei³,
Hamid Asadzadeh Aghdayi³

1. Department of immunology, School of medicine, babol University of medical sciences.

2. Department of immunology, School of medicine, shahid beheshti University of medical sciences.

3. Basic and Molecular Epidemiology of Gastrointestinal Disorders Research Center, Research institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Background: Serum starvation is a condition that has an effect on cell behavior and very little is known about this mechanism. Previous study in PBMC cells has shown that the starvation (for 96 hours) leads to down-regulation of HLA class I, increasing the relative number of T-reg cells to effective T cells and also increasing the amount of TGF- β in cellular soup. Therefore, we hypothesized that serum starvation condition can be considered as an effective tool in cells transplantation. In this project, the influence of serum starvation was evaluated on the potency of starved mouse bone marrow cells for transplantation.

Methods: Isolated bone marrow cells from C57/B6 were cultured in serum starvation condition for 24, 48, 72, 96 hours and non-starved cells were considered as control groups. Consequently, at the end of each time point, the viability of cells was determined by flow cytometry. In order to analyze the expression of foxp3, TGF β and MHC class I Real-time PCR was carried out. Also, transplantation potency of starved cells will be investigated by one way MLR test.

Results: Expression of MHC class I was declined compared with non-starved cells (24, 48, 72 hours) ($P < 0.000$); however, TGF β in starved cells were increased (48 hours) ($P < 0.000$). In both starved and non-starved groups, foxp3 did not have significant expression.

Conclusion: Although serum starvation leads to down-regulation of MHC class I, increasing of TGF β in bone marrow cells, foxp3 expression was undesirably decreased too. 48 hours starvation could result in prevention of recipient immune response to transplanted bone marrow cells by down-regulation of MHC class I and increase of TGF β , nonetheless, prevention of GVHD in this condition need to be elucidated.

Keywords: Serum starvation/ bone marrow/ MHC class I/ TGF β / foxp3

ICIA2018\Stem Cells Based Immunotherapy\Poster Discussion\8519
8519

Enrichment of Cancer Stem Cells from Murine Colorectal Primary Tumor in order to Cancer Stem Cell-based Vaccine Studies

Leila Eini¹, Zahra Madjd^{1,2*}, Roya Ghods^{1,2*}, Mahmood Bozorgmehr¹

1. *Oncopathology Research Center, Iran University of Medical Sciences, Tehran, Iran*

2. *Department of Molecular Medicine, Faculty of Advanced Technology, Iran University of Medical Sciences, Tehran, Iran*

Background: Cancer stem cells (CSC) are responsible for initiation, maintenance and resistance to chemo/radiation therapy. Inability of CSC targeting using immunotherapy may be a significant factor contributing to treatment failure. There are limited studies in murine CSCs isolation, therefore we aim to develop an animal model and isolate CSCs from CT-26 (colorectal cancer cell line) induced tumor in order to evaluate the effects of lysate vaccine on tumor growth and mice survival. Due to high cost and several limitations of NOD/SCID mice providing in Iran, also absence of adaptive immune responses in these mice, the use of CSCs in such syngeneic immunocompetent hosts permits the assessment of immune responses against these cells.

Methods: Tumor was induced in six weeks old female Balb/c mice with CT26 cells. Tumor tissues were minced and enzymatically digested. The resulting cells were seeded at a density of $1 \times 10^5/\text{ml}$ in Serum-Free Medium. Culture cells were incubated for 7 days, photographed and scored for the presence of colonospheres morphologies. Flow cytometry, was employed to characterize isolated colorectal CSCs compare to parental cells.

Results: Results showed that, cells rounded up to form independent spheroids. Majority of colonosphere cells (85%) were positive for CD133 while 15% of parental cells stained positive for the same marker. Percentage expressions for CD166 was 34% in spheroid cells versus 4% in parental cells.

Conclusion: We demonstrate that CSC populations from murine colorectal primary tumor can be enriched using spheroid formation assay, this could facilitate the establishment of immunocompetent mice models for subsequent studies such as CSCs base vaccine.

Keywords: Cancer stem cell (CSCs), vaccines, CT-26, mouse colorectal model.

ICIA2018\Stem Cells Based Immunotherapy\Poster Discussion\8610

8610

The Effects of Mesenchymal Stem cells on Clinical Scores of Multiple Sclerosis Patients

Fahimeh Lavi¹, Forouzan Yousefi¹, Mahmoud Mahmoudi¹

1. Immunology center, BuAli institute, Department of Immunology Mashhad University of Medical Science Mashhad, Iran

Background: Multiple sclerosis (MS) is the most common disabilities among CNS diseases. The lack of treatments for multiple sclerosis patients in progressive stage demonstrates a major unmet clinical need. Mesenchymal stem cells have emerged to own beneficial effects in multiple sclerosis patients and substitutive animal model of MS.

Methods : we aimed to evaluate the safety and efficacy of MSCs derived from adipose tissue as an effective treatment for secondary progressive multiple sclerosis. Eligible participants were 10 female patients, non-pregnant, unresponsive to other approved treatments, lack of chronic and acute infection or tumor. Patients are predicted to continue to have progression of disease next year. We remove abdominal fat of patients and after harvesting we expand their adipose tissue mesenchymal stem cells (AT-MSC) to reach to a range of 5×10^6 MSC/Kg of body weight. Then we infuse their expanded AT-MSC in 2 dosage during 2 injection space 7 days apart. Next we evaluate the effect of AT-MSCs on patients by assessment of EDSS on baseline, 2 weeks, 1, 3, 6 and 9 months after first injection according to MacDonald criteria.

Results: in all of patients EDSS was decreased significantly from baseline to 3 months post first injection thereafter EDSS was increases in 2 patients by 1 point at endpoint of follow-up and 6 patients were progression free and in 2 patients EDSS was reduced significantly by 1 point from baseline to 9 months after first injection.

Conclusion: MSCs were safe and effective in secondary progressive MS patients and the most pronounced influence was from baseline to 3 months after first MSC administration.

Keywords: Adipose-derived mesenchymal stem cell, secondary progressive multiple sclerosis

ICIA2018\Stem Cells Based Immunotherapy\Poster Discussion\9780

9780

Immunomodulatory Effects of Adipose Tissue-derived Mesenchymal Stem Cells in an Allograft Islets Composite Transplant for Experimental Autoimmune Type 1 Diabete

Bahare Niknam^{1,2}, Jamal Mohammadi Ayenehdeh^{1,3}, Nader Tajik¹, Seyed Mahmoud Hashemi⁴

1. Immunology Research Center, Iran University of Medical Sciences, Tehran, Iran

2. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

3. Department of Immunology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

4. Department of immunology, School of medicine, Shahid Beheshti University of Medical Sciences

Background: Allogeneic islet transplantation could be an ideal alternative therapy for Type 1 Diabetes. Adipose Tissue-derived Mesenchymal Stem Cells (AT-MSCs) characterized by immunomodulatory and protective effects may have the potential to improve the outcome of this highly immunogenic transplant. Immunomodulatory properties have been shown to be exerted through both direct contact and production of soluble markers.

Methods: Syngeneic AT-MSCs along with allograft islets embedded in hydrogelic composite and transplanted intraperitoneally in Streptozotocin (STZ) induced diabetic C57BL/6 mice.

Five groups consist of Control, Hydrogel alone, Gel+MSC, Gel+Islet and Gel+MSC+Islet, delivered into the peritoneal cavity. 32 days after transplantation, allograft composites were Collected and Real-time PCR were performed with specific primers for $\beta 2M$, IDO, iNOS, PDX1 genes.

Results: Co-transplantation of AT-MSCs significantly increased the transcript levels of IDO and iNOS in the Gel+MSC+ Islet group, in comparison with the Gel and Gel+Islet and Gel+ MSC groups. Similar results were also obtained for transcript levels of PDX1 which significantly increased in Gel+MSC+Islet group as compared to Gel and Gel+ Islet and Gel+ MSC groups.

Conclusion: According to results, AT-MSCs are capable to improving graft function with immunomodulatory effects through production of soluble markers.

Keywords: Mesenchymal stem cell, Type 1 Diabetes, Allograft islet transplantation

ICIA2018\Stem Cells Based Immunotherapy\Poster Discussion\9781
9781

Assessment of Phenotype and Function of Tregs in an Allograft Islets and Fat Derived Mesenchymal Stem Cells Composite Transplant for Experimental Autoimmune Type 1 Diabetes

Bahare Niknam^{1,2}, Jamal Mohammadi Ayenehdeh^{1,3}, Nader Tajik¹

1. Immunology Research Center, Iran University of Medical Sciences, Tehran, Iran

2. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

3. Department of Immunology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

Background: Islet transplantation is one of the most promising therapeutic approaches in Type 1 Diabetes. In order to improve the viability and function of islet transplantation, it has been proposed to associate pancreatic islets with Adipose Tissue-derived Mesenchymal Stem Cells (AT-MSCs).

Methods: Allograft islets from BALB/c mice co-embedded with syngeneic AT-MSCs from abdominal fat tissue of C57BL/6 in hydrogelic composite and delivered in to the peritoneal cavity of Streptozotocin (STZ) induced diabetic C57BL/6 mice. Five groups consist of Control, Hydrogel alone, Gel+MSC, Gel+Islet and Gel+MSC+Islet, delivered into the peritoneal cavity. 32 days after transplantation, mononuclear cells from the Mesenteric lymph nodes (MLNs) and spleen were analysis For Treg and intracellular cytokine assay by flowcytometry.

Results: Analyses showed that AT-MSCs co-transplanted with allograft significantly increased Treg ($P<0.05$) in MLNs but there is no significant variation among the groups in spleen. According to the MFI for IL-10 and TGF- β 1 in MLNs analysis, the IL-10 and TGF- β 1 significantly increased in Gel+MSC and Gel+MSC+Islet groups, in comparison with the other corresponding groups ($P<0.05$); but in spleen, TGF- β 1 significantly increased in Gel+MSC and Gel+MSC+Islet groups but variation in IL-10 was not significant.

Conclusion: These result show that AT-MSCs Can promote islet survival and function and can induced functional Treg cells. Moreover, local MSCs transplantation also exert immunomodulatory effects on infiltrating and resident immune cells.

Keywords: Type 1 Diabetes, Mesenchymal stem cells, Treg



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Tolerance and Autoimmunity

Poster Discussion

ICIA2018\Tolerance and Autoimmunity\Poster Discussion\5230

5230

Study of Therapeutic Plasmapheresis Effects on IL-6 and TGF- β and Their Receptors in Relapsing-Remitting MS PatientsAzam Jamshidian^{1,2}; Zahra Khademi²; Vahid Shaygannejad³

1. Faculty of Paramedical Sciences, Shahrekord University of Medical Sciences, Shahrekord, Iran

2. Basic Health Sciences Institute, Shahrekord University of Medical Sciences, Shahrekord, Iran

3. Neurology dept. Medical school, Isfahan University of Medical Sciences, Isfahan, Iran

Background: Multiple sclerosis (MS) is a chronic demyelinating inflammatory disease of the central nervous system. Lack of sufficient information about the effects of plasma exchange (PE) therapy in multiple sclerosis (MS), has limited this treatment to individual patients with severe refractory relapses. Th17 cells play a promoting role in MS. The cytokines, interleukin 6 and TGF- β play a critical role in the development of Th17 cells. Evaluating the effect of PE on the expression levels of IL-6 and TGF- β and their receptors was the aim of this study.

Methods: Peripheral blood samples were obtained before and after a complete course of PE therapy, from 30 Relapsing Remitting-MS patients in relapse phase. IL-6 and TGF- β mRNA levels were assayed using real-time PCR. The IL-6 and TGF- β receptors were assessed using flowcytometry.

Results: Expanded disability scale score was significantly reduced after treatment ($P < 0.01$) and this relieve of symptoms was significantly higher in males compared with females ($P = 0.039$). A significant increase in TGF- β ($p = 0.009$) and a significant decrease in IL-6 receptor were observed ($P = 0.028$). Correlation analysis showed the changes in expression levels of two cytokines are directly correlated ($P = 0.03$). mRNA level of each cytokine after treatment was conversely related to the expression level of its receptor before therapy (TGF- β , $P = 0.019$ & IL-6, $P = 0.028$). The frequency of CD4+IL6R+ cells and CD4+IL6R+ TGF- β R+ cells before PE was conversely correlated to TGF- β mRNA expression levels after PE ($p = 0.03$ & $p = 0.04$, respectively). The frequency of CD4+IL6R+ cells was also correlated with disease severity ($p = 0.001$) and the disease severity was related with symptom relief (0.009).

Conclusion: This study showed increasing TGF- β mRNA and decrease of IL-6 receptor expression can be a way in which plasma exchange improves MS relapse symptoms, and this therapy impacts on the two cytokine and their receptor in an interrelated manner.

Keywords: Multiple sclerosis, Plasma exchange therapy, IL-6, TGF- β

ICIA2018\Tolerance and Autoimmunity\Poster Discussion\5312

5312

Vitamin D Modulates the Expression of IL- 27 in the Central Nervous System in Experimental Autoimmune Encephalomyelitis (EAE)

Marziyeh mohammadi kordkheyli¹, Abdollah jafarzadeh²

1. *Tehran university of medical science, Iran, Tehran*

2. *University of medical science, Iran, Kerman*

Background: It has been reported that vitamin D has broad anti-inflammatory and immunomodulatory effects. To evaluate the effects of vitamin D on the expression of IL-in a model of experimental autoimmune encephalomyelitis (EAE).

Methods: EAE was induced in C57Bl/6 mice by immunization with myelin oligodendroglial glycoprotein mixed with complete Freund's adjuvant. The mice were administered with PBS or olive oil, intraperitoneally, in the control groups and vitamin D (200 ng every two days) in the treatment group, from day +3 to +30. At day 31, the mice were scarified and their spinal cords and brains were the expression of the IL-27 mRNA in the spinal cord was measured using real time-PCR.

Results: In PBS- or olive oil-treated EAE mice the expression of IL-27 P28 mRNA was significantly lower than that in the healthy control group ($p < 0.002$). In both PBS- and olive oil-treated EAE groups, the expression of IL-27 EBI3 mRNA was also lower than that observed in the healthy group, but the differences were not significant. In vitamin D-treated EAE group, the expression of IL- 27 P28 and IL-27 EBI3 were significantly higher compared with the olive oil-treated EAE groups ($p < 0.002$ and $p < 0.04$, respectively). The PBS- or olive oil-treated EAE mice showed the clinical symptoms of EAE at days 9 and 10, respectively. The maximum mean pathological scores were also significantly lower in vitamin D-treated EAE group, in comparison with PBS- or olive oil treated EAE mice ($p < 0.001$).

Conclusion: Vitamin D may modulate the expression of IL-27 in the spinal cord of EAE mice and also ameliorate the clinical symptoms of the disease.

Keywords: Vitamin D, EAE, IL-27

ICIA2018\Tolerance and Autoimmunity\Poster Discussion\6355
6355

Down Regulation of Neuroimmune Semaphorin 3A in Peripheral Blood Mononuclear Cells (PBMCs) and Its Serum Level of Patients with Multiple Sclerosis

Mahsa Rezaeepoor¹, Shima Shapoori¹, Mazdak Ganjalikhani hakemi¹, Masoud Etemadifar², Fereshteh Alsahebhosoul¹, Nahid Eskandari¹, Marjan Mansourian³

1. Department of Immunology, Faculty of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

2. Multiple Sclerosis and Neuroimmunology Research Center, Isfahan University of Medical Sciences, Isfahan, Iran

3. Department of Biostatistics and Epidemiology, Faculty of Medicine, Isfahan University of Medical Science, Isfahan, Iran

Background: Semaphorin 3A (Sema3A) as an immune modulator could be involve in the pathogenesis of autoimmune diseases and act as a suppressor of immune cell over-activation.

Methods: In the current study, we aimed to investigate expression of sema3A in peripheral blood mononuclear cells (PBMCs) and its serum level in relapsing-remitting multiple sclerosis (RRMS) patients. Fifteen newly determined and untreated RRMS patients were chosen and assessed in relapsing and remitting phases in compare with fifteen healthy individuals.

Results: In consistent with previous findings in other autoimmune diseases, our results revealed that serum level of Sema3A and its expression in PBMCs of RRMS patients were significantly lower than in normal subjects. We also evaluated this down regulation has predictive value with ROC analysis.

Conclusion: According to our data, we suggest that Sema3A could be participated in the pathogenesis of MS and might be a potential diagnostic biomarker for the disease.

Keywords: Semaphorin-3A, Multiple sclerosis, Autoimmune disease, Immune modulator

ICIA2018\Tolerance and Autoimmunity\Poster Discussion\7446

7446

Evaluation of the *T-bet*, *RORγt* and *Foxp3* Gene Expression in Patients with Type 2 Diabetes Mellitus

Zahra Telikani ^{1,2}, Vida Sheikh ³, Shiva Borzouei ³, Mortaza Ghasemi ³, Alireza Zamani ¹, Iraj Salehi ⁴, Mahdi Alahgholi Hajibehzad ^{1,*}

1. Department of Immunology, Hamadan University of Medical Sciences, Hamadan, Iran

2. Student Research Committee, Hamadan University of Medical Sciences, Hamadan, Iran

3. Department of Internal Medicine, Clinical Research Development Unit of Shahid Beheshti Hospital, School Of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

4. Neurophysiology Research Center, Hamadan University of Medical Sciences, Hamadan, Iran

Background: T lymphocytes are related to inflammation and insulin resistance in type 2 diabetes mellitus (T2DM) and T2DM plus nephropathy (T2DMN). The imbalance between pro-inflammatory and inhibitory lymphocytes plays a major role in development of T2DM. The objective of this study is to investigate the gene expression of *T-bet*, *RORγt* and *Foxp3* related to T helper (Th) 1, Th17 and Regulatory T (Treg) cells subsets in T2DM.

Methods: Blood samples from 39 patients diagnosed with T2DM (mean: 55.7 ± 1.12 ; female=29, men=10), 25 patients diagnosed with T2DMN (mean: 55.8 ± 6.9 ; female=19, men=6) and 26 healthy control (HC; mean: 47.3 ± 9.2 ; female=14, men=12) were collected. Peripheral blood mononuclear cells (PBMCs) were isolated from whole blood by Ficoll density gradient method. In order to determine mRNA expression of *T-bet*, *RORγt*, *Foxp3* and GAPDH (reference gene), total RNA was extracted from PBMCs and cDNA was generated. Expression of Th genes was tested/ performed with the SYBR green Real Time PCR method. The relative genes expression level for each sample were calculated by $\Delta\Delta C_t$ method.

Results: The level of *Foxp3* gene expression has shown significant decrease in T2DMN patients compared with HCs ($p=0.04$) and but there was no significant change in *RORγt* and *T-bet* gene expression. The level of *Foxp3* gene expression has demonstrated significant decrease in T2DMN patients compared with T2DM patients ($p=0.02$), whereas there was no significant change in *RORγt* and *T-bet* gene expression. No significant differences were observed in gene expression of *T-bet*, *RORγt* and *Foxp3* in T2DM patients compared with HC group.

Conclusion: This result show a decreased *Foxp3* gene expression in T2DMN, which could contribute to the pathogenesis of the disease and revealing a potential novel therapeutic target.

Keywords: T2DM, *T-bet*, *RORγt*, *Foxp3*, Nephropathy

ICIA2018\Tolerance and Autoimmunity\Poster Discussion\7460

7460

Frequency and Functional Evaluation of Treg Cells in Patients with type 2 Diabetes MellitusMahdi Alahgholi Hajibehzad ¹, Elham Mahabadi Ashtiyani ¹, Shiva Borzouei ², Alireza Zamani ¹, Vida Sheikh ²*1. Department of Immunology, Hamadan University of Medical Sciences, Hamadan, Iran**2. Department of Internal Medicine, Clinical Research Development Unit of Shahid Beheshti Hospital, School Of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran*

Background: Regulatory CD4⁺CD25⁺ T cells (Tregs) can suppress harmful immunological reactions to self or foreign antigens. The aim of this study was to evaluate the number and suppressive capacity of Tregs in patients with type 2 diabetes mellitus (T2DM).

Methods: Tregs from 19 T2DM patients and 20 healthy controls (HC) were analyzed for the number and suppression capability using flow cytometry based on surface CD4 and CD25 markers. Treg and responder T cells (Tresp; CD4⁺CD25⁻) were sorted using magnetic activated cell sorting. CFSE-labeled Tresp were cultured and stimulated with anti-CD3 and anti-CD28 for 5 days. Tregs suppression assays were performed and analyzed by flow cytometry.

Results: The addition of Tregs to the culture of Tresp cells significantly decreased the proliferation of Tresp in both T2DM and HC. The % proliferation of Tresp in the presence of Treg cells was lower in HC than T2DM patients. The % suppression of proliferation was significantly lower in patients with T2DM compared with the HC group. No significant difference was observed between proportion of CD4⁺CD25⁻, CD4⁺CD25⁺, CD4⁺CD25^{low} and CD4⁺CD25^{hi} cells in T2DM and HC subjects.

Conclusion: These data demonstrate a normal number, but a decreased functional capacity of Tregs in T2DM, which could contribute to the pathogenesis of the disease and revealing a potential novel therapeutic target.

Keywords: Treg, Tresp, T2DM, Suppression

ICIA2018\Tolerance and Autoimmunity\Poster Discussion\7475

7475

Nanocurcumin Improves Imbalance of Th17/Treg cells in Patients with Multiple Sclerosis

Sanam Dolati^{1,2}, Mehdi Yousefi^{1,2}

1. Stem Cell and Regenerative Medicine Institute, Tabriz University of Medical Sciences, Tabriz, Iran

2. Department of Immunology, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Multiple sclerosis is a chronic autoimmune disease that leads to brain inflammation. Th17 cells are considered to be important in MS pathogenesis. It was reported that a disturbance in the development and function of Tregs are associated with disability in MS patients. In this study, we aimed to identify nanocurcumin effects on Th17 and Treg cells frequency, cytokines secretion, and expression of transcription factors of these cells in patients with relapsing-remitting multiple sclerosis (RRMS).

Methods: 50 patients with RRMS were enrolled in this study in which 25 were treated for at least six months with nanocurcumin capsules while the other half received placebo capsules as the control group. At baseline and after a six-month treatment, the frequency of these lymphocytes, the expression of transcription factor and the serum levels of cytokines were assessed by flowcytometry, RT-PCR and ELISA, respectively.

Results: Nanocurcumin significantly increased the frequency of Treg cells in compared placebo group ($p=0.0027$). The levels of FoxP3, TGF- β and IL-10 mRNA increased following nanocurcumin therapy compared to the basal level ($p=0.0004$), ($p=0.0005$), and ($p<0.0002$), respectively. TGF- β and IL-10 levels were increased after treatment with nanocurcumin; from 193.7 ± 95.45 to 313.5 ± 203.6 , and from 629.1 ± 355.1 to 1038 ± 496.8 , respectively. Even though there is no significant difference in proportion of Th17 cells in placebo group compared with nanocurcumin treated group. ROR γ t and IL-17 mRNA levels were reduced in MS patients after nanocurcumin treated (0.0001) and ($p<0.04$) in compared with basal levels, IL-17A levels were decreased after nanocurcumin treated from 140.8 ± 57.24 to 88.59 ± 48.07 ($p=0.0011$) but there is no noticeable decrease in IL-23 mRNA levels and concentration in nanocurcumin group, control group.

Conclusion: The results of the current work indicate that nanocurcumin is able to restore the imbalance of Th17/Treg in MS patients.

Keywords: Multiple Sclerosis, Nanocurcumin, Th17, Tregs

ICIA2018\Tolerance and Autoimmunity\Poster Discussion\8105

8105

MiR-326 was Over-expressed in T cell-derived Exosomes of Patients with Relapsing-Remitting Multiple SclerosisMaryam Azimi Mohamadabadi ¹, Maryam Izad ^{1,2}*1. Department of Immunology, school of Medicine, Tehran University of Medical Sciences, Tehran, Iran**2. MS Research Center, Neuroscience Institute, Tehran University of Medical Sciences, Tehran, Iran*

Background: Multiple sclerosis (MS) is an immune-mediated neurodegenerative disease of central nervous system (CNS). Invasion of autoreactive CD4⁺ T cells into CNS is believed to be an underlying pathogenic mechanism in MS. CD4⁺ T cells release exosomes which are highly enriched in microRNAs, reflective of physiological or pathological condition. Exosomal microRNAs were transferred between cells and affected the physiology of target cells. Thus exosomes could be potent agents to provide quantitative and qualitative information about immune cells involved in MS. We investigated the expression of pathogenic microRNAs in T cells-derived exosomes of MS patients or healthy controls.

Methods: Conventional T cells (Tconv) derived from relapsing-remitting (RR) MS patients or healthy controls were cultured for 3 days by soluble anti-CD3/CD28. Exosomes were purified from supernatants. After RNA extraction from exosomes pellet, the expression levels of miR-146a, miR-29a, miR-155, and miR-326 were quantified by real-time PCR.

Results: A statistically significant increased expression of miR-326 in Tconv-derived exosomes was observed in RRMS patients as compared with controls (7.5 ± 1.88 vs 2.51 ± 0.9 $P=0.03$). On the contrary, no differences were found in the expression levels of miR-155, miR-146a, and miR-29a, in Tconv-derived exosomes of patients as compared with controls ($P>0.05$).

Conclusion: Our results point to an altered expression in exosome-derived microRNAs. MiR-326 was previously shown to play a role in the immunopathogenesis of MS by inducing TH17 differentiation and maturation. Therefore, miR-326 containing exosomes might also be a potential clinical target in course of MS. Moreover, the deregulation of this miRNA in exosomes may serve as a diagnostic and prognostic biomarker.

Keyword: Multiple sclerosis; Lymphocyte; Exosomes; microRNA

ICIA2018\Tolerance and Autoimmunity\Poster Discussion\8531

8531

Methylation Status of the Human Telomerase Reverse Transcriptase Gene Promoter in Patients with Multiple Sclerosis

Seyed Alireza Mesbah Namin^{1*}, Sanaz Hesami¹, Mohammad Ali Sahraian², Rozita Doosti²

1. *Department of Clinical Biochemistry, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran*

2. *Sina MS Research Center, Brain and Spinal Injury Research Center, Tehran University of Medical Sciences, Tehran, Iran*

Background: Multiple Sclerosis (MS) is a chronic inflammatory demyelinating disease of central nervous system. It is believed that both genetic and environmental factors play a role in causing the disease and shows the importance of epigenetic changes in this disease. There is also a relationship between telomere length and telomerase activity in autoimmune and inflammatory diseases which is associated with immaturity and inability of the immune system. The purpose of this study is to evaluate the promoter methylation of the human telomerase reverse transcriptase (hTERT) gene in MS patients.

Methods: In this study, the methylation status of the first CpG island of hTERT gene promoter is investigated by methylation specific PCR in two groups of normal subjects (females: 21 and males: 6) and MS patients (females: 38 and Males: 9). After DNA extraction of blood samples, the concentrated DNAs were treated and modified by sodium bisulfate. After performing of PCR using of specific primers, methylation status was analyzed by the presence of the methylated and unmethylated amplified products.

Results: The results of this study showed that hTERT gene promoter in control and patient's samples were both hypomethylated and there were no significantly differences between the hTERT promoter methylation in MS patients and normal controls, except for one case of patients.

Conclusion: It should be mentioned that study on the epigenetic modification of whole regions of the hTERT promoter is important and this field needs to be extended and further investigations would be considered in more samples.

Keywords: Multiple Sclerosis, Telomerase, Promoter, Methylation

ICIA2018\Tolerance and Autoimmunity\Poster Discussion\8665

8665

Expression Analysis of PD-1 and Tim-3 Immune Checkpoint Receptors in Vitiligo PatientsAli Rahimi¹, Hadi Hossein Nattaj¹, Zohreh Hajheidari², Abolghasem Ajami^{1*}, Hossein Asgarian Omran^{1,3*}*1. Department of Immunology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran**2. Department of Dermatology, Faculty of Medicine, Mazandaran University of Medical Sciences, Sari, Iran**3. Immunogenetics Research Center, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran*

Background: Vitiligo is a skin-related chronic autoimmune disease that degrades melanocytic cells, causing colorless and asymptomatic stains. In recent years, the contribution of immune checkpoint receptors has been addressed in the pathogenesis of multiple autoimmune diseases. In the present study, the expression profile of T-cell immunoglobulin and mucin-domain containing-3 (Tim-3) and programmed cell death-1 (PD-1) checkpoint molecules was investigated on CD8⁺ T cells of patients with vitiligo as the main pathogenic effector cells in this autoimmune complication.

Methods: A total of 30 vitiligo patients and 30 sex- and age-matched healthy controls were included in this study. To determine the frequency of Tim-3⁺/PD-1⁺/CD8⁺ T cells, PBMCs were stimulated with PHA for 72 h and a three-color flow cytometry method was applied. To measure the cytokines production, PBMCs were stimulated with PMA/ionomycin for 18 h and concentrations of IL-4, IFN- γ and TNF- α were measured in culture supernatants by ELISA. CD8⁺ T cells were then positively isolated from all participants by magnetic beads separation method and the mRNA expression of PD-1 and Tim-3 molecules was determined by TaqMan based Real-Time PCR.

Results: Vitiligo patients have significantly showed more expression of Tim-3 and PD-1 on the surface of their CD8⁺ T cells than that of normal controls. While, the production levels of TNF- α and IFN- γ were found higher by vitiligo patients than those of controls, IL-4 production was lower in patients with vitiligo. Expression analysis of Tim-3 and PD-1 mRNA confirmed the results obtained from flow cytometry and showed more expression of Tim-3 and PD-1 in CD8⁺ T cells of vitiligo patients compared to normal individuals.

Conclusion: Our results indicate that Tim-3 and PD-1 are involved in immune dysregulation mechanisms of CD8⁺ T cells in vitiligo. Further studies are needed for better understanding of Tim-3 and PD-1 roles in immunopathogenesis of vitiligo and may introduce useful biomarkers for disease progression and/or immunotherapy.

Keywords: Vitiligo, CD8⁺ T cells, Tim-3, PD-1, Flow cytometry, Real-Time PCR



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Transplantation

Poster Discussion

ICIA2018\Transplantation\Poster Discussion\7435

7435

The Study of the Relationship between Serum Levels of Soluble VEGF Receptor-1 with Delayed Graft Function after Kidney Transplantation

Zahra Hooshyar¹, Saeede Shabanitabar¹, Javad Absari², Majid Faridgozar³, Hassan Nikoueinejad^{*4}, Hossein Akbari⁵, Behzad Einollahi⁶

1. Medical Student; Student Research Committee, Kashan University of Medical Sciences, Kashan, Iran

2. Master Student; Department of Microbiology, School of Medicine, Iran University of Medical Sciences, Tehran, Iran

3. Master Student; Department of Microbiology and Immunobiology, Kashan University of Medical Sciences, Kashan, Iran

4. Medical doctor & Immunologist; Nephrology and Urology Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran

5. Statistician; Trauma Research center, Department of Community medicine, Kashan University of medical sciences, Kashan, Iran

6. Nephrologist; Nephrology and Urology Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran

Background: Delayed graft function (DGF) is a transplant complication which means needing dialysis throughout the first week after transplantation. This study aimed to ascertain the relationship between soluble VEGF receptor-1, as an immunomodulatory factor, and DGF after transplantation.

Methods: This case control study was done in a 6-month period among 2 groups of 58-member of transplanted patients with and without DGF. The control group included the patients who were operated in 2015 and didn't show DGF symptoms. Then, sFlt-1 level in all blood samples was measured by Elisa.

Results: Serum sFlt-1 levels were significantly higher in DGF group compared to those in control group ($P \leq 0.001$). sFlt-1 serum levels significantly affect DGF ($P < 0.001$) in such a way that it increases the risk of DGF (OR=1.1).

Conclusion: This study showed a significant relationship between sFlt-1 and DGF. Therefore, plasma levels of sFlt-1 may be used as a diagnostic tool to determine the risk of DGF.

Keywords: Kidney Transplantation, soluble VEGF Receptor-1, Delayed Graft Function

ICIA2018\Transplantation\Poster Discussion\7681

7681

The Absence of Immune marker HLA-DR Expression and Lack of Rejection in hEnSCs Derived Motor Neurons developed with Small Molecules (Purmorphamin) PMA

Azam Rahimi¹, Homa Mohseni Kouchesfahani¹, Somayeh Ebrahimi Barough², Jafar Ai²

1. Department of Animal Biology, Faculty of Biological Sciences, Kharazmi University, Tehran, Iran

2. Department of Tissue Engineering, School of Advance Technology in Medicine, Tehran University of Medical Science, Tehran, Iran

Background: Human endometrial stem cells (hEnSCs) are being applied in regenerative medicine. EnSCs derived motor neurons are one of the best targets for transplantation which do not stimulate immune system. Cell differentiation using small molecules provide a promising strategy for generating target cells for cell transplantation. We developed an effective and simple induction protocol for differentiating motor neurons from hEnSCs that is useful for transplantation to treat motor neuron disease. Treatments with differentiation media which contain PMA, promote the development of motor neurons.

Methods: During the first phase of differentiation cells were treated with DMEMF\12, FBS 20%, B27, FGF2, 2ME and IBMX for one day. Second environment contained DMEMF\12, N2, B27 and PMA. This phase lasted four days. Subsequent treatment with DMEMF\12, retinoic acid (RA) and PMA lasted 8 days. The last step involve adding environment contain DMEMF\12, N2, B27 and BDNF for 8 days. The whole period lasted 21 days. Then ICC test was done for the expression of Neurofilament (NF) and Acetylcholine transferase (Chat). Then Real-time PCR was performed in RNA stage for expression of neural markers such as NF, Chat, Nestin and GFAP (as glial marker). Then flow cytometry was done for the expression of the cell surface antigen HLA-DR in EnSCs and motor neurons derived from these cells.

Results: ICC results showed the expression of NF and Chat. Real-time PCR results demonstrated that the cell treated with PMA expressing neural markers NF and Chat. The results confirmed that hEnSCs and motor neurons derived from them are not expressing HLA-DR.

Conclusion: On the basis of our finding, in the present research it was stated that motor neurons developed by PMA, do not express HLA-DR. our finding indicated that EnSCs derived motor neurons developed by inducing small molecules do not stimulate immune system. Because of this unique future, they can be used in therapeutic cases such as transplantation and grafts.

Keywords: HLA-DR, hEnSCs, Motor neurons, Purmorphamin.

ICIA2018\Transplantation\Poster Discussion\7999

7999

Comparison of BASILIXIMAB Monoclonal Antibody Transient Expression in two Mammalian Cell LinesReyhaneh Hoseinpoor¹, Masoumeh Rajabibazl², Azam Rahimpour³*1. Department of Biotechnology, School of Advanced Technologies in Medicine, University of Shahid Beheshti Medical Sciences**2. Department of Medical Biochemistry, School of Medicine, University of Shahid Beheshti Medical Sciences**3. Department of Tissue Engineering and Applied cell sciences, Faculty of Advanced Technologies in Medicine, University of Shahid Beheshti Medical Sciences*

Background: Basiliximab (Simulect) is a chimeric monoclonal antibody against the alpha chain of interleukin-2 receptor (IL-2R) that is effective in reducing acute rejection in renal transplantation. The production of recombinant antibodies has been generally recognized as time-consuming and labor-intensive. For high level expression of recombinant antibodies, the most widely used mammalian expression systems are human embryo kidney (HEK-293T) and Chinese hamster ovary (CHO) cells. The aim of this study was to determine which mammalian expression cells can massively and cost effectively produce full-length antibodies.

Methods: The expression vector containing heavy and light chains of Basiliximab was constructed. HEK-293T and CHO-k1 cells were transfected for transient antibody expression using Lipofectamin LTX transfection reagent. To evaluate transfection efficiency, EGFP expression was analyzed 48 h post-transfection using the florescent microscope. The supernatant harvested from cells, was assayed to determine the level of production of antibody by ELISA. The microtiter plate wells were coated with the goat anti-human kappa light chain antibody and goat anti-human IgG (Fc specific)-peroxidase antibody was added for detection of IgG in cell culture supernatants. The amount of the antibody was assessed by comparison to a standard curve.

Results: EGFP fluorescence was observed in both cell lines. Antibody was detected in both culture supernatants. The estimate titer for CHO-K1 cells was up to 6 ng/ml while the titer for HEK293T cells was 3.5 ng/ml. Protein from transient CHO was more compared to protein produced from HEK293.

Conclusion: Production of IgG molecules with desired specificities has become an area of medical and basic research. The choice of host cell for high level expression of recombinant antibodies is critical. Our results here describe the superiority of CHO-K1 cells for transient expression compared with HEK293T cells.

Keywords: Basiliximab, transient Expression, CHO-K1, HEK293T

ICIA2018\Transplantation\Poster Discussion\8209

8209

The Impact of Killer Cell Immunoglobulin-Like Receptors/Human Leukocyte Antigen Class One Combinations on the One-Year Prognosis of Allogeneic Hematopoietic Stem Cell Transplantation in Patients with Acute Leukemia

Marzieh Mansouri,¹ Jean Villard,² Mani Ramzi³, Shirin Farjadian¹

1. Department of Immunology, Shiraz University of Medical Sciences, Shiraz, Iran

2. Immunology and Transplant Unit and National Reference Laboratory for Histocompatibility (LNRH), Geneva University Hospital, Geneva, Switzerland

3. Department of Hematology-Oncology and Stem Cell Transplantation, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Hematopoietic stem cell transplantation (HSCT) is the best therapy for hematologic malignancies such as acute leukemia. Although, graft versus-leukemia (GVL) is an important approach for prevention of relapse, it could be associated with severe graft versus host disease (GVHD). Similar to T cells, NK cells may facilitate engraftment, combat infection, and control malignant cells but do not cause GVHD. KIRs are a family of inhibitory and activating receptors that are expressed mainly by NK cells which bind HLA as ligands. Because of high variability in KIRs, careful selection of donors based on HLA and KIR is essential to optimize HSCT outcomes. Since donor KIRs are potentially encountered with non-self recipient HLA class I epitopes, this study was designed to find out the impact of KIR/HLA combinations on the outcome of HSCT in patients with acute leukemia.

Methods: From March 2015 to March 2016, a total of 31 patients with acute leukemia who received HSCs were followed for one year by chimerism analysis (30, 60, 90, 180 and 365 days after HSCT). During this period, 17 of these patients were excluded from further analysis due to the death of GVHD or GVL. KIRs genotyping of the donors and their corresponding ligands in recipients was also investigated. All alive patients were received HLA-A/B/DRB1-matched HSCs from their siblings. Three patients received HSCs from opposite sex, four patients from ABO-incompatible and one patient from Rh-incompatible donor. Chimerism was not observed in none of the recipients at the studied time points after HSCT in 14 patients who were alive at least for the first year post-HSCT.

Results and Conclusion: All donors had KIR-B haplotype but just five of them carried KIR2DS1 while seven patients were C2/C2, three patients were C1/C2, four patients were C1/C1. Our results showed that HSCT from donors with KIR-B haplotype to C2-C2 or C1-C2 bearing recipients was associated with a better outcome during the first year after transplantation. Obscure points will be cleared by analysis of chimerism and KIR/HLA combinations in the samples of expired patients.

Keywords: Leukemia, Human leukocyte antigen , Killer cell immunoglobulin like receptor

ICIA2018\Transplantation\Poster Discussion\8571

8571

Association of Epinephrine and Norepinephrine Plasma Levels and Expression Levels of β_2 -adrenergic Receptor Gene with Acute GVHD

Zahra Momeni¹, Aliakbar Movassaghpour¹, Mehdi Talebi¹, Mohammad Hossein Kazemi^{2,3}, Elham Roshandel³, Abbas Hajifathali³

1. Hematology & Oncology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

2. Department of Immunology, School of Medicine, Iran University of Medical Sciences, Tehran, Iran

3. Hematopoietic Stem Cell Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Graft versus host disease (GvHD) is the major complication of hematopoietic stem cell transplantation (AHSCT). Epinephrine (E) and norepinephrine (NE) are stress hormones recently found to have immunomodulatory properties and can decrease the probability of GvHD through interaction with adrenergic receptors. In this study, we have compared the levels of epinephrine and norepinephrine in plasma and also the expression of β_2 -adrenergic receptor gene in peripheral blood mononuclear cells (PBMCs) of patients who develop acute GvHD (aGvHD) with recipients without aGvHD.

Methods: We have studied 15 candidates for AHSCT with median age of 38 years old (range: 25-45). Blood samples were taken from patients 7 days before and after of transplantation and plasma levels of E and NE were measured using Enzyme-linked Immunosorbent Assay (ELISA). After extracting RNA from buffy coats and cDNA synthesis, the expression of β_2 -adrenergic receptor were measured using realtime-PCR. The occurrence of aGVHD were measured according patient's clinical manifestations. T-test was used to analyze data of the study.

Results: Our results showed that 5 patients out of 15 patients presented aGVHD during 100 days after AHSCT. The mean levels of E and NE in plasma and also the expression of β_2 -adrenergic receptor in PBMCs of patients who presented aGvHD were significantly lower than recipients without aGVHD (P-value <0.05).

Conclusion: According to the lower levels of E and NE in plasma and lower expression of β_2 -adrenergic receptor in PBMCs of patients with aGVHD in comparison with recipients without this complication, we concluded that these stress hormones and their receptor are associated with lower immunomodulation and consequently might have a role in developing aGVHD and could be a marker for prediction and management of aGVHD.

Keywords: Epinephrine, Norepinephrine, β_2 -adrenergic receptor, aGVHD

ICIA2018\Transplantation\Poster Discussion\9741

9741

Pre-transplant Serum Level of ferritin as Prognostic Markers of Bacterial Infection in Hematopoietic Cell Transplantation

Maryam Nikoonezhad^{1,3}, Mahshid Mehdizadeh³, Elham Roshandel^{2,3}, Abbas Hajifathali^{3*}

1. Department of Immunology, faculty of Medical Sciences, Tarbiat Modares University

2. Department of Hematology, faculty of Medical Science, Tarbiat Modares University

3. Taleghani Bone Marrow Transplantation Center, Shahid Beheshti University of Medical sciences, Tehran, Iran

Background: Although the kind of antibiotics is used to prevent bacterial infection after hematopoietic cell transplantation (HCT), due to the growth of antibiotic-resistant microorganisms, bacterial infection is one of the barriers to the success of HCT, so it is important to identify patients who are at risk of infection. Ferritin, CRP and ESR are considered, as a potential biomarker for predicting bacterial infection. Also, recently some studies have shown high levels of pre-transplant ferritin associated with mortality and morbidity after autologous or allogeneic hematopoietic cell transplantation. In this study, we have evaluated the correlation between serum level of ferritin, ESR, CRP and, prevalence of bacterial infection in patients who underwent hematopoietic cell transplantation.

Methods: We studied 64 patients with hematological malignancies who received BM transplants without any antibacterial prophylaxis. We measured the serum level of ferritin, CRP and ESR of these individuals before and after HCT. Infection in these patients was confirmed by blood culture, then studied the association of these markers with post-transplant infections.

Results: The result showed 11 patients had mild to severe bacterial infection after HCT. Serum level of pre-transplant ferritin increased in 5 patients who have bacterial infection after HCT. Also, there was a significant correlation between high levels of ferritin before HCT and infection after HCT but there was no significant relationship between the serum level of CRP and ESR with HCT.

Conclusion: These results suggest that pre-transplant serum ferritin level may be a useful marker for predicting the risk of early bacterial complications after allogeneic HCT.

Keywords: hematopoietic cell transplantation (HCT), ferritin, C-Reactive Protein (CRP), *Erythrocyte Sedimentation Rate (ESR)*, Bacterial Infections.

ICIA2018\Transplantation\Poster Discussion\9797

9797

Evaluation of the Serum Level of Interleukin-6 in Liver Ischemia-reperfusion Injury in RatAzad Bahrapour¹, Roonak Azizbeigi^{2*}*1. DVM Student, Faculty of Veterinary Medicine, Sanandaj Branch, Islamic Azad University, Sanandaj, Iran**2. Assistant of Physiology, Department of Physiology, Faculty of Veterinary Medicine, Sanandaj Branch, Islamic Azad University, Sanandaj, Iran*

Background: Liver ischemia-reperfusion injury occurs in a number of clinical settings, including liver surgery, transplantation and hemorrhagic shock with subsequent fluid resuscitation. It has been well demonstrated that there is a significant cause of morbidity and mortality which is characterized by oxidative stress accompanied with depletion of antioxidants. Interleukin-6 (IL-6) is an acute phase reactant cytokine with pleiotropic biological effects. This cytokine plays a critical role in hematopoiesis, host defense, and inflammation. The studies showed that IL-6 has a key role in anti-inflammatory properties. The aim of this study was to investigate the serum concentration of IL-6 in liver ischemia-reperfusion injury.

Methods: Adult male Wistar rats (n=10, weighing 220-250 gr) were used in this experimental. The animals were divided randomly into two groups 1) surgical control group, and 2) ischemia-reperfusion group. The control group underwent surgery but did not tolerate liver ischemia-reperfusion. The animals in the ischemia-reperfusion group were under surgical procedure and underwent ischemia for 45 minutes, then undergone liver reperfusion for 45 minutes. At the end of the study, serum concentration of IL-6 was evaluated by enzyme-linked immunosorbent assay kit (ELISA).

Results: The results showed that serum concentrations of IL-6 were 38.46 ± 3.64 pg/ml and 385.79 ± 12.96 pg/ml in the control group and ischemia-reperfusion group, respectively. There was a significant difference between the groups ($P < 0.01$).

Conclusion: It can be concluded that liver ischemia-reperfusion injury can induce significant alterations in the serum concentrations of IL-6. It could form the basis for the performance of large scale prospective randomized clinical trials of liver ischemia-reperfusion injury.

Keywords: Liver ischemia-reperfusion, Rat, Interleukin-6

ICIA2018\Transplantation\Poster Discussion\9844.doc

9844

Study of the Association between Genes Polymorphisms miRNA 146G/C and miRNA 196a2C/T and Outcome of Kidney Transplants in Iranian Patients

Anahita Abbasi,¹ Padideh Ebadi,^{2,*} Mohammad Hossein Karimi,¹ Ramin Yaghobi¹

1. Transplant Research Center, Shiraz University of Medical Sciences, Shiraz, IR Iran

2. Islamic Azad University of Shiraz, Shiraz, IR Iran

Background: Acute rejection remains a serious clinical challenge and the most serious complication in the first year of transplantation. The interaction of microRNA (miRNA) in renal transplant rejection represents an important research area for developing clinical programs. The differential expression of miRNAs in the circulation and biopsy of the organism indicates a positive association with the allograft state. Therefore, in this study the pattern of miR-146a G>C and 196a-2 C>T gene polymorphisms was evaluated in kidney transplant patients.

Methods: Tissue samples were collected from 100 renal transplant patients between years: 1386-1392. The miR-146a G>C (rs2910164) and miR-196a-2 C>T (rs11614913) gene polymorphisms were evaluated in kidney transplant patients using in-house-PCR-RFLP methods.

Results: The CC genotype, C and G alleles of the miR-146a G>C (rs2910164) polymorphism is associated with increased risk of transplant rejection in kidney transplant patients (P= 0.003, P=0.01 and P=0.01), respectively. The CC genotype, T and C alleles of the miR-196a-2 C>T (rs11614913) were also significantly more frequent in transplanted patients versus Healthy sample ones (P= 0.02, P= 0.05 and P= 0.05), respectively. But significant associations were not found between miR-196a-2 C>T (rs11614913) polymorphisms with outcomes in kidney transplant recipients.

Conclusions: The CC genotype, G and C allele of the miR-146a and also, the CC genotype, T and C alleles of the miR-196a may be genetic susceptible factors for transplant rejection especially in the patients. It is obvious that further studies are required to validate these findings in a larger population, as well as in patients with different ethnic origins.

Keywords: Kidney transplant, MicroRNA

ICIA2018\Transplantation\Poster Discussion\9938

9938

The Effect of Exercise on Immune Factors Level and FOXP₃ Gene Expression of Renal Transplanted PatientsJamshidian Tehrani Narges¹, Karimi MH^{2*}, Hemati N³*1. Transplant Research Center, Shiraz University of Medical Sciences, Shiraz, Iran**2. Transplant Research Center, Shiraz University of Medical Sciences, Shiraz, Iran**3. Research Center, Shiraz University of Medical Sciences, Shiraz, Iran*

Background: Renal transplantation is considered to treatment for most patients with end-stage of renal failure. Kidney transplanted patients are exposed to different infectious diseases because of prolonged usage of immune suppressant drugs. On the other hand, exercise may induce beneficial effects on immune system. The present study was performed to examine the effects of 12-weeks selective exercise on interleukin 4 and FOXP3 in kidney transplanted patients.

Methods: Twenty three kidney transplanted patients voluntarily and objectively were selected and randomly divided to control (n=10) and training groups (n=13). Exercise groups participated in training program for 12 weeks, three days a week each day 60–90 minutes. During this time the control group did not participate in any regular exercise. The blood samples were taken before and after 12 weeks. Cytokines expression in blood samples were measured by ELISA and gene expressions quantified with real-time PCR. Data were analyzed using t dependent test.

Results: After 12 weeks, there was no difference of IL-4 level and FOXP3 gene expression, between training group and control group was seen however IL-6 was significantly increased and TNF α was significantly decreased in trained group.

Conclusion: It can be concluded that doing exercise may activated immune system in spite of usage of immune suppressant drugs.

Keywords: Renal transplant, Interleukin 4/6, Exercise training, FOXP3, TNF α

ICIA2018\Transplantation\Poster Discussion\9952
9952

Evaluation of Lnc RNA Expression in Rejection and non-Rejection Renal Transplantation

Zahra nekooee zadeh ¹, Mohammad Hosein Karimi², Padide Ebadi³

1. Department of Biochemistry, Islamic Azad University of Shiraz
2. Department of Transplant Research Center, Shiraz University of Medical Sciences
3. Department of Biochemistry, Islamic Azad University of Shiraz

Background: Transplantation is the process of moving cells, tissues or organs from one site to another site. In spite of technical progresses in graft science and immunosuppress drugs there are some barriers to achieve a successful transplantation. One of the most important issues in renal transplantation as an ordinary treatment method is rejection. Recognize of some biomarker and biology mechanism help to reduce the possibility of rejection. Non coding RNA is a functional RNA molecule that is transcribed from DNA but not translated in proteins. Those non coding RNAs that appear to be involved in epigenetic processes can be divided in to 2 main groups the short non coding RNAs(<30nts) and the long non coding RNAs. Lnc RNAs are defined as transcripts longer than 200 nucleotides. In general Lnc RNAs function to regulate gene expression at the transcriptional and post transcriptional level. According to Lnc RNAs in regulation of the expression of the gene, the purpose of this study was using Lnc RNAs as a biomarker in renal transplantation.

Methods: This study we used 30 patients with rejection and non-rejection and then extracted RNA by using of the Trizol and convert RNA to cDNA and examined amount of Lnc RNAs expression by NGS method. We chose more patients for confirmation of changes, first all RNA was extracted by using the Trizol. The second step is to convert isolated RNA to cDNA. By using real time PCR method we measured the amount of Lnc RNAs expression.

Results: This study indicate NR-110279 is up-regulation in rejected transplantation patients and down-regulated in patients with successful renal transplant. A down-regulation of NR-120448, NR-117089 in reject transplantation and up-regulation in patients with successful renal transplant was observed.

Conclusion: According to results maybe NR-110279 upregulation and NR-120448, NR-117089 is down-regulation in reject transplantation can be used as a biomarker. But this is not definitive, must be done by more patients and real time PCR methods, so we can definitely say that is right.

Keywords: Transplantation, Kidney, Lnc RNAs, Rejection.



Vaccine

Poster Discussion

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Vaccine\Poster Discussion\7350

7350

Expression of OMP19 of *Brucella abortus* on the Surface of *Lactobacillus casei* and Evaluation of Its Immunogenicities in Laboratory Animals

Elham Mohammadi^{1, 2}, Mehdi Golchin¹

1. Department of Pathobiology, Faculty of Veterinary Medicine, ShahidBahonar University of Kerman, Kerman, Iran

2. Molecular Microbiology Research Group, Shahid Bahonar University of Kerman, Kerman, Iran

Background: Brucellosis is a zoonotic disease threatening the public health and fails the trading of animals and their products; this has a negative impact on the economic development of a country. Vaccination is the most effective action to control brucellosis. There is no human vaccine against brucellosis, and attenuated live vaccines in animal are useful but have serious disadvantages. The recombinant vaccines are promising candidates for immunization in humans and animals with more efficacies which among them, vector vaccines have significant benefits. *Brucella* cell surface antigens are the first choice to produce a vaccine because they are the primary point of contact between the pathogen and host cells. In *Brucella* genus, there is superficial antigen named OMP19 which is present in all species and biovars of the genus. Theoretically, this antigen can provide protection against all 6 species of *Brucella*. Therefore, it can be considered to produce recombinant vaccine for consumption by majority of hosts. The aim of this study was to produce a vector vaccine using *Lactobacillus casei* strains carrying OMP19 gene and assessment of its immunogenicity in mouse model.

Methods: In this study, the gene encoding OMP19 antigen was amplified and cloned into an expression vector called pT1NX and transformed to *L. casei* cell via electroporation technique. The expression was confirmed using specific antibody against the recombinant protein via immunological screening tests such as western blot, dot blot and immunofluorescence. Finally, recombinant *L. casei* was fed to mice and results were recorded.

Results: Results showed that, the mice group which had received the OMP19 through *L. casei* based vaccine represented a very good general and mucosal immune responses, similar to commercial vaccine recipient group.

Conclusion: Therefore, the vaccine produced in this study could be a very good vaccine candidate against animal and human brucellosis.

Keywords: Brucellosis, vector vaccine, *Lactobacillus casei*, OMP19

ICIA2018\Vaccine\Poster Discussion\7415

7415

Comparison of the LD50 Related to Herpes Simplex Virus type I Using Intraperitoneal and Intraocular Injection Routes in BALB/c MiceParisa Amir Kalvanagh¹, Hesam Karimi², Masoumeh Ebtokara^{1*}, Hoorieh Soleimanjahi², Parviz Kokhaei³*1. Department of Immunology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran**2. Department of Virology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran**3. Department of Immunology, Semnan University of Medical Sciences, Semnan, Iran*

Background: In order to evaluate the efficacy of prophylactic and therapeutic candidate vaccines for dealing with herpes simplex virus-related diseases in vivo, the animal model should be challenged with the relevant pathogen. To meet the challenge, it is necessary to determine lethal dose 50 (LD50) of the pathogen in the animal model affected by it. One of the important parameters in this regard is the choice of the route of pathogen injection. Our aim in this study is to compare the LD50 of herpes simplex virus type I isolated from an Iranian subject using intraperitoneal (IP) and intraocular (IO) routes.

Methods: After measuring the infectious HSV-1 titer by the TCID50 Assay, 1, 10 and 100 X TCID50 of HSV-1 injected through IP and IO injection routes to the female 4-5weeks old BALB/c mice using insulin syringe (3mice/group). Signs of disease including paralysis, infection, redness, and swelling of the eyes, anorexia, and mortality were recorded during 15 days (twice daily).

Results: The LD50 in both injections routes for HSV-1 in BALB/c mice was 1.2X TCID50. IO route practically was more difficult than IP and required experienced people, and there also was the limitation for the volume of injections, and in most cases, the virus needed to be concentrated as well.

Conclusion: Applying the IP injection route for prophylactic vaccines and IO injection route for therapeutic vaccines seem more rational.

Keywords: Herpes Simplex Virus-1, LD 50, challenge, IP injection route, IO injection route.

ICIA2018\Vaccine\Poster Discussion\7663.doc

7663

Immunogenicity of *Plasmodium falciparum* Merozoite Surface Protein-1₄₂ Antigen Formulated with poly (I: C) Adjuvant in BALB/c Mice

Niloufar Rezvani^{1,2}, Akram Abouie Mehrizi^{1*}, Sedigheh Zakeri¹, Laleh Babaeekhou²

1. Malaria and Vector Research Group (MVRG), Biotechnology Research Center, Pasteur Institute of Iran, Tehran, Iran, P.O.BOX 1316943551.

2. Islamshahr Branch, Islamic Azad University, Tehran, Iran

Background: The C-terminal region of *Plasmodium falciparum* Merozoite Surface protein 1 (PfMSP-1₄₂) is a leading malaria vaccine candidate antigen. Nevertheless, the results of clinical trials of MSP-1 based vaccines have not shown complete efficacy. In this regard, the main obstacle is the lack of potent adjuvants suitable for using in human. In this study, to induce high titer of total antibodies and polarized Th1 immune responses against PfMSP-1₄₂, poly (I: C) as a potent and human compatible adjuvant was used.

Methods: The *pfmsp-1₄₂* gene was cloned and expressed in *E. coli* M15-pQE30 expression system. Five test groups of BALB/c mice were immunized with affinity purified recombinant PfMSP-1₄₂ antigen alone or in combination with poly (I:C), poly (I:C)/Alum, Alum or CFA/IFA adjuvants. The control groups received PBS1× or corresponding adjuvants without antigen. Immunization was carried out three times, 2 weeks intervals via subcutaneous rout. Anti-PfMSP-1₄₂ IgG and IgG subclasses were measured 10 days after last immunization.

Results: The results showed that antibody response was in comparable level in the mouse groups that received rPfMSP-1₄₂ in combination with poly (I:C) or CFA/IFA adjuvants. Interestingly, among the five test groups, the ratio of IgG2a/IgG1 antibodies was highest in the mouse group that received rPfMSP-1₄₂ in combination with poly (I:C) indicating a Th1 immune response.

Conclusion: In conclusion, the results showed that in compare with CFA/IFA adjuvant, the poly (I:C) adjuvant could improve the immune responses to rPfMSP-1₄₂ antigen, which could be applied in designing in a PfMSP-1₄₂-based vaccine.

Keywords: Malaria, *Plasmodium falciparum*, PfMSP1-42, Poly(I:C)

ICIA2018\Vaccine\Poster Discussion\7940

7940

***In silico* Evaluation of RNA Based Vaccine Candidate Expressing LmSTI1 and PpSP15 Fusion Protein against Cutaneous Leishmaniasis**Nastaran Sadat Savar^{*1}, Arash Arashkia², Kenneth Lundstrom³, Ali Khamesipour⁴, Hamid M. Niknam¹

1. Immunology Department, Pasteur Institute of Iran, Tehran, Iran

2. Department of Molecular Virology, Pasteur Institute of Iran, Tehran, Iran

3. Pantherapeutics, Switzerland

4. Center for Research and Training in Skin Diseases and Leprosy, Tehran University of Medical Sciences, Tehran, Iran

Background: Leishmaniasis is among neglected diseases, 10 percent of the world population are at risk in 11 high-burden countries including Iran. However, there is still no vaccine available against any form of human disease. Self-amplifying mRNA-based vaccines are considered as the next generation vaccines. The main aims of the present study were to design a fusion form of the *Leishmania major* stress inducible protein 1 (LmSTI1) and *Phlebotomus papatasi* SP15 (PpSP15), and evaluate the stability and conformational changes of structures in RNA and protein levels compared to the native forms.

Methods: PpSP15/LmSTI1 and LmSTI1/PpSP15 fusion forms were designed by placing the PpSP15 gene sequence at the 5' or 3' ends of LmSTI1. RNA folding and minimum free energies were predicted using the RNAfold webServer. Physical and chemical parameters of proteins were computed by the ExPASy ProtParam tool and three dimensional structures of the proteins were modeled by the I-TASSER server. Modeled structures were validated and evaluated using the RAMPAGE and the ProSa web. Conformational studies on modeled fusion and native forms were performed using the Swiss-Pdb Viewer.

Results: According to the RNAfold results, PpSP15/LmSTI1 showed a higher minimum free energy compared to the LmSTI1/PpSP15 fusion form, and it was closer to the LmSTI1 native form. The ProtParam results showed that both fusion proteins were stable, although the half-life of PpSP15/LmSTI1 was estimated to be longer than that of LmSTI1/PpSP15 *in vivo*. Based on calculated RMSDs, the conformational changes of LmSTI1 and PpSP15 in PpSP15/LmSTI1 was shown to be smaller than that of LmSTI1/PpSP15 fusion form.

Conclusion: The RNA encoding PpSP15/LmSTI1 has optimal minimum free energy, at the protein level, smaller conformational changes and a longer half-life was seen. It seems reasonable to believe that this fusion construct can be used as a recombinant antigen in a self-amplifying mRNA vaccine platform against leishmaniasis.

Keywords: in silico study, RNA-based Vaccine, fusion, Leishmaniasis.

ICIA2018\Vaccine\Poster Discussion\8102

8102

Evaluation of the Adjuvant Effect of Agonists of Toll-like Receptor 4 and 7/8 in a Vaccine against *Leishmania major* in BALB/c Mice

Mosayeb Rostamian¹, Hamid M. Niknam^{1*}

1. Immunology Department, Pasteur Institute of Iran, Tehran 13164, Iran

Background: There is no effective vaccine against human leishmaniasis. Achieving successful vaccines seems to need powerful adjuvants. Separate or combined use of toll like receptor (TLR) agonists as adjuvant is a promising approach in *Leishmania* vaccine research. In present study, we evaluated adjuvant effect of separate or combined use of a TLR7/8 agonist, R848 and a TLR4 agonist, monophosphoryl lipid A (MPL) beside soluble *Leishmania* antigen (SLA) in BALB/c mice.

Methods: Mice were vaccinated three times by SLA with separate or combined TLR7/8 and TLR4 agonists and were then challenged by *Leishmania major*. Delay type hypersensitivity, lesion development, parasite load, and cytokines (interferon gamma, and interleukin-10) responses were assessed.

Results: Results showed: 1) MPL can slightly assist SLA in parasite load reduction, but it is not able to increase SLA ability in evoking DTH and cytokine responses or decreasing lesion diameter. 2) R848 does not affect the DTH response and parasite load of mice vaccinated with SLA, but it decreases/inhibits cytokine responses induced by SLA, leading to increase lesion diameter. 3) MPL neutralized inhibitory effect of R848.

Conclusion: In overall, these data emphasize that MPL slightly assists SLA to make a more potent vaccine, but R848 is not a good adjuvant to induce T cell-dependent immune response in BALB/c mice, and therefore combination of these TLR agonists in the current formulation, is not recommended for making a more powerful adjuvant.

Keywords: *Leishmania major*, Monophosphoryl lipid A, R848, Soluble *Leishmania* antigen.

ICIA2018\Vaccine\Poster Discussion\8248

8248

PPD Shifted Cellular Immunity into Th1 Pattern versus Commercial HBS Vaccine Formulated with Montanide ISA 720Bahareh Golkaran¹, Setareh Haghighat², Mehdi Mahdavi^{3,*}*1. Department of Microbiology, Pharmaceutical Sciences Branch, Islamic Azad University, Tehran, IR Iran**2. Department of Microbiology, Pharmaceutical Sciences Branch, Islamic Azad University, Tehran, IR Iran**3. Department of Immunology, Pasteur Institute of Iran, Tehran, Iran*

Background: The best way for prevention of hepatitis B infection is vaccination. Commercial HBs vaccine is formulated in alum adjuvant and does not stimulate cellular immunity in an appropriate manner. PPD is the protein extract derivate from Mycobacterium promoting the cellular immunity strongly. Studies indicate that PPD could improve the immunity responses through affecting on dendritic cells. In this study, the novel formulation of HBs Ag vaccine in companion with PPD was used to stimulate cellular immune responses.

Methods: The commercial HBs Ag vaccine was formulated in 1 and 10 µg of PPD and the mice were injected subcutaneously three times with two week interval with different formulation of vaccines. Total antibody and IgG1, IgG2a isotypes were assessed in serum samples while cytokines level including IL-4, IFN-γ, TNF-α, IL-2 was evaluated in spleen cell culture via ELISA method.

Results: Our findings show that formulation of commercial HBs vaccine in PPD improved Th1 immune response via higher IFN-γ and IL-2 cytokine release

Conclusion: The results of this study showed the ability of PPD as immunopotentiator in vaccine formulation for stimulation of better Th1 cytokine response.

Keywords: HBS Ag1, Vaccine2, PPD3, adjuvant4, montanide ISA7205.

ICIA2018\Vaccine\Poster Discussion\8358

8358

Naltrexone; as an Efficient Adjuvant in Induction of Th1 Immunity and Protection against *Fasciola hepatica* Infection

Hakim Azizi¹, Hadi Mirzaeei², Ali Akbar Nasiri³, Hajar Yaghoobi⁴

1. Department of Medical Parasitology, School of Medicine, Zabol University of Medical Sciences, Zabol, Iran

2. Department of Medical Genetics, School of Medicine, Zabol University of Medical Sciences, Zabol, Iran

3. Department of Anesthesiology, Zabol University of Medical Sciences, Zabol, Iran

4. Cellular and Molecular Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran

Background: Toxic effects of available therapeutics are major drawbacks for conventional management approaches in parasitic infections. Vaccines have provided a promising opportunity to obviate such unwanted complications. In present study, we examined immune augmenting and protective capacities of an emerging adjuvant, Naltrexone, against *Fasciola hepatica* infection in BALB/c mice.

Methods: Seventy BALB/c mice were divided into five experimental groups (14 mice per group) including 1- control (received PBS), 2- vaccine (immunized with *F. hepatica* E/S antigen), 3- Alum-vaccine (immunized with Alum adjuvant and E/S antigens), 4- NLT-vaccine (immunized with NLT adjuvant and E/S antigen), and 5- Alum-NLT-vaccine (immunized with Alum-NLT mixed adjuvant and E/S antigen). Lymphocyte stimulation index was assessed by MTT assay. Production of IFN- γ , IL-4, IgG2a and IgG1 were assessed by ELISA method.

Results: Results showed that NLT, either alone or in combination with alum, can induce immune response toward production of IFN- γ and IgG2a as representatives of Th1 immune response. Also, using this adjuvant in immunization experiment was associated with significantly high proliferative response of splenocytes/lymphocytes. Utilization of Alum-NLT mixed adjuvant revealed the highest protection rate (73.8%) in challenge test of mice infected with *F. hepatica*.

Conclusion: These findings suggest potential role of NLT as an effective adjuvant in induction of protective cellular and Th1 immune response against fasciolosis.

Keywords: Naltrexone, vaccination, Fasciolosis, *Fasciola hepatica*

ICIA2018\Vaccine\Poster Discussion\8379

8379

Immunogenicity Oral Value of *B.mellitensis* bp26 Gene Expressed in *Lactococcus lactis* as a Food Grade Vector in Mice Balb/cSeyed Davood Hosseini ^{*1}, Maryam.Azizpour.m (PhD) ², Parvaneh Jafari ², Mohadesehsadat Hosseini³

1. Razi vaccine and serum Research institute, Arak Branch, Arak, Iran

2. Department of Microbiology, Islamic Azad University, Arak Branch, Arak, Iran

3. Faculty of Biotechnology, Alzahra University, Tehran, Iran

Background: Brucellosis is still an important health problem in under developing countries. Several animal vaccines are produced but the complete protection is not achieved. A good candidate for brucellosis vaccine is *B.mellitensis* bp26 gen. Today, *Lactococcus lactis* with several positive characteristic are attractive for protein expression. These fast growing bacteria no need to aeration, easy to handling, have not exotoxin, endotoxin and protease, so the cost of culturing is inexpensive.

Methods: In this study, *B.mellitensis* bp26 gen was cloned in food grade PNZ 8149 vector and expressed in *L.lactis* NZ 3900. Fifteen BALB/c mice were divided into two groups (groups A–B). Group A mice were orally gavaged with 2.5×10^9 CFU per mL of transformed *L.lactis* with PpNZ + bp26. Group B mice were orally gavaged with 2.5×10^9 CFU per mL of *L.lactis* with pNZ. So, immunosorbent assay were conducted.

Results: Data shown the mean absorbance readings of the mice prior to and post immunization, were different significantly. In the three weeks following vaccination, the recombinant 8149 + bp26 treatment group showed an increase in antibody level whereas the non-recombinant *L. lactis* showed weak responses. The antibody levels were increased significantly in the 8149 + bp26 vaccinated group in relation to the non-recombinant *L. lactis* group ($P < 0.05$).

Conclusion: *L.lactis* can be used as a live delivery vector.

Keywords: *Lactococcus lactis*, *B.mellitensis*, bp26 gen, OMP28

ICIA2018\Vaccine\Poster Discussion\8455

8455

A Lower mRNA Expression level of TLR2, 3, 4 in PBMCs Isolated from non-responders Children to HBV Vaccine

Kousar Smailneja ganji ^{1, 2}, Mehdi Shahbazi ², Mousa Mohammadnia-Afrouzi ², Mojdeh Ghias Tabari³, Motahareh Ehsani², Majgan Bagherzadeh²

1. Student researchcommittee, Babol University of medical sciences, Babol,Iran

2. Department of Immunology, School of Medicine, Babol University of Medical Sciences, Babol, Iran

3. Department of Biochemistry, School of Medicine, Babol University of Medical Sciences, Babol, Iran

Background: Toll like receptors (TLR) have an important role in the innate immune system and also able to recognize Hepatitis B virus that exert effect on immune response. Many host related factors such as the expression of TLRs, contribute to vaccine response. This study was designed to determine the expression levels of TLR2, 3, 4 genes as indicator of innate immunity response to the hepatitis B vaccine among 3-5 years old children.

Methods: The study was carried out among 60 children with age 3-5 years old who received HBV vaccine (43 responders, 17 non responders). Blood samples were collected, then the expression of TLR2, 3, 4 genes in peripheral blood mononuclear cells (PBMCs) were analyzed by Real Time PCR method.

Results: The results of this study indicated that the average expression of TLR2, 3, 4 genes in the responder group was higher than the non-responder group. Additionally, the expression levels of TLR 3, 4 genes were significantly higher in responders compared to non-responders (P-Value \leq 0.01).

Conclusion: This study suggest that the expression of TLR2, 3, 4 could be considered as a marker of protection along with serological markers. Indeed, evaluation of the HBs antibody titer and the expression level of immune response genes together can also be used to determine efficiency of hepatitis B vaccine in responder and non-responder individuals.

Keywords: Hepatitis B vaccine, TLR2, 3, 4 genes and Real Time PCR

ICIA2018\Vaccine\Poster Discussion\9963

9963

Oral Administration of Mice with Chimeric Protein Containing Urease, Omp31 and Omp19 Induces High Protection against *Brucella melitensis* and *Brucella abortus* InfectionsMorteza Abkar ¹, Saeed Alamian ², Mahdi Fasihi Ramandi ³, Abbas Sahebghadam Lotfi ^{*4}*1. Nanomedicine and Nanobiology research center, Shiraz University of Medical sciences, Shiraz, Iran.**2. Brucellosis Department, Razi Vaccine and Serum Research Institute, Karaj, Iran.**3. Molecular Biology Research Center, System biology and poisonings institute, Baqiyatallah University of Medical Sciences, Tehran, Iran.**4. Department of Clinical Biochemistry, Faculty of Medicine, Tarbiat Modares University*

Background: Brucellosis is a group of closely related zoonotic bacterial diseases caused by members of the genus *Brucella*. It was demonstrated that *Brucella* (B) Urease, *B. melitensis* Omp31 and *B. abortus* Omp19 are promising candidates for subunit vaccines against brucellosis. It was hypothesized that application of three proteins can increase vaccine efficiency.

Methods: Three proteins were attached by hydrophobic linkers. Particle size and loading efficiency of the nanoparticles were determined. Immunogenicity of protein with Freund's adjuvant and N-trimethyl chitosan (TMC/antigen) nanoparticles as well as effect of its administration routes (i.p. and oral) on immunological responses and protection was investigated in mice. Antibody detection, cytokine measurement and protection assay were performed. Finally, immunized mice were challenged with the virulent *B. melitensis* 16M and *B. abortus* 544.

Results: Bioinformatics analysis showed chimeric protein has proper stability and conformation, and it can potentially induce humoral and cellular immune responses. After protein expression and purification, chimeric protein was loaded onto TMC nanoparticles. The results indicated that i.p. immunization of chimeric protein with Freund's adjuvant and TMC nanoparticles induces Th1-Th2 immune responses, whereas oral administration of the chimeric protein elicited Th1-Th17 immune responses. Oral administration of chimeric protein encapsulated in TMC nanoparticles displayed the highest protection level (2.58 and 2.37, respectively) against virulent strains of *B. abortus* and *B. melitensis*.

Conclusion: The chimeric protein can be a proper oral vaccine candidate against brucellosis.

Keywords: Chimeric Protein, Trimethyl Chitosan, Oral Vaccine.



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Veterinary Immunology

Poster Discussion

ICIA2018\Veterinary Immunology\Poster Discussion\7812

7812

Production, Purification and Evaluation of Specific Egg Yolk Immunoglobulin (IgY) Against *Vibrio fluvialis* of Hyper Immunized Hen Egg Yolks

Mojtaba Mirzaei¹, Mohsen Najimi², Mohammad Jahantigh³, Firoz Ebrahimi⁴, Abbas Jamshidian², Narjes Hasan Fakhrabadi⁵

1. Student of Department of Pathobiology School of Veterinary Medicine University of Zabol. I.R. Iran
2. Department of Pathobiology School of Veterinary Medicine University of Zabol I.R. Iran
3. Department of Clinical Science School of Veterinary Medicine University of Zabol I.R. Iran
4. Department of Biotechnology School of Basic Science University of Imam Hossein I.R. Iran
5. Student of Department of Pathobiology School of Veterinary Medicine University of Zabol. I.R. Iran

Background: New technologies, such as immunoglobulin egg yolk (IgY), has been in progress for therapeutic and diagnostic purposes and has been used to control many infections and results of various studies have shown that IgG immunoglobulin has inhibitory effect on the growth of many bacteria in liquid medium and animal body, the tendency to use egg yolk is increasing to produce polyclonal antibodies for application and economic purposes. *V. fluvialis* cause diseases in Human and fish. This bacterium is one of the most important factors in the development of cholera like diarrhea in human beings in underdeveloped countries.

Methods: The purpose of this study was to produce, purify and evaluate specific immunoglobulin (IgY) against *V. fluvialis* in hyper-immunized hen egg yolks. For this purpose, we immunized laying hens with formalin killed bacterial cells, IgG against *V. fluvialis* was detected in the serum of the laying hens, then IgY (IgG transferred to the yolk) using PEG 6000 from egg yolks were purified then Dialyzed. IgY evaluation was initially investigated by proteinometry to ensure produced product contains protein (IgY), then used SDS PAGE method to evaluate the purity and nature of protein. Therefore, the presence of this antibody in the purified product was confirmed. Then, to evaluate the effectiveness of this antibody, Indirect ELISA, accomplished.

Results: Concentration of 38 ng IgY/well and above, react with the antigen at Indirect ELISA. The microbial growth inhibitory assay was conducted and results showed that IgY specific anti-*V. fluvialis* with a concentration of 20 mg/ml would inhibit bacterial growth in liquid medium.

Conclusion: According to the results, it seems that the use of IgY biotechnology to produce specific antibody against *V. fluvialis* could control pathogenicity of bacterium easily and at low cost and can be a good alternative to antibiotics.

Key words: egg yolk immunoglobulin (IgY), *V. fluvialis*, Indirect ELISA, SDS-PAGE

ICIA2018\Veterinary Immunology\Poster Discussion\7978

7978

Comparison of Three Precipitation Methods for Isolation of IgY from Chicken Egg Yolk

Adibe Rahmani¹, Morad Rahimi¹, Shahrokh Navidpour², Ali Mostafaie^{3c}, Pantea Mohammadi³, Shahram Parvane³, Ahmad Bagheri³

1. Department of clinical sciences, Faculty of Veterinary Medicine, Razi University, Kermanshah, Iran

2. Razi Reference Lab. of Scorpion Research (RRLS), Razi Vaccine and Serum Research Institute, Karaj, Iran

3. Medical Biology Research Center, Kermanshah University of Medical Sciences, Kermanshah, Iran

Background: IgY, the major immunoglobulin class in egg yolk, first described in 1893 when Klemperer reported that immunized chickens produce antibodies detectable in their egg yolks as well as their blood. In this study we described three different precipitation methods to isolate IgY from egg yolk of chickens immunized against scorpion venom.

Methods: Eggs from chickens immunized with scorpion venom in poultry farm (Veterinary Faculty, Kermanshah) were collected. For extraction, egg yolks were separated and diluted in distilled water adjusted to a pH 5. Then it was subjected to a freeze-thaw cycle and centrifuged to isolate the water soluble fraction. The supernatant was divided into three identical volume that precipitated by three methods: *Polyethylene glycol, ammonium sulfate and sodium chloride precipitation*. For each precipitants, sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) was performed using 12.5% separating and 5% stacking gels. Antibody activity of different extractions were identified by an indirect ELISA assay.

Results: The total activity of IgY fractions precipitated by PEG, *ammonium sulfate* and NaCl were found to be similar and higher than untreated egg yolk. SDS-PAGE analysis indicated that purity of IgY isolated by PEG precipitation method has the highest purity of 94.7%.

Conclusion: PEG precipitation is a simple and mild technique which separates proteins by virtue of their size and solubility. Based on the present results, PEG fractionation method appeared to be more efficient in the isolation of the IgY, while there is no need for any further purification techniques.

Keywords: IgY, Purification, PEG, precipitation, Salt



ICIA2018\Veterinary Immunology\Poster Discussion\8170

8170

Evaluation of FMDV Vaccine on Immune ResponseAbbas Azadmehr¹, Reza Norian², Niloufar Sadat Miri¹, Nowruz Delirezsh²

1. Department of Immunology, Babol University of Medical Sciences, Babol, Iran

2. Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Background: Foot-and-mouth disease virus (FMDV) is a picornavirus which is highly infectious and economically devastating disease of animals. Inactivated foot-and-mouth disease (FMD) vaccines are currently used worldwide but disease still affects millions of animals around the globe.

Methods: This study measured the serum antibodies present in following vaccination with the tetravalent FMDV vaccine two times with a month interval. Serum antibody titers were measured in days 0, 7 and 14 after first vaccination and day 7 after booster, in order to determined high and low titer responder groups of antibodies. Two weeks after booster the PBMC of these groups were isolated, in order to evaluate lymphocyte proliferation at the present of mitogens and specific antigen, by MTT test. The stimulation index (SI) of proliferation in high responder group was higher than low and control groups ($P < 0.05$). On the third day of culture, the concentrations of IL-2, IL-4, IL-5, IL-6, IL-10, IL-17, IFN- γ were assayed by ELISA method.

Results: The result showed that increased in IL-2, IL-4, IL-5 and IFN- γ level of high responder group means that the immune response to vaccine bias toward Th2 cells and humoral immune response, while in low responder group, poor response of Th2 cells was shown presumably because of Th17 cells.

Conclusion: In conclusion, the low immune response to FMDV vaccine may be because of higher stimulation and cytokine release of Th1/Th2 in comparison of Th2 cell types.

Keywords: FMDV, Immune response, Antibody, Cytokine, Vaccine, Mitogen

ICIA2018\Veterinary Immunology\Poster Discussion\8362

8362

The Effect of Levamisole on the Immune System During Thiopental Sodium-induced Anesthesia in Male Mice

Mahdieh Raeeszadeh^{1*}, Javad Beheshtipour²

1. Department of Basic Sciences, Sanandaj Branch, Islamic Azad University, Sanandaj, Iran

2. DVM Student, Sanandaj Branch, Islamic Azad University, Sanandaj, Iran

Background: One of the most important concerns during surgery is anesthetic-induced immunosuppression. Immunomodulators are used to dominate this condition. Immunomodulators are compounds that are capable to regulate the immune system. It is frequently utilized to refer to substances that boost the immune response. Levamisole, a nonspecific immunomodulator, is able to amplify both humoral and cellular immune responses. It has been used to enhance immunity in various diseases. Hence, the aim of this study was to determine the effect of levamisole on the immune system during thiopental sodium-induced anesthesia in male mice.

Methods: This study was conducted on 30 male mice (30 ± 5 g). The mice were randomly divided into three groups (ten mice per group). The control group (N) received intraperitoneally (IP) normal saline. The first (T1) and second (T2) experimental groups received levamisole (20 mg/kg, IP) for 5 and 10 days, respectively. Then, all three groups were anesthetized by thiopental sodium (0.5%, 50 mg/kg, IP). In order to determination of the leukogram and neutrophil/lymphocyte ratio (NLR), blood samples were obtained from the orbital sinus before, during and after the study. After thiopental sodium administration, induction and recovery times of anesthesia were recorded. Data were analyzed using the ANOVA and Tukey's post hoc test.

Results: The leukocytosis was the remarkable leukogram outcome of the experimental groups. The NLR was significantly increased in the T2 group ($P < 0.01$). The anesthesia induction time shows no significant difference between the control and experimental groups ($P > 0.05$). The recovery time in the T2 group was significantly lower than the N group ($P < 0.05$).

Conclusion: The results of this study showed that the levamisole can be used to enhance the immune system in anesthetic condition. Clinical trials are required to characterize the impact of levamisole on the immune system.

Keywords: Levamisole, Leukogram, Anesthesia, Mice

ICIA2018\Veterinary Immunology\Poster Discussion\8476

8476

Bee Venom Melittin Effect on Human and Dog Neutrophil Respiratory Burst

Fatemeh Noparast¹; Sedigheh Nabian^{1}; Mohammad Taheri²; Mohammad Hasan Bemanian³; Abbas Gerami¹; Toran Sepehrzad³*

1. Department of Parasitology and Honey bee , Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran

2. Rastegar Reference Laboratory, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran

3. Department of Allergy and Clinical Immunology, Hazrat Rasoul Hospital, Iran University of Medical Sciences, ,Tehran,Iran

Background: The therapeutic application of bee venom has been explained in traditional medicine for treatment of some diseases. Beneficial effects of bee venom and its components have been previously described. One of the most important components of bee venom is melittin. It has been demonstrated to be considered as the anti-inflammatory, anti-viral and anti-bacterial effects, in different cell types. This study was aimed to evaluate the effect of melittin on the respiratory burst of dog and human Neutrophils using chemiluminescence assay.

Methods: Neutrophils were collected from three heparinized blood samples of dogs and human using ammonium chloride lysis. The chemiluminescence assay was performed using the luminol to compare the effect of melittin on the respiratory burst function at different melittin doses.

Results: The findings showed decreased respiratory burst process in the majority of samples at different doses of melittin, especially at dose of 2.4 ng/ml, while melittin increased the respiratory burst in dogs and human at doses of 0.3 and 0.6ng/ml.

Conclusion: Our findings revealed that melittin might affect the respiratory burst at especial doses. Further comprehensive studies are needed to provide new possibilities for treatment of inflammation related diseases.

Key words: Melittin, Respiratory burst, Neutrophils, Dog, Human

ICIA2018\Veterinary Immunology\Poster Discussion\8490

8490

Allelic Polymorphism of Ovar-DRB1 Second Exon in Iranian Lori-Bakhtiari Sheep Breed

Seyed Vahid Ganjiani¹, Elham Sadat Hosseini Nasab², Atefeh Esmailnejad³

1. Department of Pathobiology School of Veterinary Medicine Shiraz University Shiraz Iran

2. Department of Pathobiology School of Veterinary Medicine Shiraz University Shiraz Iran

3. Department of Pathobiology School of Veterinary Medicine Shiraz University Shiraz Iran

Background: The major histocompatibility complex (MHC) is the best-characterized genetic region associated with resistance and susceptibility to a wide range of diseases. Response to specific antigens is closely related to the MHC genes and their associations might therefore provide precious answers to main questions about the host–pathogen interactions. MHC of sheep is known as Ovar and is located on chromosome 20. Among Ovar MHC class II genes, DRB1 locus has been found to be highly polymorphic and associated with resistance and susceptibility to infectious diseases.

Methods: Ovar-DRB1 second exon allelic diversity was determined in 100 Iranian Lori-Bakhtiari sheep breed using polymerase chain reaction (PCR) and restriction fragment length polymorphism (RFLP) technique using *RsaI* restriction enzyme.

Results: PCR–RFLP identified seven distinct *RsaI* restriction patterns (*a*, *b*, *c*, *d*, *f*, *g*, and *h*) and 22 genotypes in Lori-Bakhtiari population. Pattern *g* had the highest (20.5 %), and Pattern *h* the lowest (1.0 %) frequency. High level of heterozygosity (82 %) and good genotype frequency fit to the Hardy–Weinberg equilibrium was observed in this population ($P=0.26$).

Conclusion: Ovar-DRB1 genotyping indicated a high level of polymorphism in the studied population. Direct sequencing is recommended for detecting allele sequences and subtypes.

Keywords: Major histocompatibility complex (MHC); Ovar-DRB1; Polymorphism; Lori-Bakhtiari

ICIA2018\Veterinary Immunology\Poster Discussion\8623

8623

Selection of High-affinity ssDNA Aptamers for Diagnostic of Infectious Hematopoietic Necrosis Virus (IHNV)Rezvan MohammadiNezhad¹, Seyed Amir Hossein Jalali^{2,3}, Hamid Farahmand⁴*1. Department of Biotechnology, College of Agriculture and Natural Resources, University of Tehran, Iran**2. Institute of Biotechnology and Bioengineering, Isfahan University of Technology, Isfahan, Iran**3. Department of Natural Resources, Isfahan University of Technology, Isfahan, Iran**4. Department of Fisheries, Faculty of Natural Resources, University of Tehran, Iran*

Background: Aptamers are single strand DNA or RNA molecules that can bind to a broad range of targets. Aptamers are selected by an iterative process known as Systematic Evolution of Ligands by Exponential Enrichment (SELEX). Due to various advantages such as high temperature-stability, animal free, cost effective production and high affinity and selectivity for the target molecule, aptamers are considered as attractive alternative to the monoclonal antibodies for diagnostic and therapeutic purposes. The objective of this study was to select DNA aptamers that can specifically bind to Infectious Hematopoietic Necrosis Virus (IHNV) nucleoprotein via SELEX method.

Methods: The selection was started with a synthetic ssDNA library of 10^{18} random molecules with 80 bp length. The purified recombinant IHNV nucleoprotein immobilized on Ni-NTA resin was used as the target protein. The recovered aptamers of each round were amplified by asymmetric PCR and used as new pool for the next round. Starting from the sixth round, counter selection with a very similar protein to IHNV-nucleoprotein was also performed to improve specificity. After 12 rounds, selection of high affinity aptamers to target protein was assessed using Enzyme Linked Oligonucleotide assay (ELONA).

Results: ELONA data from 12 rounds of selection showed that the percentage of specific aptamers to IHNV nucleoprotein was continuously increased during the first 9 rounds of selection and then reached a constant level. The binding affinity of aptamers from 9th round in ELONA assay was comparable to the binding affinity of anti-IHNV nucleoprotein monoclonal antibody in ELISA assay. No cross-reactivity was observed between selected aptamers and non-specific targets.

Conclusion: Finding of this study suggests that aptamers may be a good alternative detection probe to antibodies for rapid and specific detection of IHNV virus.

Keywords: Aptamer, SELEX, IHNV, Nucleoprotein

ICIA2018\Veterinary Immunology\Poster Discussion\9880

9880

An Immunoblot Test Based on Whole Tachyzoites Lysate for Diagnosis of Humoral Response of Domestic Cats Infected with *Toxoplasma gondii*, Rh StrainMorteza Hosseinienejad*¹, Hosseini F¹*1. Faculty of Veterinary Medicine, Shahrekord University, Shahrekord-Iran**2. DVM Student of Veterinary Medicine, Shahrekord University, Shahrekord-Iran*

Background: *Toxoplasma gondii*, an obligate intracellular protozoan parasite infects human and all warm blooded animals. Three practical ways to diagnose *T. gondii* infection in living cats include: parasite isolation, fecal examination for oocysts and serologic tests. Isolation of the parasite is difficult, costly and time consuming and it is not easy to perform in routine diagnostic laboratories. The aim of this study was to detect anti-*Toxoplasma gondii* antibodies in cats infected with Rh strain of *T. gondii* by using an immunoblotting method.

Methods: eight cats were experimentally infected using tachyzoites harvested from Vero cell cultures. Tachyzoites were then lysed and transferred to polyacrylamide gels followed by blotting to PVDF membranes. An immunoblotting was performed using these membranes to detect IgG antibodies.

Results: In the present study, positive lanes were detected in regions related to the presence of *T. gondii* surface antigens, dense granular or rhoptry proteins. No positive lanes were detected in serum samples of kittens received only Vero cell lysates (controls). Positive or negative results were in concordant to the results of IFAT.

Conclusion: The results of this study revealed that whole tachyzoites antigen based immunoblotting is an appropriate diagnostic test for serological detection of anti-*T. gondii* antibodies in recently infected kittens with this Apicomplexan parasite.

Keywords: Immunoblotting, Cats, *Toxoplasma gondii*.

ICIA2018\Veterinary Immunology\Poster Discussion\9989
 9989

Beneficial Effects of *Lactobacillus paracasei* on Isoproterenol-Induced Heart Failure in Rat

ZohrehHesari¹, Khatereh Kafshdouzan^{2*}, Parviz Kokhaei³, Bahador Bagheri ⁴, Sahar Ghaffari⁵

1. Faculty school of of Veterinary Medicine, Semnan University, Semnan, Iran

2. Department of pathobiology, School of Veterinary Medicine, Semnan University, Semnan, Iran

3. Cancer Research Center and Department of Immunology, Semnan University of Medical Sciences, Semnan, Iran

4. Cancer Research Center and Department of Pharmacology, Semnan University of Medical Sciences, Semnan, Iran

5. Veterinary Pathologist, , School of Veterinary Medicine, Tehran University, Tehran, Iran

Background: Heart failure causes extensive complications on the heart by cardiac dysfunction and hypertrophy. The emerging evidences suggest that the administration of different species of probiotics has a number beneficial therapies on heart failure. This study was performed to determine whether oral probiotic administration can attenuate or prevent the heart failure induced by isoproterenol.

Methods: Adult male Wistar rats (n=40, 185±15g), were randomly assigned to five groups, including the control group, probiotic control (0.25 mg/kg/day, oral), isoproterenol control (5 mg/kg/day, subcutaneous), pretreatment group and treatment group were divided. The groups were studied for 30 days. Serum levels of ANP and chemerin were assessed by ELIZA and the standard histological examination was done.

Results: This study showed that ANP in the treatment group compared to control group and isoproterenol control, had a significant reduction (p<0/05) and chemerin were not observed a significant reduction between groups. Pretreatment with *L. paracasei* subpp. *paracasei* 8700:2 could not prevent the development of heart failure induced by isoproterenol.

Conclusion: According to this study, receiving *L. paracasei* subpp. *paracasei* 8700:2 attenuated the serum levels of ANP, a significant improvement in cardiac hypertrophy and the diminishment inflammation.

Keywords: heart failure, isoproterenol, *Lactobacillus paracasei* subpp. *paracasei* 8700:2, ANP and Chemerin.

ICIA2018\Veterinary Immunology\Poster Discussion\10010

10010

Protein Solubilizing Strategies for Production viral Hemorrhagic Septicemia Virus (VHSV) Immunogenic Glycoprotein in *E. coli* Expression System

Marzieh Mohammadian Manesh¹, Seyed Amir Hossein Jalali^{2, 3}, mohammad zare mehrjerdi⁴, Rezvan Mohammadi Nezhad⁵

1. Department of Agricultural Biotechnology, University of Shirvan, Iran

2. Institute of Biotechnology and Bioengineering, Isfahan University of Technology, Isfahan, Iran

3. Department of Natural Resources, Isfahan University of Technology, Isfahan 84156-83111, Iran

4. Department of Agricultural Biotechnology, University of Shirvan, Iran

5. Department of Biotechnology, College of Agriculture and Natural Resources, University of Tehran, Karaj, Iran

Background: Glycoprotein of VHSV, the causative agent for viral hemorrhagic septicemia disease in rainbow trout (*Oncorhynchus mykiss*), is a membrane protein which induce protective immunity against the virus. Hydrophobicity and toxicity of VHSV glycoprotein, prevents soluble expression of this protein in *E. coli* cytoplasm. The present study aimed at developing strategies for soluble expression of VHSV glycoprotein in *E. coli* to be used as subunit vaccine.

Methods: The ectodomain sequence of VHSV-G gene was amplified with specific primers. After digestion with *Hind* III and *Bam*HI restriction enzymes, the fragment was inserted into pET41a (containing GST solubilizing tag for cytoplasmic expression) and pMAL-p4X (containing MBP solubility tag for periplasmic expression) bacterial plasmids. Recombinant constructs were then transformed to three *E. coli* expression hosts including BL21 (DE3), Rosetta (DE3) and Shuffle T7 for protein expression.

Results: Based on SDS-PAGE and dot blotting assays, both GST and MBP fusion tags resulted in soluble expression of VHSV ectodomain. Protein expression level and solubility were greater in *E. coli* cytoplasm as compared to periplasm. No remarkable difference was observed between Rosetta and Shuffle T7 strains in terms of expression level and solubility, while BL21 strain produced low amount of recombinant protein. The expressed protein was purified using Ni-NTA affinity chromatography and its accuracy was validated with western blotting.

Conclusion: The results of this study demonstrated the positive effects of solubilizing tags and engineered host for enhancing soluble expression of highly hydrophobic membrane proteins in *E. coli*. The purified VHSV glycoprotein has the potential to be used as recombinant vaccine for immunization of rainbow trout fish against VHS disease as well as to produce anti-viral antibodies.

Keywords: Membrane Protein, Periplasmic Expression, Cytoplasmic Expression, Solubilizing Tags



Allergy and Immunotherapy of Allergic Disease

Poster Presentation

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Allergy and immunotherapy\Poster\6324

6324

Epitope mapping of Hemiscorpiuslepturus scorpionvenom by means of phage display

Nastaran Mohseni¹, Fatemeh Kazemi Lomedasht^{1*}

1. Venom & Biotherapeutics Molecules Laboratory of Biotechnology Research Center of Pasteur Institute of Iran, Tehran, Iran

Fatemeh Kazemi-Lomedasht (Corresponding author) Ph.D of medical biotechnology (Assistant professor)

Background: Hemiscorpiuslepturus envenomation is recognized as a serious health concern in tropical regions. Selection of peptides mimicking antigenic and immunogenic epitopes of toxins from random peptide libraries is a novel approach for the development of recombinant toxins and poly-epitopic vaccines.

Methods: To achieve this aim, a phage display peptide library and three rounds of bio panning were performed on immobilized antibodies (IgGs).

Results: Our results showed that the highest binding of the phage to immobilized horse antibodies occurred in the third round of bio panning. The sequencing results identified unique peptides mimicking the antigenic and immunogenic epitopes of Hemiscorpiuslepturus toxins.

Conclusion: The results of this study provide a basis for further studies and the development of a putative epitopic vaccine and a re-combinant toxin.

Keywords: Epitope mapping, Hemiscorpiuslepturus scorpion, venom, Phage display

ICIA2018\Allergy and immunotherapy\Poster\6325

6325

Evaluation of food allergens in patients with Atopic dermatitis referring to Mahdiah Hospital in Tehran during the 2017Sogand Nayeri¹, Mahdiah Amiri¹, Pouya Lahoori¹, FatemehTazari^{*2}*1. Department of biology, Faculty of Basic Sciences, Higher Education Institute of AndisheSazan, Neka, Iran**2. Department of biology, Instructor of Microbial Biotechnology, Higher Education Institute of AndisheSazan, Neka, Iran*

Background: Atopic dermatitis is a chronic inflammatory disease in the skin, which often is associated with severe itching. This disease is common in children, which usually is latent in half of the cases, in the first year of life; especially in 60% of cases, those ages are under 16. The purpose of this study was to determine the prevalence of common food allergens in people with atopic dermatitis using prick test in Iran.

Methods: In this study, all individuals referred to Mahdiah Hospital in Tehran during 2017 were examined and in case of allergy were selected. After diagnosis of atopic dermatitis in these patients, Prick skin test was performed by common food allergens.

Results: Among 1100 people with allergies, 306 (approximately 31.4%) were diagnosed with atopic dermatitis. 33% of people at least were sensitive (positive prick test) to one allergen. In people with atopic dermatitis, asthma, allergic rhinitis, and urticaria were 7.1%, 10.8%, and 4.8%, respectively. There was no relationship between gender and food allergy in people with atopic dermatitis. In this study the most common food allergens were yolk (41.2%), glair (39.1%), hazelnuts (35.7%) and peanuts (29.1%), respectively. Foods like rice (5.3%), barley (6.8%), and lamb meat (7.4%) rarely caused allergies.

Conclusion: The prevalence of atopic dermatitis in patients with allergies was more than expected. According to this research, several factors contribute to the prevalence of atopic dermatitis, including the early start of auxiliary feeding, especially with milk and eggs, smoking, woolen and nylon coatings, and keeping pets in the living environment, . Thus, this research have strong suggestion on avoiding these factors.

Keywords: Atopic dermatitis, Chronic disease, Allergic rhinitis, Prick skin test

ICIA2018\Allergy and immunotherapy\Poster\6359
6359

Relationship between depression and job absenteeism in asthmatic patients

Mostafa Aalipour¹. Mohammad Ali Haghghati². Pouria Karimi³. Abolfazl Yari⁴

1. Department of Immunology, School of Medicine, Kerman University of Medical Sciences, Kerman, Iran
2. Department of Management and medical Information School of Management and medical Information Kerman University of Medical Sciences, Kerman, Iran
3. Department of Engineering environmental health, School of Health, Kerman University of Medical Sciences, Kerman, Iran
4. Department of Genetics, School of Medicine, Kerman University of Medical Sciences, Kerman, Iran

Background: Asthma is an immunological respiratory disease that has involved about 300 million people around the world in all races and age groups. The heavy economic burden caused by the disease, care and absence of work due to the illness make asthma a public issue. Previous studies have shown that asthmatic attacks are more in patients who develop both depression and asthma than who suffer from asthma only. Therefore, in this study, the relationship between depression and its impact on occupational life and absenteeism was studied in patients with asthma for the first time.

Methods: This descriptive cross-sectional study was performed on 124 patients with asthma in Kerman during the first half of 1396. The subjects were selected from among the patients referring to Besat Special Clinic and were approved before entering the study. To assess the level of depression and the amount of job absenteeism, the Persian version of the standard questionnaire PHQ-9 and WPAI-Asthma have been used, respectively. For data analysis, SPSS-24 software and Chi-square test have been used.

Results: The results showed that depression had a significant relationship with occupational absenteeism in patients with asthma ($P < 0.05$). Average weekly work in health was 48 hours in subjects under study. 50% of the subjects reported that the disease was effective in reducing their effectiveness in the workplace, and 22% had to rest at home for at least 28 hours a week. This was far higher for industrial workers and agricultural workers.

Conclusion: people who suffer from asthma and depression at the same time lose at least half of their useful time and, in addition, their effectiveness in the workplace is greatly reduced.

Keywords: Asthma, Depression, job absenteeism, PHQ-9,

ICIA2018\Allergy and immunotherapy\Poster\6360

6360

Relationship between depression and quality of life in patients with asthmaMostafa Aalipour¹, Pouria Karimi², Mohammad Ali Haghighati³, Abolfazl Yari⁴*1. Department of Immunology, School of Medicine, Kerman University of Medical Sciences, Kerman, Iran**2. Department of Engineering environmental health, School of Health, Kerman University of Medical Sciences, Kerman, Iran**3. Department of Management and medical Information School of Management and medical Information Kerman University of Medical Sciences, Kerman, Iran**4. Department of Genetics, School of Medicine, Kerman University of Medical Sciences, Kerman, Iran*

Background: Asthma and depression are among the most common diseases that threaten the human community today. Reports have shown that asthma and its deaths have increased dramatically over the past two decades. In recent years, attention has been paid to the correlation between depression and asthma. It has been shown that asthma is a disease associated with a mental condition and increase in people who suffer from depression, in addition to asthma and more asthmatic attacks occurred. This study examines the relationship between depression and quality of life in patients with asthma in Kerman, Iran.

Methods: This descriptive cross-sectional study was carried out on 200 patients with asthma (101 males and 99 females) confirmed their illness and referred to the specialized clinic of Besat in Kerman city in the first half of 1396. To assess the level of depression and quality of life, the Persian version of the PHQ-9 and ASQL questionnaires were used. Data were analyzed by SPSS software version 24 and Chi-square statistical test.

Results: The results showed that depression had a significant relationship with quality of life in patients with asthma ($P < 0.05$). 62% of asthmatic patients showed moderate to severe depressions. 95% of the patients had difficulty in their day-to-day activities, and 43% said that coexistence of depression and asthma create problems in their community activities.

Conclusion: The prevalence of depression in asthmatic patients shows a high percentage and according to the results, this factor has reduced the quality of life in patients. It seems that the study and monitoring of asthma patients with depression and their treatment can increase the quality of life in these patients and is effective in controlling the disease.

Keywords: Asthma, Depression, Quality of Life, PHQ-9, ASQL.

ICIA2018\Allergy and immunotherapy\Poster\7344

7344

Evaluation of IL-10 gene expression following rush immunotherapy in patients with allergic rhinitis

AmirabbasSalmani¹, Farahzad JabbariAzad^{2*}, Mojgan Mohammadi², Reza Farid Hosseini², Jalil Tavakol Afshari³, Maryam khoshkhooi², Ali Fouladvand¹, Sajad Dehnavi¹

1. Department of Immunology, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

2. Allergy Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

3. Immunology Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

Background: Allergic rhinitis (AR) is the most common IgE mediated hypersensitivity disease in the world. Because of high prevalence of AR, it burdens high economic & public health-related cost. Rush Immunotherapy (RIT) is a rapid treatment for atopic diseases, but its immunological mechanisms are not fully understood. In present study, we aimed to evaluate the IL-10 gene expression, as one of cytokines produced by regulatory T-cells, following the rush immunotherapy in patients with AR.

Methods: Fifteen patients with AR who had a positive Prick test for common regional aeroallergens treated with Nuevo RIT protocol (Three-day build up period) for three months. Before and after intervention, clinical symptoms was recorded by questionnaire and whole blood samples collected, RNA extraction from peripheral blood mononuclear was performed then cDNA synthesized to employ for SYBR Green real-time PCR technique. Finally gene expression of IL-10 was determined.

Results: The gene expression of IL-10 increased at three months following the RIT, but it was not significant. Significant improvement in the clinical symptoms was recorded following the RIT. ($P < 0.05$).

Conclusion: As there is no significant difference between the IL-10 gene expression following the RIT, the expression of other cytokine candidate gene of T-regs, such as TGF-beta might be suggested for evaluation in further studies.

Keywords: IL10, Gene Expression, Allergic Rhinitis, Rush immunotherapy

ICIA2018\Allergy and immunotherapy\Poster\7365
7365

Evaluation of Skin and RAST (radioallergosorbent test) test in patients with immediate hypersensitivity following betalactam(amoxicillin/penicillin) drug usage.

Zahra karami^{1,2}, Azam heidari³

1. *Shahidbeheshty university of medical sciences*
2. *Hamedan university of medical sciences*
3. *ShahidSadoughi University of Medical Sciences Yazd*

Background: Skin test is one of the most important methods for diagnosis of immediate hypersensitivity reaction in patients with following beta-lactam but also due to false positive results, in vitro tests should be done.

Methods: we decided to have skin test and radioallergosorbent test (RAST) in 50 patients who took betalactam(amoxicillin/penicillin) with other drugs and showed an immediate hypersensitivity reaction. Skin test was performed with intradermal drug solution and then we demonstrated antibetalactam (amoxicillin/penicillin) specific IgE antibodies by RAST.

Results: Out of 50 patients, 43 patients (86%) had positive skin test results (group A) and 7 patients (14%) had negative test results (group B) and of 43 patients group A, 33 patients (76.7%) have high level of allergen specific IgE (3.50-17.49 KU/L) and 10 patients (23.3%) have absent or undetectable allergen specific IgE (<0.35 KU/L) and of 7 patients(100%) group B, 7 patients have absent or undetectable allergen specific IgE (<0.35 KU/L).

Conclusions: The results show that in patients who took beta-lactam with other drugs and had immediate hypersensitivity reaction, the test results may be due to drugs other than beta-lactam.

Keyword: skin test, Radioallergosorbent test, immediate hypersensitivity reaction, betalactam

ICIA2018\Allergy and immunotherapy\Poster\7373

7373

Relationship between quality of life and Job absenteeism in patients with asthma in Kerman province

Arash Samoudi¹, Abolfazl Yari², Mohammad Ali Haghighati³, Pouria Karimi⁴, Mostafa Aalipour¹

1. Department of Immunology, Faculty of Medicine, Kerman University of Medical Sciences, Kerman, Iran

2. Department of Medical Genetics, Faculty of Medicine, Kerman University of Medical Sciences, Kerman, Iran

3. Department of Health Management, Policy and Economics, Faculty of Health Management, Kerman University of Medical Sciences, Kerman, Iran

4. Department of Environmental Health Engineering, Faculty of Health Engineering, Kerman University of Medical Sciences, Kerman, Iran

Background: Many health care centers are important indicators of life quality assessment studies and practice have understood. Asthma is one of the most common respiratory disorders that can impose serious social and economic consequences on individuals and society. So far few studies have been conducted on the consequences of occupational asthma and its impact on the quality and quantity of occupational life of individuals. The aim of this study is to determine how the impact of asthma on quality of life and ability to work of people employed in various sectors of society.

Methods: Cross-sectional and questionnaire study was carried out to determine the relationship between quality of life and job absenteeism in 124 patients with asthma referred to BESAT clinic (Kerman-Iran) in the summer of 2017. All patients were clinically confirmed by the expert and relevant tests. ASQL and WPAI-ASTHMA standard questionnaires were used to assess the quality of life and job absenteeism, which were used in similar studies, and their reliability and validity were confirmed. Finally, for analyzing the data, chi-square test was used in SPSS-16 software environment.

Results: Participants in the study included 124 patients, 83% of people aged 20-60 years. 95% of people had daily activity limitations. Also, in 50% of patients, the effect of the disease on the effectiveness of the workplace, moderate to severe, was evaluated meanwhile, there was a significant relationship between job absenteeism with different dimensions of quality of life such as age ($p < 0.015$), activity restriction ($p < 0.005$), public health ($p < 0.002$) and emotional states ($p < 0.005$).

Conclusion: Based on the results of this study, it was found that there is a significant relationship between quality of life indicators and job absenteeism in asthmatic patients.

Key words: Asthma, Job absenteeism, Quality of life, ASQL, WPAI-ASTHMA

ICIA2018\Allergy and immunotherapy\Poster\7397

7397

The anti-penicillin antibodies levels in sensitive and normal people to intradermal skin testMohammad Khosravi^{1*}, Fatemeh Boroun², Mohammad Nemat³*1. Department of Pathobiology, Faculty of Veterinary Medicine, Shahid Chamran University, Ahvaz, Iran**2. Student of Veterinary Medicine, Faculty of Veterinary Medicine, Shahid Chamran University of Ahvaz, Iran**3. Razi Vaccine & Serum Research Institute ,Agricultural Research, Education & Extension organization (AREEO)*

Background: The hypersensitivity reaction to penicillin is a public health problem. Immunological responses to penicillin and other beta-lactam antibiotics can be classified in immediate and non-immediate. The immediate hypersensitivity is mediated by IgE; however the non-immediate sensitivity facilitated by other isotypes of antibody or T lymphocytes. This research detected the non IgE antibody value against penicillin in allergic and normal people.

Methods: Thirty-eight samples from patients with positive or negative intradermal skin testing results of penicillin allergy were included in this study. The IgG and IgM levels against penicillin G were defined by in-house ELISA test.

Results: The results showed a significant ($P < 0.05$) elevation in total immunoglobulin and IgG of the sensitive groups; whoever the anti-penicillin IgM was significantly greater in non-sensitive people.

Conclusion: However the sensitized people to penicillin cannot be certainly detected with the total antibody, specific IgG and IgM value against penicillin; these values are a good indicator for prediction of immediate and late response of the immune system to penicillin.

Keywords: intradermal skin test, sensitive and normal people, anti-penicillin

ICIA2018\Allergy and immunotherapy\Poster\7455

7455

Fungal waterborne contamination as a threat for allergic infections in hot springs

Ali Ghajari*¹, Ensieh Lotfali¹

1. Department of Medical Parasitology and Mycology, School of Medicine, ShahidBeheshti University of Medical Sciences, Tehran, Iran

Background: Fungi cause allergies and many other adverse health effects. The prevalence of respiratory allergies by fungi such as *Aspergillus spp*, *Penicillium spp*, *Cladosporium spp*, *Alternaria spp*. has dramatically increased over the past decades. Hot springs are places for therapeutic purposes. Water of these pools could be carriers for transmission of various fungal diseases. The objective of this study was to survey of allergic fungal in these pools in Iran.

Methods: In this cross-sectional study, 150 samples were collected from 50 hot springs. The collected samples were filtered and cultured on SC (Sabouraud's dextrose agar with Chloramphenicol) and SCC (Sabouraud's dextrose agar with Chloramphenicol and Cyclohexamide). The plates were incubated at 25°C for 1-4 weeks. Then according to colony morphology and microscopic characteristic of fungi, the presence of allergic fungi were studied. The grown fungi were identified by macroscopic and microscopic methods.

Results: Out of 150 samples, 43 (28.6%) were positive. A total of 356 colonies observed, *Penicillium.spp* 161 colonies (45.2%), *Aspergillus.spp* 134 colonies (37.6%), *Cladosporium.spp* 44 colonies (12.3%), *Alternaria.spp* 17 colonies (4.7%). 134 colonies of *Aspergillus.spp* included *Aspergillusniger* 102 colonies (76%), *Aspergillus flavus* 17 colonies (12%), *Aspergillus fumigatus* 15 colonies (12%). The most common allergic fungi were *Penicillium.spp*, *Aspergillus.spp*, *Cladosporium.spp* and *Alternaria.spp* respectively.

Conclusion: The results of this study showed the presence of allergen fungal in hot springs. Sensitivity to *Aspergillus.spp* has been associated with severe persistent asthma. Allergic bronchopulmonary aspergillosis (ABPA) is caused by *Afumigatus*, *Alternaria.spp* and *Cladosporium.spp*, with the development, persistence, and severity of asthma. Therefore, experimental studies are essential to reduce the pollution of allergic fungal diseases in hot springs in our country.

Keywords: Fungal contamination, Allergic bronchopulmonary aspergillosis

ICIA2018\Allergy and immunotherapy\Poster\7496

7496

Seroepidemiology of Toxocariasis in Iranian asthmatic children, A study in Karaj district, IranHoma Sadri¹, Mohammad JavadGharavi²*1. Non-Communicable diseases Research Center, Alborz University of Medical Sciences, Karaj, Iran**2. Department of Parasitology, Alborz University of Medical sciences, Karaj, Iran Department of Parasitology, Iran University of Medical sciences, Tehran, Iran*

Background: The prevalence of asthma varies widely between countries and further studies are needed to evaluate risk factors for asthma in each region. A relation between asthma and toxocariasis has been reported. We aimed to investigate the possible association between Toxocara infection and childhood asthma in our region.

Methods: This was a case-control study involving 192 children aged 2 to 15 years, 92 of whom had asthma and 100 of whom did not have asthma. Blood samples were tested for the presence of Toxocara antibodies, using ELISA method.

Results: Only one patient (1.09%) had positive levels of anti-Toxocara antibodies in case group and no one in the control group. more than 90% of children in both case and control groups were residents of urban area.

Conclusion: Toxocara infection is not a common risk factor for childhood asthma in our urban area

Keywords: Asthma, Children, Toxocara

ICIA2018\Allergy and immunotherapy\Poster\7546

7546

IgG variables related to age, sex and season in patients with allergies referred to allergy clinic of Ali ibn Abitalib Hospital in Zahedan

FatemehRanjbar*¹, Mahla Jafari²

1,2 Bachelor's degree in laboratory science and member of Young and Elite Researchers Club, Department of Laboratory Sciences, Islamic Azad University, Zahedan Unit, Iran

Background: Allergic diseases are among the chronic diseases. When an extreme immune system is exposed to an allergen, the body has begun to produce a specific antibody called IgG to fight the substance and specifically secreted against antigens.

Methods: This descriptive- analytical research was done on a cross- sectional study on 88 individuals who were referred to the allergy clinic of Ali ibn Abi Talib Hospital in Zahedan. Patients reviewed included the ISAAC (International study of asthma and allergies in childhood) Standard and Global Questionnaire, Performing a Skin Prick Test and IgG was measured by ELISA method. Data was analyzed using SPSS version 21 and compiled by using Pearson correlation test.

Results: The number of patients was 88, of which 53 were females and 35 were males of these, 11.4% were less than one year, 14.8% were between one to five years, 11.4% between six to fifteen years and 62.5% of the adult population. According to the Pearson correlation test, the correlation between IgG variables with age, gender and season was performed. Results showed that this variable was not related to age and sex ($P < 0.05$) and had only a significant relationship with the season ($P > 0 / 05$).

Conclusion: There is a positive correlation between the IgE variable and the incidence of allergy. Therefore, IgE may predict a higher incidence of allergic diseases.

Keywords: Relationship, Variable, Allergy, Zahedan, Season

ICIA2018\Allergy and immunotherapy\Poster\7547

7547

Measuring Immunoglobulin Titer (IgE, IgM, IgG and IgA) in Patients with Allergy Referring to the Allergy Clinic of Imam Ali HospitalFatemeh Ranjbar², MahlaJafari^{*1}*1,2 Bachelor's degree in laboratory science and member of Young and Elite Researchers Club, Department of Laboratory Sciences, Islamic Azad University, Zahedan Unit, Iran*

Background: Allergy is a reaction to the immune system's excesses of various factors, when an extreme immune system is exposed to allergens, the body produces immunoglobulin to fight. Immunoglobulins are biological molecules that are active in the immune system and are specifically secreted against antigens.

Methods: In this study, changes in immunoglobulins in people with allergies were investigated. For this purpose, 88 patients with allergies referring to the Allergy Clinic of Imam Ali Hospital (Zahedan) in 1396, blood samples were taken and the immunoglobulin IgM, IgG, IgE and IgA were measured by ELISA method. Then, the data were analyzed by SPSS V21 software. Then, Pearson correlation test was used to determine the relationship between immunoglobulins and age, sex, and seasonal variables was investigated.

Results: 53 cases were male and 35 were female, 11.4% less than one year old, 14.8% between one and five years old, 11.4% between the ages of six and 15 years, and 62.5% of adults. The most visited was the summer season. Results showed that IgE, IgM, IgA variables had no significant relationship with age, sex and season ($P < 0.05$) and IgG had only significant relationship with seasonal variable ($P > 0.05$). The most changes were seen in IgE header, so that 34 People had abnormal IgE levels.

Conclusion: Considering the amount of IgE titer changes in the region and the severity of clinical signs in the inhabitants of this climate, further studies are needed to identify the antigenic action of allergens in different climates to reduce allergic symptoms.

Keywords: Allergy, Immunoglobulin, Imam Ali Hospital, Zahedan

ICIA2018\Allergy and immunotherapy\Poster\7553
7553

High Level of Herbal Food Additive Induces Inflammatory Responses in the Allergic Asthma Mice

Seyyede Masoume Athari¹, Bahram Yavari², Seyyed Shamsadin Athari^{3*}

1. Department of Biology, Faculty of Basic Sciences, Maragheh University, Maragheh, Iran
2. Department of Genetic and Molecular Medicine, Hamadan University of Medical Sciences, Hamadan, Iran
3. Department of Immunology, Faculty of Medicine, Zanjan University of Medical Sciences, Zanjan, Iran

Background: Herbal food additives are used routinely for rich of foods which may exacerbate asthma. The prevalence of allergic asthma has been increased over the past decades, so that prevention is an important strategy in the management of asthma. The aim of this study was to investigate the role of herbal food additive as a stimulus in asthma.

Methods: BALB/c mice were divided into four groups as followings; two OVA asthmatic and normal groups, fed with a diet containing herbal food additive and two healthy groups diet with normal food. Inflammation of airway and cytokines level in broncho-alveolar lavage fluid were measured.

Results: The group treated with both OVA and herbal food additive showed higher inflammation, interleukin-5 and 13 levels than the group sensitized only with OVA.

Conclusion: OVA and herbal food additive group showed more severe allergic asthma symptoms in comparison to the group only sensitized with OVA. Therefore, the herbal food additive can be asthmatic agents.

Key Words: Food, allergic Asthma, Immunology

ICIA2018\Allergy and immunotherapy\Poster\7554
7554

Pickles in Iranian food maybe allergic Asthma trigger in mouse model

Seyyed Shamsadin Athari¹, Seyyede Masoume Athari^{2*}

1. Department of Immunology, Faculty of Medicine, Zanjan University of Medical Sciences, Zanjan, Iran
2. Department of Biology, Faculty of Basic Sciences, Maragheh University, Maragheh, Iran

Background: Pickles is used in large amount of Iranian family that can be irritant compound for the respiratory system. The prevalence of allergic asthma is increased. The aim of this study was to investigate the role of Pickles as an asthmatic agents.

Methods: Fifteen mice were divided into three groups as followings; ovalbumin sensitized asthmatic groups, group that fed with a diet containing Pickles and normal group. Lung histopathology were examined.

Results: The group treated with Pickles showed higher lung eosinophilic inflammation and mucus hyper secretion similar to OVA (asthmatic) mice.

Conclusion: The Pickles consumption can be important trigger of allergic asthma and it should be used carefully in atopic people.

Key Words: Pickles, Allergy, Inflammation

ICIA2018\Allergy and immunotherapy\Poster\7713
7713

The Evaluation of Amount of TGF- β Produced in Supernatant of Co-culturing of DC Pulsed by Somatic Antigenic Extract of *Marshallagiamarshalli* and Peripheral T Lymphocyte in People Suffering from Asthma

Fatemeh Hemati¹, Hadi Moheb Alian², Hasan Borji³, Milad Hemati⁴, Majid Mirsadrai⁵

1. Department of Veterinary School of Ferdowsi University of Mashhad, Iran

2. Department of Veterinary School of Ferdowsi University of Mashhad, Iran

3. Department of Veterinary Faculty of Ferdowsi University of Mashhad, Iran

4. Atherosclerosis prevention research center, Medical Sciences University of Mashhad, Iran

5. Department of Medical science of Islamic Azad University of Mashhad, Iran

Background: Asthma is a chronic inflammatory disease with high prevalence and socio-economic impacts that causes respiratory dysfunction. According to the hygiene hypothesis due to decreasing of bacterial and parasite infection the prevalence of allergic asthma especially in developed countries is going up. Also dendritic cells have special potential to elect immune responses and polarizing naive T cells in to different subtypes of helper T cells. Nowadays DC based immunotherapy is a good alternative approach to treat many disease including cancer. In this study we utilized monocyte derived DC to skew naive T cells in to regulatory T cells by co culturing of DCs with somatic extract of *Marshallagiamarshalli* in order to treat allergic asthma.

Methods: In this study 10 people were chosen, 5 healthy people and 5 sick people that affected by asthma. Their blood sample were collected and their PBMCs were isolated and then monocytes separated from lymphocytes by adding of IL-4 and GM-CSF, monocytes differentiate to DC and 4th day, antigen extract of *Marshallagiamarshalli* added to culture medium. Then at 7th day of culture, these DCs cultured by autologous T cells and finally at 12th day of culture, the supernatants were collected and content of TGF- β were measured by sandwich ELISA.

Results: The average amount of TGF- β contents of supernatants in asthma group significantly increased (225 ± 6.1 pg/ml) in comparison with healthy groups (210 ± 8.5 pg/ml). ($P < 0.05$) **Conclusion:** With regards to hygiene hypothesis in which parasitic worms contain regulatory molecule of immune system, it could be possible producing of tolerogenic DCs by antigenic extract of *Marshallagiamarshalli*. *Marshallagiamarshalli* and this associated with polarizing of T cells toward regulated T cell in the blood of individuals affected by asthma; by evaluating amount of TGF- β content of supernatant of cell culture, it is clear that DCs could polarize immune responses.

Keywords: Dendritic cells, Allergic asthma, *Marshallagiamarshalli*, TGF- β

ICIA2018\Allergy and immunotherapy\Poster\7831

7831

The Specific IgE Assay to Common Respiratory Allergens in Subjects with Allergic Symptoms in Ahvaz

Maryam Moradi¹, Abbas Fayezi², Mana Momeni¹, Asyeh Javanian¹, Suzan Amini¹, Sara Yaghotian¹,
Mohammad Shahrooei^{1,3}

1. *Specialized Immunology Laboratory of Dr. Shahrooei, Ahvaz, Iran*

2. *Department of Allergy and Clinical Immunology, School of Medicine, Ahvaz University of Medical Sciences, Ahvaz, Iran*

3. *Department of Microbiology and Immunology, Experimental Laboratory Immunology, KU Leuven, Leuven, Belgium*

Background: The results of the specific IgE assay to respiratory allergens differs extensively between different geographical regions. This study was designed to assess the specific IgE to common respiratory allergens using ImmunoCAP system among allergic cases in Ahvaz province.

Methods: Four hundred and eight patients with allergic symptoms entered this cross-sectional study in Ahvaz from 2014-2017. After fulfilling a specific questionnaire and medical assessment, 5 ml sample was taken from the cases. The total IgE and specific IgE for 9 respiratory allergens was measured using the ImmunoCAP system (ThermoFisher-Phadia, Uppsala, Sweden) according to manufacturer's instructions.

Results: A total of 57% of the patients were male. The mean age of the subjects was 23 y (3 m to 68y). The majority of subjects were older than 18 years (n=195, 48%). As the results depict, the most common outdoor aeroallergens were Russian thistle (52%), Willow (47%), and Mesquite (41%), respectively, while among indoor allergens, German cockroach (38%) and *Dermatophagoides farinae* (11%) were the most prevalent. Sensitization to one allergen and more was found in 66.4% of the patients. Mono-sensitization and poly-sensitization were considered in 11.76% and 54.6% of patients, respectively.

Conclusion: The prevalence of sensitization to respiratory allergens has increased during the past decade in this city. The pattern of allergen-specific IgE showed Russian thistle as the most common aeroallergens in patients. Regarding the air pollution in some cities such as Ahvaz, the relationship between air pollution and longer pollination season and as their results increasing the allergic diseases, attention to the determination of the common respiratory allergens especially pollens seems necessary.

Keywords: Specific IgE, Respiratory allergens, Ahvaz.

ICIA2018\Allergy and immunotherapy\Poster\7837

7837

Pattern of Skin Sensitivity to Aeroallergens among Allergic Rhinitis Patients in Birjand City, Iran

Mohammad Fereidouni¹, Neda Mahavar¹, Mohammad Sadegh Naghizadeh¹, Mohammad Mofateh²

1. *Asthma, Allergy & Immunology Research Center, School of Medicine, Birjand University of Medical Sciences, Birjand, Iran (Times New Roman, font size 11 italic)*

2. *Department of Otolaryngology, School of Medicine, Birjand University of Medical Sciences, Birjand, Iran*

Background: Allergic disorders are among the most common health problems around the world and their prevalence has been increased during the last decades. Allergic rhinitis and conjunctivitis are the most prevalent forms of allergy and have negative impacts of quality of life as well as productivity of patients. Pollens and indoors allergens are the main triggers of allergic symptoms but the pattern of sensitization is varied in different parts of the world. The aim of this study was to identify the most common aeroallergens in allergic rhinitis patients at Birjand city of Iran.

Methods: patients who were referred to Birjand Allergy clinic during 2013 to 2017 because of any allergies, clinically evaluated by specialist and skin prick test with a battery of at least 20 common outdoor and indoor allergenic extracts performed based on standard method.

Results: 875 patients (mean age: 26 ± 11.9 years, range 2-64 years, M/F ratio: 0.89) who suffered from allergic rhinitis or allergic rhinoconjunctivitis enrolled in this study. Highest rate of skin sensitivity was for weeds/ grasses pollen including Salsola Kali, Amaranthus Retroflexus, Chenopodium Album and Composite family (74.3%, 62.5%, 50.9% and 39.3% respectively). Among tree's pollen; Ash (49%), Walnut (46.9%) and Mesquite (29.3%) were the most common. Less than 20% of patients showed skin reactivity to indoor allergens and Storage mites, mix of Cockroaches and house dust were the most common (17.6%, 17.3% and 9.8% respectively).

Conclusion: The results of current study confirmed the importance of weed/grass and trees pollen as the major source of allergic sensitization in our area. Interestingly the rate of sensitization to indoor allergens was low which can be explained by geo-climatic situation.

Keywords: Allergy, Rhinitis, Conjunctivitis, Skin prick test

ICIA2018\Allergy and immunotherapy\Poster\7865

7865

Evaluation of the Relationship between Exposure to Hepatitis A Virus and Allergic Disease in Birjand City in 2014

Zahra Asghari¹, Mohammad Fereidouni²

1. *medical doctor, Faculty of Medicine, Birjand University of medical Sciences*

2. *Associated Professor, Faculty of Medicine, Birjand University of medical Sciences*

Background: Childhood infections affect the immune system resulting in decrease in allergic diseases. This study aimed to evaluate the relationship between a History of Hepatitis A and Allergic Disease in Patients in Birjand Allergic Clinic of Imam Reza (AS) in 2014

Methods: This is a case control study on 352 patients had been referred to allergy and asthma clinic of Imam Reza (AS) in 2013. Sampling method was non-random in both groups (allergic 113 people and non-allergic 239 people). Both group were matched based on gender, education, age, job, income and living place. Allergic diagnosis was based on skin prick test and serum IgE beside the clinical examination. Then a questioner was completed and blood sample obtained for HAV-ab (total) anti evaluation. Data were analyzed with the software SPSS21 ($p \leq 0/05$)

Results: Overall 352 patients were enrolled in this study. Prevalence of hepatitis A and allergy was 66.5% and 32.1% respectively. Prevalence of hepatitis A were higher in allergic patients in compare to none allergic ($n=90$ 79.6% Vs $n=118$ 60.3% $p=0.0001$).

Conclusion: The results of this study, unlike most studies in this area show a higher incidence of allergic patients had a history of hepatitis. Given the high prevalence of hepatitis-A in developing countries due to the low level of health on one hand and their transition period (increasing urbanization, changing life styles and increasing the level of dealing with other environmental allergens) that independently causes atopy, it seems that a greater role these factors play in the development of atopy.

Key Words: Allergy, HAV, Hygiene hypothesis

ICIA2018\Allergy and immunotherapy\Poster\8035
8035

Increased Serum Levels of Resolvin E1 but No Difference in Serum Levels of Resolvin D1 in Allergic Rhinitis Patients Compared to Healthy Subjects

Ramin Lotfi¹, SeyedHamidreza Mortazavi², Ali Gorgin Karaji³, Farhad Salari^{3*}

1. Student research committee, School of Medicine, Kermanshah University of medical sciences, Kermanshah, Iran

2. Department of Pediatrics, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

3. Department of Immunology, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

Background: Allergic rhinitis (AR) is a common inflammatory disorder of nasal mucosa, which is characterized by pruritus, sneezing, rhinorrhoea, and nasal congestion. *Resolvins* (Rvs) are anti-inflammatory and pro-resolving lipid mediators derived from *omega-3*polyunsaturated fatty acids (PUFAs). Resolvin E1 (RvE1) and resolvin D1 (RvD1) are produced from eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA), respectively. The aim of this study was to evaluate the serum levels of RvE1 and RvD1 in AR patients and healthy subjects.

Methods: Serum samples were obtained from 37 patients with AR and 30 age- and sex-matched healthy subjects. RvE1 and RvD1 concentrations were measured by ELISA technique according to the manufacturer's instructions.

Results: RvE1 concentrations were significantly increased in patients compared to healthy subjects, but no significant difference in RvD1 concentrations between patients and healthy subjects was observed.

Conclusion: This study showed that concentrations of RvE1 were significantly increased in patients than healthy subjects, suggesting that this increment is an effort to help to timely resolution of nasal mucosa inflammation and prevention of acute inflammation progression to chronic inflammation. Our findings suggest that RvE1 or its analogues may help rational development of therapeutic approach for AR.

Keywords: Allergic rhinitis, Inflammation, Pro-resolving, Resolvin E1

ICIA2018\Allergy and immunotherapy\Poster\8061

8061

The Purification Process of *Toxocaracanis* Antigen Fragments in Order to Immunomodulate Airways Inflammation and AllergiesM.Badri¹, FGhaffarifar¹, ZS.Hasan², A.Dalimi¹*1. Department of parasitology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran**2. Department of Immunology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran*

Background: Parasitic infections are very widespread in human populations. According to WHO, about 2 billion people suffer from parasitic infections. For example, people with asthma (approximately 819 million) are more affected than asthma (approximately 300 million). The common feature of parasitic infections and allergies is often the activation of Th2 and the increase of IL4, IL5 and IL13 cytokines, which increases the eosinophils and mast cells and increases the production of IgE antibodies. In populations where parasitic infections have already been endemic and have been significantly reduced due to increased community health levels, this trend of balance adjustment of immunological responses has been impaired and consequently an increase in the incidence of diseases such as asthma, in the community. For example, the prevalence of *Ascaris* in 1976 in endemic regions of Iran was 79.71%, and a significant decline to 0.5%, while a significant increase in MS cases per 100,000 people since 2006 (26.6%) To 2011 (44.53).

Methods: The *Toxocaracanis* nematode was collected from 5 stray dogs and sonicated for isolating of antigens. The soluble antigen at first the lipid was removed by ether then we used Sephadex G200 column for purification of antigens. For evaluating the characterization of purified antigen we used SDS-PAGE.

Results: After that crossing the chromatography column the three-peak antigens were isolated and after the SDS-Page, the molecular weight (10, 27 and 50 kD) was used as a candidate. The Bradford Spectrometry method was used to determine the protein content of each fraction.

Conclusion: In these evaluation the antigenic fractions will be used for exposing to human lung cells under in vitro conditions as well as injection into laboratory mice after exposure to Ovalbumin.

Keywords: *Toxocaracanis*, Antigen fragments, Immunomodulation, Airways inflammation

ICIA2018\Allergy and immunotherapy\Poster\8179
8179

Budesonide and Fexofenadine Treatment Reduce Blood Eosinophils and Serum IgE in Allergic Rhinitis

Elham Bagherinia¹, Seyed Hamidreza Mortazavi², Farhad Salari³, Ali Gorgin Karaji^{3*}

1. Student research committee, School of Medicine, Kermanshah University of medical sciences, Kermanshah, Iran

2. Department of Pediatrics, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

3. Department of Immunology, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

Background: Allergic rhinitis (AR) is an IgE-mediated inflammation of the upper airway. The aim of this study was to evaluate the effect of treatment with Fexofenadine and Budesonide on the level of serum IgE and percent of peripheral blood eosinophil.

Methods: As part of a clinical trial, 27 AR patients (17 men and 10 women, age 18-60) were treated for one month with Fexofenadine and Budesonide. Before treatment and one month after treatment, blood was taken from AR patients. Serum IgE level was measured using IgE ELISA kit and the percentage of peripheral blood eosinophil was counted using peripheral blood smear.

Results: The results of this study showed that after one month treatment, both the percentage of peripheral blood eosinophils and serum IgE level were significantly decreased in AR patients.

Conclusion: Taken together our results showed that treatment with Fexofenadine and Budesonide can reduce serum IgE levels and blood eosinophil count.

Keywords: Allergic rhinitis, Blood eosinophil percent, Fexofenadine, Budesonide

ICIA2018\Allergy and immunotherapy\Poster\8180

8180

Treatment of Allergic Rhinitis with Fexofenadine and Budesonide reduces Clinical Symptoms and Improves the Quality of LifeElham Bagherinia¹, SeyedHamidreza Mortazavi², FarhadSalari³, Ali Gorgin Karaji^{3*}*1. Student research committee, School of Medicine, Kermanshah University of medical sciences, Kermanshah, Iran**2. Department of Pediatrics, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran**3. Department of Immunology, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran*

Background: Allergic rhinitis (AR), an inflammation of upper airway, is the most common allergic disease. AR signs and symptoms including: a runny nose, sneezing, redness, itchy, and watery eyes, and nasal congestion and obstruction, despite having a safe nature, can seriously affect patient's quality of life. Treatment of AR is a way to reduce RA symptoms and improving the quality of life. The aim of this study was to evaluate the effect of treatment with Fexofenadine and Budesonide on the quality of life and on the clinical symptoms of allergic rhinitis patients.

Methods: In this clinical trial, 30 AR patients (18 men and 12 women, age 18-60) were treated for one month with Fexofenadine and Budesonide. For each patient, before treatment and after one month treatment, two questionnaires were used, one for patient symptoms and the other for patient quality of life.

Results: Our results showed that, in AR patients, one month treatment with Fexofenadine and Budesonide significantly reduce the severity of patient's symptoms and improve the quality of patient's life.

Conclusion: Generally, our results showed that dual treatment with Fexofenadine and Budesonide could be an effective way to treat allergic rhinitis.

Keywords: Allergic rhinitis, Fexofenadine, Budesonide, Quality of life.

ICIA2018\Allergy and immunotherapy\Poster\8229 allergy
8229

The Effectiveness of Using AllergyCut® in the Allergic Rhinitis Treatment

Farid Semsarha^{1*}, Saleh Ramazani Khorshiddoost², Zahra Moeinibakhsh², Nika Khoshnevis², Maral Nazarifar², Niloofar Basrehizad³, Reza Ramazani Khorshiddoost⁴

1. MIM Daroo Pharmaceutical Co. Tehran, Iran.
2. Aramesh multi professional pain clinic, Tehran, Iran.
3. Islamic Azad university of Science and Research Branch, Tehran, Iran
4. Faculty of Systems Management, Amirkabir University of Technology, Tehran, Iran

Background: Allergic rhinitis is one of the most common chronic diseases affecting adults. AllergyCut® is a registered herbal drug with nasal spray dosage form that its efficacy has been approved during the present study.

Methods: In this study, that has been done at Aramesh clinic under supervision of shahidbeheshti university of medical sciences and health services, patients with allergic rhinitis were selected and some who were in need of further tests for diagnosis of immunological sensitivity were referred to the physician for confirming their allergy. After completing the treatment sessions, the physician re-examined the patients and their data. Then a follow-up program has been set up and the patients have used the AllergyCut® spray between mostly 60 days to 160 days (for about less than 10% of patients). The patients provided the information about their allergic attacks in a daily form, which has been designed for this study.

Results: From 110 patients who have been treated with the AllergyCut® spray, only the data of 70 patients could be cited and studied because of the accuracy in filling out the forms and the correct usage of the spray. The comparison of the data about allergy symptoms has been done according to indicators, guidelines and forms of the world allergy organizations (WAO). In first month after treatment, the condition and clinical symptoms of patients have improved significantly to 33% in average value of allergy indices, and after second month these decreased by 70% compared to the first month. The follow of patients has been done for about 2 years and this improvement maintain in this period.

Conclusion: The herbal nasal spray AllergyCut® have been used in a certain time by 70 patients who have suffered from allergic rhinitis and about 74% of the patients reported the removal of allergic rhinitis symptoms during usage and also after two years follow.

Keywords: Allergic rhinitis, AllergyCut®, Herbal medicine

ICIA2018\Allergy and immunotherapy\Poster\8277

8277

Does Air Pollution Change Common Children's Inhaled Allergens in Isfahan?Fatemeh Rezayat¹, Sadegh Feyzollahzadeh², Abbas Rezaei¹*1. Department of Immunology, School of Medicine, Isfahan University of medical Science, Isfahan, Iran**2. Department of Immunology, School of Medicine, Urmia University of medical Science, Urmia, Iran*

Background: The prevalence of respiratory allergy in children depends on their habitat. Beyond genetic factors, many environmental factors specially air pollution in large cities have been implicated in increased risk of allergy. It has believed that common inhaled allergens are different in each area. Moreover, air pollution may change the prevalence of allergens or generation of new allergens in the population. In the present study we compared prevalence of allergens in children younger than 5 years old during last year with in Isfahan province.

Methods: 75 children younger than 5 years old who were suffering from skin and respiratory symptoms, referred to Dr. Rezaei Immunology Laboratory were chosen. The total serum IgE levels were measured using ELISA. For determination of allergens, CAP RAST (radio allegro sorbent test) and membrane strip containing 50 different allergens test (EUROLINE) were performed for each patient.

Results: The Mean of total serum IgE concentration was more than normal values. Considering the results, Rough pigweed, Rose and cultivated oats were detected as the most common inhaled allergens for children in Isfahan province. IgE concentration was significantly ($P<0.05$) higher in hypersensitive children to Rough pigweed compared with Rose and cultivated oats. It was however the most number of children had allergic reaction to Rose. All of these results have been compared with existed data in earlier studies on common allergens in Isfahan.

Conclusion: In this study, three allergens including Rough pigweed, Rose and cultivated oat was detected as the most common inhaled allergens in Isfahan. Since last decade Rough pigweed was known as the most common inhaled allergen in children. Until today air pollution in this province did not have any effects on common inhaled allergens in children and its effects may involve in adult respiratory allergy.

Keywords: Respiratory allergy, Inhaled allergen, Isfahan, Children.

ICIA2018\Allergy and immunotherapy\Poster\8445

8445

Effect of treatment with Fluticasone propionate and Fexofenadine on blood Eosinophil percentage and Serum IgE level in Allergic Rhinitis

Atefe Mehrmanesh^{1,3}, Elham Bagherinia^{1,3}, Seyed Hamidreza Mortazavi², Farhad Salari³, Ali Gorgin Karaji³

1. Student research committee, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

2. Department of Pediatrics, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

3. Department of Immunology, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

Background: Allergic rhinitis (AR) is an IgE-mediated inflammation of the Upper respiratory tract and is the most common allergic disease. The goal of this study was to investigate the effects of treatment with Fexofenadine and Fluticasone propionate on the level of serum IgE and the percentage of blood eosinophil in AR patients.

Methods: In this study, 26 AR patients (12 men and 14 women, aged 18-60) were treated for one month with Fexofenadine and Fluticasone propionate. Before treatment and one month after treatment, blood was taken from AR patients. Serum IgE level was measured using IgE ELISA kit and the percentage of peripheral blood eosinophil was determined using peripheral blood smear.

Results: The results showed that after one month treatment, the percentage of blood eosinophils was significantly decreased. On the other hand, after one month of treatment, AR patient's serum IgE level was significantly increased.

Conclusion: Overall, our results showed that one-month treatment of AR patients with fexofenadine and fluticasone propionate reduces the percentage of blood eosinophil, but increases serum IgE level.

Keywords: Allergic rhinitis, Serum IgE titer, Blood eosinophil percentage, Fexofenadine, Fluticasone propionate

ICIA2018\Allergy and immunotherapy\Poster\8449

8449

Correlation of Serum *Resolvins* and the Quality of life in Allergic Rhinitis

Razieh Rezaeianpour¹, Ramin Lotfi¹, Firoozeh Khamoushi², Seyed Hamidreza Mortazavi³, Ali Gorgin Karaji⁴, Farhad Salari^{4*}

1. Student research committee, School of Medicine, Kermanshah University of medical sciences, Kermanshah, Iran

2. Student research committee, School of Nutritional Science & Technology, Kermanshah University of Medical Sciences, Kermanshah, Iran

3. Department of Pediatrics, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

4. Department of Immunology, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

Background: Allergic rhinitis (AR) is an airway inflammatory disorder with high and increasing prevalence. Although it does not have a dangerous nature, it can seriously affect the patient's quality of life (QOL). Resolvins such as resolvin E1 and D1 (RvE1 and RvD1) are omega-3 fatty acids-derived lipid mediators that have anti-inflammatory and pro-resolving effects. The major aim of this study was to investigate the relationship between quality of life and serum levels of RvE1 and RvD1 in AR patients.

Methods: This cross-sectional study included 36 AR patients with age 15–57 years. The serum concentrations of RvE1 and RvD1 were measured by ELISA technique according to the manufacturer's instructions. The Rhinitis Quality of Life Questionnaire (RQLQ) was used to assess the symptoms severity of patients. Data were analyzed by SPSS V21.

Results: The mean age of 36 patients was 34.7 ± 10.4 . 69.4% of patients were women. The mean of RvD1 and RvE1 serum concentrations in patients were 5.6 ng/ml and 242.4 pg/ml, respectively. The significant positive correlation was found between the practical problems and RvE1 serum concentrations (correlation coefficient: 0.360; $P = 0.031$; $P < 0.05$), but there was no association between the serum concentrations of RvD1 and any subscale of RQLQ.

Conclusion: Our results showed that serum levels of RvE1 might be effective in improving the quality of life of AR patients.

Keywords: Allergic rhinitis, Quality of life, Omega-3 fatty acid, Resolvin

ICIA2018\Allergy and immunotherapy\Poster\8495

8495

Determination of the Most Common Respiratory and Food Specific IgE Patients with Allergic Symptoms Using Immunoblotting method

Sepideh Moradkhani^{1,2}, Majid Khoshmirsafa^{2,3}, Maryam Mahjoub², Alireza Rezaei Manesh^{2,4}, Seyed Mehdi Ebrahimi^{2,5}, Farhad Seif³, Parisa Nourbakhsh⁶, Mehdi Shekarabi^{2,3*}

1. Department of Immunology, Faculty of Medicine, Kerman University of Medical Science, Kerman, Iran

2. Massoud Clinical Laboratory, Tehran, Iran

3. Department of Immunology, School of Medicine, Iran University of Medical Sciences, Tehran, Iran

4. Department of Immunology, Faculty of Medicine, Kermanshah University of Medical Science, Kermanshah, Iran

5. Department of Clinical Biochemistry, Faculty of Medicine, TarbiatModares University of Medical Science, Tehran, Iran

6. Departments of Immunology, Faculty of Medicine, Arak University of Medical Science, Arak, Iran

Background: The prevalence of allergic diseases, as one of the global health problems, is rising particularly in developing countries in recent years. Different allergens can cause allergic symptoms. The aim of this study was to determine the frequency of the respiratory and food IgE in patients with suspicion of allergy.

Methods: In this cross-sectional study, 162 (52% female, 48% male) patients with suspicion of allergy were investigated in Massoud Clinical Laboratory, Tehran, Iran, between May 2016 and January 2018. Specific IgE for 30 selected respiratory and food allergens was measured by immunoblotting method. Data were analyzed by SPSS-23 software. A *p* value less than 0.05 were considered as significant.

Results: The median (IQR) age of recurrence were 4 (1.5-9). Mixed grasses (GX) was the most frequent specific IgE among 30 selected respiratory allergens (42.8%) and Pistachio nut (F203) was the most frequent specific IgE among 30 selected food allergens (18.3%). In addition, the frequency (positive reaction) of respiratory and food allergens was higher in children in comparison to adults (*P*_v<0.05).

Conclusion: The results of this study indicate that evaluation of specific IgE by Immunoblotting method can be useful for precise diagnosis and follow up of desensitization therapy.

Keywords: Allergy, Food Allergens, Respiratory allergens, Specific IgE

ICIA2018\Allergy and immunotherapy\Poster\8514 allergy
8514

Association between Education and Management of Asthma Exacerbation

Babak Khodadadi^{*1}, Arash Kalantari ², MortezaSagharjoghi Farahani³

1. Young researchers and Elite Club, Khorramabad Branch, Islamic Azad University, Khorramabad, Iran

2. Assistant professor of Tehran University of medical sciences. Imam Khomeini hospital complex vali-e-asr hospital Department of Pediatrics, Tehran, Iran.

3. Young researchers and Elite Club, Islamshahr Branch, Islamic Azad University, Islamshahr, Iran

Background: Asthma is the most common chronic disorders in the world that affect children. The prevalence of childhood asthma is increasing over the last two decades, many children are able to achieve good levels of control. So this study focus on the relationship between education and asthma exacerbation control.

Methods: In this quasi-experimental study, 104 children (6-14 years old) with asthma referred to asthma and allergy clinic of Children's Medical Center in Tehran at 2011-2015, which selected by convenience sampling method and divided into two equal groups of intervention and control. Data were analyzed by SPSS version 18.0 and T-test.

Results: The results of the study indicated that reducing asthma symptoms, improving PFT(Pulmonary function test), *decreasing the frequency of exacerbations* and salbutamol spray using in the intervention group; while changes in the control group were not significantly different ($p>0.05$).

Conclusion: *The results suggest that good education have an excellent effect on asthma exacerbation control in children.*

Keywords: Asthma, Education, exacerbation,

ICIA2018\Allergy and immunotherapy\Poster\8520
8520

An Attempt to Harness the Deathful Peanut Allergy by Naturally Existing CAM Named Flavonol Quercetin

Lotfollah Behroo¹

1. Vice-chancellor for Student, Farhangian University, Nutrition, Tehran, Iran

Background: Epidemiologically, ever-growing Allergies-rates are actually/drastically baleful. Accordingly, “Peanut Allergy” is especially, of critical significance considering the pertinent fatal or life-threatening anaphylactic events. “Avoidance”- focused upon as a mainstay of remedy-, seems to be indeed a difficult labor to perform principally if the suspected food is found everywhere. As a consequence, the majority of people with food allergy seek for natural/plant-origin Complementary and Alternative Medicines (CAMs), said to be more effectual and concurrently, quite safe. A “flavonol” subgroup-member of flavonoids named Quercetin is of prime interest to investigators. To inspect the benefits of a naturally-occurring nutritional substance- Quercetin- on peanut-induced sequelae in a murine model.

Methods: Male Wistar-rats were sensitized with peanut-extract in the presence of cholera-toxin and {Al (OH) 3}. The sensitized-animals were subsequently divided into 3-group; “Positive-control”, “Quercetin-treated” and “Sham-sensitized”. One-week post-sensitization, the wistars in treatment-group were gavaged daily, with Quercetin dissolved in 5% dimethyl-sulfoxide (DMSO), at a dose of 50 mg/kg body-weight (BW)/mL/rat, for 4-week. Duly, all the study animals were challenged intragastrically, with peanut-allergens. Naive rats (n=7) served as negative-controls. Then, Plasma Histamine and Total serum IgE levels were measured.

Results: Following daily-repeated orally-administering for four weeks, Quercetin significantly lowered Plasma Histamine and Total serum IgE Levels in the Quercetin-treated wistars {(p<0.000) and (p <0.000)}, respectively.

Conclusion: It is revealed that the flavonoid “Quercetin” has suppressed thoroughly the respected IgE responses as to peanut-allergens and is efficient enough to abrogate the peanut allergy’s complications. Hence, it can be assumed/endorsed as a substitute compound against IgE mediated food allergies in human beings.

Keywords: CAMs, Flavonoid, Flavonol, Murine Model, Peanut Allergy Treatment, Quercetin

ICIA2018\Allergy and immunotherapy\Poster\8522

8522

Frequency of Common Aeroallergen in Allergic Rhinitis Patients of Gorgan in 2016

Morteza Sagharjoghi Farahani¹, Babak Khodadadi^{*2}, Arash Kalantari³, Seyed Mohammad Hadi Bagheri Rostami^{4,5}, Hosna salmani

1. Young Researchers and Elite Club, Islamshahr Branch, Islamic Azad University, Islamshahr, Iran

2. Young Researchers and Elite Club, Khorramabad Branch, Islamic Azad University, Khorramabad, Iran

3. Assistant professor of Tehran University of medical sciences. Imam Khomeini hospital complex vali-e-asr hospital Department of Pediatrics, Tehran, Iran

4. General physician, Gorgan University of medical science, Gorgan, Iran

5. Department of School of health managment and information sciences, Iran University of medical sciences, Tehran, Iran

Background: In recent years, allergic diseases have increased in the region and identifying the factors that affect these diseases is very important. The aim of this study is to determine the prevalence of common inhalers allergens based on pericardial tests in allergic rhinitis patients in Gorgan.

Methods: In this descriptive study, Prick test was performed on 270 patients who had allergic rhinitis to asthma and allergy center of Gorgan during 2016. After completing the demographic questionnaire including age, gender history of asthma, allergic rhinitis, eczema, and conjunctiva, information was recorded. Finally, the data was analyzed by SPSS16. Chi-Square test is used to evaluate the difference between the results of the test for each of the variables ($P < 0/05$)

Results: According to this study, 54 patients were seasonally and 47 persons had mixed allergic rhinitis. And disease associated with allergic rhinitis were conjugate (24%), asthma (12%) and eczema (3%) respectively. 40% of patients had allergies to Mite allergies and 40% of patients had allergic to weeds, 30% of patients were allergic to grass and allergic rhinitis, and the lowest sensitivity was related to dogs and cat's allergens. In Patients with Permanent Allergic Rhinitis (PAR), 50% of patients were sensitive to two types of mites, dermatophagouspterocnemia (DP) and dermatophagousfina (DF), and in Seasonal versus Perennial (SAR) patients, 60% of the patients were allergic to weeds and grasses.

Conclusion: Based on the findings, more than half of the patients with permanent allergic rhinitis had the highest sensitivity to the mites and then, showed the highest sensitivity to weeds and cockroaches, respectively. In patients with seasonal allergic rhinitis, the most susceptibility was allergenic to weeds and then grasses.

Keywords: Allergic rhinitis, Allergen inhalation, Prick test

ICIA2018\Allergy and immunotherapy\Poster\8529
8529

Investigating the Effect of Using Latex Gloves on Causing Skin Allergies in the ICU Ward of Kowsar Medical Center, Sanandaj

Jalil Adabi¹, Elaheh Amani¹, Sho'aub Dehghani¹, Pejman Allahvaisy, Golnaz Babamiri

1. Kurdistan University of Medical Sciences, Sanandaj, Iran

Background: Latex is recognized as a flexible substance which has been used in producing plastic products. It has been estimated that wearing gloves made of latex can cause allergies among 1% of the individuals and 3- 17 % of personnel working in medical centers. It has not been calculated that to what extent using products made of latex such as gloves can cause allergies. A stockpile of research argue that individuals might be different in terms of indicating allergies by using latex products. This study aims to investigate the effect of using latex gloves on indicating allergies among personnel of ICU ward in Kowsar Hospital of Sanandaj.

Methods: Since personnel working in ICU wards are required to use latex gloves, the subjects of this study were chosen among the personnel of ICU in Kowsar hospital of Sanandaj. Forty of them completed the checklist given to them. The collected data was analyzed through SPSS 18.

Results: The analyzed data obtained from the checklists revealed that 20% of the subjects suffer from skin redness and itching when they use latex gloves. 80% of them reported that they prefer using prep gloves.

Conclusion: Concerning the fact that latex gloves are being used increasingly specially in ICU wards, it seems significant to consider potentials of causing allergies.

Key words: Allergy, ICU Personnel, Latex Gloves

ICIA2018\Allergy and immunotherapy\Poster\8540

8540

Pathology of Patients with Allergies Referring to Ali ibn Abi Taleb Zahedan Hospital In 1396

Fateme ranjbar^{1*}, Mahla jafari², Hanie jafardokht bonjar³, Hamide danesh⁴, Fateme akbari mehni⁵

1,2,4,5. Bachelor's degree in laboratory science and member of Young and Elite Researchers Club, Department of Laboratory Sciences, Islamic Azad University, Zahedan Unit, Iran

3. Bachelor of Science in Nursing and Member of Young Researchers and Researchers Club, Islamic Azad University, Unit of Sciences, Iran

Background: Allergy is an immediate deformed reaction to the re-entry of allergenic substances that have previously been triggered by the person's immune system. Allergic diseases, the prevalence of which has increased significantly in recent years, particularly in industrialized areas, are among chronic diseases. The aim of this study was to determine the pathology of allergy.

Methods: The present study was descriptive-analytic. The statistical population consisted of allergic patients referred to allergy clinic of Ali ibn Abi Talib Hospital in Zahedan. This study included 404 patients referring to Ali ibn Abi Taleb Zahedan Hospital in 1396, based on laboratory criteria. Patients included 214 females and 190 males. Data extraction was done from the statistical sources of the hospital and then the data were analyzed by SPSS v21 software.

Results: Of the patients studied, 53% of the women and 47% of the men suffered from the disease. These patients were examined for sex, marital status, occupation, educational level, place of residence, and season of bone disease. Of these, 246 were single and married. In terms of occupational parameter, including cultural, military, administrative, etc. Cultural people earned a higher percentage. Those who were under the diploma were those who lived in Zahedan and also had more of this complication in the winter.

Conclusion: There is a significant difference between the rates of the disease and the parameters mentioned, and it is better to prevent the spread of the disease by raising awareness of the people and taking appropriate measures. How to cope with predisposing factors in the list of health education programs and it is suggested to identify the predisposing populations of the futuristic research.

Key words: Pathology, Allergy, Immune system

ICIA2018\Allergy and immunotherapy\Poster\8574

8574

Laboratory Findings in Patients with Urticaria

Maryam Moossavi¹, Mohammad Fereidouni^{2,*}, Afshin Derakhshani¹

1. Student Research Committee, Birjand University of Medical Sciences, Birjand, Iran

2. Asthma, Allergy & Immunology Research Center, School of Medicine, Birjand University of Medical Sciences, Birjand, Iran (Times New Roman, font size 11 italic)

Background: Urticaria known as hives is a common disorder with various etiologies. Studies manifested that urticaria is associated with hepatitis B and C, gastritis, thyroid malfunction and autoimmune diseases like rheumatoid arthritis. Therefore, the only screening tests that can be measured are immunologic factors. The aim of the present study was to evaluate and measure C-reactive protein (CRP), HbsAg, liver function tests, TG, thyroperoxidase (TPO), H pylori IgG, antinuclear antibody (ANA) titers, rheumatoid factor (RF), IgM and IgA in a sample of patients with urticaria in Birjand, South Khorasan province.

Methods: 337 Individuals who were referred to Birjand Allergy clinic during 2013 to 2017 because of urticaria were recruited to this study. C-reactive protein (CRP), HbsAg, liver function tests, thyroperoxidase (TPO), H pylori IgG, antinuclear antibody (ANA) titers, rheumatoid factor (RF), IgM and IgA were measured for all individuals.

Results: In our study, 69% of the individuals were women and 31% were men. Most of the urticaria individuals showed normal range of H.pylori (76.3%), IgM (92.1%), IgA (86%), TPO (90.5%), CRP (89.2%), LFT (92.4%) and TG (93.3%) followed by negative HbsAg (98%), RF (96.8%) and ANA (94.9%) test. However, some of them showed high and severe value that was seen in H.pylori (1.3%), IgM (0.4%), IgA (0.8%), TPO (0.4%), CRP (0.4%) and LFT (0.4%) tests and positive ANA (0.4%) as well.

Conclusion: In our study, it seems that high and severe value of H.pylori was the most important factor in urticaria patients.

Keywords: Urticaria, Laboratory findings, Screening

ICIA2018\Allergy and immunotherapy\Poster\8636
8636

Distribution of Different Allergic Disorders among Patients Referred to Birjand's Allergy Clinic, a Five Years Study

Afshin Derakhshani^{1,2}, Mohammad Sadegh Naghizadeh^{1,2}, Maryam Moossavi^{1,2} Mohammad Fereidouni³

1. Student Research Committee, Birjand University of Medical Sciences, Birjand, Iran

2. Asthma, Allergy & Immunology Research Center, School of Medicine, Birjand University of Medical Sciences, Birjand, Iran

3. Cellular and Molecular Research Center, Birjand University of Medical Sciences, Birjand, Iran

Background: The prevalence of allergic disorders has been increased during the recent decays. The World Allergy Organization (WAO) reports that approximately one of five people suffer from some form of allergic diseases such as allergic rhinitis (AR), asthma, conjunctivitis, eczema, food allergies and so on. High frequency of allergic disorders causes huge economic burden and negatively impact patients' quality of life and productivity. However, distribution of allergens and risk factors may be varies in different geographic areas. The aim of this study was to evaluate the prevalence of different allergic disorders among patients referred to allergy clinic during a 5 years period in Birjand, Iran.

Methods: patients who were referred to Birjand's Allergy clinic during 2012 to 2017 because of any allergy related symptoms evaluated by the specialist and relevant tests including serum total IgE, skin prick test and spirometry were performed for proper diagnosis.

Results: 2493 patients (mean age: 27 ± 15.4 years, range 1-83 years, M/F ratio: 0.77) who suffered from asthma, allergic rhinitis (AR), allergic rhino conjunctivitis (ARC), and other allergic disorders enrolled in this study. 47.4% (n=1182) of patients had allergic rhinitis or rhino conjunctivitis. 21.7 % and 16% had urticarial and atopic dermatitis respectively. Prevalence of asthma and history of drug allergy was 12% and 10% respectively.

Conclusion: The results of this study showed that allergic rhino conjunctivitis and urticaria are the most common causes of visiting specialist while asthma and drug allergy are the least.

Keywords: Allergic rhinitis, Asthma, Prevalence

ICIA2018\Allergy and immunotherapy\Poster\8639

8639

Estimating the Prevalence of Aero-allergens and Food Allergens in Patients with Atopic Eczema/Dermatitis

Mohammad Sadegh Naghizadeh^{1,2}, Afshin Derakhshani^{1,2}, Mahsa Najafzadeh^{1,2}, Mohammad Fereidouni³

1. Student Research Committee, Birjand University of Medical Sciences, Birjand, Iran

2. Asthma, Allergy & Immunology Research Center, School of Medicine, Birjand University of Medical Sciences, Birjand, Iran)

3. Cellular and Molecular Research Center, Birjand University of Medical Sciences, Birjand, Iran

Background: Prevalence of allergic diseases such as atopic dermatitis is reported to be increasing in recent decade. Identifying the many allergens is very important for diagnosis and treatment of allergic diseases. The purpose of this study was to investigate the prevalence of common allergens in patients with eczema at Birjand city of Iran.

Methods: Prevalence of 26 common aero and food allergens evaluated in atopic dermatitis patients who were referred to Birjand Allergy clinic during 2013 to 2017 by standard skin prick test (SPT).

Results: 123 patients (mean age: 28.63 ± 15.00 years, range 2-71 years, M/F ratio: 0/62) were enrolled in this study. Highest rate of skin sensitivity in food allergens were for coffee (15%), cereal mix (13%), peanut (10.4%), melon (10.3%) and black pepper (7%), and among aeroallergens, Russian thistle (53%), weed mix (48%), mugwort (38%), tree mix (36%) and chenopodium (29%) were the most common.

Conclusion: Recognition and avoidance of allergens in combination to medical therapy has an important role in management of atopic dermatitis. The rate of sensitization to indoor allergens was interestingly low which is probably because of geo-climatic situation. We conclude that the prevalence of allergens in each region is different depending on environmental conditions, food habits, ethnic diversities and life style.

Keywords: Allergy, Skin prick test, Atopic dermatitis, Food allergens, Aeroallergens

ICIA2018\Allergy and immunotherapy\Poster\8646
8646

Prevalence of Allergies and Pattern of Skin Sensitization among Patients Suffering from Otitis Externa

Kosar Salehitorabi^{1,2}, Afshin Derakhshani¹, Mahsa Najafzadeh¹, Mohammad Reza Mofateh³,
Mohammad Fereidouni^{2,4}

1. Student Research Committee, Birjand University of Medical Sciences, Birjand, Iran

2. Asthma, Allergy & Immunology Research Center, School of Medicine, Birjand University of Medical Sciences, Birjand, Iran

3. ENT department, Medical school. Birjand University of Medical Sciences, Birjand, Iran

4. Cellular and Molecular Research Center, Birjand University of Medical Sciences, Birjand, Iran

Background: Otitis externa is an inflammatory condition of external auditory canal. It is a common problem that can be seen at any ages and can be disabling for patients. The most common cause of otitis externa is bacterial infection, but fungal overgrowth, trauma and dermatologic conditions may be involved. There are some evidences that support the role of allergy in development of otitis externa. The aim of this study was to evaluate the frequency of allergies and pattern of skin sensitivity to different allergens among patients suffering from otitis externa.

Methods: Patients suffering from otitis externa evaluated for presence of any allergic disorders and skin prick test with a battery of at least 20 different allergens as well as positive and negative control performed by an allergist. 5 milliliters of venous blood was taken and after serum separation, the level of serum total IgE was measure by ELISA method.

Results: 78 patients (mean age: 31.95, in range of 4-75, M/F ratio: 0.33) were enrolled in this study. Prevalence of allergic rhino conjunctivitis, eczema, and asthma and food allergy was 24.7%, 2.6%, 1.3% and 1.3% respectively. 35.9% and 23.3% of patients had positive skin reactivity to mix of weeds and mix of trees extracts respectively.

Conclusion: The results of this study showed that except the allergic rhino conjunctivits, other allergies are not common among OE patients and weeds' pollen are the most common allergens among OE patients.

Keywords: Allergy, Otitis extern, Skin Prick Test.

ICIA2018\Allergy and immunotherapy\Poster\8647

8647

Comparison of the Pattern and Severity of Skin Prick Test Sensitivity to Different Allergens between Male and Females at Birjand

Mahsa Najafzadeh¹, Mohammad Fereidouni^{2*}

1. Student Research Committee, Birjand University of Medical Sciences, Birjand, IR Iran

2. Cellular and Molecular Research Center, Birjand University of Medical Sciences, Birjand, IR Iran
(corresponding author)

Background: Studies have shown that allergens are very important sensitizing agents in respiratory disorders such as asthma and allergic rhinitis that are common in Iran. The aim of this study was to determine the pattern and severity of skin prick test sensitivity to different allergens between male and females at Birjand, south Khorasan, Iran.

Methods: Skin Prick Test (SPT) was performed on 1484 patients for 33 common allergens. The blood sample was taken for measuring IgE concentration and eosinophil count.

Results: In the present study, the most skin reactions were observed for Russian thistle, mugwort, weed mix, lambsquarter and cat allergen which were 88.61, 64.728, 59.49, 46.326 and 39.71 mm², respectively. In contrast, the least skin reactions were seen for tree mix, peanut and betulaceae which were 12.268, 12.838 and 13.121 mm², respectively.

There were no significant association between the mean IgE levels in females (237.98 ± 12.11) and males (246.45 ± 13.64). However, the difference in the rate of eosinophilia in females ($4.13\% \pm 0.11$) and males ($4.88 \pm 0.15\%$) was statistically significant ($p < 0.001$).

In addition, a direct correlation was identified between total IgE and sum of the wheals ($p = 0.002$) and between IgE level and mean sum of the wheals ($p = 0.01$); however, there was no correlation between total IgE levels and mean of positive wheal.

Conclusion: The present study revealed the hypersensitivity to different allergens in Birjand city.

Keywords: Skin test, Allergens, Total IgE, Eosinophil

ICIA2018\Allergy and immunotherapy\Poster\8652

8652

Pattern of Allergies and Skin Prick Sensitization among Patients Suffering from Adenotonsillar hypertrophyNeda Mahavar^{1,2}, AfshinDerakhshani^{1,2}, Mohammad RezaMofatteh³, Mohammad Fereidouni^{1,4}*1. Asthma, Allergy & Immunology Research Center, School of Medicine, Birjand University of Medical Sciences, Birjand, Iran)**2. Student Research Committee, Birjand University of Medical Sciences, Birjand, Iran**3. ENT department, Medical school. Birjand University of Medical Sciences, Birjand, Iran**4. Cellular and Molecular Research Center, Birjand University of Medical Sciences, Birjand, Iran*

Background: Adenotonsillar tissue is a part of Waldeyer ring, which are responsible for antigen sampling from mouth cavity and antibody formation against antigens entering the body including allergens. Pathological growth of the adenoid and tonsil are mainly caused by antigenic stimulation as a part of inflammatory process. Adenotonsillar inflammation is one of the oldest and common problems and adenotonsillectomy is the most common operation in small children although is seen in adult as well. In spite of the high prevalence of adenotonsillar hypertrophy, the cause of this problem is unclear but some researches have shown that allergy may be a risk factor and Sensitization to a variety of allergens is believed to be involved in the development of adenotonsillar hypertrophy. The aim of this study was to evaluate the prevalence of allergic disorders in patients with adenotonsillar hypertrophy referring to Birjand's allergy clinic of Imam reza hospital.

Methods: 238 patients with adenotonsillar hypertrophy (mean age 14.12 years, range 3 to 51 years, M/F ratio: 1) were enrolled in our study. For all patients an allergy questionnaire were filled and skin prick test with a battery of 20 common allergen including tree and grass pollen, fruits, mites, molds and cockroach was done for all patients.

Results: 48.3% of patients had a positive skin reaction to at least one allergenic extract. The most common allergens were weeds including Russian thistle, Mugwort (42%), Rough pigweed (35.5%), weed mix (33.2%), Chenopodiaceae (22.31%) followed by Ragweed (14.8%) kiwi (14.3%), Melon (12.5%), coffee (12.3%) and cockroach (11.7%).

Conclusion: The results of this study show that a fairly large proportion of patients with adenotonsillar hypertrophy are sensitized to allergens particularly aeroallergens and therefore it is possible that allergy have a role in development of this problem.

Keywords: Adenotonsillar hypertrophy, Allergy, Skin prick test

ICIA2018\Allergy and immunotherapy\Poster\8661
8661

Comparison of the Skin Prick Test Wheal Size, Eosinophilia and Total Serum IgE between Allergic Male and Females at Birjand City, Iran

Mahsa najafzadeh^{1,2}, Neda Mahavar^{1,2}, Mohammadsadegh Naghizadeh^{1,2}, Mohammad Fereidouni^{*1,3}

1. Asthma, Allergy & Immunology Research Center, School of Medicine, Birjand University of Medical Sciences, Birjand, Iran)

2. Student Research Committee, Birjand University of Medical Sciences, Birjand, Iran

3. Cellular and Molecular Research Center, Birjand University of Medical Sciences, Birjand, Iran

Background: Studies have shown that allergens are very important sensitizing agents in different types of allergies especially allergic rhinitis and asthma but the pattern of sensitization and the allergenicity of different allergens are vary in different. The aim of this study was to determine the pattern of skin prick test sensitivity to different at Birjand city, Iran.

Methods: patients who suffered from any kind of allergies evaluated by an allergist and skin Prick Test (SPT) was performed.

Results: 1456 allergic patients (mean age: 26.03 ± 14.2 , range: 2-72 years, M/F ratio: 0.81) enrolled in this study. The average number of positive skin test in male and female was 1.8 and 1.7 respectively. The mean of skin wheal area because of allergenic extracts was larger in men than women but the difference was not significant (15.3 mm^2 vs 13.6 mm^2 , $P > 0.05$). There were no significant difference between the mean IgE levels in females and males (237.98 ± 12.11 vs 246.45 ± 13.64 , $P > 0.05$). However, the percentage of blood eosinophilia was significantly higher in males than females ($4.88 \pm 0.15\%$ vs $4.13 \pm 0.11\%$, $p < 0.001$).

Conclusion: The present study revealed that except for blood eosinophilia which is higher in males, there is no sex disparities in case of skin prick test reactivity to different allergens or total serum IgE.

Keywords: Skin test, Allergens, IgE levels

ICIA2018\Allergy and immunotherapy\Poster\9795

9795

The Relationship between Vitamin A and Serum IgE Level with Quality of Life in Patients with Allergic Rhinitis

Akram Davoodi¹, Firoozeh Khamoushi², Mohsen Molaei¹, Seyed Hamidreza Mortazavi³, Ali Gorgin Karaji⁴, Farhad Salari^{4*}

1. Student research committee, School of Medicine, Kermanshah University of medical sciences, Kermanshah, Iran

2. Student research committee, School of Nutritional Science & Technology, Kermanshah University of Medical Sciences, Kermanshah, Iran

3. Department of Pediatrics, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

4. Department of Immunology, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

Background: Allergic rhinitis (AR) is a common inflammatory disease of nasal mucosa that considerably affects the quality of life (QOL) of patients. Vitamin A (Vit A) and its metabolites such as retinoic acid (RA) play critical roles in development and regulation of the immune responses. Immunoglobulin E (IgE), a B cells-produced antibody, is involved in the pathogenesis of allergic diseases such as AR. The aim of this study was to investigate the relationship between dietary vit A intake and serum levels of IgE with QOL in AR patients.

Methods: This cross-sectional study included 36 AR patients with age 15–57 years. The serum level of IgE was measured by ELISA technique according to the manufacturer's instruction. The dietary vitamin A intake was measured by Food Frequency Questionnaire (FFQ) and analyzed by Nutrition 4 software. The Rhinitis Quality of Life Questionnaire (RQLQ) was used to assess the symptoms severity in AR patients. Data were analyzed by SPSS V21.

Results: The mean age of 36 patients were 34.7 ± 10.4 . 69.4% of patients were women. The mean of IgE serum concentrations in patients was 84.2 IU/mL. The daily intake of vit A was 781.8 ± 500.1 gr/day and so it was less than the recommended amount by recommended dietary allowances (RDA). Significant positive correlation was found between low intake of vit A with emotional symptoms (correlation coefficient: 0.367; $P=0.028$; $P<0.05$) and activity limitations (correlation coefficient: 0.330; $P=0.05$; $P<0.05$). Significant negative correlation was observed between IgE serum levels and practical problems (correlation coefficient: -0.376; $P=0.02$; $P<0.05$).

Conclusion: This study showed that IgE serum levels and dietary vitA intake are factors that can affect the life quality of AR patients.

Keywords: Allergic rhinitis, Immunoglobulin E, Quality of life, Vitamin A

ICIA2018\Allergy and immunotherapy\Poster\9904
9904

The Main Complaint of Patients with Allergies Referring to the Hospital of Ali ibn Abi Talib in Zahedan in 1396

Mahlajafari¹, fateme ranjbar²

1,2 - Bachelor of Science in Laboratory Sciences and Member of Young and Elite Research Club, Department of Laboratory Sciences, Islamic Azad University, Zahedan Unit, Iran

Background: Allergies are the reaction of the immune system to extremes over various factors. People who are hypersensitive have an ultrasignificant immune system against apparently harmless substances, and show an excessive response. Although the immune system protects the host against infections and alien factors, immune responses can cause injury and even disease. The aim of this study was to investigate the main complaint of patients with allergies referred to Imam Ali Hospital in Zahedan

Methods: The present study is a descriptive-analytical type. This study was performed on 404 patients referred to the Allergy Clinic of Zahedan Imam Ali Hospital in 1396. Information was collected using questionnaires completed by the patients and then analyzed by SPSSV21 software. And statistical analysis

Results: Of the patients, 53% of the patients were female and 47% were male, of whom 247 were single and the rest were married and the most frequent visits were in winter. 44.1% of patients had seasonal allergic reaction, 2.33% of cases of colds, 8.6% of cases of asthma and 9.5% of cases of febrile seizure, and 56.5% of them had no history of allergies

Conclusion: Allergic diseases are chronic diseases, affecting the quality of life of humans and are considered as the main causes of referral to health centers. It is better for authorities to prevent more social and economic harm than others by raising awareness of the people and adopting appropriate approaches

Keywords. Allergy, Complaint, Ali ibn Abi Talib, Zahedan



Cancer Immunology

Poster Presentation

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Cancer Immunology\Poster\5201

5201

Differential effects of inhibitory and stimulatory anti-HER2 monoclonal antibodies on intracellular signaling pathways

Tahereh Soltantoyeh¹, Tannaz Bahadori², Reza Hosseini Ghatar¹, Jalal Khoshnoodi¹, Azam Roohi¹, Maryam Mobini¹, Forough Golsaz Shirazi¹, Mahmood Jeddi Tehrani², Mohammad Mehdi Amiri¹, Fazel Shokri^{1,2}

1. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

2. Monoclonal Antibody Research Center, Avicenna Research Institute, ACECR, Tehran, Iran

Background: Homo- and hetero- dimerization of the receptor tyrosine kinase HER2 hyperactivate several downstream signaling pathways, leading to uncontrolled growth and proliferation of tumor cells. Anti-HER2 monoclonal antibodies (mAbs) may induce different effects on HER2 dimerization and signaling.

Methods: The effect of two inhibitory (2A8, 1T0) and one stimulatory (1H9) anti-HER2 mAbs either alone or in combination with Trastuzumab was investigated on AKT and ERK signaling pathways and HER2 degradation by Western blotting.

Results: While 1H9 mAb had no significant effect on AKT and ERK signaling pathways, 1T0 and 2A8 mAbs inhibited phosphorylation of both pathways. Combination of 1T0 mAb with Trastuzumab resulted in significant synergistic inhibition of both pathways and HER2 degradation, much more potently than the combination of Trastuzumab and Pertuzumab.

Conclusion: Our data indicate that anti-HER2 mAbs may induce different signaling pathways depending on their effect on tumor cell growth and proliferation. The significant inhibition of ERK and AKT phosphorylation by 1T0 alone or particularly in combination with Trastuzumab suggests its potential therapeutic application for targeted immunotherapy of HER2 overexpressing malignancies.

Keywords: Breast cancer, Signaling Pathways, Protein Kinase B, Extracellular Signal-Regulated Kinase, Monoclonal Antibody, Human Epidermal Growth Factor Receptor 2.

ICIA2018\Cancer Immunology\Poster\5232

5232

Effect of extremely low frequency of electromagnetic fields (ELF-EMF) on the argyrophilic nucleolar organizer regions (AgNORs) in bone marrow cellsMohammad Rafiee^{1,2}, Mohammad Abbasi³, Mohammad Yousef Alikhani⁴, Alireza Zamani^{5*}*1. Department of Medical Laboratory Sciences, School of Para medicine, Hamadan University of Medical Sciences, Hamadan, Iran**2. Department of Hematology and Blood Banking, School of Allied Medical Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran**3. Department of Internal Medicine, School of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran**4. Department of Microbiology, School of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran**5. Department of Immunology, School of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran*

Background: we are continuously exposed to increasing amount of the extremely low-frequency electromagnetic fields (ELF-EMF) in our daily life. This caused to generate concerns regarding the biological effects of such exposure. Therefore, this study was conducted to evaluate the influence of ELF-EMF on Argyrophilic nucleolar organizer regions (AgNORs) of bone marrow (BM) cells.

Methods: Eighty male rats were separated into four exposed groups (1,100,500 and 2000 μ T intensities) and a control group. The exposed groups were received ELF-EMF in solenoids which fed with the constant 50Hz sinusoidal power two hours a day for two months. The bone marrows were aspirated from bones of sacrificed rats and smears were prepared on slides. To analysis of BM cells, AgNOR staining was done on slides and Total AgNORs area (TAA), total AgNORs number (TAN) and total AgNORs length (TAL) were calculated by Scion Image Software and presented as pixels.

Results: The different intensities of ELF-EMF had no effect on the weights of rats. Significant reduction was seen in TAA, TAN and TAL in group of 1 μ T compared with groups of 2000 μ T and control. In addition, there was significant positive correlation between intensities increase of ELF-EMF and increase of TAA, TAN and TAL amounts.

Conclusion: It can be concluded that very low intensity (1 μ T) of ELF-EMF decreases AgNORs indices in bone marrow cells. But high intensity of 2000 μ T caused to return of AgNORs indices to control levels.

Keywords: Bone marrow, Argyrophilic nucleolar organizer regions, Extremely low-frequency electromagnetic fields, Proliferation, weight.

ICIA2018\Cancer Immunology\Poster\7422

7422

Cytotoxicity and Apoptosis Evaluation of *Artemisiaabsinthum*plant on AGS Gastric Cancer Cell Line

Elahe Aleebrahim Dehkordy^{1, 2}, RaziyeKarami², Amin Soltani², SadraAnsariPour³ NajmeShahinfard², Mahmoud Rafieian kopaei²

1. Young Researchers and Elite Club, Shahrekord Branch, Islamic Azad University, Shahrekord, Iran

2. Medical Plants Research Center, Basic Health Sciences Institute, Shahrekord University of Medical Sciences, Shahrekord, Iran

3. Student Research Committee, Shahrekord University of Medical Sciences, Shahrekord, Iran

Background: Gastric cancer remains the third leading cause of cancer related mortality worldwide. Natural products and derivatives of medicinal plants can play an important role to cure cancer. So Products derived from natural plants that induce apoptosis are used for cancer treatment. *Artemisia* is a genus belonging to the Asteraceae family with commonly fragrant species. This plant is widely used in Iranian traditional medicine. The current study was designed to evaluate the anticancer effect of this herb on AGS cancer cell lines.

Methods: Cytotoxic effect of hydro alcoholic extract on AGS cells was studied by MTT colorimetric assay and viability was monitored using cell counting and trypan blue. In order to investigate apoptosis, flow cytometry was performed by using Annexin V and PI apoptosis detection kit.

Results: The findings of the present study showed that the extract of *Artemisia absinthu* had inhibitory and apoptotic inducing effects on AGS cell lines. The results of MTT assay showed that the anticancer effect of *Artemisia absinthum* dependent to time and dose. The percentage of dead cell increased as dose arose. The 50% growth inhibitory concentration (IC₅₀) of extract in 48 and 72 h was 501± 21.15 and 413±16.12, respectively.

Conclusion: Our data well established the anti-proliferative effect of *Artemisia absinthum* extract, and clearly showed that the extract can induce apoptosis AGS cells. Therefore *Artemisia absinthum* could have growth inhibitory effect on AGS cell line dose and time dependent manner. It seems to come with further research, and utilizes its compound in cancer treatment.

Keywords: Cytotoxicity, Apoptosis, *Artemisia absinthum*, Gastric Cancer

ICIA2018\Cancer Immunology\Poster\7428
7428

FoxP3 Gene Variants and Breast Cancer: rs3761548 as Susceptibility not Prognosis Marker

Nadereh Naderi¹, Fahimeh Arabpour¹, Mahsa Rahimzadeh², Amin Shafizad³, Marzieh Norouzian¹

1. Molecular Medicine Research center, Hormozgan Health Institute, Hormozgan University of Medical Sciences, Bandar Abbas, Iran

2. Department of Biochemistry, Faculty of Medicine, Hormozgan University of Medical Sciences, Bandar Abbas, Iran

3. Department of Radiation oncology, Faculty of Medicine, Hormozgan University of Medical Sciences, Bandar Abbas, Iran

Background: Recent studies have characterized FoxP3 as a marker of regulatory T cells (Tregs) and an X-linked tumor suppressor gene which is involved in the pathogenesis of breast cancer. Therefore, we investigated the potential influence of a single-nucleotide polymorphism (SNPs) of FoxP3 gene on the development of breast cancer in Iranian women.

Methods: The association between FoxP3 rs3761548 polymorphism and breast cancer risk was assessed in 100 breast cancer patients and 108 healthy controls using sequence-specific primers (PCR-SSP).

Results: We identified significant difference of rs3761548 in both allele and genotype frequencies between cases and control groups. Our results showed that individuals carrying a FoxP3 rs3761548 AA genotype had about 4.3 fold increased risk of breast cancer compared with CC carriers.

Conclusion: This study has provided the first genetic data on the FoxP3 gene in south of Iran and proposes the rs3761548 polymorphism of FoxP3 gene as a risk factor in the development of breast cancer in Iranian population.

Keywords: Breast Cancer, FoxP3, Polymorphism, Treg cells

ICIA2018\Cancer Immunology\Poster\7430

7430

TGF- β Strengthens Cytotoxic effect and Senescence Induced by PI3K/ AKT/ MTOR Inhibitor in Human Colorectal Cancer Cells

Raziyeh Karami¹, Hedayatollah shirzad¹, Amin Soltani¹, Elahe Aleebrahim Dehkordy^{2,3}, Mahdi Ghatreh Samani^{1*}

1. Cellular and Molecular Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran

2. Young Researchers and Elite Club, Shahrekord Branch, Islamic Azad University, Shahrekord, Iran

3. Medical Plants Research Center, Basic Health Sciences Institute, Shahrekord University of Medical Sciences, Shahrekord, Iran

Background: Although NVP-BEZ235 and XAV-939 have been shown a promising result in the treatment of colorectal cancer (CRC), our data is little with regard to the effect of environmental cytokine on these drugs. In order to explore the reaction of CRC to this novel therapy, the current study proposes to investigate the inhibitory effect of NVP-BEZ235 and/or XAV-939 in CRC cells in combination with TGF- β as a tumor environmental cytokine to analyze their effect on cancer cell proliferation, cell death and expression of cancer promoting genes.

Methods: The cells were treated with single or combination of NVP-BEZ235, XAV-939 and TGF- β . Cell viability was evaluated by MTT assay. Flow cytometry method was used to determine cellular death. Real-time PCR was utilized to identify the effect of treatments on expression of snail, slug, c-myc and notch-1 genes.

Results: The results revealed that NVP-BEZ235 inhibited CRC cells proliferation and induced senescence cells more than XAV-939. Although, XAV-939 increased cellular death and inhibited senescence induction of CRC cells better than NVP-BEZ235. TGF- β pretreatment sensitizes CRC cells to the cytotoxic effect of NVP-BEZ235 and XAV-939. Real-time PCR analysis showed that resistance cells to NVP-BEZ235 increased expression of slug. Expression of notch enhanced remarkably when cells treated with both NVP-BEZ235 and XAV-939.

Conclusion: The results of the present study demonstrated that presence of anti-inflammatory cytokine like TGF- β augments the inhibitory effect of XAV-939 or especially NVP-BEZ235.

Keywords: TGF- β , Colorectal Cancer, NVP-BEZ235, XAV939

ICIA2018\Cancer Immunology\Poster\7457

7457

In vitro increasing cytotoxicity of CMV-specific CD8⁺ T cell in CMV-seropositive colorectal cancer patients

Shaker Ardakani M¹, Fazeli MS², Kokhaei P¹, Shakiba Y³, Tabatabaei Yazdi SM⁴, Abbasian A⁴, Emami AH⁵, Pak F¹, Nourizadeh M⁶

1. Department of Immunology, Semnan University of Medical Sciences, Semnan, Iran.

2. Department of Surgery, Imam Khomeini Complex Hospital, Tehran University of Medical Sciences, Tehran, Iran

3. Regenerative medicine Research center, Kermanshah University of Medical Sciences, Kermanshah, Iran

4. Tehran Blood Transfusion Center, Tehran, Iran.

5. Department of Hematology/Oncology, Imam Khomeini Hospital, Tehran University of Medical Sciences, Tehran, Iran

6. Immunology, Asthma and Allergy Research Institute, Tehran University of Medical Sciences, Tehran, Iran

Background: Colorectal cancer (CRC) is the third most common cancers with a high incidence of mortality and morbidity. A potential oncogenic role of HCMV in different types of cancers including CRC has been recently shown by detecting the CMV peptides in tumor tissues compared to the adjacent normal tissues. Targeting the CMV peptide-expressing tumor cells has been considered as an effective immunotherapy approach. Therefore, we tried to induce CMV-specific CD8⁺ cytotoxic T cells (CTLs) in vitro in CMV-seropositive CRC patients and compare it to healthy controls.

Methods: Sixteen CMV⁺ colorectal cancer patients and eight CMV⁺ healthy controls were enrolled the study. Patients' blood samples were obtained during tumor surgery (before chemotherapy and radiotherapy treatments). Cytotoxicity was measured by detecting the CD107a on the surface of CMV-specific CD8⁺ T cells using flow cytometry before and after 14-day culture of PBMCs in the presence of CMV peptide epitopes and IL-2 cytokine. Data were analyzed using FLOWJO software.

Results: CD107a expression on CD8⁺ T cells in both patient and control groups increased significantly after 14-day culture in the presence of CMV peptide epitopes and IL-2 which was an indicator of the potential cytotoxicity function of CTLs.

Conclusion: The results of this study specifies that in vitro stimulation of PBMC in the presence of CMV peptide epitopes and IL-2 can be an applicable method to generate potent CD8⁺ CTLs for future T cell therapy in CRC patients.

Keywords: Colorectal Cancer, CMV, CD107a

ICIA2018\Cancer Immunology\Poster\7468
7468

High Molecular weight Fractions derived from *Lactobacillus Reuteri* Cell-Free Supernatant Decrease Colon Cancer Stem- like Cell Invasion in Vitro

Faezeh Maghsood¹, Mahdi Rohani², Mehdi Kadivar^{1*}

1. Department of Biochemistry, Pasteur Institute of Iran, Tehran, Iran

2. Department of Microbiology, Pasteur Institute of Iran, Tehran, Iran

Background: Cancer stem cells (CSCs) have essential role in tumor metastasis and recurrence. Several factors are dysregulated in CSCs such as decreased expression of E-adherin, and tissue inhibitor of metalloproteinases, especially TIMP-1, and increased expression and activity of matrix metalloproteinases (MMPs), especially MMP-9 which are involved in tumor invasion and metastasis. Many studies have reported anti-cancer effects of *lactobacillus reuteri* as a probiotic bacterium. In this study, we investigated the effect of different molecular weight fractions derived from *L.reuteri* supernatant on invasion factors in a previously established human colon cancer stem-like cells derived from HT-29 cell line (HT29-CSC like).

Methods: Cell free supernatant (CFS) of *L.reuteri* was fractionated into defined molecular weight ranges (<3kD, 3-10 kD, 10-50 kD, 50-100 kD, >100 kD) using ultra-filtration tubes. HT29-CSCs were treated with different molecular weight fractions of CFS, and uninoculated bacterial growth media (MRS) as a control. Subsequently, cell invasion assessed using matrigel-coated insert plates. The expression of MMP-9, and TIMP-1 mRNAs were determined using real-time PCR, and the gelatinolytic activity of MMP-9 was evaluated by zymography.

Results: Cell invasion was significantly decreased following treatment with different molecular weight fractions of CFS in comparison with MRS-treated group ($p<0.05$). Also, 10-50 kD, 50-100 kD, and >100 kD fractions, significantly decreased activity and expression level of MMP-9 ($p<0.05$). Furthermore, expression level of TIMP-1 was significantly up-regulated by crude CFS, and >100 kD fraction ($p<0.05$).

Conclusion: Our results indicate that CFS of *L. reuteri* possesses anti-metastatic properties. Since fractions containing high molecular weight secretory factors decreased cell invasion, it suggests that the inhibitory compound(s) may fall into a macromolecule range such as a protein, nucleic acid, or a polysaccharide.

Keywords: Invasion, *Lactobacillus reuteri*, Matrix metalloproteinase, Probiotic

ICIA2018\Cancer Immunology\Poster\7498

7498

NK cell purification and expansion in the presence of IL-2, IL15 and IL18 with or without PBMCs feeder cells

Alireza Andalib, Anahita Najafalizadeh, Nafiseh Esmail, Fariborz Mokarian, Abass Rezaei

Immunology Department, Isfahan Medical School, Isfahan University of Medical Sciences, Isfahan, Iran

Background: NK cells are progressively considered as medical tools for cancer immunotherapy. The development of applicable methods to generate large numbers of functional NK cells is a crucial step to maximize the potential of this approach. So finding the appropriate condition for augment the NK cell number is part of the main aim of the study.

Methods: NK cells were isolated from PBMCs using Magni-Sort Human NK cell Enrichment methods. Purified NK cells were activated *in vitro* by IL-2 and IL-15 for two weeks. The HL-60 cell line was used as a target to assess NK *cytotoxic activity*. The cells were stained with FITC-conjugated anti-CD107a mAb, with PE-conjugated anti-CD3 mAb and PE/CY5-conjugated anti-CD56 mAb then NK cells were determined with flow cytometry method.

Results: The purity of NK cells was up to 92% by using the MACS method. Expanded NK cells were highly cytotoxic (90%) and more than 80% were viable. NK cell culture without using PBMCs as feeder cells even using IL-2+IL-15+IL18 induced no expansion of CD56+CD3- cells, but the culture including IL-15 + IL-2 has 1.3-fold expansion without using PBMCs as feeder cells.

Conclusion: NK cells did not expand significantly without feeder cells even using different cocktails of the cytokines. In contrast to T cells, NK cells cannot proliferate without feeder cells. The results using combination of IL-2, IL-15 and IL-18 were not significantly different from using IL-2 and IL-15 without using feeder cells.

Keywords: NK cells, IL-2, IL-15, IL-18, CD107a, feeder cells

ICIA2018\Cancer Immunology\Poster\7538
7538

Evaluation of TGF- β Expression in MCF-7 Cell Line after Treatment with Plant Extract

HesamBabaei Khameneh¹, Narges Amirjamshidi¹, Azin Aghamajidi¹, Abbas Azadmehr¹, Ebrahim Zabihi²,
Sadegh Fattahi³, AmrollahMostafazadeh^{*2}

1. Department of Immunology, Babol University of Medical Sciences, Babol, Iran

2. Cellular and Molecular Biology Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, Iran

3. Cellular and Molecular Biology Research Center, Babol University of Medical Sciences, Babol, Iran

Background: Breast cancer is one of the most common neoplasm and the second leading cause of cancer related deaths in women worldwide. Studies have been shown that the expression of TGF- β increased up in several subtype of breast cancer. Because of these findings, it has been suggested that, reduction of TGF- β expression can be a suitable target for prevention of tumor growth.

Methods: MCF-7 cells were cultured under cell culture conditions. MTT assay in dose and time dependent manner was conducted to find GI50 (growth inhibition 50) for each extraction. Then, the cells were exposed to different doses of plant extracts in triplicate. The level of TGF- β_1 gene expression was obtained by real-time RT-PCR.

Results: The results suggested that high concentration of *L.usitatissimum* extract significantly reduced TGF- β_1 mRNA levels and low concentration of *S.striata* extract dramatically increased the TGF- β_1 mRNA levels in MCF-7 cells.

Conclusion: According to the results, *L.usitatissimum* probably has some compounds that have negative effect on the TGF- β expression in MCF-7 cells.

Keywords: MCF-7, TGF- β , *linumusatissimum*, *scrophulariastriata*

ICIA2018\Cancer Immunology\Poster\7608

7608

NK cell purification and expansion in the presence of IL-2, IL15 and IL18 with or without PBMCs feeder cells

Alireza Andalib, Anahita Najafalizadeh, Nafiseh Esmaeil, Fariborz Mokarian, Abass Rezaei

Immunology Department, Isfahan Medical School, Isfahan University of Medical Sciences, Isfahan, Iran

Background: NK cells are progressively considered as medical tools for cancer immunotherapy. The development of applicable methods to generate large numbers of functional NK cells is a crucial step to maximize the potential of this approach. So finding the appropriate condition for augment the NK cell number is part of the main aim of the study.

Methods: NK cells were isolated from PBMCs using Magni-Sort Human NK cell Enrichment methods. Purified NK cells were activated *in vitro* by IL-2 and IL-15 for two weeks. The HL-60 cell line was used as a target to assess NK *cytotoxic activity*. The cells were stained with FITC-conjugated anti-CD107a mAb, with PE-conjugated anti-CD3 mAb and PE/CY5-conjugated anti-CD56 mAb then NK cells were determined with flow cytometry method.

Results: The purity of NK cells was up to 92% by using the MACS method. Expanded NK cells were highly cytotoxic (90%) and more than 80% were viable. NK cell culture without using PBMCs as feeder cells even using IL-2+IL-15+IL18 induced no expansion of CD56+CD3- cells, but the culture including IL-15 + IL-2 has 1.3-fold expansion without using PBMCs as feeder cells.

Conclusion: NK cells did not expand significantly without feeder cells even using different cocktails of the cytokines. In contrast to T cells, NK cells cannot proliferate without feeder cells. The results using combination of IL-2, IL-15 and IL-18 were not significantly different from using IL-2 and IL-15 without using feeder cells.

Keywords: NK cells; IL-2; IL-15; IL-18, CD107a, feeder cells

ICIA2018\Cancer Immunology\Poster\7615
7615

Thymol has selective cytotoxic effects

Pourya Tahmasebi, Seyyed Meysam Abtahi

Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Background: Thymol is a phenolic compound that possesses pharmacological activities, like anti-inflammatory, antioxidant, anticancer, and antimicrobial effects. Here, we investigated the selective cytotoxic effects of Thymol on peripheral blood mononuclear cells (PBMCs) and two cancer cell lines.

Methods: 1×10^6 live cells of PBMCs, LS174T (a colon cancer cell line) or T47-D (an invasive milk ductal carcinoma) were incubated with serial logarithmic dilutions of Thymol (0-400 μ M) for 24 h. Finally, by the time, the survivability of cells was determined by MTT method.

Results: Gained data indicated that Thymol had a profound cytotoxic effect on the LS174T and T47-D cell lines in a dose dependent manner. Interestingly, the IC₅₀ value of this compound in two cancer cell lines was lower than the IC₅₀ value in PBMCs.

Conclusion: As a result, Thymol provides favorable cytotoxicity in LS174T and T47-D cell lines without any additive cytotoxicity in PBMCs.

Keywords: Thymol, Peripheral Blood Mononuclear Cells, LS174T, T47-D, Anticancer.

ICIA2018\Cancer Immunology\Poster\7675

7675

The study on Hemiscorpiuslepturus (scorpionida: Hemiscorpiidae) venom cytotoxicity effects on K562 cells: in a in vitro studyNazanin Fathi¹, Golnaz Rashidi², Ali Khodadadi**1, 2, *.Department of Immunology, Faculty of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.*

Background: Cancer is one of the most important health problems in the world. In Iran, cancer is the third cause of death after cardiovascular disease and trauma. Evaluation of the therapeutic course of cancer is a very considered area, especially the using of natural products. Recently, using of animal venoms is to be interested by scientists to inhibit the cancerous cells. Scorpion toxins are a promising approach to confront cancer, since they have shown both *in vitro* and *in vivo* fights effects on cancer cells. The purpose of this study was to evaluate the susceptibility of *Hemiscorpiuslepturus* venom on K562 cell lines, had been derived from human chronic myeloid leukemia (CML).

Methods: Initially, calculating the concentration of protein venom by Bradford method, then K562 cells were treated with *H.lepturus* venom using an increasing rate of concentrations during a 24 hour incubation period and inhibition of CML growth was assessed by MTT assay.

Results: The amount of protein concentration in *H.Lepturus* crude venom was calculated 3.35 mg/ml. The MTT assay result showed that the IC₅₀ value was 14µg/ml after 24h incubation of K562 cells with scorpion venom.

Conclusion: The results of the current study suggest that H.lepturus venom increase the inhibition of growth activity of K562. Therefore, H.lepturus is hoped to be a step to *drug discovery* from natural toxins to treat the cancers.

Keywords: Hemiscorpiuslepturus, Venom, MTT, IC₅₀, K562 cells

ICIA2018\Cancer Immunology\Poster\7688
7688

Genistein promotes VEGF expression and Secretion in lymphoblastic leukemia cell lines

Mohsen Shahmoradi¹, Batoul Pourgheysari², Hedayatollah Shirzad¹, Fatemeh Shafiee³

1. Department of Microbiology and Immunology, Faculty of Medicine, Shahrekord University of Medical Science, Shahrekord, Iran

2. Medical Plants Research Center, Basic Health Sciences Institute, Shahrekord University of Medical Sciences, Shahrekord, Iran

3. Natural Language Processing (NLP) Research Lab, Faculty of Computer Science and Engineering, Shahid Beheshti University, Iran

Background: Genistein, a soy isoflavone, has the potential to arrest various types of cancers, partially through its anti-angiogenesis property. Vascular endothelial growth factor (VEGF), one of the angiogenic mediators, can be secreted by leukemic cells and plays an important role in tumor invasion and metastasis. In the present study, we examined the in vitro expression of VEGF mRNA and protein after treatment with genistein in two human hematopoietic tumor cell lines.

Methods: Two human lymphoblastic leukemic cell lines (Molt-4 and Jurkat) were used in this study. The cells were cultured in complete RPMI medium, and then incubated with ic_{50} concentration of genistein for 24, 48 and 72 hours. The mRNA expression of VEGF gene was determined by real-time polymerase chain reaction (RT-PCR). Furthermore, the level of VEGF secreted in the cell culture supernatants was measured using enzyme-linked immunosorbent assay kits (R and D systems).

Results: In both cell lines, VEGF protein secretion and mRNA expression were increased after treatment with Genistein in all three time periods. Likewise, genistein could act as an antineoplastic activity against jurkat and molt-4 cell lines.

Conclusion: Although the importance of angiogenic factors, like VEGF, was clearly established in solid tumors, this effect has not been fully elucidated in human hematopoietic neoplasms. In solid tumor cell lines, genistein usually decreased mRNA expression and secretion of VEGF while in our study, this pattern was different. This Difference could be useful to recognition of VEGF modulators and perhaps new treatment strategies for leukemia and other VEGF related disorders.

Keywords: Lymphoblastic cell line, Genistein, VEGF

ICIA2018\Cancer Immunology\Poster\7758
7758

Knowledge and Attitudes of Midwifery Students about Human Papilloma Virus Infection, Cervical Cancer and related Factors in Ahvaz Jundishapur University of Medical Sciences, Iran

Abdolhussein Shakurnia¹, Ata Ghadiri¹, Maede Hamidi², Samaneh Delavari¹

1. Department of Immunology, School of Medicine, Ahvaz University of Medical Sciences, Ahvaz, Iran

2. Student of Midwifery, School of Nursing and Medwifery Ahvaz University of Medical Sciences, Ahvaz, Iran

Background: Yearly, around 500,000 cases of invasive cervical cancer are diagnosed worldwide. Human papilloma virus (HPV) is a leading cause of cervical cancer and sexually transmitted infections are common health problems among young females. The knowledge and attitude of Midwifery have a considerable effect on HPV infection in women. The aim of this study was to determine the level of knowledge and attitude of Midwifery students regarding HPV infection, cervical cancer and related factors in Ahvaz Jundishapur University of Medical Sciences.

Methods: This descriptive cross-sectional study was conducted on 112 midwifery students. Data were collected using a questionnaire containing 20 items on demographic, items related to knowledge and attitudes the HPV infection and cervical cancer. Reliability and validity of the questionnaire was evaluated and confirmed. Scores higher than the average were defined as good knowledge and attitudes. The data was analyzed in SPSS 18 software by Exact Fisher and Kruskal-Wallis tests.

Results: In this study, mean age of medical students was 21.59 ± 2.41 years and all were female. The mean score of the knowledge was 26.61 ± 22.64 and 89.2% of students had poor knowledge about HPV. The mean score of attitude was 86.52 ± 10.52 and 97.3% of students had good attitudes. Only 2.7% of the students have been vaccinated against HPV. The relationship between the students' current educational year and their knowledge on HPV was not significant ($p=0.186$).

Conclusion: This study showed that the level of Knowledge of midwifery students about HPV infection was low. Therefore educational programs are needed to increase students' knowledge about HPV during their educational course. It seems that higher knowledge level of Midwifery can have significant effect on justification and informing target group for vaccination. Better knowledge and understanding of Midwifery regarding role of vaccination and decreasing mortality of cervical cancer can take a step to promote the condition of societal health.

Keywords: Human Papilloma Virus, Cervix Cancer, Knowledge, Attitude, Midwifery Students.

ICIA2018\Cancer Immunology\Poster\7780
7780

Evaluation of immune checkpoint molecules, PD-L2 and 4-1BB ligand (CD137L) expression on tumor tissue of colorectal cancer patients

Parastoo Shakerin¹, Kaveh Baghaei¹, Bijan Sedighi Moghaddam², Maryam Nourizadeh³

1. *Basic and Molecular Epidemiology of Gastrointestinal Disorders Research Center, Research institute for Gastroenterology and Liver Diseases, ShahidBeheshti University of Medical Sciences, Tehran, Iran*

2. *Semnan Medical University, Semnan, Iran.*

3. *Immunology, Asthma, and Allergy Research Institute (IAARI) Tehran University of Medical Sciences, Tehran, Iran*

Background: Immune checkpoint (ICP) molecules modulate the immune responses by either inducing or preventing T cell activation. Expressing some immune checkpoints on malignant cells has been shown to externally regulate anti-tumor immune responses.

In this study we aimed to investigate the expression levels of two immune checkpoint molecules, PD-L2 and 4-1BBL in tumor tissues of colorectal cancer (CRC) patients compared to the adjacent normal tissues.

Methods: RNA extraction and cDNA synthesis were done from tumor and adjacent normal tissues of 30 CRC patients. Expression of PD-L2 and 4-1BBL molecules was evaluated by real-time PCR. Protein levels of these molecules were assessed using immunohistochemistry (IHC) staining of the tissues.

Results: According to our IHC data, the expression of 4-1BBL ($P = 0.0006$) significantly decreased in tumor tissues of CRC patients compared to the controls, while considerable changes were not detected for PDL2 molecule. These results were in accordance to our real time PCR data.

Conclusion: Low expression of 4-1BBL, as an immune system activator, on cancer cells of CRC patients can be proposed as a target molecule to be synergized by applying agonistic anti-4-1BBL anti-bodies as an alternative immunotherapy approach for CRC patients.

Keywords: Immune Checkpoint Molecules, PD-L2, 4-1BBL

ICIA2018\Cancer Immunology\Poster\7857

7857

4-1BBL expressing B cells in breast cancer draining lymph nodesFereshteh Mehdipour¹, Mohsen Arabpour², Abdol Rasoul Talei³, Abbas Ghaderi^{1,2}

1. Shiraz Institute for Cancer Research, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

2. Department of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

3. Breast Diseases Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

Background: B cells can positively regulate immunity against tumor by increasing granzyme B expression in cytotoxic T cells via 4-1BBL/4-1BB axis. The aim of this study was to determine the frequency of 4-1BBL⁺ B cells in the tumor draining lymph nodes (TDLNs) of breast cancer patients and its change during disease progression.

Methods: Fresh axillary lymph nodes' specimens were taken from 46 patients and mononuclear cells were isolated using Ficoll-Hypaque gradient centrifugation. Cells were stimulated for 4 hours with PMA/Ionomycin. Stimulated and unstimulated cells were stained for CD19 and 4-1BBL and subjected to flow cytometry.

Results: 2.2±1.7 of unstimulated B cells were able to express 4-1BBL while 25.7±12.3% of B cells expressed this co-stimulatory molecule after stimulation. The level of 4-1BBL expression on B cells was not significantly different in the metastatic and non-metastatic lymph nodes. However, the frequency of 4-1BBL⁺ B cells in stimulated samples showed a tendency toward decrease in patients with grade III as compared with grade II+I (P=0.069). Moreover, the percentage of 4-1BBL⁺ B cells was significantly higher in TDLNs of ER⁺ or PR⁺ compared to ER⁻ or PR⁻ patients (P=0.031 and P=0.013, respectively).

Conclusion: The frequency of 4-1BBL⁺ B cells in TDLNs showed positive associations with good prognostic indicators such as ER⁺/PR⁺ and lower breast tumor grade which might be an indicative of their positive role in immunity against tumors.

Keywords: 4-1BBL, B cells, Tumor draining lymph nodes, Breast cancer

ICIA2018\Cancer Immunology\Poster

7889

Identification of antibody reactive proteins in pancreatic cancer using two-dimensional immunoblot coupled to mass spectrometry

Marzieh Rezaei^{1,2}, Saman Nikeghbalian³, Zahra Mojtahedi¹, Abbas Ghaderi^{1*}

1. Shiraz Institute for Cancer Research, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

2. Department of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

3. Department of Surgery, Nemazi Hospital, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Pancreatic ductal adenocarcinoma (PDAC) is a highly aggressive malignancy characterized by early invasiveness and resistance to treatment. Surgery in early stages is the only effective treatment, and the finding of new biomarkers for early detection of PDAC is still a major challenge. The present study aimed to compare immunoproteome between PDAC patients and healthy controls using serological proteome analysis (SERPA) method.

Methods: Cell lysates from two different pancreatic cancer cell lines were first separated by two dimensional (2D) gels, and then transferred onto the membranes probed with sera from 20 PDAC patients and 10 healthy controls. Proteins differentially reacted with autoantibodies in PDAC patients and the control groups were identified by mass spectrometry.

Results: This process led to the finding of the 18 pancreatic immunoreactive antigens can be classified into extra-cellular matrix and cytoskeletal proteins, enzymes, chaperones, signal transduction proteins and transcriptional regulators and containing of Laminin subunit alpha-5, Transcriptional regulator, Superoxide dismutase [Mn], ATP synthase subunit beta, mitochondrial ATP synthase beta subunit, Heterogeneous nuclear ribonucleoproteins C1/C2, Serine-threonine kinase receptor-associated protein, Protein SEC13 homolog, Eukaryotic translation initiation factor 3 subunit I, Chloride intracellular channel protein 1, Rho GDP-dissociation inhibitor 2, Elongation factor I-gamma, Mitochondrial Ef-Tu, chain A, Septin 2, Glyceraldehyde 3-phosphate-dehydrogenase, Phosphoglycerate mutase B isozyme, Prohibitin isoform 1, Tubulin beta 8 channel.

Conclusion: In the present study, we identified 18 immuno-reactive proteins in PDAC. While the identified proteins were critically involved in PDAC pathogenesis, further investigations in large scale population will determine the applicability of these potential biomarkers for early diagnosis or treatment of the disease.

Keywords: pancreatic ductal adenocarcinoma, biomarker, SERPA, autoantibody

ICIA2018\Cancer Immunology\Poster\7934
7934

Decrease of rankl and pthrp expression, genes involved in bone metastasis, in response to arbutin as an antioxidant in lncap cell line

Hajar Safari¹, Ebrahim Zabihi², Mahdi Pouramir³, Hamid Reza Nouri²

1. Student Research Committee, School of Medicine, Babol University of Medical Sciences, Babol, I.R. Iran.

2. Cellular and Molecular Biology Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, I.R. Iran.

3. Cancer Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, I.R. Iran

Background: Prostate cancer is the sixth highest cause of death in men worldwide. Prostate cancer lines such as LNCap express activated NF- κ B. The high levels of reactive oxygen species (ROS) lead to activation of NF- κ B, which affect expression of Receptor Activator of NF- κ B Ligand (RANKL) and parathyroid hormone-related protein (PTHrP) that are involved in osteoclast differentiation and skeletal localization of tumors. Therefore, in this study, effects of Arbutin as an antioxidant were evaluated on expression of these genes.

Methods: The human prostate carcinoma cell line, LNCaP, cultured in RPMI 1640 medium containing 5% fetal calf serum (FBS). Cells were stimulated with tert-Butyl hydroperoxide (tBHP) as ROS inducer and then treated by Arbutin in different doses. Intercellular ROS was measured by flow cytometry via a fluorogenic dye (H2DCFDA). After 12 h stimulation with Arbutin, cells were harvested and total RNAs were extracted using Trizol. The RNAs were reverse transcribed using random primers. The specific primers used to amplify RANKL and PTHrP by SYBER Green Real-time PCR. Gene expression was normalized to GAPDH by the $2^{-\Delta\Delta ct}$ method.

Results: Based on the flow cytometry obtained data, Arbutin effectively decreased intercellular ROS in dose dependent manner. It has been able to significantly reduce ROS in 1/10 and 1/100 dose of IC50. Analysis of RANKL gene expression significantly showed low level of expression in Arbutine treated group compared to tBHP as positive group. In addition, expression of PTHrP showed the same response to the Arbutin.

Conclusion: Taken together, the results of our study indicate that Arbutin in LNCap cells can decrease expression of osteoclast differentiation associated genes, thereby may be affects tumor-induced bone metastasis.

Keywords: Arbutin, ROS, Prostate cancer; RANKL; PTHrP.

ICIA2018\Cancer Immunology\Poster\7955
7955

Evaluation of prostate specific antigen (PSA) level in Suspicious Patients Who Referred to a Medical Diagnostic Laboratory in Ahvaz, Iran (2012-2016)

Abdolhussein Shakurnia¹, Asadollah Baesi², Masomeh Tehrani² Maryam Karami³

1. Department of Immunology, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

2. Department of health education, culture and research (ACECR), Branch of Khuzestan, Ahvaz, Iran

3. Veterinary Doctor, Shahid Chamran University, Ahvaz, Iran

Background: Prostate cancer is one of the most prevalent malignant cancer in men that is increasing mortality. Due to the easiest and most sensitive test for prostate cancer is serologic test, measurement of serum level of prostate specific antigen (PSA) is abundantly used for screening and diagnosis of prostate cancer. Considering the epidemiologic trend of prostate cancer, the aim of this study was to investigate the level of PSA in the males in a 5 period years from 2012 to 2016.

Methods: This study was retrospective cross sectional-descriptive. The required information of all patients who were suspected to Prostate cancer by physicians was extracted by laboratory software. The serum PSA level of 15178 patients was measured by ELISA method. According the procedure the subjects divided base on PSA values to 2 groups; lower level normal and higher level normal as health and patients men, respectively. Data was analyzed in SPSS 18 software by Chi-square tests.

Results: Mean age of the subjects was 57.56 ± 12.46 years, and the mean PSA in patients was 21.87 ± 24.35 ng/ml. The mean of PSA in patients over 50 years was 24.44 ± 25.43 ng/ml, which was significantly higher than the values of the patients under 50 years of age with 8.72 ± 10.80 ($p=0.001$). The prevalence rate of positive cases of PSA based on serological test was 2.09% (317/15178). Frequencies of positive cases of PSA has gradually increasing trend, so that the percentage of cases recorded between 2012 to 2016 was 15.4%, 17.3%, 19.5%, 22%, and 25.8%, respectively. However, this difference was not statistically significant.

Conclusion: Trends of increasing in prostate cancer in recent years are consistent with studies in other countries. Due to the aging of the Iranian population, developing one screening program for men is the way for prevention of prostate cancer progression and reducing mortality of this cancer recommended.

Keywords: Prostate cancer, prevalence, Prostate specific antigen (PSA), Ahvaz, Iran

ICIA2018\Cancer Immunology\Poster\7998
7998

Investigation of cytotoxic effect of *Tanacetumparthenium* and *Cuscutaepithymum* extracts on MCF7 human breast cancer cells

Homa Davoodi^{*1}, Mahdiah Naghavi Alhosseini², Masoumeh Mazandarani³, SeyedehNaemeh javid⁴

1. Cancer Research Center, Golestan University of Medical Sciences, Gorgan, Iran

2. Department of Immunology, School of Medicine, Golestan University of Medical Sciences, Gorgan, Iran

3. Department of botany, gorgan branch, Islamicacad university, gorgan, Iran

4. Department of Microbiology, School of Medicine, Golestan University of Medical Sciences, Gorgan, Iran

Background: Breast cancer is one of the most common carcinomas in women. Therapy on breast cancer including surgical operation, chemotherapy, radiotherapy, etc. were not completely efficient because of the serious side effect and resistance to anticancer drugs so that breast cancer still remains one of the leading causes of cancer-related death. Recently Complementary and alternative medicine for breast cancer considered to reduce or eliminate the disadvantages of conventional treatments. For centuries, traditional medicines especially natural products with herbal origin have been used for medicinal purpose. The present study was designed to explore the cytotoxic effects of *Tanacetumparthenium* and *Cuscutaepithymum* extracts on breast cancer cell line (MCF-7).

Methods: In order to evaluate the cytotoxicity properties of *T. parthenium* and *C. epithymum*, hydroalcoholic extract of these plants were prepared and then stored for further investigation. Cancer cells (MCF-7) were seeded in 96 well plates. Cultivated breast cancer cells were incubated with different concentrations (0.312-40 mg/ml) of the extracts for 48 hours and cell viability was determined using MTT assay.

Results: The results showed that in vitro exposures of the MCF-7 cells with different concentration of total extracts of these two plants significantly inhibited their growth and viability ($P<0.05$). In addition, the highest percentages of cell growth inhibition were found in concentration of 0.312 mg/ml with the values of 73.7% and 91.9% for *Tanacetumparthenium* and *Cuscutaepithymum*, respectively.

Conclusion: Hydroalcoholic extracts of *T. parthenium* and *C. epithymum* had cytotoxic effect on MCF-7 cell line and lead to cancer cell growth inhibition consequently. These findings are worth studying further and could have implications for future treatments and management of human breast cancer.

Keywords: Breast cancer, *Tanacetumparthenium*, *Cuscutaepithymum*, Cytotoxicity

ICIA2018\Cancer Immunology\Poster\8004
8004

Down-regulation of HSP40 gene family followed OCT4B1 suppression in human tumor cell lines

Shima Khajouee Ravari¹, Malek Hossein Asadi², Seyed Javad Mowla³, Mohammad Reza Mirzaei^{1*}

1. Molecular Medicine Research Center, Rafsanjan University of Medical Sciences, Rafsanjan, Iran

2. Departments of Biotechnology, Institute of Science and High Technology and Environmental Sciences, Graduate University of Advanced Technology, Kerman, Iran

3. Department of Molecular Genetics, Faculty of Biological Sciences, Tarbiat Modares University, Tehran, Iran

Background: OCT4B1, as one of OCT4 variants, were expressed in cancer cell lines and tissues more than other variants. We showed that this variant has an important role in regulation of cellular pathways as well as apoptosis and stress (heat shock protein). The aim of this study was to determine the effects of OCT4B1 silencing on expressional profile of HSP40 gene family in three different human tumor cell lines.

Methods: OCT4B1 expression were suppressed followed specific siRNA transfection in AGS (gastric adenocarcinoma), 5637 (bladder tumor) and U-87MG (brain tumor) cell lines with Lipofectamine reagent. The real-time PCR array technique was employed and the fold changes were calculated using RT² Profiler PCR array data analysis software version 3.5.

Results: Our results indicated that the expression profile of 36 studied genes from HSP40 family in studied tumor cell lines after OCT4B1 suppression showed approximately similar pattern of expression. Fifteen genes were down-regulated and tow genes (*DNAJC11* and *DNAJC5B*) were up-regulated in all three studied tumor cell line (more than 2 folds) and other studied genes (19 genes) showed different expressional pattern (up or down-expression) based tumor cell lines.

Conclusion: we suggest there is a direct correlation between OCT4B1 expression in tumor cell lines (and tissues) and HSP40 family gene expressions to escape from apoptosis and cancer expansion.

Key words: OCT4B1, HSP40 gene family, siRNA, tumor cell lines



ICIA2018\Cancer Immunology\Poster\8052
8052

Sirt1 Natural antisense transcript Down-regulates in Human Cancers

Neda Mokhberian¹, Seyed Mahmoud Hashemi², Vahid Jajarmi¹, Hossein Ghanbarian¹

1. Department of Biotechnology, School of Advanced Technologies in Medicine, ShahidBeheshti University of Medical Sciences, Tehran, Iran

2. Department of Immunology, School of Medicine, ShahidBeheshti University of Medical Sciences, Tehran, Iran

Background: Sirtuin1 (Sirt1) is a NAD-dependent deacetylase which is involved in the development and progression of human cancers. Recent studies have shown that a natural antisense transcript (NAT) is present in Sirt1 locus. Natural antisense transcripts (NATs) have recently been associated with the development of human cancers. Here, we identified the expression of Sirt1 sense and Antisense transcripts in several human cancers.

Methods: Following RNA extraction and cDNA synthesis, the expression level of Sirt1 mRNA and Sirt1-NAT was detected by quantitative real time-PCR assay in cancer cell lines and cancer clinical specimens.

Results: While Sirt1 mRNA level was upregulated in cancer cell line and cancer clinical specimens, the expression of Sirt1 antisense decreased in cancers compared to controls.

Conclusion: A converse relation was observed in the expression of Sirt1 sense and antisense transcript in normal and cancer tissues.

Keywords: Sirt1, Natural Antisense Transcript, Cancer, Cell line

ICIA2018\Cancer Immunology\Poster\8121

8121

HLA-G prognostic value in malignant liver and pancreas lesions and functional implications of HBV for HLA-G expression

Parisa Khodabandeh Shahraki,¹ Yasmin Zare,¹ Negara Azarpira,² Massood Hosseinzadeh,³ Shirin Farjadian¹

1. Department of Immunology, Shiraz University of Medical Sciences, Shiraz, Iran

2. Organ Transplant Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

3. Department of Pathology, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Human leukocyte antigen (HLA)-G is a nonclassical HLA class I molecule with modulatory effects on NK and T cells. Because HLA-G expression is frequently detected in different solid tumors, it may be involved in tumor immune evasion.

Objectives: We designed this study to elucidate the prognostic value of HLA-G in hepatocellular carcinoma (HCC) and pancreatic adenocarcinoma (PADC). The influence of hepatitis B virus (HBV) infection on HLA-G expression was also evaluated in patients with HCC.

Methods: HLA-G expression was investigated in tumor tissues from patients with HCC (n=74) or PADC (n=42) with immunohistochemical techniques. The presence of HBV genome was also examined in HCC tumor tissues by PCR.

Results: HLA-G expression was detected 66% of PADC and in 31% of HCC samples. In contrast to HCC, HLA-G overexpression was associated with advanced stages and grades in PADC. HBV genome was detected in 31% of HCC samples but we found no correlation between HLA-G expression and the presence of HBV genome in these tumors.

Conclusion: Our findings showed that HLA-G overexpression in tumor tissue correlated with poor prognosis in PADC. HLA-G expression is apparently affected more by the patient's genetic background and other epigenetic factors than by HBV infection.

Keywords: Human leukocyte Antigen-G, hepatocellular carcinoma, hepatitis B virus, pancreatic adenocarcinoma

ICIA2018\Cancer Immunology\Poster\8126

8126

Considering prevalence of B-RAF^{V600E} and PI3KCA genes' mutations in Iranian Colorectal cancer patients: Full COLD PCR and HRMAAtousa Homafar¹, NaserShagerdi Esmaeli², Shahram Samie³*1. Department of nutrition science, TabrizBranch, Islamic Azad university, Tabriz, Iran**2. Department of Hematology of ShahidBeheshti University of Medical Science, Tehran, Iran**3. Iranian high Institute for Research and Education in Transfusion Medicine, Tehran – Iran.*

Background: Colorectal cancer is the third most common cancer in the world. Mutations detection of B-RAF and PI3KCA genes is resolvable for choosing a target Anti-EGFR therapy. The aim of study is considering the mutation prevalence in B-RAF gene of by using Full-COLD PCR method and HRMA and in PI3KCA gene by using HRMA.

Methods: DNA of 27 FFPE of Colorectal Cancer was extracted. Therefore, for prevalence in exon 15 of B-RAF gene with using 2 methods combining of Full-COLD and HRMA was determined and for mutations in 2 exons 9 and 20 of PI3KCA gene by HRMA was used. Sensitivity of Full-COLD and HRMA by serial dilution of A-375 cell-line was determined. Sensitivity of this cell-line by serial dilution in Genomic DNA background with ARMS PCR method was considered. HCT-116 cell-line was chosen as positive control for PI3KCA gene mutations.

Results: Sensitivity of A-375 and HCT-116 cell-lines was 270 cell/ μ M. Sensitivity of combining 2 methods of Full-COLD and HRMA was %1. Sensitivity range of A-375 in Genomic DNA background by ARMS PCR was also %1. Although, HCT-116 cell-line was confirmed as positive control for mutations of 2 exons 9 and 20 of PI3KCA gene by using HRMA.

Conclusion: No detectable mutation in B-RAF and exon 20 of PI3KCA in extracted DNA from FFPE was found, but in exon 9 of this gene %3.7 was mutated. It seems Full-COLD plus HRMA method is more robust and sensitive method for detection or Screening of PI3KCA and B-RAF mutations.

Keywords: Colorectal cancer, B-RAF, PI3KCA, Full-COLD PCR, HRMA

ICIA2018\Cancer Immunology\Poster\8162
8162

Inhibition of the adenosine signaling in 4T1 cell lines has dose-dependent effects

Zahra Zeinaly, Seyyed Meysam Abtahi Froushani

Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Background: Adenosine plays an important role in inflammation and tumor progression, and responses to therapies. Some tumor cells like 4T1 (an invasive mice ductal carcinoma) can produce adenosine. Here, we investigated the effects of caffeine, as an adenosine antagonist and a routine drink, in mice peripheral blood mononuclear cells (PBMCs) and 4T1.

Methods: 1×10^6 live cells of PBMCs of mice and 4T1 were incubated different concentrations of caffeine (0, 0.1, 0.5 and 1 mM) for 48 hours. PBMCs were pulsed with phytohemagglutinin (final concentration of 5 μ g/ml). Finally, the survivability of cells was determined by MTT and neutral red uptake method.

Results: Attained results indicated that caffeine cannot change the rate of lymphocytes proliferation. Interestingly, caffeine at 0.5 mM significantly reduced the proliferation of 4T1 cells compared to other groups. Nonetheless, caffeine at 1 mM significantly potentiated the proliferation index of 4T1 cells.

Conclusion: The moderate dose of caffeine has beneficial anti-proliferative effect against 4T1 cells, Nevertheless higher doses of caffeine have revers and dangerous effects on 4T1 cells.

Keywords: 4T1 cell lines, Peripheral blood mononuclear cells, Caffeine.

ICIA2018\Cancer Immunology\Poster\8166

8166

Cytokine production by peripheral blood lymphocytes (PBLs) in the presence of cancer and normal adipose-derived mesenchymal stem cells (ASCs)Mahboobe Razmkhah ¹, Maryam Fakhimi ^{1,2}, Abbas Ghaderi ^{1,2}

1. Shiraz Institute for Cancer Research, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

2. Departments of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Tumors are complex tissues of different cell types, including stromal cells, cells of the immune system and mesenchymal stem cells (MSCs). MSCs play a significant role as immunomodulators in the tumor microenvironment through producing different cytokines. The present study aimed to investigate the effects of adipose derived mesenchymal stem cells (ASCs) on IL-10 and IL-17 cytokine production by peripheral blood lymphocytes (PBLs).

Methods: Human adipose-derived stem cells (ASCs) were obtained from breast adipose tissues of a breast cancer patient and a normal individual in cosmetic mammoplasty surgery. The ASCs were co-cultured with normal PBLs from 5 normal individual for 5 days. The production of cytokines including IL-10 and IL-17 in PBLs were measured by ELISPOT method.

Results: Based on the results, the production of IL-10 was increased as PBLs were cocultured with breast cancer ASCs compared to those cultured with normal ASCs or control group although this increased was not statistically significant (P value > 0.05). IL-17 production by PBLs decreased in the presence of cancer and normal ASCs compared with control. Comparing the effect of cancer-ASCs with normal-ASCs, the former had a significant effect on IL-17 production (P value $= 0.043$).

Conclusion: Based on the results of this study, the cross-talk between ASCs and the immune cells can induce the expression of anti-inflammatory cytokines such as IL-10 and reduce the expression of inflammatory cytokines such as IL-17 in the tumor microenvironment. Consequently, ASCs may promote tumor growth and progression through the immunomodulatory effects on different types of the cells from the immune system.

Key words: Adipose-derived mesenchymal stem cell, Breast cancer, Immunomodulation, IL-10, IL-17.

ICIA2018\Cancer Immunology\Poster\8175
8175

Membrane-Type-2 Matrix Metalloproteinase is selectively expressed on malignant myeloma cells

Shohreh Fakhari¹, Ali Jalili¹, Bahram Nikkhoo¹, Bayazid Ghaderi¹, Sako Mirzaie²

1. Cancer and Immunology Research Center, Kurdistan University of Medical Sciences

2. Department of biochemistry, Sanandaj Branch, Islamic Azad university, Iran

Background: Multiple myeloma (MM) is a plasma cell cancer that is considered 10% of all hematologic malignancies. Despite the development of new drugs for treatment of MM, it remains as one of the incurable cancer. Thus, performing new research is necessary to identify molecules that may play an important role in the biological behavior of myeloma cells. Matrix metalloproteinases (MMPs) are among the key players in migration and metastasis of many cancers. The role of some soluble MMPs (MMP 1,2,7,8,9,13) have been shown in hematologic malignancies, including multiple myeloma, like other cancers. However expression of Membrane Type-MMPs (MT-MMPs) is unknown in hematologic malignancies. Hence, in this study we evaluated expression of 6 MT-MMPs in myeloma cell lines, B cell lines, normal lymphocytes, mononuclear (MNC) and polynuclear cells (PMN).

Methods: We first screened expression of 6 MT-MMPs in the myeloma cell lines, B cell lines and normal lymphocytes, MNC and PMN cells by RT-PCR. Then, we performed IHC to evaluate MT2-MMP expression in 10 biopsies of normal bone marrow and 11 biopsies from patients with MM. In addition 7 bone marrow aspirated samples from MM patients MACS into 4 fractions (PMN, CD19+, CD138- , CD138+ cells) and MT2-MMP expression at mRNA and protein levels were examined by using qRT-PCR and flow cytometry, respectively.

Results: Our data showed for the first time that MT2-MMP, from among 6 MT-MMPs, is highly expressed in myeloma cells but not on other cells such as B cell lines and normal lymphocytes, MNC and PMN. Interestingly, IHC data showed that MT2-MMP was expressed in 100% (11 out of 11) MM biopsies and neither of 10 normal biopsies. Additionally, Real-time and flow cytometry data confirmed that expression of MT2-MMP is only expressed on malignant plasma cells (CD138+ cells), but not in normal fractions (CD138-, CD19+ cells and PMN).

Conclusion: In conclusion, we have shown for the first time that MT2-MMP is selectively expressed in myeloma cells and that MT2-MMP could be a suitable marker for diagnosis and possibly a novel target for treatment of MM.

Keywords: Multiple myeloma, Matrix metalloproteinase

ICIA2018\Cancer Immunology\Poster\8183

8183

H. pylori infection enhances CXCR4 and CXCR7 expression on gastric epithelial cellsAli Jalili¹, Shohreh Fakhari², Zeinab Kavyani^{1,3*}

1. Immunology Department, Medicine faculty, Kurdistan University of Medical Sciences, Sanandaj, Iran

2. Cancer and Immunology Research Center, Kurdistan University of Medical Sciences, Sanandaj, Iran

3. Maternal- Fetal and Neonatal Research Center, Tehran University of Medical Sciences, Tehran, Iran

Background: Chemokine SDF-1 is a main cytokine in recruitment and conservation of hematopoietic precursor cells in bone marrow that it's also produced in the site of tissue injury and attracts cells involving in tissue remodeling via its two specific receptors: CXCR4 and CXCR7. *Helicobacter Pylori* as a class-1 carcinogen plays an important role in development of gastric cancer by producing SDF-1 and recruitment of CXCR4/7 presenting cells. These cells can initiate dysplasia and metaplasia in gastric mucosa. In earlier study, the role of CXCR4/CXCR7/SDF-1 axis in gastric cancer and its prognosis in this cancer were evaluated so that enhanced expression of these receptors was related with metastasis; but the role of *H.pylori* in this process has not been investigated. In this study, we examined the effect of *Helicobacter Pylori* on CXCR4 and CXCR7 expression in gastric epithelial cells and gastric biopsy samples.

Methods: Gastric epithelial cell line, AGS, was co-cultured with purified colony of *H.pylori* in 100 bacteria/cell for 24h. Expressions of CXCR4 and CXCR7 mRNAs measured by RT-PCR and cell surface expression of these receptors evaluated by flow cytometry. Chemotaxis assay was performed to testing migration of treated and untreated AGS cells with *H.pylori* toward SDF-1 concentration gradient. Furthermore, expressions of CXCR4 and CXCR7 in gastric samples were assessed by Real-Time PCR.

Results: The results of the study showed that *H.pylori* exposure enhances CXCR4 and CXCR7 receptors expression both in cytoplasm and on cell surface in AGS cells. Migration of *H.pylori* treated cells toward SDF-1 gradient increased in comparison to control group. Gastric samples from *H.pylori* positive patients showed higher levels of CXCR4 and CXCR7.

Conclusion: We found that *Helicobacter Pylori* infection enhances CXCR4 and CXCR7 in gastric epithelial cells and also in gastric samples. Furthermore, enhanced migration capability to SDF-1 may be effective in gastric cancer pathology and metastasis.

Keywords: CXCR4, CXCR7, *Helicobacter Pylori*, Gastric cancer

ICIA2018\Cancer Immunology\Poster\8238
8238

IL-17 B expression in peripheral blood mononuclear cells of chronic lymphocytic leukemia patients

Mehdi Barati^{1, 2, 3}, Mehrnoosh pashaei^{2, 3}, Farahnaz Ghahremanfard^{3, 4}, Fatemeh Pak^{2, 3}, Parviz Kokhaei^{2, 3, 5*}

1. Department of Immunology, Faculty of medicine, Mashhad University of Medical Sciences, Mashhad, Iran.

2. Department of Immunology, Semnan University of Medical Sciences, Semnan, Iran.

3. Cancer Research Center, Semnan University of Medical Sciences, Semnan, Iran

4. Department of Internal Medicine, Semnan University of Medical Sciences, Semnan, Iran

5. Immune and Gene therapy Lab, Cancer Center Karolinska (CCK), Karolinska University Hospital Solna and Karolinska Institute, Stockholm, Sweden

Background: Interleukin 17B (IL-17B) as a pro-inflammatory cytokine is a member of interleukin 17 (IL-17) family cytokines (IL-17A - IL-17F). IL-17B Signal transduction mediated by a heterodimeric receptor complex that composed of IL-17RA/IL-17RB. Dis-regulation of IL-17B has been associated with a variety of diseases including cancer, autoimmune and inflammation disorders. Over expression of IL-17B in tumor, microenvironment is contributed to progression of several tumors. Therefore, the aim of this study was to investigate the expression of IL-17B in peripheral blood mononuclear cells in chronic lymphocytic leukemia (CLL) patients in compared to healthy subjects.

Methods: Initially RNA extraction and cDNA synthesis were performed on PBMCs of CLL patients (n=48) and healthy subject (n=35). RNA expression of IL-17B was quantified using Real-time PCR method.

Results: Our data indicated that IL-17B expression in PBMCs of CLL patients were significantly higher than normal subjects.

Conclusion: Overexpression of IL-17B may induce tumor progression. Therefore, targeting of IL-17B will be an effective therapeutic target in the future.

Keywords: chronic lymphocytic leukemia, IL-17B, tumor microenvironment, Real-time PCR

ICIA2018\Cancer Immunology\Poster\8274

8274

Comparison of Killing effect of synthetic and biologic nano selenium particles with on cell line derived from nervous cellsShideh Khangholi¹, Mehdi Mahdavi², Mohammadhossein Yazdi^{3*}1. *Pharmaceutical Sciences Branch, Islamic Azad University, Tehran, Iran*2. *Department of Immunology, Pasteur Institute of Iran, Tehran, Iran*3. *Department of Pharmaceutical Biotechnology and Biotechnology Research Center, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran*

Background: Today, brain tumors could be lethal without effective treatment especially in case of metastatic form. While many therapeutic approaches such as chemotherapy, radiotherapy and surgery have been utilized to cure this disease. The supplement substances such as selenium can play important role in this matter. As this respect, selenium nano particle is produced in two forms of synthetic and biologic to evaluate cytotoxic effect on nervous cell line in the present study.

Methods: A-172 nervous cell line was prepared and cultured in DMEM medium. Selenium nano particle was used in two forms of synthetic and biologic. The cytotoxicity of synthetic and biologic Nano selenium was assayed by MTT test and the result was read by ELISA reader in the wavelength of 570 nanometers.

Results: There is significant difference in group treated with biologic Nano selenium as compared to singular cell in concentrations of 100, 300 and 400 µg/ml, 400 µg/ml, 25, 100, 200, 300 and 400 µg/ml after 24, 48 and 72 hrs respectively. Also, there is remarkable difference in group treating with Synthetic Nano selenium in compared to singular cell in certain concentrations after 24, 48 and 72hrs. as a result, biologic Nano selenium has more cytotoxic effect than synthetic.

Conclusion: Nano selenium be able to suppress growth of nervous cell line via cytotoxicity function. Also biologic Nano selenium has more killing powerful than synthetic one.

Keywords: Selenium, biologic Nano selenium, synthetic Nano selenium, A-172 nervous cell line.

ICIA2018\Cancer Immunology\Poster\8284
8284

Kaempferol effect on telomerase gene expression in hela cervical carcinoma Cell Line

Saiedeh Erfanian¹, Maliheh Moradzadeh², Mohsen Farhang³

1. Department of Advanced Medical Sciences and Technologies, School of Medicine, Jahrom University of Medical Sciences (JUMS), Jahrom, Iran

2. Golestan Rheumatology Research Center, Golestan University of Medical Sciences, Gorgan, Iran

3. Student of Laboratory Sciences, Student Research Committee, Jahrom University of Medical Sciences, Jahrom, Iran

Background: Cervical cancer is one of the most common female cancers, while chemotherapy for this type of cancer is still not satisfactory. In recent years, the use of herbal medicines is expected to rise sharply. The effectiveness of these products of natural origin has urged researchers to discover new medicinal plants. Kaempferol is a natural flavonol which has therapeutic effects. Previous studies have proven the antioxidant and anti-inflammatory effects of kaempferol. In this study, the inhibitory effect of kaempferol on cervical carcinoma (HeLa) cell line proliferation and telomerase genes activity have been studied.

Method: After culturing, HeLa cells were treated by 25.50 and 75 micro-molar kaempferol concentrations, and 5-Fluorouracil was treated at a concentration of 80 mM. Also, a combination of kaempferol with concentration of 50mM and 5-Fluorouracil was used for the treatment. HeLa cells viability after proliferation was evaluated by cytotoxicity assay. Also, the polymerase chain reaction quantitative method (real time pcr) was used to evaluate the expression of TRF2 and Htert genes.

Results: Treatment with kaempferol significantly inhibited the proliferation of HeLa cells compared to the control group. Htert and TRF2 gene expression in all studied groups was significantly reduced (P-value <0.05).

Conclusion: In this study, for the first time we showed that kaempferol can be a potential inhibitor for the cervical cancer cells. Since kaempferol has lower toxicity compared to chemotherapeutic clinically used agents such as 5-Fluorouracil, it can be used as adjunct therapy in combination with 5-Fluorouracil in patients with cervical cancer.

Keywords: kaempferol, cervical cancer, inhibition of cell proliferation

ICIA2018\Cancer Immunology\Poster\8285

8285

Kaempferol effect on cell proliferation inhibition and apoptosis induction in Acute Myeloid Leukemia Cells line (HL60)Saiedeh Erfanian¹, Maliheh Moradzadeh², Mohsen Farhang³*1. Department of Advanced Medical Sciences and Technologies, School of Medicine, Jahrom University of Medical Sciences (JUMS), Jahrom, Iran**2. Golestan Rheumatology Research Center, Golestan University of Medical Sciences, Gorgan, Iran**3. Student of Laboratory Sciences, Student Research Committee, Jahrom University of Medical Sciences, Jahrom, Iran*

Background: Leukemia, one of the most common cancers in Iran, is a progressive and malignant disease of the blood-forming part of the body that is caused by the proliferation and development of white blood cells and its precursors in the blood and bone marrow. Introducing all trans retinoic acid (ATRA) for the treatment of acute promyelocytic leukemia (APL) patients could dramatically improve their survival rate. However, resistance to this drug and their toxicity are major problems in the treatment of APL. Earlier studies suggested that in haematopoietic neoplasms, the green tea polyphenol Epigallocatechin gallate (EGCG) induces cell death without adversely affecting healthy cells. This study was designed to evaluate the effect of EGCG on the level of lysis and expression of genes involved in apoptosis and cellular growth of acute myeloid leukemia cell line.

Methods: In the present study, the potential anti-promyelocytic leukemia activity of EGCG and the underlying molecular mechanisms were investigated. Acute Myeloid Leukemia Cells line (HL60) were cultured in RPMI medium and then incubated with different concentrations of EGCG (100-12.5 μ M). Cell survival (Viability) was measured by MTT assay. Real time PCR was also used to evaluate the apoptotic expression of genes and confirm the apoptotic mechanism.

Results: EGCG (100 μ M) significantly inhibited proliferation and induced apoptosis in HL-60 cells. The effect was associated with the expressions of proapoptotic genes p21 and Bax/Bcl-2 ratio were significantly increased.

Conclusion: we can conclude that the results of this study, changes in the level of expression of these genes, and the matching of the results with the findings of previous studies, can be argued that the effect of EGCG on acute myeloid leukemia cells is undeniable effects. However, further studies on these compounds are suggested to identify a more precise mechanism.

Key words: acute promyelocytic leukemia; EGCG; Apoptosis

ICIA2018\Cancer Immunology\Poster\8307
8307

Reduced Levels of T-helper 17-associated Cytokines in the Serum of Patients with Breast Cancer: Indicators for Following the Course of Disease

Mohammad Reza Atashzar ¹, Rasoul Baharlou¹, Abbas Ahmadi Vasmehjani¹, Ebarahim Rahimi², Majid Khoshmirsafa¹, Farhad Seif¹, Maryam Mahdiyar³

1. Department of Immunology and Microbiology, School of Medicine, Jahrom University of Medical Sciences, Jahrom, Iran

2. Department of Epidemiology, School of Medicine, Jahrom University of Medical Sciences, Jahrom, Iran

3. Department of Student Research Committee, School of Medicine, Jahrom University of Medical Sciences, Jahrom, Iran

Background: Interleukin (IL)-17 producing CD4+ T helper (Th17) cells that are known by producing IL-17 have recently been defined as a unique subset of proinflammatory helper cells. IL-17 is an inflammatory cytokine with robust effect on many cells and it can play important roles in pathogenesis of diverse groups of immune-mediated diseases. Therefore, the aim of this case control study was to determine serum levels of IL-17, IL-6, and transforming growth factor beta (TGF- β) in Iranian breast cancer patients.

Methods: Blood samples were collected from 55 patients with breast cancer and 34 healthy individuals with no history of malignancies or autoimmune disorders, based on simple sampling. The serum levels of IL-17, IL-6 and TGF- β were measured by the enzyme linked immunosorbent assay (ELISA).

Results: The serum level of IL-6 was significantly lower in patients with breast cancer compared with healthy individuals ($p=0.0003$), and also their IL-17 was lower in patient group than controls ($p=0.01$). Interestingly, TGF- β serum level in patients was less than controls ($p<0.0001$).

Conclusion: As most of the cases investigated in this study were in early stages, it is concluded that reduced IL-17, IL-6 and TGF- β can be used as predictors for clinical stage and prognosis of cancers such as breast carcinoma.

Keywords: Breast cancer, Interleukin-6, Interleukin-17, T-helper 17, Transforming growth factor beta

ICIA2018\Cancer Immunology\Poster\8383
8383

Study of serum IFN- γ and IL-4 levels in diffuse large B cell lymphoma (DLBCL) patients of response and non-response to Rituimab and DLBCL patients of under treatment with Rituximab

Nasrin Zare¹, NahidEskandari², Shaghayegh Haghjoo Javanmard³, Valiollah Mehrzad⁴

1. Department of Immunology, School of Medicine, Isfahan University of Medical sciences

2. Department of Immunology, School of Medicine and Applied Physiology Research Center, Isfahan University of Medical Sciences

3. Department of physiology, School of Medicine and Applied Physiology Research Center, Isfahan University of Medical Sciences

4. Department of Hematology and Medical Oncology, Isfahan University of Medical Sciences

Background: Patients with DLBCL have variable responses to rituximab. Cytokines might be deregulated after using of rituximab and also in DLBCL patients of response and non-response to Rituximab. The current study was designed to investigate and compare serum IFN- γ and IL-4 levels in DLBCL patients of response, non-response to Rituximab and DLBCL patients under treatment with Rituximab.

Methods: Serum patients were obtained from the Isfahan's Sayed Al-Shohada Hospital, Iran. Serum IFN- γ and IL-4 were measured using the Human IFN- γ ELISA MAXTM Deluxe (Biolegend) and Human IL-4 ELISA development kit (Mabtech).

Results: The IFN- γ and IL-4 were measured in serum of DLBCL patients of response (n=8), non-response (n=8) and under treatment (n=8) with Rituximab. The Mean serum values for these patients, respectively, were: for IFN- γ , 6.14 pg/mL, 8.9 pg/mL and 7.8 pg/mL; for IL-4, 4.3 pg/mL, 2.1 pg/mL and 0.005 pg/mL. DLBCL patients of under treatment with Rituximab exhibited a decrease in IL-4 levels compared to the other two groups (P=0.06). There were no significant difference in IFN- γ (P=0.3).

Conclusion: These results indicate that decrease in serum IL-4 and IFN- γ levels may be appropriate biomarkers to identify DLBCL patients with variable responses to rituximab.

Keywords: DLBCL, IFN- γ , IL-4, Rituximab

ICIA2018\Cancer Immunology\Poster\8406

8406

IL-17RB expression in peripheral blood mononuclear cells of chronic lymphocytic leukemia

Mehrnoosh pashaei^{1,2}, Mehdi Barati^{1,2,3}, Farahnaz Ghahremanfard^{2,4}, Fatemeh Pak^{1,2}, Parviz Kokhaei^{1,2,5*}

1. Department of Immunology, Semnan University of Medical Sciences, Semnan, Iran

2. Cancer Research Center, Semnan University of Medical Sciences, Semnan, Iran

3. Department of Immunology, Faculty of medicine, Mashhad University of Medical Sciences, Mashhad, Iran

4. Department of Internal Medicine, Semnan University of Medical Sciences, Semnan, Iran

5. Immune and Gene therapy Lab, Cancer Center Karolinska (CCK), Karolinska University Hospital Solna and Karolinska Institute, Stockholm, Sweden

Background: B Chronic lymphocytic leukemia (B-CLL) is the most common adult leukemia in the western world. The pathogenesis of CLL has been attributed to various forms of immune dysregulations. There are evidence that T cell dysfunction and differentiation shift to Th2 in B-CLL which may contribute to progress of the disease. The IL-17 receptor B (IL-17RB) is a receptor for IL-17E and also able to bind IL-17B. Stimulation of IL-17RB promotes Th2 responses. However, IL-17RB in B-CLL has not been investigated so far. Therefore, this study was performed to evaluate the expression of IL-17RB in peripheral blood mononuclear cells in B-CLL patients compare to healthy subjects.

Methods: Initially total RNA was extracted from peripheral Blood Mononuclear Cells (PBMCs) of CLL (n=48) and normal subjects (n=35) and cDNA were synthesized. The mRNA expression of IL-17RB was quantified using Syber Green Real-time PCR method, using ABI 7900 machine.

Results: Our data indicated that IL-17RB expression in PBMCs of CLL patients were significantly higher than normal subjects (P value <0.01, P value= 0.0090).

Conclusion: High expression of IL-17RB in B-CLL may induce tumor cells viability through its ligands. Tumor cells may express these receptors to evade from apoptosis and promote tumor cells survival. Therefore, blocking of IL-17RB could be new approach for translational investigation and clinical application.

Keywords: chronic lymphocytic leukemia, IL-17RB, Peripheral blood mononuclear cells, Real-time PCR

ICIA2018\Cancer Immunology\Poster\8407
 8407

The study of Indirubin effect on peripheral blood mononuclear cell of chronic lymphocytic leukemia patients in invitro

H. Jafarinezhad^{1, 2} (M.Sc.), B. Sedighimoghadam¹ (PhD), A. Aghaie³ (PhD), N. Ghanizadeh^{2, 1} (M.Sc.), M. Faranoosh⁴ (MD), S. Rostami⁵ (PhD) P. Kokhaei^{6, 1 *} (Professor-PhD)

1. Cancer Research Center and Immunology Department, Faculty of Medicine, Semnan University of Medical Sciences, Semnan, Iran
2. Student Research Committee, Semnan University of Medical Sciences, Semnan, Iran
3. Blood Transfusion Research center, High institute for Research and Education in Transfusion Medicine, Tehran, Iran
4. Children's Development Research Center, Endocrinology and Metabolism Research Center, Iran University of Medical Sciences, Tehran, Iran
5. Hematology, Oncology and Stem Cell Transplantation Research center, Tehran University of Medical Sciences, Tehran, Iran
6. Laboratory of Immunology and Gene Therapy, Karolinska Cancer Center, Karolinska University Hospital, Stockholm, Sweden

Background: Chronic lymphocytic leukemia (CLL) is a malignancy of naive B-cells. Indirubin is an active constituent of a traditional Chinese herbal medicine. Indirubin or its derivatives had the ability to induce apoptosis in cancer cell. Therefore, the aim of this study is evaluation of indirubin effect on peripheral blood mononuclear cell of chronic lymphocytic leukemia patients compared with healthy donor in invitro.

Methods: Peripheral blood samples (5 mL) obtained from 14 CLL patients and 8 age-matched healthy subjects. Peripheral blood mononuclear cells (PBMC) were isolated from blood patients and healthy subjects by using Ficoll technique. PBMC cultured with indirubin (in concentration of 0.1 to 80 μ M) for 24, 48 and 72 hours. The cytotoxicity percentage of indirubin was measured by MTT technique and also apoptosis percentage was detected by AnnexinV Assay. Finally BAX, Bcl2, CDK1 and CDK2 mRNA expression were investigated by Real time PCR using SYBR Green method.

Results: The MTT assay showed that indirubin has high cytotoxicity percentage (20 μ M, 48h) in the PBMC of CLL patients compared with control group ($P < 0.05$). The Annexin v assay indicated that the indirubin has high apoptosis percentage (20 μ M, 48h) in the PBMC and B cells (CD19⁺ cells) compared with control group ($P < 0.05$). The expression level of Bcl2, CDK1 and CDK2 mRNA in PBMC of CLL patients that treated with indirubin compared with control group decreased significantly ($p < 0.05$) and BAX mRNA level showed no significant differences in PBMC of CLL and healthy donors ($p > 0.05$).

Conclusion: It seems indirubin induces apoptosis in tumor cells in dose and time dependent manner. This herbal component causes apoptosis by down regulation of Bcl2 level mRNA and induces cell cycle arrest by down regulation of CDK1 and CDK2 level mRNA in PBMC of CLL patients.

Keywords: Chronic lymphocytic leukemia, CDK1, CDK2, Bcl2, BAX

ICIA2018\Cancer Immunology\Poster\8421
8421

Lower circulating levels of interleukin-10 in patients with bladder cancer

Taybeh Tavakoli¹, Hossain Khorramdelazad¹, Mohammad-Taghi Rezayati¹, Somayeh Abolhassani¹, Somayeh Pourtalebi¹, Abdollah Jafarzadeh²

1. Department of Immunology, Medical School, Rafsanjan University of Medical Sciences, Rafsanjan, Iran

2. Department of Immunology, Medical School, Kerman University of Medical Sciences, Kerman, Iran

Background: Interleukin 10 (IL-10) is considered an immune-regulatory cytokine, exerting both antitumor and pro-tumor properties. The role of IL-10 in tumor development is very controversial. The aim of this study was to investigate the serum levels IL-10 in patients with bladder cancer (BC).

Methods: The blood samples were collected from 28 patients with bladder cancer and 23 age and gender-matched healthy subjects as a control group. The serum levels of IL-10 were measured by ELISA.

Results: The mean serum levels of IL-10 was 8.39 ± 3.17 Pg/ml in BC group and 10.72 ± 3.60 in healthy subjects. The mean serum levels of IL-10 in BC patients was significantly lower than healthy group ($P < 0.02$). The serum levels of IL-10 in BC men was significantly lower than healthy men (7.96 ± 3.44 Pg/ml vs 12.26 ± 3.60 Pg/ml, $P < 0.01$). However, IL-10 levels similarly expressed in BC women and healthy women.

Conclusion: These results showed lower levels of IL-10 in patients with BC. The IL-10 levels were affected by gender of patients. The clinical values of IL-10 in BC need more consideration.

Keywords: Bladder cancer, Interleukin-10, Serum, Gender

ICIA2018\Cancer Immunology\Poster\8424

8424

The Hepatocyte Growth Factor (HGF) can act as immune modulator by increasing IL10, TGF β and IFN γ cytokines in Mesenchymal Stem Cells: in vitro studyKavehTari¹, Ali Jalili¹*1. Cancer and Immunology Research Center, Kurdistan university of medical science*

Background: The Hepatocyte Growth Factor (HGF) has important role as a modulator of immune system. This factor produce mainly by Bone Marrow Mesenchymal Stem Cells (BM-MSCs.) The purpose of this study was evaluation of the effect of HGF on human mesenchymal stem cells to produce the gene of IL10, TGF β and IFN γ cytokines and suppression of the proliferation of Peripheral blood mononuclear cell(PBMCs.)

Methods: In this study, the BM-MSCs were Isolated, cultured, characterized and then treated with HGF at dose of 20 ng/ml in different times. IL10, TGF β and IFN γ gene were measured by qRT-PCR, Also, the ability of the HGF in suppression of the proliferation of PBMNC cells were evaluated by using MLR. Results were analyzed by SPSS software and T-test.

Results: The results showed significant production of IL10, TGF β and IFN γ upon HGF treatment in RNA level (P<0.05). Also a significant decrease of PBMNC proliferation co-cultured with HGF treated- MSCs were seen (P<0.05).

Conclusion: The findings showed that using of HGF could decrease proliferation of HSCs by altering expression of key cytokines in the process of modulating of immune system.

Keywords: Hepatocyte Growth Factor, Mesenchymal Stem Cells, Immune Modulating.

ICIA2018\Cancer Immunology\Poster\8439
8439

CD25 expressing B cells in head and neck squamous cell carcinoma draining lymph nodes

Marzie Norouzian^{1,2}, Fereshteh Mehdipour², Sima Balouchi Anaraki^{1,2}, Bijan Khademi³, Abbas Ghaderi^{1,2}

1. Department of Immunology, School of Medicine, Shiraz University of Medical Sciences

2. Shiraz Institute for Cancer Research, School of Medicine, Shiraz University of Medical Sciences

3. Department of Otolaryngology, Khalili Hospital, Shiraz University of Medical Sciences

Background: Tumor-draining lymph nodes (TDLNs) are the main site in which tumor-specific immune responses are initiated. B cells are among main residents of TDLNs but compared to T cells, little is known about phenotype and function of B cell subsets repopulated in the TDLN. It has been suggested that B cells expressing CD25 have different phenotypical and functional properties as compared to CD25⁻ B cells. CD19⁺CD25^{hi} B cells have also been reported to have regulatory capacity in humans. The aim of this study was to assess the frequency and phenotype of CD25 expressing B cells in the TDLNs of head and neck squamous cell carcinoma.

Methods: Freshly isolated lymph node cells were directly stained with fluorochrome conjugated anti-CD19, -CD25, -CD27, and -IgM antibodies or their isotype controls and analyzed by flow cytometry.

Results: Flowcytometric analysis showed that 21.7±17.4% and 6.9±2.1% of CD19⁺ cells in TDLNs were CD25⁺ and CD27^{hi} CD25⁺, respectively. Moreover assessment of CD27 by CD25⁺ B cells revealed that 68.4± 20.2% of CD25⁺ B cells expressed CD27. In addition, 24.1± 16% of CD25⁺ B cells showed naïve (CD27-IgM⁺) phenotype while 22.7±15.3% and 40.5± 15.5% of them had unswitched memory (CD27⁺ IgM⁺) and switched memory (CD27⁺ IgM⁻) phenotypes, respectively.

Conclusion: A subset of B cells expressing CD25 in TDLNs of patients with head and neck squamous cell carcinoma was detected. These CD25⁺ B cells had either memory or naïve phenotypes, most of them however, showed memory phenotype. Further functional and phenotypical investigations are needed to understand the function of these B-cell subsets in TDLNs of head and neck squamous cell carcinoma.

Keywords: Head and Neck squamous cell carcinoma, CD25⁺ B cell, tumor draining lymph nodes

ICIA2018\Cancer Immunology\Poster\8477

8477

Investigation of the gene polymorphism of *CCL22* 16C/A in patients with Basal Cell CarcinomaMaryam Fakhimi¹, Sara Ghafarpour³, Mohamad reza Haghshenas¹, Faride Jowkar³, Nasrollah Erfani^{1,2}

1. Shiraz Institute for Cancer Research, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

2. Departments of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

3. Department of Dermatology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

Background: The CC-motif ligand 22 (*CCL22*) is a chemokine with crucial role in tumor-associated immunosuppression. *CCL22* is constitutively produced by macrophages, monocytes and dendritic cells and is inducible in many other immune cells such as Th2 and B cells. *CCL22* production has also been reported by several types of tumor cells. This chemokine is a functional ligand for *CCR4* which is commonly expressed on Th2 and Treg cells, and preferably recruits them to tumor microenvironment. The aim of this study was to investigate the association of 16C/A genetic variation (rs4359426) in *CCL22* gene with susceptibility of Iranian patients with Basal Cell Carcinoma (BCC).

Methods: 154 pathologically confirmed BCC patients (mean age: 60.67±11.57) and 154 age-sex-matched healthy individuals were recruited. Genomic DNA was extracted from white blood cells, and genotyping was performed using Polymerase Chain Reaction following by Restriction Fragment Length Polymorphism (PCR-RFLP) method.

Result: Arlequin analysis showed no deviation of genotype frequencies from Hardy-Weinberg equilibrium among both patients and controls. The frequencies of genotypes among patients and controls were 85.7% vs. 81.8%, 13.6% vs. 16.9%, 0.6% vs. 1.3% for CC, CA, AA genotypes respectively. Statistical analysis revealed no significant differences in the frequencies of genotypes and alleles between patients and controls ($P>0.05$). Although statistical analysis revealed no significant differences in the frequencies of genotypes and alleles between two groups, the A allele (CA or AA) significantly decreased in male patients comparing to male controls ($P_{\text{value}}=0.03$). No association was also observed between these genotypes and clinicopathological parameters in the patients ($P>0.05$).

Conclusion: Results of this investigation do not support the association of 16 C/A in *CCL22* gene with susceptibility and progression of BCC in Iranian patients. However, investigation of other *CCL22* gene variations and haplotypes may be required to completely roll out the association of *CCL22* gene variants with BCC.

Keywords: Basal Cell Carcinoma, Genotypes, Haplotypes, CC Chemokine ligand 22 (*CCL22*).

ICIA2018\Cancer Immunology\Poster\8492
8492

Investigating the effect of interaction between IL-17A, IL-17E, TNF- α and rs3783605 on the activity of VCAM-1 promoter in the human umbilical vein endothelial cells (HUVECs)

Fatemeh Dadgar Pakdel¹, Javad Dadgar Pakdel³, Mohammad Keramatipour^{*2}, Mohammad Vodjgani^{*1}

1. Department of Medical Immunology, School of Medicine, Tehran University of Medical Sciences (TUMS), Tehran, Iran

2. Department of Medical Genetics, School of Medicine, Tehran University of Medical Sciences (TUMS), Tehran, Iran

3. Department of pharmaceutical biotechnology, Pasteur Institute, Tehran, Iran

Background: Vascular Cell Adhesion Molecule-1 has a key role in the cell adhesion and transendothelial movement. Different factors such as gene promoter and its single nucleotide polymorphisms affect the VCAM-1 expression. The previous reports proved an important role of rs378605 in -420 position in the pathogenesis of diseases connecting with VCAM-1. Present study was designed to investigate the effect of rs3783605 and its interaction with IL-17A, IL-17E and TNF- α cytokines on the activity of VCAM-1 gene promoter in HUVEC.

Methods : Two vectors with different alleles of rs3783605 were constructed to express the GFP. GFP expression level was measured by real-time PCR.

Result : Results showed that -420 G led to about 2-fold higher transcriptional activity compared with the A allele. HUVECs stimulated with mentioned cytokines alone showed a decrease of expression. The stimulation with IL-17A in addition to TNF- α showed an increase of expression in G allele vector and the stimulation with IL-17E in addition to TNF- α led to an increased expression in the cells containing A allele. The stimulation by IL-17A in addition to IL-17E led to decrease of expression in the cells containing G allele and stimulation with three cytokines simultaneously showed a decrease of expression in G allele vector.

Conclusion: In general, the results can be interpreted in three states of stimulation with one, two and three cytokines and on that basis, the results are different. Data from this study can be useful in the treatment of diseases in the future.

Keywords: SNP; promoter; VCAM1; HUVEC; IL-17E

ICIA2018\Cancer Immunology\Poster\8546
8546

Expression of the level of CD44 protein in various breast cancer cell lines

Fatemeh Vahidian¹, Elham Safarzadeh¹, Behzad Baradarn¹

1. Department of Immunology research center, Faculty of medicine, Tabriz university of medical sciences, Iran

Background: Breast cancer is the most common cancer diagnosed in women. Breast cancer can occur in both men and women but it's far more common in women. This cancer is a heterogeneous disease and can be divided into in situ and invasion modes and the most common cause of death in this cancer is metastatic. CD44 is a superficial cellular glycoprotein that plays an important role in the cell-matrix and cell-cell-interaction of the migration and invasion of cancer cells and also is a medium to cell environment, which helps with the survival and migration of cancer cells such as breast cancer. Also the expression of CD44 increases in cancer cells and plays an important role in helping the growth and metastasis of breast cancer cells, and can serve as a therapeutic goal. Interestingly, CD44, as a key marker of cancer on the surface of cancer cells in various malignancies, includes breast cancer.

Methods: Breast cancer has different grades and cell lines. In this study we examined the expression of CD44 in four cell lines including MDA-MB231, MDA-MB468, MCF-7 and SKBR3. We cultured each of these cells and then extracted their RNAs and transferred to their related cDNAs then, we did real time PCR on them.

Results; By analyzing the data, we noticed that the highest expression of CD44 was in turn in MDA-MB468, SKBR3 and MDA-MB231 which are metastatic cell lines. Further, we recognized that the expression of this protein was very low in MCF-7 which is a non-metastatic cell line.

Conclusion: We concluded that the expression of CD44 in metastatic cell lines of breast cancer is more than non-metastatic cell lines, and more importantly, the expression of this protein in MDA-MB468, which is more metastatic than the others, was more.

Keywords: breast cancer, CD44, cell lines, metastasis

ICIA2018\Cancer Immunology\Poster\8550
8550

Expression of the inhibitory molecule, Tim-3 on the cells of the breast cancer draining lymph nodes

Fereshteh Mehdipour¹, Atri Ghods^{1,2}, Sahar Shariati^{1,3}, Abdol Rasoul Talei⁴, Abbas Ghaderi^{1, 2}

1. Shiraz Institute for Cancer Research, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

2. Department of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

3. Department of Immunology, School of Medicine, ShahidSadoughi University of Medical Sciences, Yazd, Iran

4. Breast Diseases Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

Background: T cell immunoglobulin and mucin domain-3 (Tim-3) is expressed on different cell types including T cells, dendritic cells, macrophages and natural killer cells. Tim-3 has been suggested to act as an inhibitory molecule and have a negative impact on anti-tumor immunity due to its expression on exhausted or regulatory T cell. The aim of this study was to determine the frequency of Tim-3⁺ lymphocytes and CD8⁺T cells in the tumor draining lymph nodes (TDLNs) of breast cancer patients and its relation with disease parameters.

Methods: Axillary lymph node specimens were taken from 22 breast cancer patients and mononuclear cells were isolated using Ficoll-Hypaque gradient centrifugation. Cells were then stained for CD8 and Tim-3 and subjected to flow cytometry.

Results: The percentages of Tim-3⁺ cells were determined in lymphocytes and CD8⁺ T cells respectively. Our results showed that 5.8±2.4 of lymphocytes and 6.1±2.8 of CD8⁺ T cells expressed Tim-3 without significant differences in the metastatic and non-metastatic lymph nodes, however the geometric Mean Fluorescent Intensity (gMFI) of Tim-3 expression was significantly higher in lymphocytes of the metastatic lymph nodes (P=0.033). The percentage of Tim-3⁺ lymphocytes were significantly higher in the TDLNs of ER- compared to ER+ patients (P=0.047). A similar non-significant trend was seen in the frequency of TIM-3⁺CD8⁺ T cells. Moreover, the frequency of Tim-3⁺ lymphocytes and CD8⁺ T cells showed significant and non-significant increases in the TDLNs of patients with grade III as compared with grade II (P=0.020 and P=0.10, respectively). The frequency of Tim-3⁺ lymphocytes or CD8⁺ T cells did not show significant associations with tumor size or stage of the disease.

Conclusion: Our results suggested a link between higher frequency of Tim-3⁺ lymphocytes and poor prognosticators like ER- and higher tumor grade. Further functional and phenotypical analysis of this lymphocyte subset is warranted.

Keywords: Tim-3, Lymphocyte, CD8⁺ T cells, Tumor draining lymph nodes, Breast cancer

ICIA2018\Cancer Immunology\Poster\8563

8563

The Increased level of Osteopontin and its receptor in patients with benign and malignant bone tumorAmeinh Hosseini¹, Alireza Mirzaei², Khodamorad Jamshidi², Masoumeh Tavakoli Yaraki¹*1. Department of Biochemistry, School of Medicine, Iran University of Medical Sciences, Tehran, Iran**2. Bone and Joint Reconstruction Research Center, Shafa Orthopedic Hospital, Iran University of Medical Sciences, Tehran, Iran*

Background and aim: Osteopontin has emerged as an active player in many physiological and pathological processes, including biomineralization, tissue remodeling and inflammation. As an extracellular matrix protein and proinflammatory cytokine, osteopontin is thought to facilitate the recruitment of monocytes/macrophages and to mediate cytokine secretion in leukocytes. Because OPN acts through several receptor mechanisms including both integrins and CD44, targeting these receptor ligand interactions as already under investigation in cancer therapy. The aim of the present study was to investigate the expression level of OPN and CD44 in tumor tissue of patients with benign and malignant bone tumor comparing to normal bone tissue.

Materials and Methods: 50 patients of Shafa Orthopedic Hospital in Tehran with benign and malignant bone tumor have participated in this case-control study. The tumor tissues and margins were used for mRNA extraction and cDNA construction, and to determine the expression level of CD44 gene, a Real-Time PCR-based Cyber Green method was used and data were analyzed using $\Delta\Delta CT$ method. The level of circulating OPN was determined using ELISA. Finally, statistical analysis was performed using Graph Pad Prism software version 5 and independent t-test.

Results: Measurement of CD44 expression level in tumor tissues of patients with bone tumor revealed that the level of this gene was significantly increased in patients comparing to healthy subjects and normal tissues. Also, the increased level of CD44 was associated with elevated level of tumor grade and stage ($p < 0.05$). The plasma level of OPN was increased in patients with bone tumor comparing to healthy controls which was correlated with severity of disease.

Conclusion: The results of the current study have shown that the OPN and its receptor (CD44) can account as a circulating also local bio marker in patients with bone tumor due to the significant differences of the CD44 and OPN level in patients comparing to controls, it can be noticed as a possible biomarker for controlling disease.

Keywords: OPN, CD44, gene expression, Bone Tumor.

ICIA2018\Cancer Immunology\Poster\8601
8601

Evaluation of tissue expression of BTLA and HVEM molecules and soluble form of HVEM in gastric cancerpatients

Maryam Azarafza¹, Mohsen Tehrani¹, Reza Valadan¹, Iradj Maleki², Omid Emadian³, Abolghasem Ajami¹

1. *Department of Immunology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran*

2. *Department of Gastroenterology, Gut and Liver Research Center, Mazandaran University of Medical Sciences, Sari, Iran*

3. *Department of Pathology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran*

Background: Gastric cancer is the second leading cause of cancer death and the fourth most commonly diagnosed cancer worldwide. Identifying and subsequently targeting pathways by which tumors inhibit activity and cytotoxicity of immune cells has been regarded as a novel strategy to improve clinical approaches and survival outcome. The BTLA/HVEM pathway is one of the inhibitory mechanisms, playing an important role during the inflammation and tumorigenesis and has been recently recognized as a tumor escape strategy. There is limited information regarding precise level of BTLA and HVEM expression in gastric cancer. The aim of this study was to investigate the expression of BTLA, HVEM and soluble form of HVEM in gastric cancer specimens and serum, respectively.

Methods: The expression level of BTLA and HVEM was evaluated by immunohistochemistry and mRNA expression was examined using real-time PCR in gastric tissue specimen of gastric cancer, metaplasia and control groups. sHVEM in serum was assessed by ELISA assay kit.

Results: Our results indicated that the expression of BTLA, HVEM in tissue and sHVEM in serum of patients with gastric cancer and metaplasia were high compared to control groups. The increased expression of BTLA, HVEM and serum level of sHVEM was significantly correlated with poor prognosis of gastric cancer and clinicopathological features such as TNM staging and lymph node metastasis.

Conclusions: Our data indicated that increase of BTLA and HVEM in patients with gastric cancer may have correlated to tumor progression and might be considered as an independent prognostic factor. Altogether, this inhibitory pathway could be a promising candidate for immunotherapy of gastric cancer.

Keyword: Gastric cancer, B and T lymphocyte attenuator, Herpesvirus entry mediator

ICIA2018\Cancer Immunology\Poster\8606
8606

+49A/G (rs231775) and -318C/T (rs5742909) Polymorphisms of *CTLA-4* Gene Showed No Association with Patients with Prostate Cancer

Mahyar Malekzadeh¹, Parvin Nikaein¹, Ali Ariaifar², Abbas Ghaderi¹

1. Institute for Cancer Research (ICR), Shiraz University of Medical Sciences, Shiraz, Iran

2. Department of Urology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Prostate cancer is the fourth most common cancer in both sexes and the second most common cancer in men which lethality is normally associated with occurring the metastasis rather than primary tumor. Therefore, targeting the molecular pathways that promote tumor formation and progression could assist better prognosis. CTLA-4 is a protein receptor that functions as an immune checkpoint which its blockage is regarded as an effective mechanism in treating many cancer types.

Methods: 5ml peripheral blood from 100 prostate cancer patients and 100 age matched normal men were taken. DNA extraction was done by Proteinase K method. PCR-RFLP and AS-PCR were performed to determine the genotypes of +49A/G (rs231775) and -318C/T (rs5742909) sites, respectively.

Results: There was no significant correlation between these sites in patient and control groups. Moreover, no significant relation between clinicopathological findings either the mean age in patient group was found.

Conclusion: Our data suggested no association between *CTLA-4* gene variants at positions +49A/G(rs231775) and -318C/T(rs5742909) in patients with prostate cancer compared with healthy controls. To conclude the role of *CTLA-4* gene variants in prostate cancer, investigation of other known polymorphisms reported in *CTLA-4* gene and also reports from other ethnic populations are warranted.

Keywords: CTLA-4 Polymorphism, +49A/G, -318C/T, Prostate, Cancer

ICIA2018\Cancer Immunology\Poster\8619

8619

Investigation of CTLA-4 Gene +49A/G (rs231775) and -318C/T (rs5742909) Polymorphisms in Patients with Thyroid Cancers

Shabnam Abtahi¹, Fatemeh Izadi Jahromi¹, Mahyar Malekzadeh¹, Mohammad Javad Fattahi¹, AmirReza Dehghanian², Mohammad Hossein Dabbaghmanesh³, Abbas Ghaderi¹

1. Institute for Cancer Research (ICR), Shiraz University of Medical Sciences, Shiraz, Iran

2. Department of Pathology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

3. Department of Endocrinology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Cytotoxic T Lymphocyte Antigen 4 (CTLA-4 or CD152) is among the immune checkpoint molecules and its abnormal expression in genetically and environmentally cancer predisposed individuals might make the person more susceptible to disease initiation and progression. The rs5742909 (-318C/T) and rs231775 (+49A/G) are among the most commonly studied SNPs and have been considered as genetic factors related to thyroid diseases.

Methods: We aimed to elucidate the association between these SNPs and susceptibility to various types of thyroid cancers, by comparing the frequency distribution of the genotypes/alleles in these regions with a group of healthy individuals.

Results: A statistically significant association was observed between the presence of G allele in rs231775 locus and thyroid carcinoma, when comparing cases and controls (OR=2.4; CI: 1.54 to 3.78; P<0.001) and the frequency of heterozygotes (AG) was higher than homozygotes for allele A (AA), in thyroid cancer patients. However; we observed no difference among allele/genotype frequencies in regards to locus rs5742909.

Conclusion: In this study we observed that G allele in rs231775 and possibly lower expression and affinity of mutated CTLA-4 molecule, is associated with higher susceptibility to thyroid carcinoma, especially papillary subtype.

Keywords: CTLA-4, Polymorphism, rs231775, rs5742909, Thyroid, Neoplasms

ICIA2018\Cancer Immunology\Poster\8637
8637

Autophagy, using Atmospheric Cold Plasma (CAP) method, is an alternative therapeutic mechanism for the treatment of melanoma cancer

Mina Alimohammadi¹, Alireza Refeie¹, Monireh Golpour¹

1. Mazandaran University Of Medical Sciences , Mazandaran , Iran

Background : Autophagy is a pathway to tolerate stress that preserves survival under adverse conditions. However, long autophagy prevents the cell from coping with stress due to degradation of cellular structures, leads to autophagy cell death. ROS induction by starvation, involved in stimulating autophagy during the onset and progression of cancer. The new treatment with cold atmospheric plasma (CAP) promotes the production of ROS in the cell and induces oxidative stress. It leads to autophagy in the melanoma cells without affecting normal cells.

Methods : In this study, by starving to induce autophagy in B16 melanoma cells and L929 in Invivo and Invitro, the CAP effectiveness measured compared to chemotherapy. Using the MTT assay, we showed the survival rate of CAP-treated cells in comparison with the chemotherapy group and the combination of both treatments significantly decreased.

Result : Also, with flowcytometry, we showed that the percentage of cell death in tumor cells in the CAP group was significantly higher than other groups. Also, by examining the mRNA expression of LC3 and ATG5 which involved in autophagic, CAP increased the expression of these proteins. Also, in Invivo study, with B16 inoculation in Balb/C mice, we observed that tumor size in the CAP-treated group was significantly decreased compared to the chemotherapy. Then, to confirm the selective CAP effect on the toxicity and metastasis and the cellular attachments of surrounding healthy tissues, by staining H&E on vital tissues such as the liver, kidneys, lymph nodes and spleen, has showed that CAP does not have any toxicity and deformity in the tissue And cellular connections, while other groups were strongly associated with serious injury, high intracellular toxicity, and loss of intercellular connections.

Conclusion: This study showed that CAP did not cause damage to the normal tissue and also caused a high percentage of cell death autophagic in cancer cells .so It can be a good alternative for the treatment of malignant melanoma

Keywords:Ros , CAP , Autophagy ,Melanoma

ICIA2018\Cancer Immunology\Poster\8645
8645

Down-regulation of E-Cadherin in Acute Lymphoblastic Leukemia

Mohammad Mahdi Mohammadi¹, Amin Mirzaeian¹, Shahrbanoo Rostami², Fatemeh Mirzaeyan³

1. Department of Immunology, Faculty of Medicine, Kerman University of Medical Sciences, Kerman, Iran

2. Department of Hematology-Oncology and Stem Cell Transplantation, Tehran University of Medical Sciences, Tehran, Iran

3. Department of Hematology and Blood Banking, Faculty of Allied Medicine, Tehran University of Medical Sciences, Tehran, Iran

Background: E-cadherin is one of the adhesion molecules which contribute to the interaction of haematopoietic progenitors with their niche. While the pivotal role of the loss of E-cadherin expression in tumourigenesis has been well characterized, little is known about its expression level in Acute Lymphoblastic Leukemia.

Methods: RT-qPCR was carried out on bone marrow aspirates from 89 ALL patients at the time of diagnosis and 33 age-matched healthy individuals. The ALL patients were categorized into B-ALL (69.7%) and T-ALL (30.3%), and the B-ALL cases were further classified into pro-B, early pre-B, pre-B and immature/mature-B based on immunophenotypic characteristics.

Results: Compared with controls, E-cadherin was significantly downregulated in ALL ($p < 0.001$). There were no correlation between ALL subtypes or leukocyte count with gene expression level of E-cadherin. However, there was significantly negative correlation between E-cadherin expression and the number of bone marrow blasts as a disease-related prognostic factor ($r = -0.26$, $p = 0.01$) in ALL Patients.

Conclusion: Our results showed decreased expression of E-Cadherin in ALL patients. Since alterations of adhesion behaviour of haematopoietic stem cells may alter the homeostasis of haematopoietic cells within bone marrow microenvironment and ultimately lead to transforming ability and vigorous growth of leukaemic cells, we suggest that E-Cadherin signaling pathway should be further explored to reveal disease biology and potential therapeutic targets.

Keywords: Acute Lymphoblastic Leukemia, E-cadherin, RT-qPCR.

ICIA2018\Cancer Immunology\Poster\8669
8669

SNP variation of immune system gene in medulloblastoma

FatemeShaabanpour Aghamaleki¹, Shirin Farivar²

1. Department of molecular and cellular Biology School of Biotechnology and Life Science University of ShahidBeheshti

2. Department of molecular and cellular Biology School of Biotechnology and Life Science University of ShahidBeheshti

Background: medulloblastomas are the most frequently diagnosed embryonal tumors of the central nervous system. Immunotherapy might be key to improve survival and to avoid morbidity. The efficient killing of tumor cells using immunotherapy requires overcoming cancer-associated strategies to evade cytotoxic immune responses. Single nucleotide polymorphisms have the most effect on immune genes in medulloblastomas that can be used as a specific target in immunotherapy of medulloblastomas.

Methods: Genome variation profiling by SNP array of 7 primary medulloblastomas and 10 metastatic medulloblastomas patients was extracted from Geo datasets. SNPs are compared with logfc for their frequency between two groups and sorted by their p-value. Also, the related gene for an SNP, the most related pathway, location, function, and protein networks of this SNPs were identified with NCBI and GeneCard databases.

Results: several SNPs were identified for medulloblastomas that may associate with it. Some these SNPs related to an immune system and have a high frequency in metastatic medulloblastoma. One of these SNPs were in PTPRS gene (rs886932). The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. rs886932 is an intron variant with C/T polymorphism. This SNP has a high frequency in metastatic medulloblastoma.

Conclusion: identification of critical SNP of immune system gene in two kinds of medulloblastoma patients can lead to appropriate and specific chemotherapy and immunotherapy by targeting specific SNP.

Keywords: medulloblastoma, SNPs, PTPRS gene, cancer immunology and immunotherapy

ICIA2018\Cancer Immunology\Poster\8679
8679

The effects of *Aloe vera* leaf exudate on the proliferation of K562 human tumor cell line

Hamid Shamsi¹, Payman Tavakoli¹, Minoo Shaddel², Meysam Abtahi Foroushani³, RoyaShamsi⁴

1. Department of parasitology University of Urmia

2. Department of parasitology Faculty of Medicine University of Aja University

3. Department of Microbiology Faculty of Veterinary University of Urmia

4. Department of Clinical Nutrition and Dietetics, Faculty of Nutrition and Food Technology, ShahidBeheshti University of Medical Sciences, Tehran, Iran.

Background: Leukemias are a group of blood malignancies which are classified as lymphoid and myeloid based on the origin of the malignant cells. K562 is a leukemic cell line which was first isolated from a 53 years old woman suffering from chronic leukemia. The present study was carried out to investigate the probable inhibitory effect of *Aloe Vera* on the proliferation of K562 cell line.

Methods: The inner parts of the plant including the gelatinous substance were emptied and the outer layer was used for the preparation of the lyophilized extract. The extract was dissolved in DMSO to be used in all experiments. The leukemic cells were cultured in RPMI 1680 medium containing 10% Fetal Bovine Serum (FBS) and were then treated with increasing doses of the extract. The viability of the cells was studied using the MTT test. Peripheral Blood Mononuclear Cells (PBMC) were isolated from the peripheral blood of normal donors using ficoll-hypaque and were subsequently treated with the same doses of the extract. After the incubation time, the IC50 was determined for both the K562 and normal PBMCs using the MTT assay. Doxorubicin and Phosphate-buffered Saline (PBS) were used as the positive and negative controls, respectively. The results were analysed in SPSS software using One-way ANOVA.

Results: Statistical analysis of the obtained data showed that the viability of cells in both the leukemic cell line and normal PBMCs decreased following 24, 48, and 72 hours treatment with the extract in a dose-dependent manner.

Conclusion: The results of the present study showed that *Aloe Vera* outer layer extract significantly decreases the number of viable cells in both K562 cell line and normal PBMCs with less selectivity in comparison to the conventional anti-leukemic doxorubicin.

Keywords: *Aloe Vera* leaf exudate; K562; MTT

ICIA2018\Cancer Immunology\Poster\8681

8681

Expression analysis of activatory and inhibitory receptors of NK cells in patients with chronic lymphocytic leukemia

Leila Hadadi¹, Ali Akbar Amirzargar², Ramazan Ali Sharifian³, Saied Abediankenari^{1,4}, Hossein Asgarian Omran^{1,4*}

1. Department of Immunology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

2. Department of Immunology, Faculty of Medicine, Tehran University of Medical Sciences, Tehran, Iran

3. Clinic of Hematology and Oncology, Vali-Asr Hospital, Faculty of Medicine, Tehran University of Medical Sciences, Tehran, Iran

4. Immunogenetics Research Center, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

Background: Natural killer (NK) cells have pivotal roles in both innate and adaptive immune responses to many inflammatory and chronic infectious diseases. Initial activation and function of NK cells are mediated by signals received from multiple activating and inhibitory receptors. NK cell dysfunction and imbalance in these regulatory receptors have been reported in various solid and hematologic malignancies. In this study, the expression pattern of NKp30 and T cell immunoglobulin and mucin domain containing molecule-3 (Tim-3), as candidates of activating and inhibitory receptors, were evaluated in patients with chronic lymphocytic leukemia (CLL).

Methods: Twenty five untreated CLL patients and fifteen sex- and age-matched healthy controls were enrolled in this study. Fresh peripheral blood mononuclear cells were collected from all subjects. CD56⁺/CD3⁻ cells in lymphoid region were defined as NK cells and the frequency of CD56⁺/CD3⁻/NKp30⁺ and CD56⁺/CD3⁻/Tim-3⁺ cells was determined by three-color flow cytometry method.

Results: Our Results revealed that Tim-3 expression, as an inhibitory receptor, was significantly increased on NK cells of CLL patients in comparison to normal healthy controls. NK cells from CLL patients showed lower expression of NKp30 activatory receptor compared to controls, but the difference was not statistically significant.

Conclusion: Altogether our results confirm that NK cells in CLL patients show dys-regulation in their activatory and inhibitory receptors and may be functionally exhausted.

Keywords: Chronic lymphocytic leukemia, NK cell, Tim-3, NKp30

ICIA2018\Cancer Immunology\Poster\8693

8693

Relationship between helicobacter pylori and TIMP-1 in gastric adenocarcinoma

Rana Ezzeddini¹, Mohammad Taghikhani¹, Mohammad Hossein Somi², Amir Salek Farrokhi³,
Amirtaher Eftekharrosadat⁴

1. Department of Clinical biochemistry, Faculty of Medical Sciences, TarbiatModares University, Tehran, Iran

2. Liver and Gastrointestinal Diseases Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

3. Department of Immunology, Faculty of Medical Sciences, TarbiatModares University, Tehran, Iran

4. Department of Clinical biochemistry, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Various proteins can be found in adjustment of inflammatory immune response in H. pylori (H.P)-infected in gastric mucosa. Tissue inhibitor matrix metalloproteinase 1 (TIMP-1) plays a vital role in carcinogenesis and is over-expressed in many human malignancies. Although, the value of TIMP-1 is debated as a biomarker in gastric cancer (GC) patients. Overexpression and therefore, increased serum level of TIMP-1 can be related to the pathogenesis of H.pylori-associated stomach adenocarcinoma. The objective of this study is to investigate the association between H.pylori with TIMP-1 in adenocarcinoma cases.

Methods: In order to carry out this experiment, 85 gastric adenocarcinoma patients and 130 healthy controls selected. Blood samples collected and TIMP-1 concentration determined by ELISA method. Also, Crystal violet fast (CVF) staining performed for H.pylori detection on endoscopic biopsies. We used T-test for data analysis and P values less than 0.05 considered statistically significant.

Results: Our findings indicated significant increased level of TIMP-1 in gastric adenocarcinoma patients comparing control subjects ($P \leq 0.05$). Also, our results showed that 39% of gastric adenocarcinoma patients were positive H.pylori. In addition, we demonstrated that TIMP-1 concentration significantly up regulated in H.pylori positive adenocarcinoma cases compared to negative patients.

Conclusion: Altogether, the present survey showed that increased level of TIMP-1 in gastric adenocarcinoma has a correlation with involving H.P in stomach

Keywords: Gastric adenocarcinoma, TIMP-1, Helicobacter pylori



ICIA2018\Cancer Immunology\Poster\8712
8712

Correlation among Age and Genetics and Regulatory T cells in Chronic Myeloid Leukemia Patients

Zahra Shaker¹, Zohair Mohammad Hasan², Davar Amani³, Masume Ebtekar²

1. *AJA University of Medical Sciences, TarbiatModares University, ShahidBeheshti University of Medical Sciences*

2. *TarbiatModares University*

3. *ShahidBeheshti University of Medical Sciences*

Background: Chronic myelogenous leukemia(CML, Chronic Myeloid Leukemia)Can be created at any age, but most adults over age 50 have the disease, and these patients about 70% of patients are included.Hmchnbn the disease occurs in men than in women.Prevalence in children is rare and is reported almost 4 years.Due to the large variable called way of life that is influenced by climate (air, water and soil), economy, customs, traditions and religious beliefs are different age.

Method: This is a historical cohort study of 15 patientsCMLFor a period of two years, a peripheral blood sample was taken and testedRT-PCR in real time, To determine the presence of Philadelphia gene and flow cytometerFACS CaliburTo assess the number of cellsTSetting was used .Information obtained by the test-GEE;Generalized Estimation EquationWere analyzed .

Result:Some of the results show that when the oncoproteinbcr / ablThe amount of cells has increasedTThe regulation for each year increases the age of the patient by 0.729 , which is significant($P_v = 0.000$, $P_v < 0.05$). In this study, it was determined that genetic aging also increased

Conclusion: Has a relationship .These findingsDThe prescribed medications and follow up of the treatment, as noted by the experts is very effective.

Keywords:CML,Treg,bcrlabl, Age

ICIA2018\Cancer Immunology\Poster\9851
9851

Increased CD8⁺/PD-1⁺/TIGIT⁺ T-cells in Patients with chronic lymphocytic leukemia (CLL)

Reza Hajiasghar Sharbaf¹, Hossein Asgarian Omran^{1,2}, Hadi Hossein Nattaj¹, Ramin Shekariz³,
Mohsen Tehrani^{1,3}

1. Department of Immunology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran.

2. Immunogenetic Research Center, Mazandaran University of Medical Sciences, Sari, Iran.

3. Gastrointestinal Cancer Research Center, Mazandaran University of Medical Sciences, Sari, Iran.

Background: Tumors can escape immune response via upregulation of immune-checkpoint receptors, such as CTLA-4, PD-1, Tim-3, LAG-3, and TIGIT, leading T-cells to a state of exhaustion. In this study, the frequency of CD8⁺/PD-1⁺/TIGIT⁺ T-cells, as exhausted T-cells, was measured in patients with CLL.

Methods: PBMCs were isolated from blood samples from CLL patients and healthy controls. Cells were cultured in the presence of phytohemagglutinin (10 µg/mL) for 72 h. Frequency of CD8⁺/PD-1⁺/TIGIT⁺ T-cells was then measured using three-colored flow cytometry.

Results: Frequency of CD8⁺/PD-1⁺/TIGIT⁺ T-cells were higher in patients with CLL (mean, 18.45) than in healthy controls (mean, 7.76 %).

Conclusions: Our data indicate that a high number of CD8⁺ T-cells in CLL patients express immune checkpoint receptors, PD-1 and TIGIT. Therefore, reversal of these inhibitory pathways could be a promising therapeutic approach for CLL.

Keywords: Exhausted T-Cell, PD-1, TIGIT, chronic lymphocytic leukemia (CLL)

ICIA2018\Cancer Immunology\Poster\9882

9882

Melphalan cytotoxicity increased in combination with Arsenic trioxide in MT-2 lymphoma cells

Faeze Khodadadi¹, Mohammad Hadi Sadeghian^{1*}, Houshang Rafatpanah², Fatemeh B. Rassouli³,
Mehrdad Iranshahi⁴

1. Cancer Molecular Pathology Research Center, Department of Hematology and Blood Bank, Faculty of Medicine, Mashhad University of Medical Sciences, Iran

2. Inflammation and Inflammatory Diseases Research Center, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

3. Cell and Molecular Biotechnology Research Group, Institute of Biotechnology, Ferdowsi University of Mashhad, Mashhad, Iran

4. Department of Pharmacognosy and Biotechnology, Biotechnology Research Center, Faculty of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

Background : Adult T cell leukemia lymphoma (ATLL) is a CD4+ lymphoproliferative malignancy resulting from human T-cell leukemia virus type 1 (HTLV1) infection. Iran is known as one of the endemic regions for HTLV-1. Despite advances in treatment of ATLL, the average survival rate of patients is low. Melphalan is one of the alkylating agents, the first types of drugs to be used in chemotherapy of malignant disease, are still an important group of drugs in clinical cancer chemotherapy. To introduce more effective anticancer strategies for the treatment of ATLL, here we investigate whether melphalan could enhance the efficacy of arsenic trioxide, a chemical drug prescribed for ATLL.

Methods: In this regard, MT-2 cells, an ATLL cell line, were treated with increasing concentrations of melphalan and Arsenic for 24, 48 and 72 hours, and the IC50 value of drugs was determined. Next, cells were treated with combination of melphalan (0.5 µg/ml) and Arsenic (1 and 2 µM). Then, cell viability was evaluated using Alamar Blue 14 g% reagent.

Results: Obtained results indicated that 0.5 µg/ml melphalan increased the toxicity of Arsenic 2 µM, and thus, had synergic effects with this anticancer drug.

Conclusion: The results of the present study indicated that combinatorial use of melphalan and arsenic trioxide can be considered in new therapeutic approaches to ATLL.

Keywords: melphalan, Adult T-cell leukemia/lymphoma, Arsenic trioxide, HTLV-1

ICIA2018\Cancer Immunology\Poster\9883

9883

Effects of melphalan on the expression of NF- κ B in human lymphoma cells

Faeze Khodadadi¹, Mohammad Hadi Sadeghian^{1*}, Houshang Rafatpanah², Fatemeh B. Rassouli³, Mehrdad Iranshahi⁴

1. Cancer Molecular Pathology Research Center, Department of Hematology and Blood Bank, Faculty of Medicine, Mashhad University of Medical Sciences, Iran

2. Inflammation and Inflammatory Diseases Research Center, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

3. Cell and Molecular Biotechnology Research Group, Institute of Biotechnology, Ferdowsi University of Mashhad, Mashhad, Iran

4. Department of Pharmacognosy and Biotechnology, Biotechnology Research Center, Faculty of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

Background : Melphalan is a widely used classical chemotherapeutic agent in the group of alkylating nitrogen mustard agents that was developed more than 50 years ago, and substantial clinical experience has been accumulated. Adult T-cell leukemia/lymphoma (ATLL) is an aggressive T-cell malignancy caused by human T-cell leukemia virus type 1 (HTLV-1) infection and often occurs in HTLV-1 endemic areas. Iran especially Khorasan province is known as endemic regions for this virus. In spite of medical advances, the prognosis of ATLL is still very poor. HTLV-1 Tax oncogene activates NF- κ B signaling pathway. Herein we studied the effects of melphalan on the expression of NF- κ B in ATLL cells.

Methods: In this study, we showed the melphalan effects on the expression of NF κ B (REL-A) in HTLV-1 infected cell line. Accordingly, MT-2 cells were treated with 0.5 μ g/ml melphalan for 72 hours, while control cells were treated with DMSO 0.4% (used as melphalan solvent). Next, total cellular RNA was extracted and treated with DNase I and then reverse transcribed into cDNA. For confirmation of cDNA fidelity PCR using GAPDH primers was conducted. Further Real-time PCR was performed using Taq man probe and specific primers for NF- κ B (REL-A).

Results: Compared to relevant control, MT-2 cells treated with melphalan (in non-toxic concentration) exhibited significant down regulated levels of NF- κ B (REL-A) ($p < 0.05$).

Conclusion: These *in vitro* results suggest that the alkylating agent melphalan could be considered as an effective agent to alleviate malignant properties of ATLL cells in future *in vivo* studies.

Keywords: melphalan, Adult T-cell leukemia/lymphoma, HTLV-1, NF- κ B

ICIA2018\Cancer Immunology\Poster\9900
 9900

Interaction of Perforin and Granzyme Band HTLV-1 Viral factors Is Associated with ATLL development

Mohammad Mehdi Akbarin¹, Sadegh Farhadi², Abolghasem Allahyari³, Mohammad Mehdi Koshayar³, Abbas Shirdel², SeyedAbdolrahim Rezaee¹, Zahra Mozaheb³, Asadollah Mohamadi¹, Alireza bari², SeyedehTahereh Mohaddes², Hossein Rahimi², Houshang Rafatpanah¹

1. Inflammation and Inflammatory Diseases Research Centre, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran
2. Hematology Department, Ghaem Hospital, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran
3. Hematology Department, Imam Reza Hospital, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

Background: Human T lymphotropic virus (HTLV-1) is a human retrovirus which is associated with adult T cell leukaemia (ATLL). ATLL is a malignancy lymphoproliferative disease which infects CD4 T cells. It is not clear why the majority of HTLV-1 infected individual's remains healthy carries and minority develop ATLL. Cellular immune response has a critical role in ATLL and destroys malignant and HTLV-1-infected cells. Perforin and granzyme have important functional role in apoptosis and destruction of infected cells. In the present study we examined the role of perforin and granzyme in ATLL patients and HTLV-1 asymptomatic carries (HAC).

Method: Peripheral blood mononuclear cells (PBMCs) were isolated from ATLL patients and asymptomatic carriers by using Ficoll-hypopaque density centrifugation. RNA was extracted and cDNA was synthesised. A real-time PCR TaqMan method was designed and optimized for evaluation of perforin, granzyme, Tax and HBZ gene expression. HTLV-1 proviral load was quantified in patients with ATLL and carriers.

Results: The mRNA expression of TAX and HBZ was significantly higher in ATLL patients than HACs ($p=0.011$) and the HTLV-1 proviral load was higher in ATLL patients compared to HAC group ($p=$). There was a significant increase in perforin gene expression in HACs than ATLL patients ($p=0.002$). Furthermore the expression of granzyme was also higher in HTLV-1 carries compared with ATLL patients and significant differences was observed between two groups ($p=0.036$).

Conclusion: Low expression of perforin and granzyme in ATLL patients seem influences the efficiency of CTL function and destruction of HTLV-1-infected cells which might contribute to the disease pathogenesis.

Keywords: HTLV-1, ATLL, perforin, granzyme, proviral load

ICIA2018\Cancer Immunology\Poster\9947
9947

The survey of Interleukin-35 gene expression in tissue and blood of colon cancer

Milad Ameri¹, Vedad Alkhamis²

1. Department of Pathobiology, University of Bu Ali Sina Hamedan

2. Department of Microbiology University of Shahid Chamran Ahvaz

Background: Colon cancer is the third most common cancer Worldwide. Molecular methods such as Real Time PCR have been used to determine the expression of IL-35 as a screening test for the detection of colon cancer or diagnosis in different stages. The primary purpose of this study was to investigate the expression of IL-35 (IL-35) as a suppressor of activity and increase apoptotic sensitivity in cancer cells in the intestinal cancer tissue.

Methods: Due to certain limitations, considering the nature of the IL-35 which have heterodimers of both IL-12 and EBI-3 components, this study was conducted on IL-12 α as a marker for the presence and expression of IL-35 In the tissue and peripheral blood of patients with colon cancer. In this study, 20 samples of paraffinaized tissue and 20 blood samples from people with colon cancer and 10 healthy samples were collected from the Khomeini hospital in Tehran. After de-paraffinization, RNA extraction was performed with RNX Plus solution. The cDNA was synthesized by reverse transcription using the MMULV enzyme. Gene expression was measured by the Real time PCR method. Glyceraldehyde phosphate dehydrogenase gene (GAPDH) was also used as internal control.

Results: Results showed that only 3 samples with an expression rate (RQ) greater than 1 out of 43 samples, 3 of samples were normal and used to verify the accuracy of the tests, suggesting a decrease in IL 12-R gene expression in most of the samples.

Conclusion: Due to staging of the disease, it can be concluded that with the decreased IL-12R expression gene, the colon cancer has progressed.

Keywords: Intestinal Cancer, Interleukin 35 Gene, Real Time PCR

ICIA2018\Cancer Immunology\Poster\10054

10054

Highly Reliable Simultaneous Biosensing of CA125 and CA15-3 Tumor Markers and selective tracking and imaging over MCF-7 and OVCAR-3 cells lines via FRET phenomenon based on CDs-MnO₂ heterojunction in single Excitation WavelengthSomayeh Hamd Ghadareh¹, Abdollah Salimi^{1,2*}, Fardin Fathi^c, Sara Parsa¹

1. Department of Chemistry, University of Kurdistan, 66177-15175, Sanandaj, Iran

2. Research Center for Nanotechnology, University of Kurdistan, Sanandaj, Iran

3. Cellular and Molecular Research Center, Kurdistan University of Medical Sciences, Sanandaj, Iran

Background: A fluorescence resonance energy transfer (FRET) strategy designed for simultaneous detection of two tumor markers based on coupling carboxylic acid functionalized carbon dots (CDs-COOH) and Rhodamine B (RB) conjugating antibodies. The strong green fluorescent-CDs were synthesized by a simple one-pot hydrothermal method. Monodispersed MnO₂ nanosheets were functionalized with ssDNA related aptamers and applied as quencher. More importantly, the solution containing both of the CDs and RB conjugating to CA125 as Ab1 and CA15-3 as Ab2 (CD1-Ab1, RB-Ab2) could be excited by single wavelength excitation of 490 nm. Inspired by these facts, we fabricated a selective and sensitive fluorescence nanosensor for simultaneous targeting of tumor markers.

Method: In the absence of CA125 and CA15-3 target antigens the fluorescence of the antibody labeled CD1 and RB, quenched with MnO₂ due to energy transfer (FRET) process, while with addition of target antigens, the formation of immunocomplex between CDs-Ab probe, antigen and ssDNA leading to the release of MnO₂, resulting in the distinct changes in fluorescence response. The response of the architecture nanoassembly enhanced gradually with the increase in concentration of targets in the range of 80 fg/mL to 5 pg/mL with 15.0 fg/mL as calculated limit of detection for CA125, and the response was found to be linear in the concentration range of 1.0 to 25 pg/mL for CA15-3 and the calculated limit of detection was 0.7 pg/mL. This strategy revealed a stable and good analytical performance for the detection of OVCAR-3 and MCF-7 cells lines ranging from 1000 to 10000 cells/mL with detection limit 2-3 cells in 10 µL of injected sample.

Result: Finally, due to low cytotoxicity of the applied CDs, the CDs-antibody hybrids were used in selective tracking and imaging of the cancer cells over MCF-7 and OVCAR-3 cells lines. The selectivity and sensitivity of this fluorescent nanosensor over other markers were carefully investigated. The obtained results implying the potential applications of presented assay for pathological diagnosis, biomedicine research and management of cancer diseases.

Keywords: Fluorescence resonance energy transfer, carbon dots, MnO₂ nanosheets, Simultaneous detection, CA125, CA15-3, Bioimaging

ICIA2018\Cancer Immunology\Poster\10061

10061

Mesenchymal stem cells derived from stage II of human breast tumor tissue promote the proliferation of the PBMCs :A in vitro assay

Koushansineh sepehr¹, Alireza Razavi¹, Zuhair Mohammad Hassan², AbdolReza Fazel³,
Majid Mossahebi Mohammadi⁴, Seyed Mahmoud Hashemi⁵

1. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

2. Department of Immunology, Faculty of Medical Sciences, TarbiatModares University, Tehran, Iran

3. Cancer research center, Golestan University of Medical Sciences, Gorgan, Iran

4. Sarem Cell Research Center(SCRC),Sarem Women's Hospital,Tehran,Iran

5.Department of Immunology, School of Medicine, ShahidBeheshti University of Medical Sciences, Tehran, Iran

Background: Mesenchymal stem cells (MSCs), as a subpopulation of stroma cells found in the tumor microenvironment,are critical in promoting tumor progression and possess immunomodulatory properties. Therefore, the purpose of this study was to isolate MSCs from primary human breast tumor tissue, and to study the effect of soluble factor of MSC on the peripheral blood mononuclear cell (PBMC) proliferation.

Methods: The tumor tissues (n=8) obtain from patients with pathological stage II of breast tumor. MSCs were isolated by explant culture method and identified. The Mitogen-induced PBMC was treated with 25 and 50% MSC conditioned medium (CM) for 72h, and changes in proliferation evaluated by BrdU ELISA kit.

Results: We successfully isolated and identified MSCs from primary breast tumor tissues. Flowcytometry analysis demonstrated that isolated cells were positive for, CD73, CD44, CD29, CD105 and CD90 but negative for, CD11b, CD45, CD34, and HLADR. In addition, cells possessed the capability of multipotential differentiation into osteoblasts and adipocytes.

The results of theBrdU ELISA assay showed that 50% CM concentrations significantly increased the proliferation of PBMCs.

Conclusion: Collectively, our study showed that MSCs were confirmed to exist in stage II of human breast tumor and significantly promote the proliferation of PBMCs In Vitro.

Keywords: Breast Cancer, Mesenchymal Stem Cells, Isolation, Immunomodulatory

ICIA2018\Cancer Immunology\Poster\11064
11064

Evaluation of NK(CD16+CD56+CD3-), NKT(CD16+CD56+CD3+), T(CD16-CD56-CD3+) cells cytotoxic activity in PBL of breast cancer patients by using CD107a expression assessment by flow cytometry (before and after co-culture with MCF-7 & PMA/I)

Minoo Adib^{1*}, Nasim Samarghandi¹

1. Department of Immunology, Faculty of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

Background: Invasive ductal carcinoma is the most prevalent type of breast cancer. Because of attention in recent years to innate immunity against tumors, assessment of NK cell function as an important component of the innate immune response is essential to find precise information about these cells. A lack of available assays for the detection of NK cells functional activity is an important limitation in the field of NK cells researches.

Methods: There are several methods to estimate the cytotoxic activity of NK cells. A new method has been described by using CD107a expression on the cell surface as a marker of cytotoxic NK & CD8⁺ T-cell degranulation.

Results: There was a significant increase in CD107 expression on NKT cells (with PMA/I stimulation) of control group comparing with patients group. The mean percent of T cells decreased significantly in grade III compared to grade I, II and control group.

Conclusion: In this study CD107 expression, the initial marker of cytotoxic activity, on the three populations of cells is investigated in two groups and resulted in expression of this marker on NKT cells is higher than NK and T cells in both groups. Just as many studies indicated a decrease in NK cell cytotoxicity in any grade of disease, this study showed that decrease of NK cells cytotoxicity only occur in grade I & II, but as a striking point, in grade III of disease cell cytotoxicity of these cells increased. Eventually, in this study expression of CD107 on T cells of both groups is equal. Several studies to confirm of our results reported major reasons for the observed decreased cellular immune responses in patients with breast cancer were associated with defective presentation of antigens by dendritic cells, not function of effector T cells.

Keywords: NK cell, CD107a, Breast Cancer, flow cytometry, MCF-7

ICIA2018\Cancer Immunology\Poster\11078
11078

Comparison of miR146a-5p expression in 4T1 and MKN cancer cell lines

Atefe Alirezaee¹, Ahmad Zavaran Hosseini¹, Sara Soudi¹

1. Department of Immunology, Faculty of Medical Sciences, TarbiatModares University, Tehran, Iran

Background: Different studies demonstrated that MicroRNA (*miR*) are participated in tumor formation and progression by regulation of cell proliferation, apoptosis and differentiation. *miR-146a* are among the important microRNAs that control cell apoptosis. Upregulation of *miR-146a* will terminated to downregulation of apoptotic pathways. In this study we want to know if there is a difference in the expression of miR-146a in two cell lines.

Methods: For this purpose the amount of miR-146a-5p expression were measured in 4T1 and MKN-45 cancer cell lines as represented the related cells to breast cancer and gastric cancer, respectively. So 3×10^5 cells of 4T1 and MKN-45 were cultured in 12 well plates. 48 hours after culture, the cells harvested and assessed for miR-146a-5p expression by Real time PCR.

Result: According to obtained result, MKN-45 cancer cell line demonstrated two fold decrease in miR-146a-5p expression compared with 4T1.

Conclusion: As a result, we confirmed the miR-146a-5p expression in both cell lines. However there is a difference in miR-146a-5p level between different cancer cell lines. So miR-146a-5p, has a different therapeutic value dependent to kind of cancers.

Keywords: miR146a-5p, 4T1, MKN-45

ICIA2018\Cancer Immunology\Poster\11097

11097

The anti-tumor effects of arsenic trioxide in combination with an EGFR inhibitor on breast cancer cell linesElham Boustanipour¹, Zahra Sadr¹, Ali Zekri², Seyed H. Ghaffari³

1. *Department of Medical Genetics, Faculty of Medicine, Tehran University of Medical Sciences (TUMS), Tehran, Iran*
2. *Department of Medical Genetics and Molecular Biology, Faculty of Medicine, Iran University of Medical Sciences (IUMS), Tehran, Iran*
3. *Hematology, Oncology and Stem Cell Transplantation Research Center, Tehran University of Medical Sciences, Tehran, Iran*

Background: Breast cancer (BC) is the most prevalent malignancy found in women worldwide. Despite extensive advances in diagnosis, surgical techniques, and treatment, BC is among the leading causes of cancer-related deaths in women. Metastasis is the main challenges in the BC management and it accounts for majority of deaths in breast cancer patients. Therefore, understanding the cellular and molecular factors responsible for the metastasis and developing novel therapeutic approaches against metastasis of breast cancer is urgently needed. Previous studies showed that epidermal growth factor receptor (EGFR) plays key roles in breast cancer initiation, growth, dissemination and cancer immunotherapy. Arsenic trioxide (ATO) as a multifunctional drug has been investigated for the treatment of leukemia and several solid tumors. EGFR inhibitor could influence anti-tumor immune responses by changing immune gene expression.

Methods: Different breast cancer cell lines were obtained and cultured in RPMI 1640 medium supplemented with 10% fetal bovine serum. The effects of ATO as a single and in combination with EGFR inhibitor were assessed on the viability and migration of BC cell lines using MTT, apoptosis and wound-healing assays in different time intervals.

Results: ATO or EGFR inhibitor as a single treatment did not have a considerable effect on the viability and wound-healing rate of BC cell lines, but the combination of the ATO with EGFR inhibitor significantly reduced metastasis effect of the cell lines.

Conclusions: Present study uncovered that combination of ATO with EGFR inhibitor could overcome metastasis of the breast cancer. Overall, combination of cytotoxic treatment with targeted therapy may represent a promising strategy for treatment of the breast cancer.

Keywords: Arsenic trioxide, EGFR inhibitor, Breast cancer, Metastasis.



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Cancer Immunotherapy

Poster Presentation

ICIA2018\Cancer Immunotherapy\Poster \5236 cancer immunotherapy
5236

Evaluation of Optimal Pattern of Injection and Dosage of Dendritic Cell Cancer Immunotherapy Using the Mathematical Model Based on Artificial Neural Network Model

Zahra Mirsanei¹, Sima Habibi¹, Nassim Kheshtchin¹, Jamshid Hadjati*¹

1. Department of Immunology, School of Medicine, Tehran University of Medical Science, Tehran, Iran

Background: Previous studies have demonstrated that maturation of dendritic cells by *Listeria monocytogenes* lysate or CpG DNA can enhance the efficiency of the cancer vaccines in experimental models. A mathematical model based on artificial neural network of those results, proposed a new exponential pattern for vaccination. According to the model, injection of dendritic cells matured by CPG (CpG-DCs) and *Listeria monocytogenes* (List-DCs) at the same time and with decreasing and increasing dose of each vaccine in a group of mice would induce optimal antitumor immune response and minimum immunosuppression in the tumor microenvironment.

Method: 164 cells were implanted subcutaneously in right flank of Balb/C mice in four groups: vaccinated with *Listeria monocytogenes* matured DCs, CpG, both and none as control group. DC vaccines were injected according to proposed mathematical model. Tumor growth in the course of treatment and survival of mice after that was followed by measuring tumor size. GrBELISA test for assessing the activity of CTL, and real time PCR test for evaluation of T-bet (Th₁ transcription factor) and FOXP₃ (Treg transcription factor) gene expression were used.

Result: Results from the group received combination therapy with *Listeria monocytogenes* and CPG matured DCs, compared with other groups that have received only one vaccination alone, showed that growth of tumor is slower and declining, as well as higher activity of CTLs in this group. Gene expression results also confirm the increase of immune response in the tumor microenvironment.

Conclusion: Based on the results it can be concluded that mathematical models can be useful tools for predicting the tumor behavior and more efficient treatments, so the effort to improve them should be considered.

Key word: Dendritic cells, Cancer vaccine, Mathematical model, Immunotherapy.

ICIA2018\Cancer Immunotherapy\Poster \6330 cancer immunotherapy
6330

MiR-34a Replacement Effect on Growth and Migration Inhibition in HCT 116 Colon Cancer Cell

NazaninJafarpour¹, Habib Onsori², Behzad Baradaran³

1. Department of Genetics, East Azarbaijan Science and Research Branch, Islamic Azad University, Tabriz, Iran

2. Department of Genetics, Tabriz Branch, Islamic Azad University, Tabriz, Iran

3. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran (Corresponding Author)

Background: Colorectal cancer is the most common cancer of the digestive tract and is also the fourth leading cause of death in the world. Colorectal cancer is the third most commonly reported case in women, and is the second leading cause of cancer deaths in women. In this study, we increased the miRNA-34a substitution in colon cancer cells by transferring the miRNA-34a gene and expressing this miRNA in these cells and examined the effects of this miRNA on growth and migration inhibition.

Methods: Cancer cells were cultured in HCT-116 in RPMI1640 culture medium and all cellular experiments were performed in a logarithmic growth phase. For the transfer of miRNA-34a into colon cancer cells, the JetPAI reagent (poly plus) was used. The expression of miR-34a was induced by qRT-PCR after miRNA induction. To investigate the effect of this miRNA on the cellular migration status after induction of Wound healing, ultimately the cells were induced by untreated cells were compared and significant amounts were determined.

Results: MTT assay showed that miR-34a induction induced death of colorectal gradient cells. The results of the qRT-PCR test showed a significant increase in the expression of miRNA-34a in transfected HCT116 cells. Ultimately, the results of the Wound Healing test showed reduced migration of transfected cells compared to control cells.

Conclusion: The results showed that increase of miRNA-34a induced cell death and decreased metastasis and migration of HCT-116 cell line colon cancer cells.

Key words: miRNA, Colon cancer, Metastasis, Transfection

ICIA2018\Cancer Immunotherapy\Poster \7339 cancer immunotherapy
7339

Selection and Evaluation of Human Recombinant Antibodies against ErbB2 Antigen for Breast Cancer Immunotherapy

ElhamNadimi^{1,2}, Foroogh Nejatollahi^{1,2}, Setareh Moazen³

1. *Recombinant Antibody Laboratory, Department of Immunology, Shiraz University of Medical Sciences, Shiraz, IR Iran*

2. *Shiraz HIV/AIDS Research Center, Institute of Health, Shiraz University of Medical Sciences, Shiraz, IR Iran*

3. *Faculty of Science, University of British Columbia, Vancouver, Canada*

Background: Breast cancer is the most common cause of cancer-related death in women worldwide. ErBb2/HER2+ breast cancer accounts for 25% - 30% of all cases of breast cancer. Human recombinant single-chain antibodies, which are produced by antibody engineering technologies, are new and effective antibodies in cancer immunotherapy.

Methods: Panning process was performed on the clones of a human scFv library using two ErbB2 epitopes and Polymerase chain reaction (PCR) and DNA fingerprinting were done on the selected clones. The clones were evaluated via phage ELISA in terms of reactivity and specificity to epitopes.

Results: Single-chain antibodies, scFvI and scFvII, the anti-ErbB2-specific scFvs with 40% and 45% frequencies, were selected against epitopes I and II, respectively. The results of phage ELISA demonstrated a significant difference in the optical density (OD) of scFvs in reaction with the related peptides.

Conclusions: Targeted cancer therapy, which acts on a specific molecule in cancer cells, minimizes the side effects of immunotherapy. Due to the unique properties of scFvs, these antibodies have been used in targeted therapy of several cancers. In this study, 2 specific scFvs were selected against 2 ErbB2 epitopes, which contained trastuzumab and pertuzumab binding sites. The results of the panning process demonstrated the selection of 2 specific scFvs (with frequencies of 40% and 45%, respectively), which significantly reacted with the corresponding epitopes in phage ELISA assay. These small, high-affinity, human antibodies, which were selected against regions containing the binding sites of 2 food and drug administration (FDA)-approved monoclonal antibodies for breast cancer immunotherapy, have the potential to be considered for breast cancer targeted therapy. However, in vitro and in vivo tests should be performed to evaluate the antitumor effects of these scFvs.

Keyword: Breast cancer, scFv, ErbB2

ICIA2018\Cancer Immunotherapy\Poster\7437 cancer immunotherapy
7437

Anti-tumor activity of Neurokinin 1 receptor inhibitor, in multiple myeloma

Elham Razani, Davood Bashash*

Department of Hematology and Blood banking, School of Allied Medical Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: The substance P/neurokinin 1 receptor (SP/NK1R) cascade has been demonstrated to play a considerable role in development of human cancers. In spite of therapeutic improvements in treatment strategies over the past decades, multiple myeloma (MM) still remains as one of the leading causes of person-years of life lost all over the world. On the basis of the pathogenic role of SP/NK1R pathway in cancer, it was of great interest to investigate the anti-cancer effect of aprepitant, an oral competing non-peptide antagonist of NK1R, in MM cells.

Methods: To evaluate the cytotoxic and anti-proliferative effects of aprepitant in multiple myeloma, KMM cells were treated with increasing doses of the inhibitor. Cell viability and metabolic activity were investigated by trypan blue and MTT assays, respectively. Afterwards, Annexin-V/PI staining and DNA content analysis were performed to evaluate whether aprepitant-induced cytotoxic effect could be attributed to either apoptosis induction or cell cycle arrest. Additionally, transcriptional alteration of apoptosis-related target genes was also studied using real-time RT-PCR.

Results: We found that treatment of KMM cells with aprepitant resulted in inhibition of both viability and metabolic activity in a concentration-dependent manner. Moreover, flow cytometric analysis delineated a considerable pro-apoptotic potential of the inhibitor in MM cells, as evidenced by increased externalization of phosphatidylserine, and elevated cell population in sub-G1 phase. Noteworthy, the results of real-time PCR revealed that aprepitant-induced cytotoxicity is mediated through shifting the ratio of death promoters to death repressors via alteration of Bax and Bcl-2 expression.

Conclusions: Based on the pharmacologic safety of aprepitant and its broad clinical application in chemotherapy-induced nausea and vomiting prevention, our study suggests this inhibitor as a promising agent for the treatment of multiple myeloma. However, further investigation, including clinical trials will provide valuable clues to add this inhibitor for treatment of MM patients.

Keywords: Multiple myeloma, Aprepitant, Neurokinin-1 receptor.

ICIA2018\Cancer Immunotherapy\Poster\7530 cancer immunotherapy
7530

Native human PD1: reducer of PD1⁺FOXP3⁺CD8⁺T cells

Samane Mohamadzade¹, Alireza Andalib¹, Nafise Esmael¹, Nahid Eskandari¹, Abbas Rezaei¹,
Hosein Khan Ahmad², Ilnaz Rahimmanesh²

1. Department of Immunology, School of Isfahan Medical, Isfahan University of Medical Sciences

2. Department of Genetics and Molecular Biology, School of Isfahan Medical, Isfahan University of Medical Sciences

Background: PD1 is one of the immune check point receptors which expresses on the activated T cells and suppress stimulatory signaling of TCRs. In the tumor microenvironments, continuous stimulation of CD8⁺T cells increases PD1 expression, PD1⁺CD8⁺T cells, and PD1 signaling. PD1/PD-L1 engagements induce FOXP3 expression and regulatory T cell differentiation, FOXP3⁺PD1⁺CD8⁺T cells, which are inhibitor cells in the tumor microenvironments. Therefore, FOXP3⁺PD1⁺CD8⁺T cells are the progenitor population of FOXP3⁺PD1⁺CD8⁺T cells. Inhibition of PD1 receptors on the T cells could decrease PD1/PD-Ls interactions and CD8⁺ regulatory T cells induction. In this study we constructed a native human PD-1, shPD1, and assessed its effect on FOXP3⁺PD1⁺CD8⁺T cells.

Methods: We constructed shPD1 which was contained extracellular domain of human PD1 and GFP. We co-cultured concanavalinA stimulated T cells and MDA-MB-231 cell line with or without shPD1 (groups 1 and 2). After 24 hours, PD1 and FOXP3 expression by CD8⁺T cells was assessed with anti CD8, anti PD1 and anti FOXP3 antibodies using flow cytometry.

Results: Population of FOXP3⁺PD1⁺CD8⁺T cells in group 1 (6.4%) was significantly, $P < 0.05$, lower than group.2 (8.12%).

Conclusion: Stimulation of lymphocytes with concanavalinA induced PD1 expression on the surfaces of T cells; therefore PD1 signaling increased FOXP3⁺PD1⁺CD8⁺T cells which were progenitors of FOXP3⁺PD1⁺CD8⁺T cells. However, shPD1 in co-culture of stimulated T cells with MDA-MB-231 cell line, inhibited PD1/PD-L1 interaction and decreased PD-1 signaling and PD1 expression. Therefore FOXP3⁺PD1⁺CD8⁺T cells were decreased and accordingly FOXP3 induction was reduced. We concluded that our shPD1 could be used as a reducer of inhibitory PD1⁺CD8⁺T cells and preventer of FOXP3⁺PD1⁺CD8⁺T cells in the tumor microenvironments. Therefore inhibitory molecules secretion could be decreased using shPD1 in the tumor microenvironments.

Keywords: shPD1, PD1⁺ CD8⁺ T cells, FOXP3.

ICIA2018\Cancer Immunotherapy\Poster \7535 cancer immunotherapy
7535

Immunogenicity of Rituximab, Trastuzumab and Bevacizumab Monoclonal Antibodies in patients with malignant diseases

Fatemeh Saffari¹, Abdollah Jafarzadeh^{1, 3, 4}, Kalantari-Khandani B³, Farzaneh Saffari²,
Soleimani S¹, Mohammadi MM¹

1. Department of Immunology, Medical School, Kerman University of Medical Sciences, Kerman, Iran
2. Department of Laboratory Sciences, Para-Medicine School, Shiraz University of Medical Sciences, Shiraz, Iran
3. Department of Hematology-Oncology and BMT, Kerman University of Medical Sciences, Kerman, Iran
4. Molecular Medicine Research Center, Rafsanjan University of Medical Sciences, Rafsanjan, Iran.
5. Department of Immunology, Medical School, Rafsanjan University of Medical Sciences, Rafsanjan, Iran

Background: One of the most important limiting efficacy of monoclonal antibodies (mAb) is their immunogenicity that may influence the diagnostic and therapeutic process of them. The aim of this study was determine the presence of antibody against some mAbs including Rituximab, bevacizumab and trastuzumab in malignant patients.

Methods: Blood samples were collected from 32 patients with lymphoma and leukemia, 43 patients with breast cancer and 23 patients with adenocarcinoma during treatment with Rituximab, trastuzumab and bevacizumab, respectively. The serum levels of human antibodies against mentioned mAb were determined by designing a standard of sandwich ELISA method.

Results: The presence of human antibodies against mentioned antibodies was detected in 4 of 32 (12.5%) of Rituximab-treated patients and 7 of 43 (16.3%) of trastuzumab- treated patients with mean titer of 2.33 ± 0.37 AU/mL and 1.2 ± 0.21 AU/mL, respectively. The seropositivity for the presence of anti-mAb in patients treated with Rituximab alone was significantly higher than patients administrated both Rituximab and chemotherapeutic agents (26.6% versus 0.0%; $P < 0.02$). None of bevacizumab-treated patients were developed antibody against administrated mAb.

Conclusion: In conclusion, these results indicate the production of antibody against therapeutic mAbs rituximab and trastuzumab were detected in a number of treated patients that may influence their efficacy. The production of the human anti-mAb antibody may influenced by chemotherapeutic drug in rituximab-treated patients. Bevacizumab did not show immunogenicity in our treated patients.

Keywords: Immunogenicity, Malignant patients, Rituximab, Trastuzumab, Bevacizumab

ICIA2018\Cancer Immunotherapy\Poster \7555 cancer immunotherapy
7555

Effect of miR-200c replacement on growth and migration of prostate cancer cells (pc3)

Roghayeh Pirzadeh¹, Habib Onsori², Behzad Baradaran³

1. Department of Genetics, East Azarbaijan Science and Research Branch, Islamic Azad University, Tabriz, Iran

2. Department of Genetics, Tabriz Branch, Islamic Azad University, Tabriz, Iran

3. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran (Corresponding Author)

Background: Prostate cancer is one of the most common cancers among men around worldwide. In advanced countries, prostate cancer is the most common cancer after skin cancer and the most deadly cancer after lung cancer. Studies show that mir-200c expression in prostate cancer is reduced. In this study, we increased mir-200c level in the prostate cancer cells and examined its effect on cell growth and migration. Also the effect of mir-200c replacement on ZEB1 and ZEB2 expression was evaluated.

Methods: The pc3 cells cultivating in RPMI-1640 medium, the IC₅₀ of antibiotic (G418) geneticin was obtained by using the MTT assay. Using jetPEI transfection reagent, miR-200c was transmitted to prostate cancer cells. The qRT-PCR test was used to prove the action of transferring this miRNA into cells. The effect of mir-200c on target genes ZEB1, ZEB2 was evaluated by qRT-PCR. Wound healing assay was performed to examine the migration status of transfected cancer cells.

Results: Using the MTT test, an antibiotic Geneticin, IC₅₀ 200 μ M was obtained. The results of qRT-PCR showed increased expression of miRNA-200c in PCR cells transfected with miRNA-200c. The results of Wound healing showed reduced migration of transfected cells with miRNA-200C compared to untreated control. The result of qRT-PCR showed the expression level of ZEB1 and ZEB2 was decrease after replacement of mir-200c.

Conclusion: The results showed that increasing miR-200C plays an important role in inhibiting the growth and migration of PC3 cell line prostate cancer cells.

Keywords: miRNA, Prostate Cancer, Immigration, Transfusion

ICIA2018\Cancer Immunotherapy\ Poster \7703 cancer immunotherapy
7703

Leukemia cell growth inhibition by silencing of Mcl-1 gene

Shaghayegh Askarian amiri¹, Abbas Azadmehr¹, Farzane Ordoni aval¹, Morteza Oladnabi²

1. Cellular and Molecular Biology Research Center, Department of Immunology, Health Research Institute, University of Medical Sciences, Babol, Iran

2. Gorgan Congenital Malformations Research Center, Department of Human Genetics, School of Advanced Technologies in Medicine University of Medical Sciences, Gorgan, Iran

Background: T cell acute lymphoblastic leukemia (T-ALL) is one of the most malignant hyperplastic diseases of the hematopoietic system. This tumor is most common tumor diseases in children and youths. Mcl-1 is a pro-survival protein of the Bcl2 family member. Mcl-1 was found as an important factor to conventional cancer therapies. Down regulation of this gene can be potential therapeutic target in multiple types of cancers.

Methods: The aim of this study was to investigate the effect of specific Mcl-1 small interfering RNAs (siRNAs) on the survival and induction of apoptosis on jurkat cells. Specific Mcl-1 siRNA was transfected. Then, relative Mcl-1 mRNA expression level was measured by quantitative real-time PCR. Moreover, cell survival was determined using colorimetric MTT assay and apoptosis was analyzed using flow cytometry.

Results: Our findings showed that 48 hours after transfection with Mcl-1 siRNA the expression of mRNA levels was effectively reduced in dose dependent manner. Furthermore, treatment with Mcl-1 siRNA significantly reduced tumor cell survival and significantly increased the extent of T-cell acute leukemia cell apoptosis.

Conclusion: Our results suggest that down-regulation of Mcl-1 by specific siRNA in T-cell acute leukemia cells can effectively reduce cell survival and induces apoptosis cell death. Therefore, Mcl-1 siRNA may be a complementary agent in treatment of T-ALL.

Key words: Mcl-1, siRNA, Jurkat cell line, T-ALL, Apoptosis

ICIA2018\Cancer Immunotherapy\Poster\7704 cancer immunotherapy
7704

BACH1 as inducer marker for prostate cancer cells metastasis

Neda Shajari^{1, 2}, Behzad Baradaran^{1, 2}

1. Department of Immunology, School of Medicine, University of Tabriz

Background: Metastasis to distant organs is a hallmark of many tumor cells. BACH1 (BTB and CNC homology 1) has been shown to transcriptionally regulate expression of a range of genes that are associated with breast cancer metastasis. The aim of this study was to investigate the effect of BACH1 silencing on prostate cancer cells in vitro.

Methods: BACH1 expression was silenced in DU145 prostate cancer cell line using siRNA approach. Quantitative RT-PCR was used to detect BACH1 expression and other metastasis-related genes following siRNA knockdown.

Results: Quantitative RT-PCR analysis revealed that the expression levels of BACH1 mRNA in DU145 cells were significantly suppressed after transfection. Also, the CXCR4 and MMP1 expression levels decreased following BACH1 knockdown in DU145 cells.

Conclusion: Our results indicated that BACH1 down-regulation in DU145 cells decrease the expression levels of some metastasis-related genes. These results suggest that BACH1 may function as an oncogenic driver in prostate cancer and may represent as a potential therapeutic target for metastatic prostate cancer.

Keywords: BACH1, siRNA, Prostate Cancer

ICIA2018\Cancer Immunotherapy\Poster\7745 cancer immunotherapy
7745

Inhibition of glioblastoma cell growth by silencing of TGF β RII

Farzane Ordoni aval¹, Abbas Azadmehr¹, Shaghayegh Askarian amiri¹, Payam Saadat², Somayeh Fallahnejad³, Morteza Oladnabi⁴, Behzad Baradaran⁵, Behzad Mansoori⁵

1. Cellular and Molecular Biology Research Center, Department of Immunology, Health Research Institute, University of Medical Sciences, Babol, Iran
2. Mobility Impairment Research Center, Health Research Institute, University of Medical Sciences, Babol, Iran
3. Obstetrics and Gynecology department, Shahid Beheshti University of Medical sciences, Tehran, Iran
4. Gorgan Congenital Malformations Research Center, Department of Human Genetics, School of Advanced Technologies in Medicine University of Medical Sciences, Gorgan, Iran
5. Immunology Research Center, University of Medical Sciences, Tabriz, Iran

Background: Glioblastoma (GBM) is the most malignant and aggressive type of glioma, associated with a high rate of mortality. The transforming growth factor- β receptor II (TGF β RII) is involved in glioma initiation and progression. On the other hand, TGF β RII blockade is critical to the inhibition of GBM. Therefore, we aimed to determine the effects of specific TGF β RII siRNA on the survival of U-373MG cells.

Methods: TGF β RII siRNA was transfected and qRT-PCR was performed to examine TGF β RII mRNA expression. Cell survival was determined using colorimetric MTT assay, and platelet-derived growth factor-BB (PDGF-BB) level was measured in the culture supernatant using ELISA assay.

Results: Our findings indicated that specific siRNAs could dose-dependent suppress TGF β RII mRNA expression after 48 hours. In addition, treatment with TGF β RII siRNA significantly reduced tumor cell survival and decreased the amount of PDGF-BB proteins in the cell culture supernatant.

Conclusion: Our results suggest that TGF β RII blockade can be a promising complementary treatment for glioma.

Keywords: TGF- β RII, siRNA, U-373 MG cell line, PDGF-BB, Glioblastoma

ICIA2018\Cancer Immunotherapy\Poster\7826 cancer immunotherapy
7826

Enhancement of 5-fluorouracil-induced cytotoxicity by LPS in MCF7 human breast cancer cells

Negin Nokhandani¹, Mahdieh Naghavi Alhosseini², Seyedeh Naemeh Javid³, Homa Davoodi⁴

1. Laboratory Sciences Research Center, Golestan University of Medical Sciences, Gorgan, Iran
2. Department of Immunology, School of Medicine, University of Golestan Medical Sciences, Gorgan, Iran
3. Department of Microbiology, School of Medicine, University of Golestan Medical Sciences, Gorgan, Iran
4. Cancer Research Center, University of Golestan Medical Sciences, Gorgan, Iran

Background: 5-Fluorouracil (5-FU) is a widely used anticancer drug in patients with breast and other solid cancers. Despite its many advantages, clinical applications were limited because of the drug resistance. On the other hand, the chemotherapy has shown common side effects include myelosuppression, dermatitis, cardiac toxicity, diarrhea, and mucositis. Therefore, new strategies and tumoricidal agents for therapy and resistance reversal are urgently needed. Lipopolysaccharide (LPS) is one of the major components of the outer layer of gram-negative bacterial membrane. The Stimulation of toll-like receptor 4 (TLR4) through LPS activates immune responses. In the previous study, Davoodi et al. showed that LPS increased the cytotoxic effect of 5-FU on colon cancer cells. In the current study, the effect of LPS pretreatment on cytotoxicity induced by 5-Fu in breast cancer cell line MCF-7 was investigated.

Methods: The effect of LPS pretreatment on the cytotoxicity of 5-FU was tested in a breast cancer cell line, MCF7. Cell viability was analyzed using MTT assay. Cancer cells were seeded in 96 well plates. On the following day cells were exposed to LPS (1µg/ml) for 4 hours and then were treated with different concentration of 5-FU (0-500µg/ml). After 24-72 h, the cells were collected and cell viability was determined by MTT assay.

Results: The results showed that, 1µg/ml LPS pretreatment increased the cytotoxicity effect of different dose of 5-FU in MCF-7 ($P < .05$). We showed that 4 hours pretreatment of LPS has desired effect in comparison with simultaneous treatment of LPS and 5-FU.

Conclusion: LPS augmented the tumoricidal effects of the chemotherapy by 5-FU in breast cancer cells. LPS can increase the anti-cancer effect of 5-FU and therefore reduce advert side effects of chemotherapy and also reduce the rate of resistance to 5-Fu in patients.

Keywords: Breast cancer, Lipopolysaccharide, 5-fluoracil

ICIA2018\Cancer Immunotherapy\Poster \7987 cancer immunotherapy
7987

Overexpression of microRNA-193a-5p, as a novel prognostic marker, inhibits migration of human HT-29 colon cancer cells via down-regulating metastasis-related genes expression

Naghmeh Shirafkan^{1,2}, NavidShomali^{1,2}, Behzad Mansoori¹, Ali Mohamadi¹, Behzad Baradaran^{1*}

1. Immunology Research Center (IRC), Tabriz University of Medical Sciences, Tabriz, Iran

2. Student Research Committee, Tabriz University of Medical Sciences, Tabriz, Iran

3. Correspondence*: Behzad Baradaran, Immunology Research Center, Tabriz University of Medical Sciences

Background: Colorectal cancer (CRC) is among three leading reasons of cancer-related mortality in all over the world. MicroRNAs are important post-transcriptional regulators which dysregulated in cancers and affect the expression of different genes in cancer progression. Down-regulation of miR-193a-5p has been found in some types of cancer. However, effects and potential mechanisms of miR-193a-5p remains elusive in CRC.

Methods: Quantitative RT-PCR was performed to detect expression of miR-193a-5p and metastasis-related genes before and after mimic transfection in CRC cell lines. Wound healing assays measured migration capacity of HT-29 cell lines after miRNA transfection. MTT and Annexin V staining were used to determine the role of miR-193a-5p in the regulation of HT-29 proliferation and apoptosis, respectively.

Results: In this study, we found that miR-193a-5p was down-regulated in CRC cell lines. And the restoration of miR-193a-5p in human HT-29 cell line could inhibit cell migration. But, reduced viability and apoptosis induction did not occur significantly after miR-193a-5p overexpression. Also, the quantitative RT-PCR analysis showed that the expression levels of miR-193a-5p mRNA were increased after mimic transfection. Furthermore, the Vimentin, E-cadherin and CXCR4 mRNA expression levels decreased after overexpression of miR-193a-5p with mimic in HT-29 cell line.

Conclusion: Our results showed that miR-193a-5p may function as a tumor suppressor and play an important role in metastasis and suggest it may be a potential therapeutic target for applying in CRC therapy.

Keywords: Colorectal cancer, microRNA, Metastasis, Apoptosis

ICIA2018\Cancer Immunotherapy\Poster \8077 cancer immunotherapy
8077

Investigation of anti-tumor and anti-inflammatory effects of *Nigella sativa* on 4T1 and CT26 cell lines

Fatemeh Hosseini¹, Hadi Hassannia²

1. Department of Immunology, School of medicine, Semnan University of Medical Sciences, Semnan, Iran

2. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

Background: Cancer is the second leading cause of death globally and cancer-related inflammation plays an important role in malignant progression of several cancer types. Current therapeutic approaches for cancer therapy are not completely effective and are associated with many side effects. In recent years, there has been growing interest in the application of medicinal herbs, since they have fewer side effects than chemical drugs. Therefore, the present study aimed to investigate of anti-tumor and anti-inflammatory effects of *Nigella sativa* in *in vitro* conditions.

Methods: In this study, 4T1 and CT26 cell lines were treated with different concentrations of hydro alcoholic extracts of *Nigella sativa* (50, 250, 500, 1000 µg/mL). After 48h, the effects of the extract on tumor growth and COX-2 gene expression were analyzed by XTT and Real time PCR, respectively.

Results: The results revealed that 4T1 and CT26 cell lines growth significantly decreased in a dose dependent manner in compared to the control ($p < 0.05$). Also, Real time PCR results showed that the expression of COX-2 significantly decreased in a dose dependent manner even at the low concentration of extract in compared to the control ($p < 0.05$).

Conclusion: Based on these results, *Nigella sativa* extract not only suppressed the growth of cancer cell lines but also significantly reduced cancer-related inflammation. Therefore, it may be utilized as a novel therapeutic in several types of cancer. However, this claim needs further investigations.

Keywords: *Nigella sativa*, Cancer, Inflammation, COX-2

ICIA2018\Cancer Immunotherapy\Poster \8163 cancer immunotherapy

8163

Effect of conditioned medium of mesenchymal stem cells pulsed with different doses of caffeine on murine breast cancer cells (4T1).

Zahra Zeinaly, Nouroz Delerizh, Seyyed Meysam Abtahi Froushani

Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Background: Conditioned medium of Mesenchymal Stem Cells (MSCs) have potent immunomodulatory and anticancer benefits. It is clear that environmental factors modulate the functions of MSCs. This study was set out to evaluate the effects of the conditioned medium of MSCs (CM) of caffeine (as a very popular drink) treated MSCs on the proliferation of 4T1, a breast adenocarcinoma cell line.

Methods: MSCs were isolated by flashing the tibia and femur bones of mice. Third Passage of MSCs incubated with different doses of caffeine (0, 0.1, 0.5 and 1 mM) for 24 hours. Then cells were washed and cultured for 24 hour with fresh medium. The CM was collected and used in the culture medium of 4T1 (50%).

Results: Neutral red (NR) uptake and MTT proliferation test indicated that the CM could significantly decrease the proliferation of 4T1 cells. NR uptake and MTT test showed that CM pulsed with caffeine at 0.1 mM and CM pulsed with caffeine at 0.5 mM could significantly increase the anticancer effects of CM, respectively. Data from both tests indicated that CM pulsed with caffeine at high dose (1 mM) could significantly regress these beneficial effects of CM.

Conclusion: It seems that caffeine at the low to moderate doses can potentiate the cytotoxic effect of CM. Nevertheless, the high dose had trouble effects on the beneficial effects of CM on the proliferation of cancer cells.

Keywords: Caffeine, Conditioned medium, Mesenchymal stem cell.

ICIA2018\Cancer Immunotherapy\Poster \8417 cancer immunotherapy
8417

Immunotherapy in ependymoma brain tumor

Fateme Shaabanpour Aghamaleki¹, Shirin Farivar²

1. Department of molecular and cellular Biology School of Biotechnology and Life Science University of Shahid Beheshti

2. Department of molecular and cellular biology school of biotechnology and Life Science University of Shahid Beheshti

Background: Approximately 50% of children with ependymoma will suffer from tumor recurrences that will ultimately lead to death. Children with the recurrent disease commonly experience cumulative neurological morbidity from repeated surgical and adjuvant therapy interventions, and almost universally suffer from tumor progression. Accordingly, conceptually new treatment approaches are needed, both to decrease the risk of tumor recurrence and to enhance disease control in those children who experience recurrent disease. One of this approaches is immunotherapy, also called biologic therapy, and is a type of cancer treatment that boosts the body's natural defenses to fight cancer. It uses substances made by the body or in a laboratory to improve or restore immune system function.

Methods: Gene's expression profiling by array of 20 ependymoma patients with recurrent experience and 45 primary ependymoma patients was extracted from Geo datasets. Genes are compared with logfc for their expression between two kinds of patients and sorted by their p-value. Also, the most related pathway, location, function, and protein networks of this gene were identified with string, GeneCard and DAVID databases.

Results: Several important immune genes that associated with recurrent ependymoma tumor are identified such as CCL19, CXCL9, NCF1, and IGSF6. These genes have hyperexpression in ependymoma patients with recurrent experience. Among these genes, CXCL9 is a small cytokine belonging to the CXC chemokine family. CXCL9 is a T-cell chemoattractant, which is induced by IFN- γ . It is associated with CCL19 and CXCL2 and plays roles in Toll-like Receptor Signaling Pathway and ERK Signaling. It has hyperexpression in ependymoma patients with recurrent experience.

Conclusion: According to CXCL9 function and expression changes, it can play an important role in ependymoma with recurrent experience. Therefore, it can be used as an immunotherapeutic target for cancer therapy.

Keywords: Ependymoma, Immunotherapy, Gene expression, CXCL9

ICIA2018\Cancer Immunotherapy\Poster \8434 cancer immuno
8434

Using CRISPR/Cas9 system for immunotherapy of brain tumors

Fateme Shaabanpour Aghamaleki¹, Shirin Farivar²

1. Department of cellular and molecular Biology School of biotechnology and life science University of shahid beheshti university

2. Department of cellular and molecular biology Faculty of biotechnology and life science University of shahid beheshti

Background: Brain tumors are the leading cause of cancer mortality in children and remain difficult to cure despite advances in surgery and adjuvant therapy. Most current brain tumor research is focused on the molecular and cellular analysis of the tumor mass. There are wide variety of immune genes that play an important roles in brain tumors pathogenesis and progression. Identification of these genes can be used as therapeutic target in CRISPR/Cas9 system. CRISPR/Cas9 system has been modified to edit genomes, by delivering the Cas9 nuclease complexed with a synthetic guide RNA (gRNA) into a cell, the cell's genome can be cut at a desired location, allowing existing genes to be removed and/or new ones added.

Methods: Genes expression profiling by array of 117 brain cancer patients and 13 healthy individuals were extracted from Geo datasets. Genes are compared with logfc (fold change) for their expression in healthy and patients group, and then sorted by their p-value. Also the most related pathway, genes function and ontology are identified by DAVID and GeneCard databases. Finally genes with most modification and related to critical pathway in immune system are selected and their best gRNAs (guide RNA) in CRISPR system are identified for editing by CHOPCHOP datasets.

Results: TNFRSF19 or TNF receptor superfamily member 19 has hyperexpression among brain tumor patients. It plays an important roles in immune system and inflammatory response. It's important pathway and function are Wnt and TNF Superfamily Pathway. The best gRNAs for this gene is identified through CHOPCHOP database that haven't any secondary structures.

Conclusion: Many experiences showed that immune genes play an important roles in cancer progression and metastasis, therefore they can be as a therapeutic target for CRISPR/Cas9 system and gene therapy.

Keywords: Brain Tumor, Immune System, CRISPR/Cas9, Gene Therapy

ICIA2018\Cancer Immunotherapy\Poster \8435 cancer immuno
8435

MiR-199a-5p may sensitize the triple-negative breast cancers to paclitaxel chemotherapy

Navideh Haghnava^{1,2}, Faezeh Asghari^{1,2}, Najibeh Shekari^{1,2}, Behzad Baradaran^{1,2}, Tohid Kazemi^{1,2*}

1. Department of Immunology, Faculty of Medicine, Tabriz University of Medical Science, Tabriz, Iran

2. Immunology Research Center, Tabriz University of Medical Science, Tabriz, Iran

Background: Paclitaxel is a chemotherapeutic agent that is used widely in the treatment of breast cancer and acts by arresting in cell cycle. We aimed in this study to investigate the effect of paclitaxel treatment on the expression level of miR-199a-5p, cell cycle arresting miRNA in breast cancer cell lines.

Methods: MCF-7, MDA-MB-231 and SKBR-3 cell lines were cultured and MTT assay was performed to determine IC₅₀ of paclitaxel. After RNA extraction and cDNA synthesis, expression level of miR-199a-5p was quantitatively evaluated using Real-Time PCR.

Results: IC₅₀ concentration for MCF-7, MDA-MB-231 and SKBR-3 was 3.5 μ M, 0.3 μ M and 4 μ M, respectively. Our results showed increased expression level of miR-199a-5p in MDA-MB-231 cell line (1.27 fold, P=0.001) and decreased expression level in two other cell lines including MCF-7 and SKBR-3 (125 fold, P< 0.0001 and 29.4 fold, P< 0.0001, respectively) in response to paclitaxel therapy.

Conclusion: Altered expression level of miR-199a-5p could be related to the IC₅₀ concentration of paclitaxel. More sensitive cell line, MDA-MB-231, showed increased expression level of miR-199a-5p after treatment with paclitaxel. In contrast, other two cell lines with higher IC₅₀ concentration showed decreased expression level of miR-199a-5p. In conclusion this miRNA could be involved in resistance/sensitivity of different types of breast cancer to paclitaxel therapy.

Keywords: miR-199a-5p, paclitaxel, breast cancer

ICIA2018\Cancer Immunotherapy\Poster \8448 cancer immuno
8448

Fas expression on CD4⁺ T cells in tumor draining lymph nodes of patients with breast cancer

Zahra Faghih¹, Yasmin Vahidi¹, Abbas Ghaderi^{1,2}

1. Shiraz Institute for Cancer Research, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

2. Department of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Fas receptor, also known as CD95 and APO-1, is a member of tumor necrosis factor α -family of death receptors which modulates T-cell based immune responses. Fas-mediated apoptosis is an important contributor in the contraction phase of antigen driven T cell responses. Fas receptor directly binds to and activates TCR components in a stimulus-dependent manner. Additionally, the interaction of CD95 (Fas) and CD95L (Fas ligand) leads to apoptosis of thymocytes with incorrect TCR rearrangement and of those that recognize self-antigens, a process called negative selection. Moreover, Fas–FasL interaction could also promote activation-induced cell death, a form of apoptosis induced by repeated TCR stimulation, responsible for the peripheral deletion of activated T cells. Recent studies revealed that most cancer cells are resistant to CD95-mediated apoptosis and this resistance play a crucial role in the pathogenesis of several malignancies. In the present study, we aimed to investigate Fas expression on CD4⁺ helper cells in tumor draining lymph nodes of breast cancer (BC) patients as the main site of immune response formation.

Methods: Mononuclear cells were isolated from lymph nodes of 53 untreated BC patients and stained for CD4 and CD95 antibodies to detect Fas expressing cells. The cells were then acquired on four-color flowcytometer and data were analyzed by CellQuest Pro software.

Results: More than 43% of CD4⁺ lymphocytes in draining lymph nodes of BC patients expressed CD95 marker though no significant difference was found in the frequency of cells with Fas expression in the patients with different clinicopathological parameters.

Conclusion: Our results indicated no direct association between Fas expression on CD4⁺ lymphocytes and breast cancer prognosis. However, its expression in respect to expression of Fas ligand on tumor cells and other immune cells should also be investigated.

Key words: Breast cancer, CD4⁺ T cells, Fas expression

ICIA2018\Cancer Immunotherapy\Poster \8586 cancer immuno
8586

Evaluation of MicroRNA181a and MicroRNA181b after suppression of PTPN22 gene in T-cell leukemia cell line (Jurkat)

Elham Baghbani¹, Behzad Baradaran¹

1. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

Background: T-cell acute lymphoblastic leukemia (T-ALL) is one of the most frequent malignancies associated to T-cells. Effectiveness interference in cancer medical care has been characterized by high efficiency and potential, induction of silencing within the advanced stages of growth, low price in compared to the opposite strategies of gene therapy, and high specificity compared to the opposite strategies of cancer therapy like chemotherapy. PTPN22 gene (protein tyrosine phosphatase, non-receptor type 22), encodes a protein, lymphoid tyrosine phosphatase (Lyp) in human. MiR-181 was concerned in regulation of the differentiation of B cells, T cells and natural killer cells during normal hemogenesis. In this study, we specifically investigated the effects of PTPN22 silencing in human acute T-cell leukemia cell line (Jurkat) and its affect in the expression of miR181a and miR181b. We have shown that PTPN22 can be considered as a potent adjuvant in T-ALL therapy.

Methods: Jurkat cells were grown in RPMI-1640 supplemented with 10% (v/v) FBS, penicillin (100 U/ml), streptomycin (100 µg/ml), and maintained at 37° C in humidified 5% CO₂ atmosphere. PTPN22 siRNA transfection was performed and then total RNA was isolated from cells. Quantitative Real-time PCR was performed by using a standard SYBR Green PCR master mix.

Results: Results of Quantitative Real-time PCR showed that downregulation of PTPN22 leads to downregulation of miR-181a and miR181-b in jurkat cell line.

Conclusion: The results showed that the expression of miR-181a and miR181-b in jurkat cell line are affected by PTPN22. Deregulation of their expression is related to development of T-cell malignancies and pathological process of leukemia.

Keywords: Jurkat, PTPN22, miR181, T-ALL

ICIA2018\Cancer Immunotherapy\Poster \8662 cancer immuno
8662

Restitution of miRNA-145 expression inhibits migration in MKN-45 gastric cancer cell line

Tahereh Zeinali¹, Dariush Shanehbandi¹, Naiyer Hosseini¹, Ali Mohammadi¹, Behzad Mansoori¹, Elham Baghbani¹, Behzad Baradaran¹

1. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Although gastric cancer (GC) is third most common cause of cancer-related death, major therapeutic advances have not been made, because GC is a polygenic disease arising as the result of multiple gene dysregulations. In addition to other protein-coding oncogenes and tumor suppressor genes, microRNAs play an important role in gastric cancer tumorigenic progression. The tumor suppressor miR-145 is ubiquitously expressed in normal tissues and body fluids but is lost or expressed at reduced levels in various forms of cancer including GC. In this study, we aim to reconstitute the miR-145 expression in MKN-45 cells and investigate the role of miR-145 in gastric cancer metastasis.

Methods: We transfected MKN-45 cells through PCMV-miR-145 plasmid vector delivery. The effects of the ectopic expression of miR-145 were examined by performing wound healing assay in vitro. Also, alterations in K-RAS and MMP-9 mRNA levels, which play significant roles in invasion, were evaluated by qRT-PCR analysis, respectively.

Results: Consequently, the results indicated that forced expression of miR-145 decreased cell migration in MKN-45 cells, with concomitant repression of K-RAS and MMP9 as putative targets of miR-145.

Conclusions: Replacement of exogenous miR-145 regain control over multiple metastasis pathways that are regulated by the endogenous miR-145, it may have a therapeutic potential to suppress gastric cancer metastasis.

Keywords: gastric cancer; restitution; miR-145 replacement therapy; metastasis

ICIA2018\Cancer Immunotherapy\Poster \8664 cancer immuno
8664

UCB CD34+Expansion and Differentiation into NK cells for immunotherapy approaches

Gholamreza Anani Sarab¹, Alireza Shekohifar²

1. Department of Immunology, Birjand University of Medical Sciences. Birjand-Iran

2. Department of Molecular Medicine., Birjand University of Medical Sciences. Birjand-Iran

Background: Human Immune system has some essential reflectors called natural killer cells which are differentiated from CD34⁺ HSCs and are active in fighting and killing infectious agents and tumor cells.

Methods: In this study we examined the serum-free expanded CD34⁺ cells and their differentiation into NK cells. This procedure was 5 weeks long and at the end of this period, the differentiated NK cells which were then characterized through immunophenotyping analysis. IFN-gamma secretion assay was also performed.

Results: According to the performed analyses the CD34⁺ differentiated NK cells were reported to be acceptable as normal NK cells based on immunophenotypical properties.

conclusion: Concludingly, the expanded CD34⁺ cells in the serum-free medium were proved to have the capability of being differentiated into NK cells and providing a good source for anti-cancer immunotherapy.

Keywords: NK cells, immunotherapy, UCB CD34⁺

ICIA2018\Cancer Immunotherapy\Poster \8666 cancer immuno
8666

Restoration of miR-143 expression increases apoptosis in MKN-45 gastric cancer cell line

Naiyer Hosseini¹, Dariush Shanebandi¹, Tahereh Zeinali¹, Nasrin Hosseini², Ali Mohammadi¹, Behzad Mansoori¹, Elham Baghbani¹, Behzad Baradaran¹

1. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

2. Azarbaijan Higher Education and Research Complex, Tabriz, Iran

Background: Gastric cancer (GC) with high incidence and mortality is a major health concern especially in the developing countries. Involvement of multiple genes in initiation and progression of GC leads to poor prognosis in this cancer. miRNAs are a class of small non-coding RNAs which negatively control the expression of messenger RNAs (mRNA). MiR-143 is a tumor suppressor miRNA which its aberrant expression has been shown in a variety of human cancers including GC. In the present study, we have tried to restore the miR-143 expression in MKN-45 cells and examine the role of miR-143 in gastric cancer apoptosis.

Methods: We transfected MKN-45 cells by pCMV-miR-143 plasmid vectors. Effects of exogenous expression of miR-143 on cell apoptosis were assessed by MTT, respectively. Furthermore, DAPI staining assay was utilized for detection of apoptosis induction. After miR-143 grafting, alterations in Caspase-3, Caspase-9 and Bax mRNA levels, which play key roles in apoptosis, were evaluated by qRT-PCR analysis.

Results: According to the outcomes, exogenous expression of miR-143 induced apoptosis in MKN-45 cells. Moreover, increased expressions of Caspase-3, Caspase-9 and Bax, as putative targets of miR-143, were observed.

Conclusion: It could be concluded that, reversing the miR-143 expression, by methods such as miRNA replacement, could be considered as an effective strategy to increase apoptosis in gastric cancer.

Keywords: gastric cancer, apoptosis, restoration, miR-143

ICIA2018\Cancer Immunotherapy\Poster \8723 cancer immunotherapy
8723

Evaluating the expression of M1 phenotype inducing miRNAs in metastatic breast cancer cell line

Maryam Moradi Chaleshtori¹, Seyed Mahmoud Hashemi², Sara Soudi³, Mojgan Bandehpour¹,
Samira Mohammadi Yeganeh¹

1. Department of Biotechnology, School of Advanced Technologies in Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

3. Department of Immunology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

Background: Breast cancer is the most frequent carcinoma in females and the second most common cause of cancer-related mortalities in women. In recent years, tumor microenvironment (TME) has attracted a lot of attention for developing more effective treatments. Among the immune cells in TME, macrophages are particularly abundant. In breast cancer, more than 50% of the tumor mass includes tumor-associated macrophages (TAMs). In the early stages of tumors, TAMs adopt the antitumor phenotype (M1) and in advanced tumors TAMs shift to a tumorigenic phenotype (M2). The aim of this study was to identify the specific miRNAs in metastatic breast cancer cells carried in exosomes and induce M1 phenotype in macrophages.

Methods: We used bioinformatics software and programs such as TargetScan, miRWalk, miRanda, as well as microarray data from GEO to find miRNAs differentially expressed between M1 and M2 macrophages. Real-Time PCR was performed to evaluate the expression of candidate miRNAs in breast cancer cell line which probably carry by exosomes to TAMs.

Results: The results of bioinformatic studies proposed candidate miRNAs with differentially expressions between M1 and M2 macrophages and also target the key genes in signaling pathways of macrophage polarization. miR-33 and miR-130 were selected among all candidate miRNAs. The results of Real-Time PCR indicated increased expression of miR-130 and decreased expression of miR-33 in metastatic breast cancer cell line.

Conclusion: According to our findings, targeting of these candidate miRNAs can be considered for reprogramming of M2 macrophages into antitumor M1 macrophages as a therapeutic strategy in breast cancer.

Keywords: Breast cancer; Tumor-Associated Macrophages; miRNA

ICIA2018\Cancer Immunotherapy\Poster \9746 cancer immuno
9746

The role of alopelopin in expressing the DNMT gene and inhibition of cancer cells

Amirhossein Alimohammadian¹, Negin Maleki²

1. Member of the brilliant talent of young and elite peoples youth club of Ardabil Islamic Azad University, Iran

2. Department of medical laboratory, Ardabil branch, Islamic Azad University, Ardabil, Iran

Background: The cytosine methylation in the CpG islets in the promoter region is an important mechanism for regulating the expression of genes, and this arrangement can play a role in the developmental and evolutionary stages and, by binding to TLR9, produces IL12, and results in gamma IFN production, which results in stimulation cytotoxicity of NK cells or can be a factor in silencing the expression of genes, especially in some cancer cells.

Methods: In breast cancer cells, the most important marker for methylation is the Her2-neu gene promoter. In this study, different concentrations of nano-alopelopin on the expression of the gene expression in DNMT were investigated using time real time PCR method.

Results: The results showed that the relative expression of DNMT1 gene was significantly reduced by applying 30ppm nano-alopelopin in breast cancer cell line.

Conclusion: Obviously, by reducing the expression of the MT gene, methylation can be reduced, and the expression of exogenous genes may be expressed again.

Keywords: CpG, alopelopin, Cancer, Methylation

ICIA2018\Cancer Immunotherapy\Poster \9827 cancer immuno
9827

Comparative study of the effects of oshexaenoic acid (DHA), Linoleic acid (LA) and Taxol in decreasing the expression of miR-10b and miR-20a oncomirs in triple-negative metastatic breast cancer cell line

Sepideh Maralbashi^{1,2}, Farhad Salari¹, Mahsa Javadian³, Najibeh Shekari³, Cynthia Aslan³, Milad Asadi³, Tohid Kazemi³

1. Department of Immunology, Faculty of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran
2. Student Research Committee, Kermanshah University of Medical Sciences, Kermanshah, Iran
3. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Molecular signatures of the effects of widely used omega-3 and omega-6 fatty acids as dietary supplements and even in cancer therapy have been studied. Of them, microRNAs, have been shown to undergo altered expression after treatment with oshexaenoic acid(DHA) and linoleic acid (LA) in different cancer types. In this study, the expression level of two well-known oncomirs, miR-10b and miR-20a, and also MHC class I polypeptide-related sequence A (MICA) as the target for miR-10b was investigated after treatment with DHA, LA, and common therapeutic agent taxol.

Methods: MDA-MB-231 cells were cultured and MTT assay was performed to determine IC50 of DHA, LA and taxol. Cells were treated by DHA(100μM), LA(50μM) and Taxol(3μM) alone or in different combinations. After RNA extraction and cDNA synthesis, the expression level of miR-10b, miR-20a and MICA were determined by quantitative real-time PCR.

Results: Our findings showed that DHA, LA, and taxol, and their combinations were led to the significantly decreased expression of miR-10b and miR-20a (all $p < 0.0001$). In contrary, MICA as the target of miR-10b was also down-regulated significantly ($p < 0.0001$).

Conclusion: Useful effects of DHA, LA and their combination in breast cancer could be interpreted in part by decreasing the expression level of oncomirs. Decreased expression of MICA in contrast to the decreased expression of its controlling miRNA, miR-10b, could be the result of altered expression of other miRNAs in treated cells.

Keywords: Breast cancer, oshexaenoic acid, miR-10b, miR-20a, MICA

ICIA2018\Cancer Immunotherapy\Poster \9832
9832

Synergistic effect of oshexaenoic acid (DHA) and Linoleic acid (LA) with Taxol in up-regulating the expression of miR-101, but not miR-342 tumor suppressor microRNAs, in HER2-positive breast cancer cell line

Cynthia Aslan^{1,2}, Mahsa Javadian¹, Najibeh Shekari¹, Sepideh Maralbashi³, Navid Shomali¹, Tohid Kazemi¹

1. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

2. Student research committee, Tabriz University of Medical Sciences, Tabriz, Iran

3. Department of Immunology, Faculty of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

Background: Breast cancer is the most common malignancy that occurs in women globally. The effect of diet and mainly dietary fat on breast cancer risk has been widely studied. The ω -3 (e.g. DHA) and ω -6 (e.g. LA) polyunsaturated fatty acids (PUFAs) are two key families of PUFAs that their usefulness as dietary supplement especially in cancer has been interested. Several studies have shown that DHA and LA alter breast cancer related microRNAs (miRs). In this study, we investigated the expression of miR-342 and miR-101 as tumor suppressor miRs in human breast cancer cell line BT-474 under treatment with DHA, LA, and also paclitaxel.

Methods: Human breast cancer BT-474 cell line was grown and treated by DHA (1, LA (50 and taxol (3 for 24h. Total mRNA was isolated and cDNA was synthesized using standard protocols. Then, expression of miR-101 and miR-342 in all treated and untreated (as control) cells was determined by quantitative real-time PCR.

Result: Our results showed that DHA, LA and taxol and also their different combinations caused significant up-regulation of the expression of miR-101 ($p < 0.0001$). In contrary, such treatments led to the significant decrease in the expression of miR-342.

Conclusion: Our results showed useful effect of DHA and LA on increasing the expression of tumor-suppressor miRNA, miR-101. As we failed to find similar results about miR-342, another tumor suppressor miRNA, it could be concluded that more studies on the wider range of tumor-suppressor miRNAs and also oncomiRs are needed to find more about molecular signatures of the useful effects of dietary fats in cancer setting.

Key words: Breast cancer, DHA, LA, microRNA

ICIA2018\Cancer Immunotherapy\Poster \9848 cancer immuno
9848

Synergistic effect of oshaxaenoic acid (DHA) with etaxel in decreasing the expression level of miR-20a, a MICA-regulating oncomiR, in metastatic gastric cancer cell line

Najibeh Shekari^{1, 2}, Mahsa Javadian¹, Sima Ghaffari³, Maedeh Moradi¹, Masoud Shirmohammadi¹, Tohid Kazemi^{1, 3*}

1. Immunology Research Center, Tabriz University of Medical Science, Tabriz, Iran
2. Student Research Committee, Tabriz University of Medical Sciences, Tabriz, Iran
3. Infectious and Tropical Diseases Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Gastric cancer is one of the most prevalent and lethal cancers all over the world. MICA (MHC class I chain-related gene A) is induced upon stress, damage or cancer and recognized by the activating receptor NKG2D on NK cells. MicroRNAs play critical roles in carcinogenesis and their relation with MICA expression has been shown. Here, we aimed to investigate the effect of etaxel, DHA and LA on the expression level of miR-20a oncomir and MICA in gastric cancer cell line, MKN45.

Methods: MKN45 cells were cultured and MTT assay was performed to determine IC₅₀ of etaxel, DHA and LA. After RNA extraction and cDNA synthesis, the expression level of miR-20a and its target MICA were determined by quantitative real-time PCR for both treated and untreated cell lines.

Results: Analysis of results showed that after treatment, expression level of MICA in all treated cell lines decreased significantly ($p < 0.0001$). While, miR-20a expression level significantly decreased in treated cells ($p < 0.05$), especially in etaxel/DHA/LA ($p < 0.0001$, and fold change = 0.08). LA led to significant up-regulation of miR-20a ($p < 0.05$, fold change = 1.12).

Conclusion: Decreased expression level of miR-20a as OncomiR in all treatments showed the efficacy of etaxel, DHA and LA therapy, especially when combined together which indicated their synergistic effects. Down-regulation of MICA, independent of miR-20a, may be the effect of other factors which might regulate its expression in MKN45 cell line.

Keywords: MICA, miR-20a, etaxel, DHA.

ICIA2018\Cancer Immunotherapy\Poster \9849

9849

Investigating the effects of osahexaenoic acid (DHA), Linoleic Acid (LA) alone or in combination with etaxel on the expression of MMP-2 and Talin-2, and their regulating microRNAs in metastatic gastric cancer cell line

Najibeh Shekari^{1, 2}, Mahsa Javadian¹, Motahhareh Ghasemi³, Parviz Farzamifar¹, Masoud Shirmohammadi¹, Tohid Kazemi^{1, 3*}

1. Immunology Research Center, Tabriz University of Medical Science, Tabriz, Iran

2. Student Research Committee, Tabriz University of Medical Sciences, Tabriz, Iran

3. Infectious and Tropical Diseases Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Gastric cancer (GC), as one of the most lethal cancers, afflicts many people every year. Efforts to improve the current therapies have been continued, and efficacy of using natural supplements including osahexaenoic acid (DHA) and Linoleic Acid (LA) has been suggested. MicroRNAs play critical roles in GC carcinogenesis. Here, we aimed to evaluate the effects of DHA, LA and etaxel on the expression level of metastasis-related genes including; MMP-2 and talin-2, and their controlling miRNAs in GC cell line, MKN45.

Methods: MKN45 cells were cultured and determination of IC50 of etaxel, DHA and LA was done by MTT test. After RNA extraction and cDNA synthesis, the expression level of miRNAs and target genes were determined by using quantitative real-time PCR.

Results: Analysis of results showed that MMP-2 expression level was decreased significantly in all treated MKN45 cells ($p<0.05$), except of those treated by LA. Expression level of talin-2 was increased significantly ($p<0.05$), but not in cells treated with etaxel and its combinations that showed decreased expression (all $p<0.001$). On the other hand, level of their related tumor-suppressor miRNAs, miR-194 and miR-30 were increased in all cells except of those treated with DHA and LA. Mir-126b increased significantly in cells treated by etaxel, DHA and etaxel/DHA ($p<0.05$).

Conclusion: Regardless of the effects of DHA and LA alone, combined usage of DHA, LA and etaxel showed better results and led to the down-regulation of MMP-2 and talin-2 in mRNA level. Also, miRNA expression was significantly up-regulated in combination setting. Increased expression level of miR-194, miR-30 and miR-126b supported useful combinatory effect of DHA, LA and etaxel in inhibition of metastasis.

Keywords: Gastric cancer, metastasis, miRNAs, DHA, LA, etaxel.



Computational Immunology and Systems Biology

Poster Presentation

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Computational Immunology and Systems Biology\Poster\5190
5190

Application of *In Silico* Approaches in the Development of New Drugs Against Infectious Diseases

Garshasb Rigi¹, Ibrahim Toktaz²

1. Department of Genetics, Faculty of Basic Sciences, University of Shahrekord, Shahrekord, Iran

2. National Institute of Genetic Engineering and Biotechnology (NIGEB), Tehran, Iran

Background: The extension of computational approaches and their usage for the bioinformatics investigation of immunity has prepared new insight into the engineering, variety, and progress of immune repertoires.

Methods: In this study we used libraries for ligand screenings were obtained a subset from drug like category from ZINC database. AutoDock vina software was used as the virtual screening software. Top successive hits were then analyzed regarding attraction, dispensation, metabolism, and expulsion confidants by FAFDrugs3 webserver and ADME and toxicity analysis. Moreover probable accessorial human protein target of successive hits and oral toxicity was checked with PROTOX web server.

Results: Four new anti-Ebola drugs were introduced by inhibition of VP40 proteins, also four chemical glycoprotein1 (GP1) inhibitors which theoretically prevented Ebola virus entrance were found.

Conclusion: We presented a ligand for inhibition of sortaseA (srtA) and two compounds against Listeriolysine-O of *Listeria monocytogenes* using molecular docking based screening.

Keywords: Computational Immunology, Drug Design, Anti Infectious Diseases



ICIA2018\Computational Immunology and Systems Biology\Poster\6322
6322

Cloning, expression and purification of humanized format of anti-VEGF single domain antibody

Nastaran Mohseni¹, Fatemeh Kazemi Lomedasht^{1*}

1. Venom & Biotherapeutics Molecules Laboratory of Biotechnology Research Center of Pasteur Institute of Iran, Tehran, Iran

Background: Over expression of Vascular Endothelial Growth Factor in tumor cells angiogenesis has been proven. Therefore, blocking and inhibiting of its function by monoclonal antibodies could be effective way to inhibit angiogenesis. Nanobodies have very high binding affinity and specificity to their target. In addition, nanobodies are able to detect hidden antigen efficiently. The camel origin of nanobodies concerns their use in human therapy. Humanization is effective way to reduce immunogenicity problems.

Methods: Here, We used *in silico* approach to design humanized format of nanobody. The 3-Dstructure of camel and humanized nanobody predicted by I-TASSER. The affinity, specificity and bioactivity of both nanobodies were evaluated.

Results: Our *in silico* results showed that both nanobodies had similar structures. The significant differences between camel and humanized nanobody was not observed.

Conclusion: All in all, the humanized Nanobody could be developed as promising candidate for use in human therapy to target pathologic angiogenesis.

Keywords: Nanobody, Antibody engineering, Humanization.

ICIA2018\Computational Immunology and Systems Biology\Poster\6327
6327

Analysis of transcription factors involved in responses of plasmacytoid dendritic cells to influenza virus

Fatemeh Hadi^{1,2}, Rasoul Godini¹, Hossein Fallahi¹

1. Department of Biology, Faculty of Sciences, Razi University, Kermanshah, Iran

2. Department of Stem Cells and Developmental Biology, Brain and Cognitive Center, Royan Institute for Stem Cell Biology and Technology, ACECR, Tehran, Iran

Background: Flu, caused by several types of influenza virus, is a contagious disease of respiratory organs which infects millions of people worldwide yearly. Plasmacytoid dendritic cells are from defense lines protecting lung against viral infections, including influenza. These cells stimulate immune responses by secretion of substantial concentrations of interferon- α . Herein, to reveal regulatory network governing their defense mechanisms, we analyzed transcriptome profile of plasmacytoid dendritic cells responding to influenza.

Methods: To do this study, we used freely available microarray dataset, GSE68849. Differentially Expressed Genes (DEGs) were determined using GEO2R tool of NCBI. We detected TFs governing the DEGs by ChEA, a database for protein-DNA interactions. We also determined all experimentally validated human TFs using TFcheckpoint database. Protein-protein interactions (from STRING database) and network analysis were applied for constructing core regulatory network and detecting hub TFs. DAVID database was utilized to reveal biological roles of the TFs.

Results: Seeking for all possible TFs, we found 75 DE-TFs in 1127 DEGs, 27 down-regulated and 28 up-regulated. However, ChEA database determined statistically significant protein-DNA interactions for only 11 TFs which based target genes FLI1, SPI1, ATF3, and KLF6 are from most important TFs. By constructing core regulatory network, we determined the TFs, such as JUN, FLI1, and RUNX1, controlling most number of the other TFs. On the other hand, annotation of the TFs showed that immune and inflammatory responses are included for some TFs.

Conclusion: Using protein-DNA and protein-protein interactions and rigorous statistical analysis, present study revealed critical TFs involved in plasmacytoid dendritic cells responding to influenza virus. The results can help to a better understanding of defense system against viral infections and might assist to find better way for treatment.

Keywords: Influenza, Network, Plasmacytoid dendritic cells, Transcription factors

ICIA2018\Computational Immunology and Systems Biology\Poster\7560
7560

Prediction and Analysis of Antigenic Determinants In HLA-G Molecule

Motahareh Ehsani¹, Mehdi Shahbazi¹, Mousa Mohammadnia afrrouzi¹

1. Department of Immunology, School of Medicine, Babol University of Medical science, Babol, Iran.

Background: Antigenic determinant is an important structural part of an antigen molecule that is also highly accessible to solvent. Generally when an antigen is recognized by immune cells including dendritic cell, its antigenic determinants are processed and presented to T lymphocyte which finally initiates immune response. In this study, Antigenic properties of HLA-G were predicted and the assessment of motifs in the aspect of antigenic potential was done.

Methods: using Predicting Antigenic Peptides (<http://imed.med.ucm.es/Tools/antigenic.pl>), the antigenic properties of HLA-G molecule were assessed and antigenic epitopes were determined. Consequently these epitopes could induce antibody response.

Results: The sequence of HLA-G with 414 amino acids has 17 antigenic determinants. According to an antigenic plot, the average antigenic propensity of this molecule was 1.0130 and only a small sequence in the N-terminal region of HLA-G showed a high average antigenic propensity.

Conclusion: As abnormal expression of HLA-G on normal renal cells is representative of renal cell carcinoma (RCC), Predicting and analyzing of its antigenic properties could open a way for designing of monoclonal antibodies in order to promote immunotherapy in the field of cancer .

Key words: HLA-G, Dendritic cell, T lymphocyte, Antigenic determinant

ICIA2018\Computational Immunology and Systems Biology\Poster\7590
7590

***In silico* study of miRNA involved in TGF- β pathway to prevent tissue fibrosis**

Faride Sadeghkhani¹, Ehsan Arefian²

1. Department of Bioengineering, School of Science and Technology, University of Tehran.

2. Molecular virology lab, Department of Microbiology, School of Biology, College of Science, University of Tehran

Background: Transforming growth factor-beta (TGF- β) isoforms are one of the most important and well known multifunctional cytokines which play the main role in wound healing and tissue repairing. It is produced by most of the parenchymal cell types but it also released by infiltrating cells such as lymphocytes, monocytes/macrophages, and platelets after wounding or inflammation. Increased TGF- β can lead to restore normal tissue architecture or may cause tissue fibrosis. Progressive fibrosis in organs like liver, kidney, lung or heart can lead to organs dysfunction and even death. Skin fibrosis caused Scleroderma and other disorders. So development in therapeutic intervention which could involve in this pathway is required to downstream fibrosis-specific genes regulation. One of the most important strategies is study and use miRNAs that have been specialized for regulating this pathway during evolution. In this study, *in silico* methods have been used to find the best interfering miRNAs.

Methods: For bioinformatics study, “KEGG” algorithms were used to identify proteins involved in TGF- β pathway including SMAD and then “Target Scan” data help us find miRNAs which have been involved in the TGF- β pathway and sort them in a score table. The most frequent and best ranking miRNAs in score table were selected.

Results: according to the score table, *hsa-miR-3613-3p* (32times) and *hsa-miR-4668-5p* (14times) have the most ability to inhibit this pathway.

Conclusion: In order to inhibit the progressive fibrosis, current study shows *hsa-miR-3613-3p* and *hsa-miR-4668-5p* could inhibit the pathway with the best total score and highest chance.

Keywords: TGF- β pathway, fibrosis, *In silico* study, Target scan.

ICIA2018\Computational Immunology and Systems Biology\Poster\7638
7638

Immuno-informatics based approaches to design a novel multi epitope-based vaccine for immune response reinforcement against Leptospirosis

Ahmad Karkhah¹, Majid Validi², Hamid Reza Nouri³

1. Student Research Committee, School of Medicine, Babol University of Medical Sciences, Babol, I.R. Iran
2. Clinical Biochemistry Research Center, Basic Health Sciences Institute, Shahrekord University of Medical Sciences, Shahrekord, I.R. Iran
3. Cellular and Molecular Biology Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, I.R. Iran

Background: Leptospirosis is known as a zoonotic disease of global importance originated from infection with the spirochete bacterium leptospira. Although several leptospirosis vaccines have been tested, the vaccination is relatively unsuccessful in clinical application despite decades of research. Therefore, this study was conducted to construct a novel multi-epitope based vaccine against leptospirosis by using Hap1, LigA, LAg42, sphH and HSP58 antigens.

Methods: T cell and IFN gamma epitopes were predicted from these antigens. In addition, to induce strong CD4⁺ helper T lymphocytes (HTLs) responses, Pan HLA DR-binding epitope (PADRE) and helper epitopes selected from Tetanus toxin fragment C (TTFrC) were applied. Moreover, for boosting immune response, Heparin-Binding Hemagglutinin (HBHA), a novel TLR4 agonist was added to the construct as an adjuvant. Finally, selected epitopes were linked together using EAAAK, GPMPG, AAY and HEYGAEALERAG linkers.

Results: Based on the predicted epitopes, a multi-epitope vaccine was constructed with 490 amino acids. Physico-chemical properties, secondary and tertiary structures, stability, intrinsic protein disorder, solubility, and allergenicity of this vaccine construct were assessed using immunoinformatics analysis. A soluble, and non-allergic protein with a molecular weight of 53.476 kDa was obtained. Further analyses validated the stability of the chimeric protein and the predicted epitopes in the chimeric vaccine indicated strong potential to induce B-cell and T-cell mediated immune response.

Conclusion: Immunoinformatics analysis showed that the modeled multi-epitope vaccine can properly stimulate the both T and B cells immune responses and could potentially be used for prophylactic or therapeutic usages.

Keywords: Leptospirosis; Immunoinformatics; Multi-epitope vaccine; Reverse vaccinology.

ICIA2018\Computational Immunology and Systems Biology\Poster\7756
7756

Evaluation of Azacitidine-Incorporated SNA as Targeted Delivery to TSDR of FoxP3 Gene by Bioinformatics Tools

Mrs.Zahra Sasanipour¹, Dr.Maryam Darayee²

1. Department of Nanotechnology, School of Advanced Medical Sciences and Technologies, Shiraz University of Medical Sciences

2. Department of Molecular medicine, School of Advanced Medical Sciences and Technologies, Shiraz University of Medical Sciences

Background: FoxP3 is a transcription factor that is responsible for the evolution of regulatory T-cells (Treg cells) and make it to perform better. Treg-specific demethylated region (TSDR) is a highly conserve locus on the FoxP3 gene that is fully demethylated in natural Tregs. Azacitidine is a chemical drug that inhibits DNA methylation. Naturally, the fetus is an external body and must be rejected by the mother's immune system but during the pregnancy, the expression level of this factor is increased in these cells that resulted in fetus keepin.

Methods: GC percent of this factor in promotor region was investigated by Endmemo and Sciencebuddies datasets. An azacitidine-incorporated SNA (spherical nucleic acid) was designed through AutoDock to bind to the TSDR of FoxP3 gene specifically. Specific binding of SNA to the TSDR was simulated by PyMOL software.

Results: Bioinformatics analysis indicated that TSDR was GC rich (67.5 %). Furthermore, azacitidine incorporation to the SNA particles was possible and this particle could bind to TSDR through targeted delivery.

Conclusion: Because of high percent of GC in TSDR, this region is susceptible to methylation and gene suppression and consequently the number of Treg cells should be decreases in *pregnant person's body* and caused to the fetus rejection and abortion as an external body. Therefore through targeted delivery of azacitidine to this region, demethylation would be occurred and lead to prevention of abortion.

Keywords: FoxP3, SNA, Azacitidine, TSDR, Drug delivery

ICIA2018\Computational Immunology and Systems Biology\Poster\7918
7918

Designing an *in Silico* chimeric multi epitope-based vaccine targeting LLO, GAPDH and tP60 for robust immune response against Listeria

Maryam Rostamtabar¹, Hamid Reza Nouri²

1. Student Research Committee, School of Medicine, Babol University of Medical Sciences, Babol, I.R. Iran

2. Cellular and Molecular Biology Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, I.R. Iran

Background: *Listeria monocytogenes* as an intracellular pathogen mainly infects pregnant and immunocompromised individuals. Therefore, a subunit vaccine targeting multiple antigens could be considered as an ideal approach for prevention and treatment of listeriosis. Currently, the most effective immunogenic antigens of *Listeria* are included Listeriolysin-O (LLO), glyceraldehyde-3-phosphate-dehydrogenase (GAPDH) and truncated form of the listerial P60 antigen (tP60).

Methods: The T_{CD8}⁺ helper lymphocytes and IFN gamma epitopes were predicted from these antigens. Moreover, Heparin-Binding Hemagglutinin (HBHA), a novel TLR4 agonist was added to the construct as an adjuvant. Then, a multi-epitope chimeric protein was constructed from these dominant epitopes. The chimeric gene structure, its mRNA, and deduced protein were analyzed. Then, the three dimensional (3D) structure was predicted using Modeller. Finally, validation of the predicted protein was assessed by Ramachandran plot statistics.

Results: Based on the predicted epitopes, a chimeric construct with 356 amino acids length was prepared by applying GPGPG linker. Physico-chemical properties, stability, intrinsic protein disorder, secondary and tertiary structures, of this construct were assessed. A soluble and non-allergic protein with a molecular weight of 31.4 kDa was obtained. The predicted 3D structure of the chimeric protein showed that most of the dominant epitopes were folded individually. The validation experiment showed that more than 88% residues of chimeric protein are located in favorable regions of the Ramachandran plot. Further analyses indicated T cell epitopes have enough potency for MHC binding and strong potential to induce the T-cell mediated immune response.

Conclusion: Immunoinformatics analysis showed that the modeled chimeric vaccine can properly stimulate the CD4⁺ and CD8⁺ immune responses and could potentially be used for prophylactic or therapeutic usages.

Keywords: Immunoinformatics, Listeriosis, Listeriolysin-O, Multi-epitope vaccine.

ICIA2018\Computational Immunology and Systems Biology\Poster\8401
8401

***In silico* determination of superficial clefts in antigen binding fragment of human IgG**

Fatemeh Hajighasemi ^{*1}, Soheila Rohani¹, Fatemeh Sefid²

1. Department of Immunology, Faculty of Medicine, Shahed University, Tehran, Iran

2. Department of Biology, Faculty of Basic Science, Shahed University, Tehran, Iran

Background: Immunoglobulins (Igs) are a group of plasma proteins consisting of heavy and light chains. Each chain contains variable and constant regions. Idiotypic determinants are epitopes sited on Igs variable region. Private idiotypic is a marker exclusive to a singular clone of B cell. As epitopes are commonly positioned close to surface clefts of proteins, recognition of Igs clefts could be very helpful in Igs epitopes determination. In current study superficial clefts in antigen binding fragment of human IgG have been defined by computational immunology.

Methods: Amino acid sequence and third structure of reference human IgG were found in PDB databank. The second IgG structure was determined by Phyre 2 software. Surface clefts in IgG antigen binding fragment have been distinguished by Isocleft finder software.

Results: Three clefts were recognized by Isocleft finder software. These clefts were located in Fab region of human IgG. First two clefts were in variable and constant domains and third one was in constant domains of Fab region. First, second and third clefts contained 93, 60 and 32 amino acids, respectively. The biggest and deepest cleft was in amino terminal (antigen binding site) of IgG molecule.

Conclusion: In present study three clefts in human IgG Fab were identified by computational immunology. The first two clefts contained many amino acids of variable domains. Thus it is very probable that private idiotypes be located in these clefts. So these clefts are useful for private idiotypes determination and generating specific anti-idiotypic monoclonal antibodies to monitor/ target clonally expanded malignant B cells.

Keywords: *In silico*, Human IgG, Clefts

ICIA2018\Computational Immunology and Systems Biology\Poster\8423

8423

Assessment of flexible regions in human IgG light chainsby computational analysisFatemeh Hajighasemi *¹, Soheyla Rohani¹, Fatemeh Sefid²1. *Department of Immunology, Faculty of Medicine, Shahed University, Tehran, Iran*2. *Department of Biology, Faculty of Basic Science, Shahed University, Tehran, Iran*

Background: Immunoglobulins (Igs) are glycoproteins produced by plasma cells, play an essential role in protecting the body against infections and destroying them. Immunoglobulin G (IgG) has an important role in defense against pathogens. The serum IgG level differs in several diseases such as infections and immunodeficiencies. Thus IgG has a very high diagnostic importance. For exact measurement of IgG, we need subtle diagnostic tools such as anti IgG- epitope specific monoclonal antibodies (MAbs). Immunogenic epitopes are beneficial for generating very proficient MAbs. More flexible regions in a molecule have more immunogenicity. Computational immunology helps in well identification of immunogenic epitopes through definition of their physiochemical properties such as flexibility. The aim of this study is evaluation of human IgG light chains flexibilityby computational immunology.

Methods: The amino acid sequence and third construction of reference human IgG was obtained in PDB database. The second IgG structure was specified by Phyre 2 software. IgG light chain flexible regions were distinguished by IEDB software.

Results: The greatest flexible positions were situated in 150 – 160 and in 165-170 amino acid sequences of IgG light chains as was determined by IEDB software.

Conclusion: According to results of this study the amino acid sequences located in 150 – 160 and in 165-170 positions which are placed in constant domain of human IgG light chains, organize the most flexible sites and hence are very useful tools for recognition of more immunogenic epitopes to producing greatly sensitive and specific anti- IgG MAbs.

Keywords: Human IgG, computational analysis, flexibility

ICIA2018\Computational Immunology and Systems Biology\Poster\8686
8686

Evaluation of operational amino acids in human immunoglobulin G light chain by computational Interpro surf software

Fatemeh Hajighasemi ^{*1}, Soheila Rohani¹, Fatemeh Sefid²

1. Department of Immunology, Faculty of Medicine, Shahed University, Tehran, Iran

2. Department of Biology, Faculty of Basic Science, Shahed University, Tehran, Iran

Background: Immunoglobulins (Igs) are bifunctional serum glycoproteins protect the body against foreign substances. Igs consist of light and heavy chains. Human immunoglobulin G light chain involves variable (VL) and constant (CL) domains. The variable domain takes part in antigen recognition and the constant domain influence on the antigen binding site conformation. Operational amino acids have a key role in function of proteins. Computational biology softwares are very useful in solving immunologic problems and well understanding of immune cells responses. Recognition of IgG operational amino acids would be useful in determination of vital amino acids in IgG defensive activities. In this study the human IgG light chain functional amino acids were identified by computational Interpro surf software.

Methods: Amino acid sequence and third structure of reference human IgG were attained from PDB database. Second IgG structure was determined via Phyre 2 software. Human IgG light chain operational amino acids were defined using Interpro surf software which is available in <http://curie.utmb.edu/prosurf.html> database.

Results: According to our data **the** operational amino acids of human IgG light chains are generally situated in 1-111 amino acids sequence exist in VL domain.

Conclusion: Our results showed that human IgG light chains functional amino acids are mostly situated in VL domain. This is consistent with well-known amino terminal function of IgG (antigen binding). **These** functional amino acids would be helpful in recognition of IgG idiotypic determinants which are located in its variable regions as well as detection of some IgG dysfunction and related immune disorders.

Key words: IgG, computational software, operational, amino acids

ICIA2018\Computational Immunology and Systems Biology\Poster\9752
9752

***In silico* analysis of FoxP3's isoforms and its effects**

Maryam Darayee^{1,2}, Amir Rahimi^{1,2}

1. Bioinformatics and Computational Biology Research Center, Shiraz University of Medical Sciences, Shiraz, Iran
2. Department of Molecular Medicine, School of Advanced Medical Sciences and Technologies, Shiraz University of Medical Sciences, Shiraz, Iran

Background: FoxP3 is a transcription factor responsible for the evolution and performance of regulatory T cells (Treg) which have pivotal role in the maintenance of homeostasis. T helper 17 cells (Th17) are a subset of pro-inflammatory T helper cells which related to T regulatory cells and the signals that cause Th¹⁷s to differentiate actually inhibit Treg differentiation. According to NCBI-Gene database, in human FoxP3 has two different isoforms. One isoform comprises 431 amino acids and the other has 396 amino acids which the smaller form lacking exon 2.

Methods: Protein sequencing and related analysis were collected by uniprot database. Pairwise sequence alignment of Foxp3 isoforms was done and compared with each other and the interactions between FoxP3 and other proteins was investigated in STRING database.

Results: Collected data from STRING data base showed that FoxP3 interacts with ten proteins including CTLA-4, jun, IL-17A, IL-2RA, NFATC2, IL-2, IL-6, IL-10, TGF- β , ROR γ t and ROR γ t among them binds to the second exon of full-length FoxP3. Protein sequence alignment showed that amino acids from 71 to 106 of the full-length (431 amino acids) were excluded in the shorter isoform (396 amino acids).

Conclusion: As the biological information revealed, the missed region in shorter isoform is located on the repressor domain of FoxP3 protein and attaches to transcription factors (such as ROR γ T) that are responsible for T-cells differentiation to Th17 and suppresses them. This occurrences resulted in inflammatory condition and may induce allergy or autoimmune diseases.

Keywords: FoxP3 isoforms, Th17, ROR γ t, *in silico* study

ICIA2018\Computational Immunology and Systems Biology\Poster\9760
9760

An *in silico* assay of the structure and function of HA22 anti-cancer drug

Nadia Manavi¹, Bahareh Hasani Sabzevar^{2*}, Nazanin Gholampour Faraji², Aliakbar Haddad-Mashadrizheh³, Behrouz Jalali Ghassam¹

1. Pharmaceutical Research Group, Khayyam Bioeconomy Institute, Mashhad, Iran

2. Structural biology and Bioinformatics Research Group, Khayyam Bioeconomy Institute, Mashhad, Iran

3. Cell and Molecular Biotechnology Research Group, Institute of Biotechnology, Ferdowsi University of Mashhad, Mashhad, Iran

Background: The high incidence of lymphoma and associated mortality among societies has led to the development of novel diagnostic and therapeutic methods. In this regard, In 1997 the BL22 was introduced as an immunotoxin drug to treat this type of patients, and in 2002, its optimization resulted in the production of a new form called HA22. Accordingly, structural modeling and functionality assay of this drug in quasi-physiology situation could provide approaches for more optimization which are considered in this study.

Method: NCBI, Uniprot, Protein Atlas and RCSB databases were used to retrieve desired protein sequences and structures of drug as well as the amount and hosting tissues which express antigen. Modeller software was used for modeling and assembling the fragment sequences of drug via homology modeling method. ERRAT, Verify 3D and RAMPAGE programs were employed to determine the quality of 3D structures of protein. On the other hand, GROMACS, HADDOCK and IEDB were utilized to assay the structural stability, functionality features and immunogenicity properties of drug, respectively. The PYMOL software was used to visualize structures.

Result: Assessment of the primary structure of the HA22 immunotoxin revealed the presence of RFB4 monoclonal antibodies with the ability to target CD22 antigen. Meanwhile, truncated part of *Pseudomonas* exotoxin was revealed in its context. On the other hand, the fragments assembled has led to the production of 7 models with various quality. The best model from a sustainability perspective, showed desirable curves of RMSD and RMSF after being placed in a quasi-physiological condition. However, several epitopic regions were revealed in the structure of the drug which indicating the irritation of the immune response after injection.

Conclusion: Totally, the results of this study provide a model of HA22 immunotoxin with stability structure in quasi-physiological condition, which in turn could lead to introduction of a novel isoform of drug, after immunogenicity optimization, which are considered in group.

Keywords: lymphoma, BL22, HA22, *In-silico* biology

ICIA2018\Computational Immunology and Systems Biology\Poster\9804
9804

Production of cytokinesuper agonist by mutagenesis (growth hormone as an example)

Shima Bahador, Sayyed Hamid Zarkesh Esfahani, Rahman Emamzadeh, Samira Mirbaha

Department of Biology, Faculty of Sciences, University of Isfahan, Isfahan, Iran

Background: Human growth hormone (hGH) is a single-chain peptide containing 191 amino acids. Structurally it belongs to class one cytokines. Cytokines are now widely used as modulators of the immune systems and for therapeutic purposes. High does multiple injections are usually needed to achieve and maintain the therapeutic plasma levels of cytokines which incur discomfort and high cost. Long acting cytokine super agonists which need less dose and less administration intervals are highly desirable. The aim of this study was to produce a cytokine super agonist through genetic manipulation. Human growth hormone molecules was used as a model because of its simple and well known cytokine structure and high market value.

Materials and methods: Molecule structure and binding site of the growth hormone and its receptor (GHR) and amino acids involved were analyzed using available soft-wares. Eight amino acids in site1of binding site were targeted for replacement. The new construct was designed and synthesized and then the PCR product of mutated gene was cloned into pCold expression plasmid and was transfected in *E. coli*. The mutant rhGH was produced in *E. coli* and purified using its His-tag by Nickel column. The expression and properties of the mutant protein was analyzed using ELISA, SDS page, dot and western blotting techniques and its final concentration was determined by Bradford test and then Biological activity of the mutant rhGH was tested in cell culture.

Results: High levels of mutant rhGH were produced using *E.coli* prokaryotic protein production system. Mutant rhGH showed a much higher biological activity compared to wild GH.

Conclusion: The affinity of GH to GHR and its biological activity can be increased by introducing appropriate mutations in the GH molecule. This approach can be utilized to make other cytokine super agonists.

Keywords: growth hormone, agonist, cytokine, mutation



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Diagnostic Methods in Immunology

Poster Presentation

ICIA2018\Diagnostic Methods in Immunology\Poster\11085
11085

The Changes of Hepatic Leptin, Resistance, and Adiponectin in Streptozotocin-Induced Diabetic Rats at Translational and Transcriptional Levels.

Hussam Al-Humadi^{1*}, Rafal Al-Saigh², Ahmed Al-Humadi³

1. Pharmacology and Toxicology department, Pharmacy College, University of Babylon, Babylon, Iraq

2. Clinical Laboratory sciences department, Pharmacy College, University of Babylon, Babylon, Iraq

3. Department of Pharmacology, Medical School, National and Kapodistrian University of Athens, Athens, Greece

Background: Liver is considered as a one of endocrine organs that secrete a variety of metabolically important substances including cytokines and adipokines which may represent a link between the insulin resistance and diabetes (DM). This study was to investigate hepatic leptin (LP), adiponectin (AD), and resistin (RS) in streptozotocin (STZ)-induced DM rats with the immunohistochemical (IHC) expression and real-time polymerase chain reaction (PCR).

Methods: Sixteen albino wister male rats with DM induced by single STZ administrated through intra peritoneal injection compared with control group for 30-days duration of experiment and assessing the changes in weight, insulin level (IN), blood sugar (BS), in addition to IHC and gene expression (GEX) of hepatic LP, AD, and RS; (N=8) for each group.

Results: There was a non-significant increase in weight with significant increase in BS levels of DM group ($P<0.001$) with a non-significant decrease in IN; serum liver enzymes were demonstrated a significant increase in DM group ($P<0.05$) except in ALT. Non-HDL lipid profile assay (TC, TG and LDL) revealed a significant increase ($P<0.05$) while HDL revealed no significant changes in comparison with control. Hepatic LP and RS by IHCs were significantly increased in DM group ($P<0.05$ vs. control) while LP and GEXs correlated and RS GEX was not vs. control. Although IHC expression of hepatic AD was significantly decreased in DM group ($P<0.05$) with high GEXs correlation.

Conclusions: The significant increase in hepatic IHC expression of LP and RS and decreased in AD level with highly correlation with their gene expression in STZ induced DM are augmented the features of insulin resistance with progressive dyslipidemia and abnormal liver function tests. These adipokines have very important role in drawing the scenario of diabetic pathological process and complication and ultimately in choosing an appropriate pharmacological and therapeutic approach.

Keywords: Adipocytokines, Leptin, Resistin, Adiponectin, Streptozotocin-Induced Diabetic Rats

ICIA2018\Diagnostic Methods in Immunology\Poster\11085
7612

Ethics in the Laboratory of Immunology

Tajaddin akbarzadeh khiavi

Ardabil University of Medical Sciences

Background: With the increasing number and variety of laboratory tests in recent years, the role of the immunology lab in identifying diseases is very important. Accuracy and use of modern methods of the world, as well as attention to the maintenance of rights, respect for lending and human dignity, are among the priorities that have contributed to increasing the satisfaction of clients and the characteristics of an optimal laboratory. In the Laboratory of Immunology, medical ethics is an absolute principle and professional ethics and commitment to the laboratory as expertise are influential. Undoubtedly, scientific advancement and professional ethics lead to the growth and improvement of the treatment process and the prevention of disease and the health of families and society. Morality is like a specialty with a variety of levels that can be used to provide better laboratory services in more effective ways.

- 1) Pre-test ethics include: patient's well-being and satisfaction, non-discrimination between patients, correct collection and standardization of the sample, and obtaining information and identifying patients.
- 2) Ethics in carrying out the test: All tests should be based on standard and professional skills and qualifications, and qualitative methods should not be reported in place of quantitative methods.
- 3) Post-test ethics: The most important principle in the immunology lab is the confidentiality of test results, accurate reporting of results, and the keeping of records and documentation of tests.

Keywords: Laboratory, Immunology, Ethics



Immune Cell Therapy

Poster Presentation

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Immune Cell Therapy\Poster\7483
7483

The Effect of Lipopolysaccharide on the Expression Level of Immunomodulatory and Immunostimulatory Factors of Human Amniotic Epithelial Cells

Hossein Motedayyen¹, Somayeh Ghotloo², Mohsen Masjedi¹, Nariman Mosaffa³, Abbas Mirshafiey⁴,
Mahmood Saffari^{2*}

Background: Human amniotic epithelial cells (hAECs) are a novel source of stem cells and have immunomodulatory effects on both the innate and adoptive immune systems. hAECs can differentiate into multiple cell lineages that make them a suitable cell source for regenerative medicine. These cells express multiple toll-like receptors (TLRs) and respond to various TLR ligands. This study aimed to evaluate the effect of lipopolysaccharide (LPS), a TLR4 ligand, on the level of immunomodulatory and immunostimulatory factors of hAECs.

Methods: hAECs were isolated from term placentas obtained from 10 healthy women during uncomplicated elective cesarean deliveries. hAECs were cultured in the presence and absence of LPS (5 µg/ml) at 37 °C with 5% CO₂. After 6 hours of incubation, the cells were collected and RNA extraction and cDNA synthesis were performed. Afterwards, the expression level of protagelandin E2 synthase, transforming growth factor-beta1 (TGF-β1), interleukin-10 (IL-10), interleukin-1beta (IL-1β), interleukin-6 (IL-6) and tumor necrosis factor-alpha (TNF-α) in hAECs were analyzed by q-PCR.

Results: Our results indicated that LPS had the ability to up-regulate the expression of protagelandin E2 synthase and TGF-β1 in hAECs. However, there was no change in the level of IL-1β, IL-6 and IL-10 in hAECs when were stimulated with LPS. In addition, we observed TNF-α was only expressed at very low level in some of hAECs samples which its expression was independent of the effects of LPS.

Conclusion: These findings suggest that LPS may enhance the immunomodulatory effects of hAECs through up-regulating immunosuppressive factors and down-regulating pro-inflammatory cytokines in hAECs that may represent an advantageous cell source with potential applications for immunotherapy of disease with immune pathophysiology.

Keywords: Human amniotic epithelial cells (hAECs), Immunomodulatory effect, Toll-like receptors (TLRs), Lipopolysaccharide, Regenerative medicine.

ICIA2018\Immune Cell Therapy\Poster\8057

8057

Effect of STAT3 Specific Inhibition on Dendritic Cells Maturation and Function

Jahangiri A.H., Eskandaryan M., Yaftiyan A., Moazzeni S. M.

Immunology Department, Faculty of Medical Sciences Tarbiat Modares University, Tehran, Iran

Background: Dendritic cells (DCs) are the most potent antigen-presenting cells of the immune systems with apparent influences on outcome of innate and adaptive immune response. Signal transducer and activator of transcription-3 (STAT3) is a transcription factor that modulates the expression of some important genes involved in the immunostimulatory functions of DCs. STAT3 signaling is also essential for NF- κ B mediated IL-12 expression. In this study, a small molecule (S31-201) was used as the specific inhibitor of STAT3 signaling to investigate its effects on dendritic cells differentiation and function.

Methods: DCs were differentiated from mouse bone marrow (BM) cells in the presence of DCs differentiation cytokines, GM-CSF and IL-4 and their maturation was induced using the LPS as DCs maturation factor. S31-201 was added to select DCs cultures during the maturation period. The DCs maturation state was assessed by the expression of MHCII, CD40 and CD86. The expression of IL-10 and IL-12 cytokines genes were also determined by RT-PCR. The allogenic T cell response induction potency of S31-201 treated and non-treated DCs was measured using the MLR assay.

Results: Specific inhibition of STAT3, increased the expression of DCs maturation markers. Increment T-cell stimulation was observed in MLR assay using the S31-201 treated DCs. We also observed increase in IL-12 and decrease in IL-10 production after STAT3 inhibition.

Conclusion: Our results suggest that specific inhibition of STAT3 cases a clear change in DCs maturation state and produces cells with profound mature DCs phenotypic and functional characteristics which could be useful in DC based vaccination of tumors and infectious diseases.

Keywords: Dendritic cells, STAT, Cell signaling, small molecules

ICIA2018\Immune Cell Therapy\Poster\9769
9769

Mycoplasma Contamination in cell based investigation: avoidance, Detection and Eradication

Hedieh Rahmati¹, Meysam Ganjibakhsh¹, Parvaneh Farzaneh¹, Masoumeh Asadi¹, Zahra Elyasi Gorji¹, Seyed Abolhassan Shahzadeh Fazeli^{1, 2}

1. Human and Animal Cell Bank, Iranian Biological Resource Center (IBRC), ACECR and Tehran, Iran

2. Department of Molecular and Cellular Biology, Faculty of Basic Sciences and Advanced Technologies in Biology, University of Science and Culture, Tehran, Iran

Background: Contaminations in primary cell culture can make so many problems in cell culture. Therefore, an appropriate method for removal of tissue contamination is needed. It is obvious that Mycoplasma contamination causes genetic and epigenetic changes and also disturbances in metabolism pathway of cells. So, prevention and detection of mycoplasma contamination is an important issue in cell culture study.

Methods: The tissue specimen obtained and transported under sterile conditions. In this method, the contamination from the tissues (human placenta, chorionic, amnion and human oral cancer) was removed by washing the tissue using the PBS in several times. After that, tissues were immersed in antibiotics including 200 µg/ml Penicillin, 200 µg/ml Streptomycin, 100 µg/ml Gentamicin, 100 µg/ml Ciprofloxacin and 100 µg/ml Amphotericin B at 4 °C for 1 hour in sterile conditions. Next, the cells were isolated by enzymatic digestion and cultured using ciprofloxacin 100 µg/ml for 3 weeks. After that, the antibiotic was removed and these cells were analyzed for mycoplasma contamination by DNA staining, PCR and microbial direct culture.

Results: Our results showed 69% of oral cancer tissues and 50% of embryonic membranes that transported to lab had mycoplasma contamination. The results obtained from quality control laboratory showed that utilization of this method helped us to decrease the tissue contamination in less than 10 %.

Conclusion: Employing proper aseptic technique in laboratory, and occasionally utilization of antibiotic treatments can lead to save valuable cell lines which it should not be lose. Therefore, assessment of the contamination rate before employing of the cells is a practical technique to preserve costly cells.

Key word: Contamination, Cell culture, Mycoplasma

ICIA2018\Immune Cell Therapy\Poster\9729

9729

The Effect of Silymarin on the Expression of Chemokine Receptors in T Helper 1 (Th1) CellsMohadeseh Toghyani Khorasgani¹*1. Department of Immunology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran*

Background: Silymarin, a polyphenolic flavonoid derived from milk thistle (*Silybum marianum*), is known to have anti-inflammatory, hepatoprotective, and anticarcinogenic effects. The aim of this study was to investigate the effect of silymarin on the chemokine receptor of Th1 (T helper 1) cells.

Methods: Peripheral blood mononuclear cells (PBMCs) from 8 healthy individuals were activated with Concanavalin 'A' and treated with silymarin (100 μ M) or dimethyl sulfoxide (DMSO), as negative control, in a standard condition (RT: 37 and CO₂: 5%). Cells were incubated (72 hours) and then examined for the flowcytometric evaluation of chemokine receptor CXCR3 expression. Peripheral blood lymphocyte subpopulations were identified and evaluated by two-color flowcytometric analysis. A non parametric paired samples Wilcoxon test was applied to compare the grouped data. Results were expressed as the mean \pm standard deviation (SD). P-values < 0.05 were considered to indicate significant differences.

Results: Silymarin increased the expression of CXCR3 on Th1 cells. These results were significant (P = 0.017) in all samples.

Conclusion: This study provided evidence of effectiveness for silymarin on the expression of CXCR3. Therefore, in future, silymarin, regarding its lower side effects, could be used instead of other drugs such as immunosuppressive ones in autoimmune diseases.

Keywords: Silymarin, Chemokine receptor, T helper cells, Flow cytometry



14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

Immunogenetics

Poster Presentation

ICIA2018\Immunogenetics\Poster\5197

5197

Evaluation of the Situation and Function of Immunoglobulin Binding Domains from Full length and Truncated Forms of Protein AGarshasb Rigi¹, Gholamreza Ahmadian²

1. Department of Genetics, Faculty of Basic Sciences, University of Shahrekord, Shahrekord, Iran

2. Department of Industrial and Environmental Biotechnology, National Institute of Genetic Engineering and Biotechnology (NIGEB), Tehran, Iran

Background: The Staphylococcal protein A (SpA) is widely used in biotechnology to purify polyclonal and monoclonal IgG antibodies. It is also as a very important protein in medicine. The most recent use of protein A is the application of this protein as a drug against autoimmune diseases especially, Rheumatoid Arthritis (RA) and Idiopathic Thrombocytopenic Purpura (ITP).

Methods: During our study, recombinant full-length and truncated forms of this protein were expressed in *Escherichia coli*. The bioinformatics review of the <http://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi> and <http://www.wwpdb.org> databases. Molecular modeling using the I-TASSER software was done. The function of these two forms of protein A was studied using ELISA.

Results: *In silico* analyses confirms that protein A is a 42-56 kDa protein containing five homologous domains with about 58 amino acids for binding to immunoglobulin. The Immunoglobulin binding domains are displayed from N terminal of the protein, with the letters E, D, A, B and C. The results of molecular modeling on Protein A represented the IgG-connected domains and the alpha-helix and beta-sheet secondary structures. The ELISA test showed a higher activity and ability for the truncated form for binding to IgG, compared to the full-length protein. Probably, in the truncated form, due to its proper folding and the lack of extra and hydrophobic regions and the lack of incorrect accumulation, the IgG binding domains are more likely to interact with IgG and thus exhibit more activity.

Keywords: Immunoglobulin Binding Domains, Protein A, Autoimmune Diseases

ICIA2018\Immunogenetics\Poster\7368
7368

rs1883025 and rs2230806 Polymorphisms of ABCA1 Gene are Associated with Myocardial Infarction in Southern Iranians

Hamed Fouladseresht¹, Sahar Karimzadeh¹, Shahdad Khosropanah², Mehrnoosh Doroudchi¹

1. Dept. of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

2. Cardiovascular Research Center, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Coronary artery disease (CAD) is a common disease and a major cause of mortality and morbidity worldwide. Accumulation of lipids in macrophages to become foam cells is a preliminary step in CAD development. ATP-binding cassette protein A1 (ABCA1) is responsible for lipid and cholesterol efflux to extracellular environment, thereby decreasing foam cell formation. The purpose of our study was to investigate the association between ABCA1 genes intronic +1594G/A (rs1803025) and exonic +656G/A (rs2230806) polymorphisms with Myocardial Infarction (MI).

Methods: 282 MI patients and 266 healthy blood donors were included in this case-control study. The DNA extracted and the genotype of rs1883025 and rs2230806 SNPs were determined by restriction fragment length polymorphism-polymerase chain reaction (RFLP-PCR). The analysis of genetic data was performed using the SPSS19.0 software.

Results: We found a significant association between MI and different genotypes of ABCA1 gene. In patients with MI, the frequencies of AA genotype and A allele of rs1883025 were significantly higher as compared to healthy controls (AA vs. GG: OR=2.44, 95%CI=1.146-5.196, p=0.018; A vs. G: OR=1.339, 95%CI=1.028-1.745, p=0.03). Also, the frequencies of GG genotype and G allele of rs2230806 in patients were significantly higher as compared to healthy controls (GG vs. AA: OR=1.865, 95%CI=1.124-3.096, p=0.015; G vs. A: OR=1.329, 95%CI=1.041-1.696, p=0.022).

Conclusion: Our study shows for the first time, that intronic polymorphism of ABCA1 gene is associated with Myocardial Infarction. The G allele of the rs1883025 polymorphism is known to result in higher levels of HDL-C in European-American, African-American, American-Indian, Hispanic, Japanese and East-Asia population. Our results also confirms a previous report from Tehran concerning the association of rs1883025 with CAD.

Keywords: ABCA1, rs1883025, rs2230806, Myocardial Infarction

ICIA2018\Immunogenetics\Poster\7369

7369

-586G/C and -872C/T Polymorphism of ABCA1 Gene are Associated with Predisposition to Myocardial Infarction in Southern Iranians

Hamed Fouladseresht¹, Kazem Nemati¹, Sahel Khazaei¹, Mohammad Hossein Nikoo², Mohammad Javad Zibaenezhad², Mehrnoosh Doroudchi¹

1. Dept. of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

2. Cardiovascular Research Center, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

Background: ATP-binding cassette transporter 1 (ABCA1), a member of the ATP-binding cassette family, plays a critical role in the development of atherosclerosis (AS). ABCA1 plays a pivotal role in the reverse cholesterol transport from intra to extra cellular environment. The aim of this study was to evaluate the single nucleotide polymorphisms (SNPs) in ABCA1 gene, [-586G/C (rs1800976) and -872C/T (rs2422493)], in patients with Myocardial Infarction (MI).

Methods: Blood samples were collected from 282 individual with MI before receiving medical treatment and 266 age-matched healthy blood donors as a control group. The DNA extracted and the SNPs were determined by amplification refractory mutation system-polymerase chain reaction (ARMS-PCR) technique. The analysis of genetic data was performed using the SPSS19.0 software.

Results: The rs1800976 and rs2422493 polymorphisms of ABCA1 were statistically associated with increased MI risk in the studied population. In patients with MI, the frequencies of GG genotype and G allele of rs1800976 were significantly higher as compared to healthy controls (GG vs. CC: OR=2.204, 95%CI=1.275-3.809, p=0.005; G vs. C: OR=1.410, 95%CI=1.099-1.809, p=0.007). Also, the frequencies of TT genotype and T allele of rs2422493 were significantly higher as compared to healthy controls (TT vs. TC: OR=1.934, 95%CI=1.217-3.074, p=0.005; T vs. C: OR=1.489, 95%CI=1.009-2.196, p=0.044).

Conclusion: This investigation show that G allele of -586G/C position and T allele of -872C/T position of ABCA1 gene are associated with MI in southern Iranian individuals.

Keywords: ABCA1; rs1800976; rs2422493; Myocardial Infarction

ICIA2018\Immunogenetics\Poster\7383

7383

Decreased levels of Bax and Bcl-2 mRNA Expression in Peripheral Blood from Patients with Multiple Sclerosis

ArezouDabiri¹, Meysam Amidfar², Seyed Javad Hashemini³, VahidShaygannejad⁴

1. M.Sc of Immunology, Shahid Sadoughi University of Medical Sciences, International Campus, Yazd, Iran
2. Department of neuroscience, school of advanced technologies in medicine, Tehran University of medical sciences, Tehran, Iran
3. Department of immunology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran
4. Isfahan neuroscience research center, Alzahra research institute, Isfahan University of medical sciences, Isfahan, Iran

Background: Apoptosis plays an important role in normal function of immune system and impairments of apoptotic pathways especially Bcl-2 family members could be associated with multiple sclerosis (MS).

Methods: In the present study we investigated the mRNA expression of Bax and Bcl-2 in peripheral blood cells of 25 MS patients and 25 control subjects.

Results: We found a significant down regulation in the relative gene expression of Bax ($p=0.001$) and Bcl 2($p=0.02$) in white blood cells from MS Patients when compared to healthy individuals. we also observed an increased Bcl-2/Bax expression ratio in peripheral white blood cells from MS patients in compared to healthy subjects but that was not significant($p=0.437$).

Conclusion: Our data suggest the reduced gene expression patterns of pro-apoptotic Bax and anti-apoptotic Bcl-2 in white blood cells of MS patients as a peripheral marker of MS.

Keywords: Multiple sclerosis, Peripheral blood, Gene expression, Bax, Bcl-2

ICIA2018\Immunogenetics\Poster\7395

7395

Association between TRAF3IP2 rs33980500 Polymorphism and Plasma Levels of CXCL1 in Acute Myocardial Infarction

Pordel, S., SajediKhanian, M., Nikoo H., Doroudchi, M.

1. M.Sc., Department of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran
2. Department of Cardiology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran
3. Karimi, M.H., Associate Professor, Fars Blood Transfusion Center, Shiraz, Iran
4. Associate Professor, Cardiovascular Research Center, Shiraz University of Medical Sciences, Shiraz, Iran
5. Associate Professor, Department of Immunology, Shiraz University of Medical Sciences, Shiraz, Iran

Background: IL-17A and the key adaptor (Act1) in its signaling pathway play important roles in inflammatory responses in Myocardial infarction (MI). CXCL1 chemokine is one of the target genes of the IL-17A signaling pathway. Certain genotypes of rs33980500 from TRAF3IP2, which encodes Act1 adaptor, reinforce the stability of CXCL1 mRNA, which may result in higher production of the protein. We investigated association between plasma levels of CXCL1 and TRAF3IP2-rs33980500 variations in acute myocardial infarction (AMI).

Methods: Confirmed AMI patients (n=126) referring to the hospitals affiliated to the Shiraz University of Medical Sciences in a one year period were included in this study. DNA was extracted from PBMCs of venous blood. Control individuals (n=50) were recruited from among healthy blood donors of the same age range and gender. Plasma levels of CXCL1 in patients and controls were measured by ELISA test. Also TRAF3IP2-rs33980500 variation was identified using PCR-RFLP method.

Results: Statistical analysis showed a significant difference in the plasma levels of CXCL1 chemokine between MI patients and healthy controls (P=0.0006). Also we found a significant correlation between plasma contents of CXCL1 and TT genotype of TRAF3IP2-rs33980500 polymorphism in patients (P=0.04).

Conclusion: Our results indicated an association of TT genotype of TRAF3IP2-rs33980500 with the plasma levels of CXCL1 in AMI. Alteration of C to T leads to the replacement of Aspartic to Asparagine that increases the activity of TRAF2/TRAF5 pathway which causes augmentation of the stability of CXCL1 mRNA and increase in the plasma levels of CXCL1.

Keywords: Myocardial infarction, Single nucleotide polymorphism, TRAF3IP2, CXCL1

ICIA2018\Immunogenetics\Poster\7429
7429

STAT3 Polymorphisms and Acute Kidney Injury After Cardiopulmonary Bypass

Nadereh Naderi^{1, 2}, Mahsa Rahimzadeh³, Sara Aghakhani³, Mahmoud Khayatian³, Hossein Montazerghaem⁴, Farzaneh Dasturian³

1. Department of Immunology, Faculty of Medicine, Hormozgan University of Medical Sciences, Bandar Abbas, Iran
2. Food health research center, Hormozgan University of Medical Sciences, Bandar Abbas, Iran
3. Department of Biochemistry, Faculty of Medicine, Hormozgan University of Medical Sciences, Bandar Abbas, Iran
4. Cardiovascular Research Center, Hormozgan University of Medical Sciences, Bandar Abbas, Iran

Background: Cardiopulmonary bypass -associated acute kidney injury (CPB -AKI) is clearly linked to increased morbidity and mortality. We explored the association between *inflammation-associated transcription factor*, STAT3, polymorphism and the risk of CPB -AKI.

Methods: In this study, STAT3 rs744166 polymorphism were analyzed in 102 patients undergoing coronary artery bypass grafting in Jorjani heart center, Bandar Abbas, Iran. The genotypes were determined using sequence-specific primers (PCR-SSP).

Results: 39 patients met the criteria for AKI after cardiac surgery (AKI group). The remaining *patients* did not develop AKI (non-AKI group). *Our results did not show any difference* in rs744166 frequency between AKI and non-AKI groups but there was an association between rs744166 CC genotype and reduced risk of CPB -AKI in older patients (Age \geq 60).

Conclusion: The rs744166 polymorphism is significantly associated with a decreased risk of CPB -AKI in old subjects.

Key words: Acute kidney injury; Cardiac surgery; Cardiopulmonary bypass, Polymorphism, Stat3

ICIA2018\Immunogenetics\Poster\7743

7743

Association between Vitamin D Receptor Gene ApaI Polymorphism and Vitamin D Status with Hashimoto's Thyroiditis in Urmia

Amir Hossein Faghfour¹, Rasoul Zarrin¹, Morteza Bagheri², Alireza Mehdizadeh³, Parvin Ayremlou⁴

1. Department of Nutrition, School of Medicine, Urmia University of Medical Sciences

2. Cellular and Molecular Research center, Faculty of Medicine, Urmia University of Medical Sciences

3. Department of Endocrinology, Imam Khomeini Teaching Hospital, Urmia University of Medical Sciences

4. Clinical Researches Development Unit of Imam Khomeini Teaching Hospital, Urmia University of Medical Sciences

Background: Hashimoto's Thyroiditis (HT) is one of the most frequent autoimmune diseases caused by autoantibodies production against thyroglobulin and thyroid peroxidase resulting in progressive destruction of thyroid gland. Some genetic and environmental backgrounds have been linked to HT. Vitamin D and vitamin D receptor may contribute to the immune system regulation and thereby autoimmune processes in HT. The aim of this study was to examine vitamin D receptor gene ApaI (rs7975232) polymorphism and vitamin D status in HT patients in Urmia population.

Methods: A total of 105 patients with HT (16 men and 89 women; mean age=38.7±11.8 years old) and 117 healthy controls (93 men and 24 women; mean age=39.5±10.49 years old) matched by age, sex, vitamin D intakes from foods and supplements and sun-exposure habits were enrolled in this case-control study. ApaI polymorphism were genotyping by polymerase chain reaction and restriction fragment length polymorphism and serum levels of 25-hydroxyvitamin D were measured by electro chemiluminescence immunoassay.

Results: The frequencies of AA, AC, CC, A and C genotypes/alleles at ApaI marker were 31.4%, 47.6%, 21%, 55.2% and 44.8% in HT and 35.9%, 41.9%, 22.2%, 55.6% and 44.4% in controls, respectively. Genotype and allelic distribution of ApaI polymorphism were not different in two groups ($P = 0.68$ and $P = 0.95$; respectively). Vitamin D status (deficiency= ≤ 20 ng/ml; insufficiency= 20-30 ng/ml; sufficiency= ≥ 30 ng/ml) did not differ between two groups ($P=0.06$), but vitamin D deficiency were associated with increased risk of HT compared with vitamin D sufficiency ($P = 0.02$; OR = 2.32 CI 95%, 1.11-4.83). Serum levels of 25-hydroxyvitamin D in patients with "AA" genotype were higher compared with "AC" and "CC" genotype (40.03±15.45 ng/ml vs 30.39±15.43 ng/ml and 35.27±16.53 ng/ml; respectively).

Conclusion: Vitamin D status can be associated with HT susceptibility. ApaI may alter 25-hydroxyvitamin D levels in HT patients.

Keywords: Hashimoto's Thyroiditis, Vitamin D Receptor, Gene Polymorphism, Vitamin D Status

ICIA2018\Immunogenetics\Poster\7947
7947

Association of Osteopontin Gene Promoter Polymorphism (-66T>G) with Susceptibility to Multiple Sclerosis in Iranian Population.

Fahimeh Dadkhah¹, Fereshteh Alsahebhosoul², Nahid Eskandari³, Mansour Salehi⁴,
Masoud Etemadifar⁵, Mohammad Kazemi⁶

1. MSc student, Department of Immunology, School of Medicine, Isfahan University of Medical Science, Isfahan, Iran.
2. Department of Immunology, school of Medicine, Isfahan University of Medical Science, Isfahan, Iran
3. Department of Immunology, school of Medicine, Isfahan University of Medical Science, Isfahan, Iran
4. Department of Cellular & Molecular Biology, school of Medicine, Isfahan University of Medical Science, Isfahan, Iran
5. Department of Neurosciences, Al-Zahra Hospital, Isfahan University of Medical Science, Isfahan, Iran
6. Department of Cellular & Molecular Biology, school of Medicine, Isfahan University of Medical Science, Isfahan, Iran

Background: Multiple sclerosis (MS) is a chronic autoimmune disease of the central nervous system (CNS). Various factors including genetic and environmental factors are involved in the prevalence of MS. The Osteopontin (OPN) has important role in immune response, so its encoding gene could be considered as a candidate for susceptibility to MS diseases. In the present study, we evaluated the frequency of OPN (-66T>G- rs2835709T > G) polymorphism in MS patients compared with healthy controls in Isfahan, Iran population.

Methods: Blood samples were collected from 100 patients with multiple sclerosis and 100 healthy controls. After DNA extraction, OPN (rs28357094) polymorphism was detected with high resolution melt real time polymerase chain reaction (HRM Real Time PCR) technique. The results were analyzed with the SPSS software.

Result: In this study, frequency of TT genotype (OPN-rs28357094) (P = 0.020, OR = 1.85) in Patients with multiple sclerosis was higher than the control group significantly.

Conclusion: The results of this study show that rs28357094 T>G polymorphism (-66T>G) on OPN gene promoter was associated with susceptibility to multiple sclerosis in Isfahan population.

Keywords: Multiple sclerosis, Polymorphism, OPN gene

ICIA2018\Immunogenetics\Poster\8287

8287

-308 G/A Tumor Necrosis Factor Alpha Polymorphism and Coronary Artery Disease risk in Hemodialysis PatientsSaiedeh Erfanian¹, Maliheh Moradzadeh², ElhamKashafi Jahromi³*1. Department of Advanced Medical Sciences and Technologies, School of Medicine, Jahrom University of Medical Sciences (JUMS), Jahrom, Iran**2. Golestan Rheumatology Research Center, Golestan University of Medical Sciences, Gorgan, Iran**3. Food and Drug Department, Jahrom University of Medical Sciences, Jahrom, Iran*

Background: Prevalence and incidence of chronic renal failure is increasing and it has been considered as a threatening factor in the worldwide. Cardiovascular diseases has been considered as the cause of more than half of death in hemodialysis patient. It has been shown that systemic inflammation has a key role in this issue. TNF α is a mediator cytokine of inflammation response which has a key role in pathophysiology of vascular disease caused by metabolism of lipids and obesity and insulin resistance. The aim of this study was to assess the association of polymorphism -308G/A gene TNF alpha and severity and vulnerability of cardiovascular disease in hemodialysis patient.

Methods: In this case control study, 41 hemodialysis patient with cardiovascular risk factors were considered as the case group and then 41 hemodialysis patient without cardiovascular risk factors were considered as the control group. DNA extraction was done with the specific Kit for the extraction of DNA. PCR was used for detection of polymorphism -308G/A and then NcoI enzyme was added to each polymorphism for slicing and then electrophoresis was used for confirmation of enzyme slicing. Data were gathered and analyzed with SPSS v 15.

Results: there is no significant difference between the 2 group regarding genotype AA distribution and frequency of Allel A (p value>0.05).

Conclusion: The results of this study demonstrate that there is not any association between this polymorphism and cardiovascular risk in Iranian population. Other studies with larger samples are beneficent in order to detect the role of this polymorphism.

Keywords: Polymorphism -308G/A, TNF alpha, Atherosclerosis, Hemodialysis patient

ICIA2018\Immunogenetics\Poster\8354

8354

Evaluation of MiR-146a and MiR-193a Expression in Colorectal Cancer Cell Lines

Saeed noorolyai^{1,2}, Elham baghbani², Ahad Mokhtarzadeh^{2,1}, Behzad baradaran²

1. Department of Biochemistry, Higher Education Institute of Rab-Rashid, Tabriz, Iran

2. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Colorectal cancer (CRC) is a major cause of cancer related deaths, and the third most commonly diagnosed cancer worldwide. Although the exact mechanisms have not been clearly identified yet, many predisposing conditions are known to be related to the occurrence and progression of CRC, such as the lack of physical activity, diet containing few vegetables and fruits, tobacco smoking, alcohol abuse, overweight, obesity and a family history of cancer. MicroRNAs (miRNAs) are a family of endogenous, small, noncoding RNA molecules that involved in a wide range of biological processes including differentiation, proliferation, and apoptosis. Previous studies have confirmed that miR-193a and miR-146a exhibit pathogenic effects in carcinogenesis, especially in the colorectal adenocarcinoma. In this study, we evaluated the expression of miR-193a and miR-146a levels in three colorectal cancer cell lines.

Methods: HCT-116, SW480 and HT-29 cell lines were grown in RPMI-1640 supplemented with 10% (v/v) FBS, penicillin (100 U/ml), streptomycin (100 µg/ml), and maintained at 37°C in humidified 5% CO₂ atmosphere. Total RNA was extracted from cells. We used a QRT-PCR to evaluate of miR-193a and miR-146a expression levels. Quantitative Real-time PCR was performed by using a standard SYBR Green PCR master mix.

Results: Results of Quantitative Real-time PCR showed that the expression levels of miR-193a and miR-146a in HT29 cell line is lower than SW480 and HCT116 cell lines.

Conclusion: The results showed that expression of miR-193a and miR-146a in different colorectal cell lines are different and may have diverse effects on the cell lines.

Keywords: microRNA, Colorectal cancer, HT29, Quantitative Real-time PCR

ICIA2018\Immunogenetics\Poster\8494

8494

Establishment of Lymphoblastoid Collection from Multiple Sclerosis Patients

Mehrnaz Izadpanah^{1,2}, Samaneh Mahmoud Aghdam¹, Masoumeh Asadi¹, Parvaneh Farzaneh¹, Mansoureh Farhangnia¹, seyedAbolhassanShahzadeh Fazeli^{1,3}

1. Human and animal cell bank, Iranian Biological Resource Center (IBRC), ACECR, Tehran, Iran

2. School of advanced technologies in medicine, Tehran University of Medical Sciences, Tehran, Iran

3. Department of Molecular and Cellular Biology, Faculty of Basic Sciences and Advanced Technologies in biology, University of Science and Culture, Tehran, Iran

Background: Multiple Sclerosis (MS) is an inflammatory disease of the nervous system and pathologic effect of the immune system. Genetic factors play an important role in the development of the disease. The main source for the MS study and treatment is the DNA sequences analysis of related genes and their function. Preparing a human genome for basic and applied studies is not always feasible, and also, these samples do not respond to ongoing research and researchers need to re-sample. The unrestricted way to access the genome sample without re-sampling of human beings is to store unlimited storage and proliferation cells such as immortalized lymphocytes. Thus, large samples of human cells can be replicated and exploited forever, especially for DNA extraction and genetic analysis in following steps.

Methods: The blood sample is taken from the patient for MS diseases, and the lymphocyte cells are isolated with Ficholl 1.077g/m. Cells immortalized by the Epstein-Barr virus and sub cultured during 6-8 weeks. Then Lymphoblastoid cells stored at -196 ° C after identity, observing the principles of the cell bank and, if necessary, re-cultivated and exploited.

Results: 50 blood samples from MS disease patients were collected. Lymphocyte cells were isolated, immortalized with EBV, expanded and stored finally. MS Lymphoblastoid cells appeared singly or in small and large clusters. All cells were analyzed for quality control tests which are critical in banking process including bacterial, fungal, viral contamination in addition to genetic identification such as Short Tandem Repeat (STR) test.

Conclusion: By creating immortalized cell bank from MS diseases in the country, the possibility of further studies and screening will be facilitated even for the discovery of new and not discovered genes involved. In addition, the relationship between different genetic factors and polymorphisms can be reviewed and analyzed by researchers in the field of molecular and medical biology.

Keywords: Banking, Lymphoblastoid cell, Multiple Sclerosis

ICIA2018\Immunogenetics\Poster\8544

8544

Association Study between TLR4 Gene Variant (rs4986790) and Toxoplasmosis in Iran

Elnaz Asghari¹, Seyed Shapoor Reza Shojaei², Hamid Reza A. Naghash²

1. PhD students of Parasitology, Karaj Branch, Islamic Azad University, Karaj, Iran

2. Karaj Branch, Islamic Azad University, Faculty of Veterinary Medicine Karaj, Iran

Background: Variety and genetic variation are very important in the components of the human immune system that recognize and fight against foreign agents and antigens. As a result, humans differ widely in how to detect and respond to pathogens, and some people may be resistant to certain sick strains. *Toxoplasma gondii* is an obligate intracellular parasite that can invade and replicate in almost all nucleated cells of warm-blooded animals. It has a world-wide geographic distribution and is known to infect many species of birds and mammals, including approximately one third of humans. TLR4 plays a key role in the process of the innate immune response, and activates the inflammatory cell via the NF- κ B pathway by inducing the expression of a variety of cytokines and other molecules crucial to immune responses.

T. gondii infection induced the production of interferon (IFN)- β through TLR4 and MyD88 signaling. The glycosyl phosphatidylinositol (GPI) of *T. gondii* was demonstrated to trigger TLR4 signaling Pathways. The purpose of this investigation was the determination of the distribution of genotypes at single nucleotide polymorphism (SNP) of the toll-like receptor 4 (TLR4) in humans infected with *Toxoplasma gondii* and the identification of genetic changes predisposing to infection. We decided to describe in this study the prevalence rates of the genotype and alleles at the TLR4, 896 A>G SNP (rs4986790) in women infected with *T. gondii* in IRAN and compare them to the prevalence rates observed in women uninfected controls.

Methods: 35 peripheral blood samples were collected from 35 women with *Toxoplasma* infection (positive serum test) and the blood of non-sick women was also used as a control group (negative).

Results and conclusion: The results were analyzed for SPSS software for statistical analysis. GG genotype (0.070) has been obtained, resulting in a 95% confidence level for this allele, there is no significant difference between the patient and the healthy person.

Keywords: *Toxoplasma gondii*, TLR4, SNP, rs4986790, Genetic variation

ICIA2018\Immunogenetics\Poster\9774
9774

IL-33 Polymorphism rs192992 and its Association with Susceptibility to Different Pattern of Multiple Sclerosis

Mahdavi R¹, Jamali M², Rostami Rad M³

1. Department of Immunology, Birjand University of Medical Sciences, Birjand, Iran

2. Department of Immunology, Medical School, Kerman University of Medical Sciences, Kerman, Iran

3. Department of Microbiology, Medical School, Tehran University of Medical Sciences, Tehran, Iran

Background: Multiple sclerosis is the most common autoimmune demyelinating disease of the central nervous system (CNS) characterized by destruction of the myelin sheath, gliosis with progressive neurological dysfunction. Interleukine-33 (IL-33) is a cytokine with both pro- and anti- inflammatory activities that implicated in the pathogenesis of some auto immune diseases .the aim of this study was to determine single nucleotide polymorphism (SNP) of IL-33, rs1929992, in its gene in patients with multiple sclerosis (MS) and investigation of this polymorphism with susceptibility to multiple sclerosis.

Methods: In this case-control study, peripheral blood samples were collected from 140 MS patients from January 2016 to February 2017 in the Imam Reza Hospital and Neurology Department of Vali-e-Asr Hospital in Birjand and blood sample of 140 healthy subjects (people referred to the Blood Transfusion Organization) as a control group. Patients with multiple sclerosis and healthy control groups were matched for age and sex. SNP at rs1929992 was determined by polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) method.

Results: There was significant difference between both, healthy control group and patient with multiple sclerosis. The frequencies of AA genotype at SNP rs1929992 were significantly higher in patients with SPMS and PPMS as compared with control group ($P < 0.05$). However, the frequencies of AG genotype was significantly lower in patients with RRMS in comparison to the healthy group ($P < 0.05$). In patients with RRMS, PPMS and SPMS patterns, the frequencies of A allele was significantly higher than that in control group ($P < 0.003$, $P < 0.001$, $P < 0.0001$). In patients with RRMS, PPMS and SPMS pattern, the frequency of G allele was significantly lower than control group ($P < 0.003$, $P < 0.001$, $P < 0.0001$).

Conclusion: These results of the present study suggest that the SNP rs1929992 in IL33 gene, may be associated with different pattern of MS susceptibility.

Keywords: Multiple sclerosis, Interleukine-33, gene polymorphism

ICIA2018\Immunogenetics\Poster\9802
9802

Bioinformatics Study of Homologous Photolyase Enzyme in Humans and Immune Responses against Skin Cancer

SeyedSaadihosseini^{1,2*}, Elmira shiuokhi^{2,3}, Hashem Marawne¹, Ehsan Sohrabi¹

1. Department of Biotechnology Faculty of Bioscience University of ShahidBeheshti

2. Lab. of Regenerative Medicine and Biomedical Innovations, Pasteur institute of Iran

3. Department Science and Research .University of Islamic Azad .Central Tehran

Background: With attention to the damage of ozone layer, the impact of UVB rays on DNA is increasing day by day. Ultraviolet (UV) radiation is one of the main causes of skin cancer in the world. The aim of bioinformatics study of Photolyase is to find its homologues in humans and their role in DNA repair and stimulation of immunological responses.

Methods : Bioinformatics analyzes with such software F genes H, BioEdit, Mega7, Multi Align, PROTPARM, PROTSKALE, Inter Pro, UNIPROT, Signal P, Inter Pro Scan, MMDB& Cn3D, STRING and R Was performed.

Results: Analyzes showed that PHR2 is an enzyme with 579 amino acid and protein PI 8.91, molecular weight of 64304.67. Instability index (71.54) Indicate the low half-life of the protein and its expression in a specific condition in the cell and whit two cofactors, FAD cofactor and diverse classes of antenna chromophores likes 5, 10- methenyltetrahydrofolate (MTHF), 8-hydroxydeazaflavin, FAD is essential for the activity of the enzyme photolyases. Very high homology with the CRY gene in humans , That CRY has more features than PHR2 , which can be used as a genes for the damage caused by ultraviolet beam projection in the future, according to predictions about ozone depletion, and on the other hand, this gene with many genes Key involved in skin diseases and immune responses, The gene has almost all of the characteristics of PHR2, which can be used as a genes in relation to the damage caused by ultraviolet light ejection of the skin in the future, according to predictions about ozone depletion, and this gene with many key genes, including (CLOCK, PER, ARNTL, CSNK1E, FBXL3, CDKN1A, SMARCB (BRG1), which are linked to skin disease and immune responses.

Conclusion: This study showed that CRY in human has a high homology in its structure and its properties with PHR in other organisms. Also, these genes were correlated with a number of mechanisms involved in immunological responses.

Keywords: Photolyase, Skin cancer, CRY gene, Immunology

ICIA2018\Immunogenetics\Poster\9936

9936

Vitamin D Receptor (VDR) Gene Polymorphisms are not Associated with Susceptibility to Endometriosis in Iranian Women Population

Ali Akbar Delbandi^{1, 2}, Sepideh Khodaverdi³, Mohammad Jafari^{1, 2}, Zahra Moradi², Tahereh Mohammadi², Zhima Akhavansales⁴

1. Immunology Research Center, Immunology and Infectious Disease Institute, IRAN University of Medical Sciences

2. Immunology Department, School of Medicine, IRAN University of Medical Sciences

3. Endometriosis Research Center, IRAN University of Medical Sciences

4. Biotechnology Research Center, International Campus, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

Background: Susceptibility to endometriosis is influenced by a combination of genetic, hormonal, immunological, and environmental factors. Vitamin D is known to reduce inflammatory responses, and it has been shown to have regulating effects on the immune system. This effect may be influenced by genetic polymorphisms in the vitamin D receptor (VDR) gene. The aim of this study was to analyze of association between VDR gene polymorphisms with susceptibility to endometriosis in Iranian women population.

Methods: In a case control study, we evaluated the distribution of four genetic polymorphisms of VDR gene [defined by the presence of restriction endonuclease sites for FokI (F/f), BsmI (B/b), TaqI (T/t), and ApaI (A/a)] in 118 women with endometriosis and 116 matched healthy women controls. Simultaneous genotyping of polymorphisms was performed by Polymerase Chain Reaction Sequence-Specific Primers (PCR-SSP) technique that previously reported by our laboratory.

Results: No significant difference observed in the distribution of alleles and genotypes between patients and controls for any of the VDR polymorphisms.

Conclusion: Our results demonstrated that VDR polymorphisms are not determinative Immunogenetic marker for endometriosis.

Keywords: Endometriosis, Vitamin D, Vitamin D receptor, Genetic polymorphism



ICIA2018\Immunogenetics\Poster\9975

9975

The Relationship between *FGF1-1385A/G* Polymorphism and Endometriosis in an Iranian Population

Fariba Mirjalili, Ehsan Zare Mehrjardi, Seyed Morteza Seifati

Medical Biotechnology Research Center, Ashkezar Branch, Islamic Azad University, Ashkezar, Yazd, Iran

Background: Endometriosis is a chronic disease which endometrial tissue are found outside the uterine cavity. It is a polygenic and multifactorial disease that has a strong genetic base. Angiogenesis appears to be an important event in the pathophysiology of endometriosis (EM). One of the angiogenic factors, fibroblast growth factor (FGF1), play a central role in the initiation of angiogenesis. In this study, the possible association between *FGF1 -1385A/G* polymorphism and susceptibility to endometriosis in an Iranian population was investigated.

Methods: Seventy six unrelated premenopausal women with endometriosis and 106 unrelated healthy premenopausal women without endometriosis were enrolled in the study. Genomic DNA was extracted from Peripheral blood in all subjects. After extracting DNA, *FGF1 - 1385A/G* polymorphism was analyzed by PCR-RFLP.

Results: There was significant difference between frequencies of *FGF1 -1385A/G* polymorphism in case and control groups ($p=0.025$). So, this polymorphism was associated with endometriosis in our sample.

Conclusion: Further studies involving gene-environment and gene-gene interactions, particularly combination of *FGF1 -1385A/G* polymorphism and other *FGF1* gene family polymorphisms are needed.

Keywords: Endometriosis, *FGF1*, *-1385A/G* polymorphism

ICIA2018\Immunogenetics\Poster\10006

10006

TSLP and TSLPR Gene Expression Levels in Different Phases of Menstrual Cycle in Women with Endometriosis

SoolmazHabibi¹; ElhamAmirchaghmaghi^{2*}; Fariba Ramezanali.³; RezaAflatoonian³; ParvanehAfsharian¹; Maryam Shakhoseini^{1*}

1. Department of Genetics, Reproductive Biomedicine Research Center, Royan Institute for Reproductive Biomedicine, ACECR, Tehran, Iran

2. Department of Regenerative Biomedicine, Cell Science Research Center, Royan Institute for Stem Cell Biology and Technology, ACECR, Tehran, Iran

3. Department of Endocrinology and Female Infertility, Reproductive Biomedicine Research Center, Royan Institute for Reproductive Biomedicine, ACECR, Tehran, Iran

Background: Endometriosis is an Estrogen-dependent and inflammatory disease that characterized by the presence of endometriotic tissue outside the uterine cavity. It has been suggested that immunological factors play roles in the pathogenesis of endometriosis. Thymic stromal lymphopoietin (TSLP) is one of cytokines that triggers the dendritic cell-mediated Th2 response and has been suggested to promote Endometriosis. TSLP Receptor, also known as cytokine receptor-like Factor 2 (CRLF2) that forms a functional hetero dimeric complex with Interlukine-7 receptor (IL-7R) to bind TSLP. The present study aimed to elucidate *TSLP* and *TSLPR* genes expression in endometrial tissues of patients with endometriosis compared to women without endometriosis in different phases of menstrual cycle (proliferative phase /secretory phase).

Methods: This study consisted of two groups: women with endometriosis (n=15) and healthy fertile women (n=16). Women with any other uterine abnormalities were excluded. Informed written consent was obtained from all women. Endometrial samples were obtained from eutopic and normal endometrial biopsies were obtained by Pipelle ($p < 0.05$).

Results: The expression levels of *TSLP* and *TSLPR* (*IL7R* and *CRLF2*) genes in ectopic and eutopic endometrium of endometriosis group was significantly increased in comparison to normal endometrium in both proliferative and secretory phases of menstrual cycle. In addition, expression levels of *TSLPR* (*IL7R* and *CRLF2*) gene in ectopic and eutopic endometrium was significantly increased in proliferative phase in comparison to secretory phase.

Conclusion: Overexpression of *TSLP* and its receptor in endometriotic tissues may have essential role in development of endometriosis through drive Th2 immune response. Also a higher expression of *TSLPR* observed in endometriotic tissues in proliferative phase, may be secondary to higher estrogen level in this phase as endometriosis is estrogen dependent disease.

Keywords: Endometriosis, Th2 Immune Response, Thymic Stromal Lymphopoietin (TSLP), Thymic Stromal Lymphopoietin receptor (TSLPR). Menstrual cycle.



Immunohematology

Poster Presentation

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Immunohematology\poster\7504
7504

Volumetric Fluid System and Direct Absolute Counting for Platelets Micro Particles Detection in the Patients with Diabetes Mellitus type 2

Razie Mahmoodian^{1*}, Morteza Salimian², Mohsen Hamidpour³, Ahmad Gharehbaghian⁴

1. MSc, Department of Hematology and Blood Banking, School of Allied Medical Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. PhD, Paramedical Faculty, Kashan University of Medical Sciences, Kashan, Iran.

3. PhD, Department of Hematology and Blood Banking, School of Allied Medical Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran

4. PhD, Department of Hematology and Blood Banking, School of Allied Medical Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Platelet micro particles (PMPs) are submicron particles released from the membrane of activated platelets through shedding. They are involving in thrombotic problems of the diabetes mellitus type 2. As high clinical relevance, standardization of PMPs detection is required. Flow cytometry methods based on flow-rate and based on volumetric fluidic system are used as calibration factor to turn cytometer events to absolute count.

Methods: 40 patients with diabetes mellitus type 2 and 35 healthy individuals volunteers were under study. 5 ml of anti-coagulated (sodium citrate 3.8%) blood of patients and controls were collected. After sampling, Platelets Rich Plasma was separated. Absolute counting of CD41⁺- Annexin-v⁺ micro particles was evaluated by bead-based strategies and volumetric system. Isotype controls were used for each flow cytometry run.

Results: Direct Absolute Counting for Platelets Micro particles with both methods were recorded and compared. The volumetric Fluid system counts micro particles as accurate as flow-rate beads. In comparison with to healthy individuals, Diabetic group showed a significant increase in shedding of platelet micro particles in all of states with ADP stimulation. ($P < 0.001$, SD: ± 74.52)

Conclusion: The flow cytometric data show that the cytometer armed with a volumetric fluid system provides direct absolute counting and fixing the need for counting beads. The counting of this method may be even more precise and more reliable that it may result in difference in refractive index between beads and membrane vesicles. Platelets in patients with diabetes mellitus type 2 patients are pre active and they are more susceptible to released micro particles that it can be associated to metabolic environment in this patient.

Keywords: flow cytometry, Volumetric Fluid System, PMPs, diabetes mellitus type 2

ICIA2018\Immunohematology\poster\7523

7523

The effect of *Tribulusterretris* aqueous extract on production and secretion/activity of platelets related factors in healthy Subjects

Soodabeh Aghapour^{1,3}, Zahra Koolivand¹, Mehrdad Payande², Kamran Mansouri¹, Fariba sohrabi¹, Ahmad Tajemiri¹, Mohammad Alimoradi¹, Ali Mostafaie¹

1. Medical Biology Research Center, Kermanshah University of medical sciences, Kermanshah, I.R.Iran.

2. Department of Hematology and Medical Oncology, Kermanshah university of medical sciences, Kermanshah, I.R.Iran

3. Department of Immunology, Babol university of medical sciences, Babol, I.R.Iran

Background: Platelet has essential role in immune regulation, tissue repair and homeostasis. The plants of the zygophyllacea family have many effects on the immune system. The aim of this study was to investigate the effect of the aqueous extract of *Tribulusterretris* on the production and function of platelets in healthy individuals.

Methods: Aqueous extract of *Tribulusterretris* was prepared and dried and the final powder was distributed twenty healthy subjects to take as pills three times a day for fourteen consecutive days. Blood samples were then taken from individuals on before and days 1, 14 and 28 after stopping taking the pills. The platelet count (in CBC) and secretion of platelet factor-4 (PF4), Von Willebrand factor (VWF) and serotonin (assessed by Enzyme Linked ImmunoSorbent Assay) were used to analyze production and function of platelet, respectively.

Results: The Concentration of VWF and serotonin was significantly higher in 1, 14 and 28 days after than before extract usage. Additionally, the concentration of PF4 was significantly lower in 1 day after using extract. However, significant difference was not seen at 14 and 28 days after extract usage. Moreover, there was not any significant difference in platelet count before and after extract usage.

Conclusion: Findings of this study suggest that *Tribulus Terrestris* seems to be remarkable safe choice for prevention and treatment of heart disease and Idiopathic thrombocytopenic purpura (ITP).

Key words: *Tribulusterretris*, Platelet, serotonin, VWF, PF4

ICIA2018\Immunohematology\poster\8356
8356

Incidence of Factor VIII Inhibitors in Patients with Hemophilia A

Akbar Dorgalaleh¹, Shadi Tabibian¹, Yadollah Farshi², Esmail Saneei Moghaddam³, Morteza Shamsizadeh⁴

1. Department of Hematology and Blood transfusion, School of Allied Medicine, Iran University of Medical Sciences, Tehran, Iran

2. Department of Hematology and Blood transfusion, School of Allied Medicine, Tehran University of Medical Sciences, Tehran, Iran

3. Iranian Blood Transfusion Organization

4. Hamedan University of Medical Sciences

Background: Hemophilia A (HA) is an autosomal recessive X-linked disorder resulting from decrease of factor VIII (FVIII) level. In affected patients with replacement therapy, inhibitor development is induced that complicating patients management.

Methods: This study was performed in 54 patients with different forms of HA. Patients including 17, 25 and 12 patients with severe, moderate and mild forms of HA, respectively. Activated Partial Thromboplastin Time (aPTT) mixed and Bethesda tests were performed for each patient.

Results: Among these 54 patients, 12 patients had high titers FVIII inhibitor and severe form of HA and none of the patients with mild and moderate forms of HA had FVIII inhibitors.

Conclusion: It seems that FVIII inhibitor is higher in patients with severe form of HA and it could be the reason behind neutralization of FVIII and development of complications in the treatment process.

Keywords: Hemophilia, Factor VIII, Inhibitor, Replacement therapy

ICIA2018\Immunohematology\poster\8420
8420

Introduction of leuko-reduction filters as a biological research source of immunologic cells

Maryam Valizadeh¹, Maryam Ajami¹, Ali Akbar Pourfatollah^{1,2}

1. Department of immunology school of TarbiatModares university of medical faculty

2. Department of immunology school of Iranian blood transfusion organization

Background: monocyte cells and monocyte-derived dendritic cells (Mo-DCs) are one of the efficient tools for research purpose. Traditionally, peripheral blood mononuclear cells (PBMCs) isolated by density gradient centrifugation on Ficoll-Histopaque, and then differentiated to DCs.

Today, because of leukocytes side effects on the blood products, leukocytes remove from blood prior to entering the process of blood products and then use in transfusion. In order to solve this issue, now a day's leukocyte depletion by filtering has been implemented in blood banks.

After filtration process, leukoreduction filters are known as waste. In this study, we intended to retrieve viable peripheral blood leukocytes from a leukoreduction filters that currently are implemented in blood banking. If we succeed to elute high quality leukocyte, introduced to others as a big and efficient source of leukocyte cells for using in research.

Methods: After Blood sampling from eligible donors, leukofilters were removed from the blood bag and back-flushed by optimized washing buffer for leukocyte elution. Preparation of PBMCs from Filter Buffy Coats is the next step; monocyte cells isolation was done using immune-magnetic cell sorting (MACS) method. Differentiation of Mo-DCs accomplished by using cytokine treatment. Finally DCs were characterized for specific surface marker by flow cytometry.

Results: leukocyte from filter were recovered at high number (about $5-6 \times 10^9$) and viability (>95%). Phenotypic and functional assay showed no difference between filter and buffy coat derived Mo-DCs.

Conclusion: Leukoreduction Filter can be introduced as a big and efficient source of immunologic cells for using in research.

Keyword: Leukoreduction filter, Leukocyte, Dendritic cell, MACS

ICIA2018\Immunohematology\poster\8504
8504

Survey of relationship between the severity of anemia and blood transfusion requirements

Elham Khalaf Adeli¹, Ali Akbar Pourfathollah¹

1. Blood Transfusion Research Center, High Institute for Research and Education in Transfusion Medicine, Tehran, Iran

Background: Conventionally, allogeneic blood transfusion is a supportive care for anemic patients in surgical producers. However, that is related to adverse events such as transfusion-related immunomodulation (TRIM), infection, etc. We surveyed transfusion requirement considering with severity of anemia in patients undergoing cardiac surgery.

Method: 181 adult patients scheduled for elective cardiac surgery were included in a descriptive study. Preoperative Hb<13g/dl for men and Hb<12g/dl for women considered as anemia. Anemia were classified as mild (11-12.9 g/dl for male, 11-11-11.9 for female) moderate (8-10.9 g/dl), or severe (<8g/dl) based on WHO criteria. Transfusion requirements were recorded during and 24 hours after surgery.

Results: Overall, 54(29.8%) patients were anemic. Anemia was mild in 58%, moderate in 28%, and severe in 14% of patients. 81% of patients with mild anemia and all of patients (100%) with severe anemia were transfused. The prevalence of transfusion was 57% in moderate anemia condition. The severity of anemia significantly was correlated with transfusion requirements ($P=0.03$).

Conclusion: The need for blood transfusion is prevalent in severe anemia. To avoid of allogeneic blood transfusion, anemia should be correct before surgery. Furthermore using blood conservation strategies and autologous transfusion could be reduce the need for allogeneic blood transfusion.



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Immunology of Chemical Victims and Environmental Pollution

Poster Presentation

ICIA2018\Immunology of Chemical Victims and Environmental Pollution\Poster\7354
7354

Evaluation of some of the factors associated with immune system following effect of magnetic field in rats

Amin Lafzi¹, Mansour Khakpur²

1. Dvm, Veterinary Medicine Clinician , Tabriz ,Iran

2. Department Of Pathobiology, Tabriz Branch, Islamic Azad University, Tabriz, Iran

Background: This study was designed to investigate the effects of electromagnetic fields (EMF) uniformly on the immune system of 12 Norwegian male Wistar rats were used as experimental model.

Method: These rats were divided into three control, treatment one and treatment two groups. The first treatment group was subjected to electromagnetic field for four hours and fifteen days. The second group received this treatment for eight hours and fifteen days. At the end of the fifteenth day, blood samples from rats were taken and were prepared to be analyzed. Then, blood cells were counted and after conducting biochemical tests, the data were analyzed using SPSS computer application, one-way ANOVA and Tukey range tests.

Result: The results revealed that the level of WBC increased in the first experimental group compared to control group; yet, it decreased in the second experimental group. The level of total protein relatively decreased compared to control group; however, level of albumin decreased significantly and at the same time Gamma globulin increased.

Conclusion: Eventually, the increase in the number of lymphocytes and decrease in the number of neutrophils and monocytes are the most significant effects of electromagnetic field on immune system.

Keywords: Immune system; Magnetic field; Rats.

ICIA2018 Immunology of Chemical Victims and Environmental Pollution\Poster\7564
7564

The Immune statues of Regulatory T Cells in Chronic Myeloid Leukemia Chemical Victims Exposed to Sulfur Mustard

Zahra Shaker¹, Zohair Mohammad Hasan², Masume Ebtekar², Davar Amani³

1. Tarbiat Modares University, Shahid Beheshti University of Medical Sciences, AJA University of Medical Sciences
2. Tarbiat Modares University
3. Shahid Beheshti University of Medical Sciences

Background: The importance of the role of cell balances T in their health and imbalance in various diseases, especially in various types of cancers, it seems that the number of these cells increases in chronic myeloid leukemia, and in different stages of the disease, this increase is different. The suppression of the immune system in chemical injuries causes this increase and exacerbates the disease in these people.

Methods: In order to determine the number of T cells, The regulation of flow cytometry and the evaluation of the function of these cells are the amount of production of two cytokines IL-10 And TGF- β By the same cells using the technique RT-PCR in real time Done.

Results: As the findings show, in cases of forming Philadelphia Gene or oncoprotein Bcr / Abl, the number of regulatory T cells has increased significantly ($P_v < 0.05$, $P_v = 0.015$). In cases of patients who were both philadelphia positive and exposed to chemical gas of mustard, the percentage of T-reg is higher than non-exposed Philadelphia positive cases. This increase is not significant. ($P_v = 0.311$, $P_v > 0.05$) In the exposed group with mustard, the incidence of Bcr / abl onco protein has effect positive on the number of T-regs and this increases significant ($P_v = 0.001$, $P_v < 0.05$), and in non- exposed group with mustard, the occurrence of Bcr/Abl oncoprotein has no meaningful relationship ($P_v = 0.398$, $P_v < 0.05$).

Conclusion: The imbalance of T-regs number CML Can be a reliable indicator in tracking the process of treatment and the disease process. so that, compared with cytogenetic tests, the follow up and comparison of the number of these cells at different times of the disease according to peripheral blood sampling compared with the sampling Bone marrow in genetic tests is both cost-effective and time-consuming and it inflicts fewer illnesses on the patient .

Keywords: Chemical Sulfur Mustard, CML, Regulatory T cell

ICIA2018\Immunology of Chemical Victims and Environmental Pollution\Poster\7621
7621

Effect of extremely low frequency electromagnetic fields (ELF-EMFs) on serum levels of IL-6, IL-21 and expression of BCL-6, AID genes in spleens of rats

Mohammad Gholamian Hamadan,^{1,2} Mahdi AlahgholiHajibehzad,¹ Hanie Mahaki^{1,3}, Zohreh Zareighane,¹ Elaheh Talebi Ghane,⁴Alireza Zamani^{1,3}

1. Department of Immunology, School of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran.
2. Student Research Committee, Hamadan University of Medical Sciences, Hamadan, Iran
3. Molecular Immunology Research Group, Research Center for Molecular Medicine, Hamadan University of Medical Sciences, Hamadan, Iran.
4. Department of Biostatistics, School of Paramedical Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Background: The spleen is the important organ in humoral immune system against blood antigens. Some environmental factors such as extremely low frequency electromagnetic fields (ELF-EMFs) may cause harmful effects on it. The purpose of this study was assessing the role of long-term exposure of ELF-EMFs in different intensities on expression of B-cell lymphoma 6 protein (BCL-6) gene, and Activation-induced cytidine deaminase (AID) gene.

Methods: Eighty adult male rats were exposed to ELF-EMFs in intensities of 1,100,500 and 2000 μ T at frequency of 50 Hz for 2 hours daily for up to 60 days.

Results: The results showed that AID gene expression was declined in low intensity of 1 μ T compared with control group, while no changes were observed after ELF-EMFs exposure on expression levels in other intensity and other gene. Serum level of IL-6 increased in 500 μ T group in compared with other groups. Also, serum level of IL-21 have no significant change in all groups.

Conclusion: ELF-EMFs exposure does not alter differentiation of T_{FH} (Tcell Follicular Helper), the most important factor for initiation specific humoral immune system but can effect on Centrobasts.

Keywords: BCL-6, AID, IL-6, IL-21, T_{FH}, ELF-EMFs.

ICIA2018 Immunology of Chemical Victims and Environmental Pollution\Poster\7725
7725

Maternal exposure to A.lwoffii F78 prevents allergic asthma in offspring

Najmefrozannia¹

1. Department of basic science School of Medicine University of mashhad Islamic azad university

Background: Asthma is a chronic inflammatory disease. Epigenetic factors affecting the asthma phenotype have been lately investigated. Cowshed, farm animals and their products, which are found in farming environment, must have an important role in preventing asthma.

A.lwoffii f78 is a gram-negative cowshed bacterium, which causes allergoprotective effect in prenatal model of transmaternal experimental asthma protection.

Methods: For this review we searched for asthma in titles and abstracts and for A.lwoffii F78 in whole papers at PubMed and Elsevier and Google scholar. We found 25 articles 12 of which were irrelevant to our subject or had a low impact.

Results: Maternal exposure to A.lwoffii F78 prevents allergic asthma in the mice with homozygous asthmatic mother. The Exposure first causes a transient increase in proinflammatory cytokines and toll like receptor messenger RNA but later it has shown lower amounts in cord. Its effect is dependent on maternal toll like receptor signaling.

Dendritic cells stimulated by A.lwoffii f78 cause to induce T_H1 polarising gene in naïve T-cells .In the mice treated by A.lwoffii F78 T_H1 immune response shifting and decrease of T_H2 antibodies titration rather than T regulatory cells response were observed.

Exposure to A.lwoffii causes immune suppression and regulatory mechanisms by IL8, TNF, IL-12 cytokine.

Conclusion: The articles proved that the strong influence of prenatal exposure to A.lwoffii F78 would prevent allergic asthma as it was claimed in hygiene hypothesis.

Keywords: Allergy protection, A.lwoffii F78 Hygiene hypothesis, T_H1 response shifting, epigenetic

ICIA2018\Immunology of Chemical Victims and Environmental Pollution\Poster\8460
8460

Study the Effect of Genetics on MMP2 and Regulatory T cell in Chronic Myeloid Leukemia patients who Exposure to Sulfur Mustard

zahra Shaker¹, Zohair Mohammad Hasan ², EbtekarMasume ³, Davar Amani ⁴

1. Tarbiat Modares University, Shahid Beheshti University of Medical Sciences, AJA University of Medical Sciences

2, 3 Tarbiat Modares University

4. Shahid Beheshti University of Medical Sciences

Background: Philadelphia chromosome and expression of an encryption Bcr / Abl that happen in stem cells and comet impacts on the production of molecules involved in processes such as matrix metalloproteinase and cancer of blood cells, including lymphoid cell lines and T cells.

The above changes of cells and molecules in some cases even produce an oncoprotein known genetic and is not related to examine the relationship between cancer chronic myeloid leukemia in this study means CML (Chronic Myeloid Leukemia) More than 90% and in many cancers that particular oncogene or cancer is found only extends.

Methods: In this study, on patients with CML The two victims exposed to mustard and other chemical exposure cases with mustard has been done, the duration of disease was 8.6 on average during the investigation and treatment of the disease process is followed. Cytogenetic tests include karyotyping, FISH, RT-PCR in real time to obtain findings and test GEE the data used for the analysis.

Result: Some of the results from the analysis of the data suggests that in some 55/3707 and 12/3089, such as the concentration of ng Lynr MMP2 That $P < 0.05$, MMP2 Along with the occurrence of Philadelphia gene in the mustard exposure group, significantly increased the number of cells T Regulation is effective. The concept of this finding is to increase the number T-reg with variables MMP2. In some experiments, the presence of Philadelphia, will be one of the main causes of cancer metastasis and Mytlayan to CML.

Conclusion: Therefore, one of the most important therapies for lowering the concentration MMP2 or deactivating the desired molecule. Approach to the prevention of metastases could be blocking the progression of the disease.

Keywords: MMP2, Leukemia, Sulfur Mustard

ICIA2018\Immunology of Chemical Victims and Environmental Pollution\Poster\11171
11171

Evaluation of serum levels of Pro-inflammatory cytokines(TNF- α , IL1- β)in Sulfur Mustard(SM)exposed patients with long term pulmonary complications

Ashraf Vafa, Tooba Ghazanfari, Naghizadeh MM

Immunoregulation Research Center, Shahed University, Tehran, Iran

Background: Exposure to mustard gas leads to acute and chronic toxic effects on the pulmonary system. The results of various studies have been suggested the role of Pro-inflammatory cytokines including IL1- β and TNF- α in the pathogenesis of Pulmonary inflammatory diseases. IL- β and TNF- α are recognized as a key mediator of systemic inflammatory reactions and numerous changes including altered cell signaling, migration, changing on cytokine production and fever. The aim of this study is the evaluation of serum levels of IL-1 β and TNF α in SM-exposed patients with long-term pulmonary complications.

Materials: In this study, 65 SM-exposed patients as exposed group and 69unexposed people as control group were studied. The both exposed and control groups were homogeneous in terms of gender, age and body mass index. Pulmonary function test was performed by spirometry. The serum levels of IL-1 β , TNF- α were measured by ELISA method.

Results: The serum levels of TNF- α was high in SM-exposed group compared to control group. However, there were no significant differences between the serum levels of IL-1 β in SM exposed group and the control group.

Conclusion: Due to increased serum levels of TNF- α in exposed group compared to the control group, It may play an important role in pulmonary inflammatory complications. It is a proof of TNF α effects on pulmonary system as a mediator of inflammatory reactions. Although there were no significant differences in the serum levels of IL-1 β in SM-exposed group compared to the control group, It doesn't rollout the role of IL-1 β in pathogenesis of long term pulmonary complications, because it may have local changes in pulmonary system and long term complications. Therefore, locally assessment of IL-1 β would be valuable.

Keywords: Mustard Gas; TNF α ; IL-1 β ; Pulmonary Complications;



Immunology of Exercise, Aging, and Nutrition

Poster Presentation

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Immunology of Exercise, Aging, and Nutrition\Poster\6364
6364

Effects of Endurance Training on the Serum Levels of Tumour Necrosis Factor- α and Interferon- γ in Sedentary Men

Abdolreza Sotoodeh Jahromi¹, Abdossaleh Zar², Fatemeh Ahmadi³, Peter Krstrup⁴, Khosrow Ebrahim⁵, Friborz Hovanloo⁵, Davar Amani⁶

1. Research Center for Non-communicable Diseases, Immunology Department, Jahrom University of Medical Sciences, Jahrom, Iran
2. Department of Physical Education & Sport Science, University of Jahrom, Jahrom, Iran
3. Department of Physical Education & Sport Sciences, Shiraz University, Shiraz, Iran
4. Sport and Health Sciences, College of Life and Environmental Sciences, University of Exeter, Exeter, UK
5. Department of Exercise Physiology, G.C. Shahid Beheshti University, Tehran, Iran
6. Department of Immunology, Medical School, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Physical activity could be considered one of the factors that affect the immune system status and function. To find the relation between exercise and cytokines, we examined the possible effects of an 8-week endurance training program on the serum levels of cytokines, including tumour necrosis factor-alpha (TNF- α) and interferon-gamma (IFN- γ) in sedentary men.

Methods: A total of 30 healthy young male volunteers were randomly divided into an endurance training group and a control group. The training group followed a specific exercise protocol (running on a treadmill for 15~30 min at 50~70% maximal heart rate) for 8 weeks and the control group did not participate in any exercise program. Venous blood samples were collected from both the groups 24 h before and 24 h and 48 h after the exercise. Repeated ANOVA was used for statistical purposes. The serum levels of TNF- α and IFN- γ were determined by ELISA.

Results: Significant ($p < 0.05$) and non-significant ($p > 0.05$) decreases were observed in the serum levels of IFN- γ and TNF- α , respectively, after the 8-week endurance training program.

Conclusion: Our findings indicated that an 8-week endurance exercise may affect the serum levels of some inflammatory cytokines, suggesting the beneficial role of this training protocol in elderly population and people with certain conditions (inflammation of the vertebrae or other inflammatory diseases).

Keywords: Endurance Training, TNF- α , IFN- γ , Sedentary Men

ICIA2018\Immunology of Exercise, Aging, and Nutrition\Poster\7861
7861

Effect of Probiotics on Beta-Interferon Gene Expression

Mohaddase Jarrahi¹, Roya Bishe Kolaei^{1*}, Rooholla Abdollahpour²

1. Department of Biology, Islamic Azad University of Qaemshahr, Qaemshahr Iran

2. Animal Science Department, Islamic Azad University of Qaemshahr, Qaemshahr Iran

Background: Probiotics, as useful microorganisms for the health of host animals, can affect the function of digestive and immune system. However, the mechanism of beneficial effects of probiotics on immunity is still not fully understood. Since Beta-Interferon gene expression is important for improving immunity function, the present study was designed to investigate the effect of kefir Probiotic compounds on Beta-Interferon gene expression in broiler chickens intestine.

Methods: This research was conducted in a completely randomized design in four replications and two stages. The first stage was carried out at a research farm for 80 broiler chickens for 42 days. Treatments in the diet of broiler chickens included the following: 1. The first treatment without any additive in the diet as control. 2. The second treatment kefir was added to the diet of broiler chickens. In the second stage, after 42 days of chickens breeding, their samples were taken from the Jejunum tissue of the treated chicks and were transferred to the lab for molecular testing. After that RNA was extracted from the Jejunum tissue sample and the synthesis of cDNA was carried out by PCR. Finally, the samples were applied to examine gene expression by Real-time PCR using the GAPDH gene.

Result: The results showed that kefir fed to boiler chickens had significant effect on the immunity system ($P \leq 0.05$) compared with the control group. Application of kefir did not reduce beta- interferon gene expression. Therefore resistance to gastrointestinal infections did not increase.

Key words: Beta-Interferon, Probiotics, Kefir, Real time PCR.

ICIA2018\Immunology of Exercise, Aging, and Nutrition\Poster\7863
7863

Effect of Organic Acid on Beta-Interferon Gene Expression

Mohaddase Jarrahi¹, Roya Bishe Kolaei^{1*}, Rooholla Abdollahpour²

1. Department of Biology, Islamic Azad University of Qaemshahr, Qaemshahr Iran

2. Animal science Department, Islamic Azad University of Qaemshahr, Qaemshahr Iran

Background: Today healthy breeding of poultry or improving immune system against infectious diseases, antibiotics are used at therapeutic level. At the same time as they are fed, antibiotics are usually applied as growth promoters at lower levels than the treatment dose. Due to immune system resistance of the body to some antibiotics and damages to intestinal microbial balance suitable alternative has been introduced. Feeding some for improving animal immune system resistance is an alternative method. One of the most important protein produced against the immune system is Beta-Interferon, which improves cell resistance to viral infection. The purpose of this project was to investigate the effect of Organic acid on Interferon gene expression.

Methods: This research was performance as a completely randomized design in four replications and two steps. The first step was carried out in a research farm for 80 broiler chickens for 42 days; as follow: 1. The first treatment without any additive considered in diet of the control. 2. In the second treatment the organic acid (Acid fires) was added to the broiler chickens diet. In the second step, after 42 days sampling was undertaken from Jejunum tissue of the chickens under treatment and were send to the laboratory for molecular testing. RNA was extracted from the Jejunum tissues and the synthesis of cDNA was carried out by PCR. Finally the samples were applied to examine gene expression by Real-time PCR using GAPDH gene.

Result: Result showed that application organic acid (Acid fires) to feeding diet of broiler chickens significantly reduced Beta- Interferon gene expression compared to control group.

Key words: Beta-Interferon, Organic acid, Acid fires, Real time PCR

ICIA2018\Immunology of Exercise, Aging, and Nutrition\Poster\8361

8361

Effects of aerobic exercise training on serum Leptin level and insulin resistance in overweight breast cancer womenMahnaz Sourani¹, Amin Isanejad¹, Mohamadreza Khosravi Farsani²1. *Department of physical education, Shahed University*2. *Department of Medical Hematology-Oncology, Isfahan University of Medical Sciences*

Background: Because both breast cancer and the hormone leptin are associated with obesity and insulin resistance in women and with regard to the positive effects of aerobic exercise in breast cancer patients. The purpose of this study was to investigate the effect of 10 weeks aerobic exercise on serum leptin level and insulin resistance of overweight women with breast cancer.

Methods: Twenty- five overweight women with breast cancer based on inclusion criteria for participation in the study selected and randomly assigned to an exercise (n=15) or control (n=10) group. The exercise group trained aerobic exercise three times per week for 10 week at 40 -70 % maximal heart rate. The outcomes were changes in VO_{2Max} , Body composition, fasting blood sugar, insulin, leptin from baseline to post intervention. The serum leptin level was measured by ELISA. Data analyzed by independent t-test and ANCOVA ($P < 0.05$).

Results: Body mass was decreased in exercise group ($P = 0.033$), VO_{2max} increased in the exercise group ($P = 0.001$). Fasting blood sugar didn't change after 10 week of exercise training, however the insulin resistance was decreased in exercise group ($P = 0.001$). The serum leptin level decreased significantly in exercise training group ($P = 0.001$).

Conclusion: Aerobic exercise had beneficial effects on VO_{2Max} , insulin resistance, leptin in overweight women with breast cancer. Therefore aerobic exercise recommended as safe rehabilitation for breast cancer survivors.

Keywords: breast cancer, aerobic exercise, leptin, insulin resistance

ICIA2018\Immunology of Exercise, Aging, and Nutrition\Poster\8564
8564

Relationship between BMI with IL-17 and IL-23 in renal transplant patients following 10 weeks of exercise training

Somaye pooranfar¹, Mohsen Salesi², Omidreza Salehi³

1. Department, Tehran markazi Branch, Islamic Azad University, Tehran, Iran.

2. Department of Shiraz Branch, University, shiraz, Iran

3. Department of Marvdasht Branch, Islamic Azad University, Marvdasht, Iran

Background: Chronic renal failure has been considered as one of the health problems in world. In recent studies of immunology, exercise has been emphasized on the components of the immune system. The aim of this study was investigation relationship between BMI with IL-17 and IL-23 in renal transplant patients following 10 weeks of exercise training

Methods: 36 kidney transplant patients were selected and randomly divided into two groups of exercise (n=18) and control (n=18). All the patients took blood and Anthropometric test before and after training. Training group subjects participated in a 10-week training program including a 60-90 minutes training sessions three times a week. Pearson correlation (IL-23) and Spearman (IL-17). Moreover, SPSS software were used for statistical analysis of the findings ($P_{\text{value}}=0/05$).

Results: Findings showed that there was no significant relationship between the levels of BMI with IL-23 ($r = -0.19$; $r = -0.26$) and IL-17 ($r = -0.55$, $p = 0.75$) after 10 weeks of selected exercise

Conclusion: It can be concluded that there is no significant relationship between BMI with IL-17 and IL-23 following exercise in renal transplant patients.

Keywords: BMI, IL-17, IL-23, Kidney transplant patients, Exercise training.

ICIA2018\Immunology of Exercise, Aging, and Nutrition\Poster\8599
8599

An in vitro evaluation of anti-aging effect of guluronic acid (G2013) based on enzymatic oxidative stress gene expression using healthy individuals PBMCs

Mahsa Taeba, Seyed Shahabeddin Mortazavi Jahromib, Abdollah Jafarzadeha, Mohammad Reza Mirzaeic, Abbas Mirshafieyd,

1. Department of Immunology, School of Medicine, Rafsanjan University of Medical Sciences, Rafsanjan, Iran
2. Department of Cellular and Molecular Biology, Kish International Campus, University of Tehran, Kish, Iran
3. Molecular Medicine Research Center, Rafsanjan University of Medical Sciences
4. Department of Immunology, School of Public Health, Tehran University of Medical Sciences

Background: Aging is usually associated with increased levels of oxidants, and may result in damages caused by oxidative stress. There is a direct relationship between aging and increased incidence of inflammatory diseases. The present research intended to study the anti-aging and anti-inflammatory effects of the drug G2013 (guluronic acid) at low and high doses on the genes expression of a number of enzymes involved in oxidative stress (including SOD2, GPX1, CAT, GST, iNOS, and MPO) in peripheral blood mononuclear cells (PBMCs) of healthy individuals under in vitro conditions.

Methods: Venous blood samples were taken from 20 healthy individuals, the PBMCs were isolated and their RNAs extracted and their cDNAs were synthesized, and the genes expression levels were measured using the qRT-PCR technique.

Results: Our results indicated that this drug could, at both low and high doses, significantly reduce the expression of the genes for SOD2, GPX1, CAT, and GST compared to the LPS group ($p < 0.0001$). Moreover, it was noticed that the drug is able to significantly reduce gene expression levels at the high dose and at both doses (low and high), for iNOS and MPO compared to the LPS group ($p < 0.0001$), respectively.

Conclusions: The present research showed that G2013, as a novel NSAID drug with immunomodulatory properties, could modulate the expression levels of the genes for SOD2, GPX1, CAT, GST, iNOS, and MPO, to the level of healthy gene expression, and possibly it might reduce the pathological process of aging and age-related inflammatory diseases.

Keywords: G2013 Guluronic acid Anti-aging Oxidative stress NSAID Aging

ICIA2018 Immunology of Exercise, Aging, and Nutrition\Poster\9876
9876

The effect of naringin on serum TNF- α level in male rats after swimming

Vahid R Haghaniat¹, Ali R Khoshdel¹, Mohammad R Hajizadeh¹, Mehdi Mahmoodi¹,
Giti Farsi¹, Mohammad Ayubi Rad¹, Mohammad Zamanian²

1. *Department of Clinical Biochemistry, Faculty of Medicine, Rafsanjan University of Medical Sciences, Rafsanjan, Iran*

2. *Physiology-Pharmacology Research Center, Rafsanjan University of Medical Sciences, Rafsanjan, Iran*

Background: TNF- α is one of the most known circulatory inflammation marker. Exercise is considered to be a safe activity that causes moderate inflammation, the purpose of this study was to investigate the effects of a moderate-intensity exercise like swimming on the level of TNF- α in serum of male wistar rats with naringin as supplement. Naringin belongs to the flavonoid family the flavonoid naringin is found naturally in citrus fruits, especially in grapefruit, where as naringin is responsible for its bitter taste.

Methods: Twenty- four male wistar rats were randomly divided into 3 groups including; a group of exercised rats with supplement of naringin, and 2 control groups containing non-exercised control without naringin supplement and normal control rats. 160 mg/kg naringin was used as supplement. Naringin was administered orally (with use of gavage syringe) daily for 20 days. On the 20th day, the rats were killed and the level of TNF- α in the collected serum were measured.

Results: The level of TNF- α in exercised control group had significantly increased compared to non-exercised control group. Naringin supplement significantly decreased serum TNF- α level in treated group in comparison to exercised control group.

Conclusion: According to our study, swimming exercise in male rat models increase inflammation and Naringin can significantly delay that by decreasing the level of TNF- α in blood circulation.

Keywords: Naringin, Swimming, TNF- α , Inflammation

ICIA2018\Immunology of Exercise, Aging, and Nutrition\Poster\9889
9889

The effect of eight weeks aerobic and anaerobic swimming on some factors of cardiovascular disease in middle men

Mahdi Ghasemiartiyan¹, Seyed Mahmud Hejazi²

1. Assistant professor in sport physiology, Department of physical education, Mashhad branch, Islamic Azad University, Mashhad, Iran

2. Institute of Higher Education hakim Nezami, Ghuchan, Iran

General practitioner Department of Education Mashhad, Area 6, Iran

Background: The aim of this study was to compare the effects of aerobic exercise aerobic swimming on the surface of IL-6 and TNF- α and HS.CRP serum in young women.

Method: This semi-experimental study population consisted of 100 male swimmer in Mashhad that were in the age range 25 to 30 years and have had sufficient skills in 4 main swimming. 28 patients were randomly assigned to 2 groups (two groups of 14 each aerobic and anaerobic workout swim) and 10 patients in the control group. Blood samples were taken to measure the factors considered and analyzed the data using descriptive and inferential statistics, Kolmogorov-Smirnov test and ANOVA (ANOVA) was performed and the following results were obtained.

Result: Serum levels of IL-6 subjects is a significant difference. ($0.019/0 = P$). Between the levels of serum TNF- α subject is a significant difference. ($0.005/0 = p$). Between the level of HS.CRP subject is a significant difference in serum participants. ($0.001/0 = P$)

Conclusion: Based on this study, aerobic exercise and anaerobic swimming reduces the level of IL-6 and TNF- α and HS. CRP levels in subjects that this reduction was statistically significant.

Keywords: IL-6 and TNF- α and HS CRP

ICIA2018\Immunology of Exercise, Aging, and Nutrition\Poster\9999
9999

Comparison of 6-month endurance and resistance exercise on the balance of TH1, TH2, cortisol and leptin in young people

Mahdi Ghasemi artiyani¹

1. Faculty Member of Sanabad Golbahar Institute of Higher Education.
2. Hakim Nezami Institute of Higher Education, Ghuchan, Iran
3. teacher of Education Physical Education Department of Education Mashhad, Area 6, Iran

The aim of this study was to compare the effects of 12 weeks of aerobic and resistance exercise on th1 and th2 balance and serum leptin and cortisol in young men this study used a quasi-experimental data have been collected .Among healthy people and 52 volunteers in carrying out research in the age range 24 to 35 years old, Of the 100 subjects were randomly selected, Demographic information, health history, physical activity and smoking Will complete. The sample consisted of 52 young men, 20 patients who will do aerobic training (Experimental) And 20 other people will doing strength training (Experimental) 12 patients in the control group. Blood samples for measurement of these factors on ELISA (ELISA) of the three groups before and after exercise will take. Moreover, to analyze the data using descriptive statistics and Kolmogorov-Smirnov test and t-test statistical method the student will perform.



Immunology of Infectious Diseases

Bacteria & Fungi

Poster Presentation

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Immunology of Infectious Diseases Bacteria & Fungi\Poster\6357

6357

Studying the Polymorphism in the Gene Encoding Interleukin- 17 [rs4711998 (A/G)] with PCR-RFLP in Patients with Brucellosis Compared with Healthy Subjects: A Case-Control Study

Hamed Farhadi Kohan¹, Fariba Keramat^{2,3}, Hassan Mahmoudi⁴, Alireza Zamani⁵, Massoud Saidijam⁶, Mohammad Ebrahim Ghaffari⁷, Sima Kazemi⁴, Mohammad Yousef Alikhani^{2,8}

1. MSc in Microbiology, Faculty of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran
2. Brucellosis Research Center, Hamedan University of Medical Sciences, Hamedan, Iran
3. Professor, Department of Infectious Diseases, Faculty of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran
4. PhD Candidate in Microbiology, Faculty of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran
5. Professor, Department of Immunology, Faculty of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran
6. Professor, Research Center for Molecular Medicine, Hamadan University of Medical Sciences, Hamadan, Iran
7. PhD Candidate in Biostatistics, Faculty of Health, Hamadan University of Medical Sciences, Hamadan, Iran
8. Professor, Department of Microbiology, Faculty of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

Background: Brucellosis is a systemic infection caused by gram-negative coccobacilli and facultative intracellular bacteria of the genus *Brucella*. Interleukin-17 is one of the important cytokines that plays a role in controlling host immune response in patients with brucellosis. The aim of this study was to investigate polymorphism of the genes encoding IL-17 in patients with brucellosis compared to healthy subjects

Methods: This case-control study included 86 patients with brucellosis who were selected based on clinical symptoms, serology, culture and PCR results. The control group composed of 86 healthy people. The polymorphism gene encoding Interleukin-17 was evaluated in both groups by PCR-RELP method.

Results: Current study showed that the frequencies of AA [OR=0.047 (95%CI: 0.01-0.12)] and GG [OR=337.20 (95%CI: 20.49-5541.39)] were significant at position -1998 (G/A) in both cases and controls [P-values (AA) and (GG) = 0.001]. But, the frequency of AA genotypes in the control group was greater than the frequency of GG genotype in patients. The odds ratio for catching brucellosis in people who have genotype GG was 41 times higher than those who have genotype AA.

Conclusion: Findings showed that, AA and AG genotypes at -1998 (A/G) position are more important. So, the risk of brucellosis in people with GG genotype at position -1998 was higher than that in people with AA genotype. In other words, people with AA genotype are more protected against brucellosis at this position

Keywords: Interleukin-17, Cytokines, Brucellosis, Polymorphism.

ICIA2018\Immunology of Infectious Diseases Bacteria & Fungi\Poster\7762
7762

The Survey of Typhoid Fever in Suspicious Patients Who Referred to a Medical Diagnostic Laboratory in Ahvaz, Southwest of Iran (2012-2017)

Abdolhussein Shakurnia¹, Sara Iranparast¹, Asadollah Baesi², Samaneh Salehi pour², Mehri Ghafourian¹

1. Department of Immunology, School of Medicine, University of Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.

2. Department of health education, culture and research (ACECR), Branch of Khuzestan, Ahvaz, Iran

Background: Typhoid fever is a systemic infection caused by *Salmonella species* and is an important cause of illness and death, particularly among children and adolescents. Typhoid fever remains an important public health problem in the world especially in the population from developing countries. The aim of this study was to determine the prevalence of Typhoid fever amongst suspicious patients who referred to a medical diagnostic laboratory in Ahvaz, Southwest of Iran in a six years' period from 2012 to 2017.

Methods: This study was cross sectional-descriptive. In this epidemiological study, the required information of all patients who were suspected to Typhoid fever by physicians was extracted over the last 6 years by laboratory software. The diagnosis of Typhoid fever was performed using serological Widal test; and raised anti-O antibody level over titer of >1:80 was considered significant of having typhoid fever. The data was analyzed in SPSS 18 software by Chi-square tests.

Results: Of 6870 surveyed patients with the mean age of 36.83±16.74 years, 4435(64.6%) were male. The prevalence rate of typhoid fever based on serological test was 2.2%. The trend of prevalence rate of typhoid fever has increased from 0.9% to 5.1% in 2012 and 2017, respectively. (p=0.001). It was observed that prevalence of typhoid fever was higher in females with 106 (2.4%) than males 44 (1.8%) patients (p=0.066). Similarly, that prevalence of typhoid fever was observed higher in spring months (p=0.129). It was also observed that prevalence of typhoid fever was significantly higher among the patients <15 years old (p=0.034). It was concluded that prevalence of typhoid fever has increased among the youths who consume unsafe drinking water and food from outside source.

Conclusion: Typhoid fever is an important public health concern in Ahvaz, southwest Iran. This surveillance tool could have wide applications for surveillance for Typhoid fever in developing countries.

Keywords: Prevalence, Typhoid fever, Ahvaz, Southwest Iran

ICIA2018\Immunology of Infectious Diseases Bacteria & Fungi\Poster\7798
7798

Assessment of Serum Level of CXCL10 and CCL11 in Patients with Sepsis at Admission and Discharge

Mahsa Rahgoshay¹, Gholamhossein Hasanshahi², Maryam Kafizadeh³

1. Department of Paramedical Sciences, School of Paramedical Sciences, Kerman University of Medical Sciences

2. Department of Medicine, School of Medicine, Rafsanjan University of Medical Sciences

3. Department of Medicine, School of Medicine, Rafsanjan University of Medical Sciences

Background: Sepsis is a leading cause of death and the most common cause of death in the intensive care unit. CXCL10 plays a critical role in the recruitment of leukocytes and has angiostatic properties. In addition CCL11 has a potent eosinophil chemo attractant in allergic inflammation and asthma. This study aimed to compare the level of these chemokines in patients with sepsis at admission and discharge based on age and gender studies

Material and Method: This study was performed in a cohort of 33 patients with sepsis who were admitted to the Hospital. The peripheral blood sampling was performed (during admission and discharge) twice. Demographic data were collected through clinical history. The serum concentrations of CXCL10 and CXCL11 measured by ELISA. Results obtained along with demographic data were analyzed using the software SPSS, and ANOVA statistical methods

Results: The 54 patients participating in the project, 22 were male and 11 females respectively. The minimum age of participants was 45 years and a maximum of 92 years. The mean values of serum chemokines in patients with sepsis at the time of hospital discharge was significantly reduced and the differences were statistically significant ($P < 0.0001$) While CCL11 chemokine showed increased level during discharge. Compared to the overall, mean concentration of serum chemokines in patients with sepsis at admission by sex and age were approximately equal in terms of differences not statistically significant ($P = 0.193$ $P = 0.816$)

Conclusion: Our study showed that serum level of CXCL10 and CCL11 increased significantly in septic patients. After the phase of antibiotic therapy, Serum level of CXCL10 was decreased while serum level of CCL11 were remained elevated. Overall, the changes in concentrations of chemokines were not directly associated with age and sex. Finally, our study proposed that chemokines may use for the early diagnosis and treatment of patients with sepsis as a new treatment.

Keywords: Sepsis, SDF-1 α (Cell- inflammatory chemokine and angiogenesis construction)

ICIA2018\Immunology of Infectious Diseases Bacteria & Fungi\Poster\7896
7896

Improving the Delivery of HCV NS3 Protein Using Cady-2 Penetrating Peptide in HEK-293T Cell Line

Sina Alizadeh¹, Azam Bolhassani^{2*}, Shiva Irani¹, Seyed Mehdi Sadat²

1. Department of Biology, School of Basic Science, Science and Research Branch, Islamic Azad University, Tehran, Iran

2. Department of Hepatitis and AIDS, Pasteur Institute of Iran, Tehran, Iran

Background: Hepatitis C virus (HCV) is associated with several hepatic and extrahepatic disorders, including several malignancies. Many studies showed that T-cell responses to HCV nonstructural protein 3 (NS3) play an important role in clearing acute HCV infections, thus NS3 can be considered as a suitable candidate antigen for development of a HCV therapeutic vaccine. However, there are limitations for penetration of NS3 protein into the cells without a suitable delivery system. In this study, we used Cady-2 cell penetrating peptide (CPP) which harbors highly hydrophobic and positively charged residues for the controlled delivery of HCV NS3 antigen, *in vitro*.

Methods: The expression of HCV NS3 was performed using prokaryotic pET expression system, and detected by SDS-PAGE and western blotting. Its purification was carried out using affinity chromatography under denaturing condition. The NS3/ Cady-2 complexes with different molar ratios were individually formed at room temperature. The presence of the complexes was confirmed by SDS-PAGE electrophoresis. Moreover, their size and morphology were analyzed with a scanning electron microscope. The efficiency of HCV NS3 transfection using Cady-2 was evaluated by SDS-PAGE and western blotting in HEK-293T cell line.

Result: Generally, our data indicated a clear band of ~ 32 kDa for NS3 protein in SDS-PAGE and western blot analysis. SEM data confirmed the formation of discrete nanoparticles with a size of ~200-250 nm. The dominant band of ~ 32 kDa was detected in the transfected cells with NS3/Cady-2 nanoparticles using the anti-NS3 polyclonal antibody. The corresponding band was not detected in the un-transfected cells and transfected with NS3 protein alone.

Conclusion: These data suggest Cady-2 CPP as a possible effective protein delivery system for development of therapeutic HCV vaccine.

Keywords: Hepatitis C virus, Nonstructural protein 3, Cell penetrating peptide

ICIA2018\Immunology of Infectious Diseases Bacteria & Fungi\Poster\7970
7970

The Survey of Brucellosis in Suspicious Patients Who Referred to a Medical Diagnostic Laboratory in Ahvaz, Southwest of Iran (2012-2017)

Abdolhussein Shakurnia¹, Asadollah Baesi², Sara Iranparast¹, Masomeh Tehrani²

1. Department of Immunology, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

2. Department of health education, culture and research (ACECR), Branch of Khuzestan, Ahvaz, Iran

Background: Brucellosis is a bacterial disease caused by pathogenic *Brucella* species. It is a zoonotic disease causing serious public health and economic problems in many parts of the world. To address the lack of recent data on human brucellosis in the region and explore changes in the epidemiology presentation, we conducted a survey on suspected brucellosis patients. The aim of this study was to survey the Brucellosis in Suspicious Patients Who Referred to a Medical Diagnostic Laboratory in Ahvaz, Southwest of Iran in a six years' period from 2012 to 2017.

Methods: This study was retrospective cross sectional-descriptive. In this epidemiological study, the required information of all patients who were suspected to Brucellosis by physicians was extracted over the last 6 years by laboratory software. The diagnosis of Brucellosis was performed using serological Wright test; and raised antibody level over titer of >1:80 were considered significant of having Brucellosis. The data was analyzed in SPSS 18 software by Chi-square tests.

Results: Of 17872 surveyed patients with the mean age of 39.14 ± 16.43 years, 4938(27.6%) were male. According to Wright's test, 422 patients were diagnosed, the prevalence of brucellosis was 2.42% in this study. Patients had the titers of 1.80, 1.40 and 1.40 for Wright, 2-Mercapto ethanol (2ME) and Coombs tests respectively. Most participants were in the age group of 15-24 years and 69.4% were male. The lowest disease occurrence observed in winter (18%) and the highest frequency observed in autumn (28%). Frequencies of Brucellosis has gradually decreasing trend, so that the percentage of cases recorded during 2012 to 2016 was 4.2% to 1.05%, respectively. This difference was not statistically significant.

Conclusion: This study revealed that brucellosis is not prevalent in Ahvaz city. The incidence of brucellosis in the Ahvaz was less than the national average, and Khuzestan province classified to have very low incidence.

Keywords: Brucellosis, Prevalence, Epidemiology, Ahvaz, Iran

ICIA2018\Immunology of Infectious Diseases Bacteria & Fungi\Poster\8050
8050

A Survey on Goat and Sheep Milk Contamination with *Brucella Melitensis* from Urmia

Javad Alizadeh^{1*}, Amir Tukmechi²

1. M.Sc. student, Department of Microbiology, Division of Bacteriology, Faculty of Veterinary Medicine, Urmia University

2. Associate Professor, Department of Microbiology, Division of Bacteriology, Faculty of Veterinary Medicine, Urmia University

Background: Brucellosis is an important zoonosis in animal and human. The prevalence of brucellosis is high worldwide and it's endemic in some country such as Iran. Milk, milk products and meat are the most important source for Brucella. The aim of this study was to determine goat and sheep milk contamination with *Brucella melitensis* from different zone of Urmia (North, South, West and East) by milk ring test technique.

Methods: So, 202 samples were taken and transferred to the laboratory on ice immediately. At first milk samples were stored at 4 °C and their contamination with *B. melitensis* were screened by ring test.

Results: Results showed all of 202 milk samples (120 samples for sheep and 82 samples for goat) sixty one samples were contaminated with *B. melitensis*. Also, result documented that contamination in sheep and goat milk were 21.6 (26 samples) and 42.6 (35 samples) respectively.

Conclusion: It's concluded that the prevalence of *B. melitensis* in goat and sheep milk from Urmia are very high and for confirmation complementary test such as culture and polymerase chain reaction should be used.

Keywords: *Brucella melitensis*, Goat, Milk, Sheep, Urmia

ICIA2018\Immunology of Infectious Diseases Bacteria & Fungi\Poster\8107
8107

Effect of Stimulation with *Helicobacter pylori* and *Lactobacillus Acidophilus* Antigens on the Production of Various Cytokines by Patient's PBMCs with Abdominal Aortic Aneurysm in Co-culture with Endothelial Cells

Hamid Aria^{1,2}, Mehdi Kalani³, Hossein Hodjati⁴, Mehrnoosh Doroudchi¹

1. Department of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

2. Bioinformatics and Computational Biology Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

3. Prof. Alborzi Clinical microbiology research center, Nemazee hospital, Shiraz University of Medical Sciences, Shiraz, Iran

4. Department of Vascular Surgery, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Probiotics are of great interest in the prevention and treatment of metabolic and inflammatory diseases. In this study we investigated the effect of *Lactobacillus acidophilus* on the pattern of cytokine production by PBMCs in the co-culture with endothelial cells.

Methods: PBMCs were isolated from 5 men with diagnosis of AAA and 5 men with normal/insignificant angiography, CT-Scan and Ultrasonography results. EC cells were extracted from umbilical cords by collagenase method (HUVEC). PBMCs were cultured in plates with bacterial extract of CagA+ *Helicobacter pylori* with and without *Lactobacillus* extract in co-culture with HUVEC cells for 48 hrs. Then, the supernatant were removed to measure IL-2, IL-4, IL-5, IL-6, IL-9, IL-10, IL-13, IL-17A, IL-17F, IL-21, IL-22, IFN- γ and TNF- α using a commercial fluorescent-labeled bead assay.

Results: CagA+ *Helicobacter pylori* as well as *Lactobacillus acidophilus* induced higher levels of IL-9 by controls' PBMCs in co-culture with HUVECs than patients (P=0.05 and P=0.01). *Lactobacillus* induced higher production of IFN- γ by PBMCs of controls but co-culture condition and/or presence of CagA increased its production by patients' PBMCs as well. Both CagA+ and *Lactobacillus* extracts decreased IL-21 production by PBMCs' of patients in the co-culture with HUVECs but not in single culture

Conclusion: IL-21 is known to enhance survival of endothelial cells but in patients with AAA, its production by stimulated PBMCs was decreased in the co-culture. The significance of the decrease in IL-21 production in AAA development needs further investigation.

Keywords: Abdominal aortic aneurysm, Cytokines, *Helicobacter pylori*, *Lactobacillus acidophilus*

ICIA2018\Immunology of Infectious Diseases Bacteria & Fungi\Poster\8108
8108

Elevated levels of IL-6 and IL-9 in the Sera of Patients with AAA do not Correspond to their Production by Peripheral Blood Mononuclear Cells

Hamid Aria^{1,2}, Mehdi Kalani³, Hossein Hodjati⁴, Mehrnoosh Doroudchi¹

1. Department of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran
2. Bioinformatics and Computational Biology Research Center, Shiraz University of Medical Sciences, Shiraz, Iran
3. Prof. Alborzi Clinical microbiology research center, Nemazee hospital, Shiraz University of Medical Sciences, Shiraz, Iran
4. Department of Vascular Surgery, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Abdominal Aortic Aneurysm (AAA) is the local dilatation of abdominal aorta. AAA is an inflammatory condition in which cytokines may play a pathogenic role.

Methods: Peripheral Blood Mononuclear Cells (PBMCs) and serum were isolated from 5 men with diagnosis of AAA and aortic dilation greater than 5.5 centimeters, and 5 men with normal/insignificant angiography, CT-Scan and Ultrasonography results. The supernatant of PBMCs and sera were used to measure IL-2, IL-4, IL-5, IL-6, IL-9, IL-10, IL-13, IL-17A, IL-17F, IL-21, IL-22, IFN- γ and TNF- α using a fluorescent-labeled bead assay.

Results: The mean serum IL-6 and IL-9 levels were significantly higher in patients than controls ($P=0.007$ and $P=0.007$, respectively). While serum TNF- α level was not different between groups, its production by patient's PBMCs was significantly lower than controls ($P=0.047$). The mean serum levels of IL-10 and IFN- γ in patients were marginally higher than controls ($P=0.055$, $P=0.055$, respectively). Mean serum IL-2 level was not different between the groups but its production by PBMCs of patients was significantly higher than the control group ($P=0.047$).

Conclusion: Our study showed alteration in the levels of cytokines from inflammatory, Th1, Th2 and Th17 subtypes in the sera of AAA patients. The production of IL-6, IL-9, IFN- γ and IL-10, however, was not solely attributed to the PBMCs. Therefore, participation of other cells in the tissue or blood should be considered.

Keywords: Abdominal aortic aneurysm, IL-9, IL-10, Serum

ICIA2018\Immunology of Infectious Diseases Bacteria & Fungi\Poster\8177
8177

Frequency of *Staphylococcus Aureus* Nasal Carriers in Patients with Allergic Rhinitis and Healthy People

Sahar Mohammadi¹, Habib Soheyli², Ehsanollah Ghaznavi Rad^{3,1,*}

1. Department of Microbiology and Immunology, School of Medicine, Arak University of Medical Sciences, Arak, Iran

2. Department of pediatrics, School of Medicine, Arak University of Medical Sciences, Arak, Iran

3. Molecular and Medicine Research Center, Arak University of Medical Sciences, Arak, Iran

Background: Allergic rhinitis is the most common allergic disease in the breathing organ of atopic patients due to an IgE release during inflammatory dysfunction in the nasal mucosa. The most common symptoms of the disease include nasal congestion, itching and nasal congestion, sneezing and runny nose and eye pruritus. There are evidences that some microorganisms, such as *Staphylococcus aureus*, contribute to the development of allergic rhinitis due to the ability to produce toxins, enzymes and virulence factors. Therefore, the aim of this study is to evaluate the frequency of *Staphylococcus aureus* nasal carriage in patients with allergic rhinitis and healthy people.

Methods: A sample of 100 patients with allergic rhinitis who referred to the allergic clinic in Arak city was subjected to swab specimens that were taken from the anterior part of the nose and then identified by standard methods of microbiology. Also, 100 healthy people who did not have any type of allergy diagnosis were included in the study as control group.

Results: Of the 100 patients with allergic rhinitis, 17 (17%) and 20 healthy subjects (20%) were carriers of *Staphylococcus aureus*.

Conclusion: Although there was no difference in the frequency of *Staphylococcus aureus* in patients with allergic rhinitis and healthy individuals, it is necessary to evaluate the typing and virulence factors of isolated *S. aureus* for any correlation.

Keywords: *Staphylococcus aureus*, Nasal carriers, Allergic rhinitis

ICIA2018\Immunology of Infectious Diseases Bacteria & Fungi\Poster\8240
8240

The Survey of Typhoid Fever in Suspicious Patients who Referred to a Medical Diagnostic Laboratory in Ahvaz, Southwest of Iran (2012-2017)

Abdolhussein Shakurnia¹, Sara Iranparast¹, Asadollah Baesi², Samaneh Salehi pour², Mehri Ghafourian¹

1. Department of Immunology, School of Medicine, University of Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.

2. Department of health education, culture and research (ACECR), Branch of Khuzestan, Ahvaz, Iran

Background: Typhoid fever is a systemic infection caused by *Salmonella species* and is an important cause of illness and death, particularly among children and adolescents. Typhoid fever remains an important public health problem in the world especially in the population from developing countries. The aim of this study was to determine the prevalence of Typhoid fever amongst suspicious patients who referred to a medical diagnostic laboratory in Ahvaz, Southwest of Iran in a six years' period from 2012 to 2017.

Methods: This study was cross sectional-descriptive. In this epidemiological study, the required information of all patients who were suspected to Typhoid fever by physicians was extracted over the last 6 years by laboratory software. The diagnosis of Typhoid fever was performed using serological Widal test; and raised anti-O antibody level over titer of >1:80 was considered significant of having typhoid fever. The data was analyzed in SPSS 18 software by Chi-square tests.

Results: Of 6870 surveyed patients with the mean age of 36.83 ± 16.74 years, 4435 (64.6%) were male. The prevalence rate of typhoid fever based on serological test was 2.2%. The trend of prevalence rate of typhoid fever has increased from 0.9% to 5.1% in 2012 and 2017, respectively. ($p=0.001$). It was observed that prevalence of typhoid fever was higher in females with 106 (2.4%) than males 44 (1.8%) patients ($p=0.066$). Similarly, that prevalence of typhoid fever was observed higher in spring months ($p=0.129$). It was also observed that prevalence of typhoid fever was significantly higher among the patients <15 years old ($p=0.034$). It was concluded that prevalence of typhoid fever has increased among the youths who consume unsafe drinking water and food from outside source.

Conclusion: Typhoid fever is an important public health concern in Ahvaz, southwest Iran. This surveillance tool could have wide applications for surveillance for Typhoid fever in developing countries.

Keywords: Prevalence, Typhoid fever, Ahvaz, Southwest Iran

ICIA2018\Immunology of Infectious Diseases Bacteria & Fungi\Poster\8265
8265

First-line Treatment of Helicobacter Pylori in Lebanon: Comparison of Bismuth-containing Quadruple Therapy Versus 14-days Sequential Therapy

Mahdi Tarhini*¹, Mohammad Fayyad Kazan*², Mahmoud Mokbel⁴, Ghada Kchour³

1. Faculty of Health Sciences V, Lebanese University, and director of Clinical Pathology and Blood Banking Department, Raee Hospital, Saida, Lebanon

2. Institut de Biologie et de Médecine Moléculaires, Université Libre de Bruxelles, 6041 Gosselies, Belgium

3. Laboratory of Cancer Biology and Molecular Immunology, Faculty of Sciences I, Lebanese University, Hadath, Beirut, Lebanon

Gastroenterology Department, Raee Hospital, Saida, Lebanon

Background: Helicobacter Pylori (H. Pylori) is the most common cause of peptic ulcer disease (PUD) and represents a strong risk factor for gastric cancer. Treatment of H. Pylori is, therefore, a persistent need to avoid serious medical complications. Resistance to antibiotics remains to be the major challenge for H. Pylori eradication.

Methods: In this study, we determined the prevalence of H. pylori infection and evaluated H. pylori eradication efficacy of bismuth-containing quadruple therapy (Pylera) versus 14-days sequential therapy in treatment naïve-Lebanese patients. 1030 patients, showing symptoms of peptic ulcer (PU) and gastritis, underwent 14C-Urea Breath Test and esophagogastroduodenoscopy to examine H. Pylori infection and gastrointestinal disorders. Among the H. Pylori-positive patients 60 individuals were randomly selected, separated into two groups (each consisting of 30 patients) and treated with either bismuth-containing quadruple therapy or 14-days sequential therapy.

Results: We show that of the 1050 patients tested: 46.2% were H. pylori-positive, 55% had gastritis, 46.2% had both gastritis and H. pylori infection, 8.8% had gastritis but no H. pylori infection, 44.9% had neither gastritis nor H. pylori infection. Following the 14-days sequential therapy, the eradication rate was significantly higher than that obtained upon using bismuth-containing quadruple therapy [80% (24/30) versus 50% (15/30), $\chi^2=5.93$, $P=0.015$].

Conclusion: we determined H. pylori and gastritis prevalence among Lebanese PU-patients and showed that 14-days sequential therapy is more efficient than bismuth-containing quadruple therapy in terms of H. Pylori-eradication.

Keywords: H. pylori, Peptic ulcer, Pylera, Sequential Therapy.

ICIA2018\Immunology of Infectious Diseases Bacteria & Fungi\Poster\8270
8270

***Clostridium Difficile* Toxin Assay in Patients Referred to Mahdیه Hospital, Tehran, Iran**

Ezzati Mehmandust Olya Mohammad¹, Delavari Mahdi¹, Arbabi Mohsen¹, Fatemeh Goudarzi², Ebadi Maryam³

1. Department of Parasitology and Mycology, School of Medicine Kashan University of Medical Sciences, Kashan, Esfahan, Iran

2. Department of Parasitology and Mycology, School of Medicine Shiraz University of Medical Sciences, Shiraz, Iran

3. Department of Parasitology and Mycology, School of Medicine Qazvin University of Medical Sciences, Qazvin, Iran

Background: *Clostridium difficile* colonizes the large bowel of patients undergoing antibiotic therapy and produces toxin A (Tissue Damaging Enterotoxin) and toxin B (Cytotoxin). *C. difficile* is the leading cause of hospital-acquired diarrhea, known as *C. difficile*-associated disease. Diagnosis of *C. difficile* is based on Glutamate Dehydrogenase (GDH) as antigen and has been shown to be very effective because all strains produce high amount of this enzyme.

Method: In this study, *clostridium difficile* toxin assay was performed for 55 patients referred to Mahdیه Hospital, Tehran in 2016-2018. Toxin assay conducted by Clostridium K-set (Corisbio-Belgium) using membrane technology with colloidal gold contains antibodies against GDH antigen.

Results: Fecal samples from 55 patients (26 females: 29 males) were collected and *clostridium difficile* toxin assay performed. Mean age of patients was 51.8 years. There was no significant relation between age (P value=0.45), sex (P value: 0.8) and positive results. 20 patients (36.3%) were positive for *clostridium difficile* toxin Antigen vs. 35 patients (63.7%) negative. The most important risk factor associated with the infection was previous use of antibiotics.

Conclusion: Positive results for *clostridium difficile* toxin assay is rather high in patients referred to Mahdیه Hospital, south of Tehran. Clostridium difficile infection should be considered in patients with diarrhea and traditional risk factors associated with this disease.

Keywords: *Clostridium difficile*, Diarrhea, Toxin assay.

ICIA2018\Immunology of Infectious Diseases Bacteria & Fungi\Poster\8545
8545

Induction of Regulatory T and B cells in Patients with Brucellosis

Sanaz Jourghasemi¹, Fariba Keramat², Ghasem Solgi¹, Mehrdad Hajilooi^{1*}

1. Immunology Department, Faculty of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

2. Department of Infectious Diseases; Department of Pharmacology, Hamedan University of Medical Sciences

Background: Brucellosis is the most common bacterial zoonotic infection which caused by intracellular gram-negative bacteria of the genus *Brucella*. The aim of this study was to analyze the frequencies of regulatory T cells and regulatory B cells in brucellosis patients compared to healthy controls.

Method: Twenty seven patients with brucellosis and twenty healthy controls were included in this case-control study. Frequencies of CD4⁺ CD25⁺ FOXP3⁺ T cells and CD5⁺ CD19⁺ CD38^{Hi} CD1d^{Hi} B cells in peripheral blood samples of both groups of the study were determined by flow cytometric analysis.

Results: We observed a higher significant frequency of CD4⁺CD25⁺FOXP3⁺ T cells in the brucellosis patients compared to healthy controls ($2.130 \pm 0.1609\%$ vs $0.2123 \pm 0.02288\%$, $P=0.0001$). Similarly, percentage of CD5⁺CD19⁺CD38^{Hi}CD1d^{Hi} B cells was significantly higher in patients compared to healthy controls ($6.313 \pm 0.6823\%$ vs $0.7700 \pm 0.1139\%$, $P=0.0001$).

Conclusion: Our results suggested that *Brucella Species* may induce regulatory responses to evade immune system and survive in the host.

Keywords: Regulatory T cells, Regulatory B cells, *Brucella*

ICIA2018\Immunology of Infectious Diseases Bacteria & Fungi\Poster\8588
8588

Comparison of Naloxone and Propanol on the Immunity Responses of Wistar Rats Challenged with S19 Vaccine

Samira Zand, Meysam Abtahi Foroushani, Masumeh Ahanjan, Sara Golbahari, Nasrin Nourollahzadeh

Department of Microbiology, Faculty of Veterinary Medicine, University of Urmia

Background: Alum as a most popular human vaccine can only produce humoral immunity. The immunomodulatory effects of Naloxone and Propanol reported in some pervious study. This study was designed to investigate the Adjuvant potential of naloxone and propanol on the immunity responses of Wistar Rats challenged with live *brucella abortus bovis* vaccine (S19).

Materials and Methods: In this experimental study, 15 Wistar rats were divided into 3 groups according to the following conditions: 1) Naloxone treatment group; 2) Propranolol treatment group; 3) Control group. In the two treatment groups, on day zero, the treatment solution and the vaccine containing 2% brucella was injected into the peritoneum of the mouse. On the 15th day of injection, the vaccine S19 was injected to the foot of each rat. In control group, on day zero, the vaccine containing 2% rubella was injected into the peritoneum of the mouse and on the 15th day of injection, the vaccine S19 was injected to the foot of each rat. Rats were kept for 20 days and the event of symptoms of the disease and their survival were examined.

Results: DTH, specific antibody titer and respiratory burst were more prominent in naloxone treated group compared to control group. Nevertheless, there was no significant difference between propranolol and control groups.

Conclusion: The use of naloxone is a promising strategy to develop more favorable immune stimulation.

Keywords: Respiratory burst, Cell-mediated immunity, Propranolol, Naloxone,

ICIA2018\Immunology of Infectious Diseases Bacteria & Fungi\Poster\8605
8605

Antimicrobial Effects of Acetone Extract Rasha Gavan (*B.M.V*) in Kurdish Ethnomedicine on *E. Coli* & *S. Aureus*

Mehrdad Pooyanmehr¹, Amir Amiri Paryan^{2*}, Erfan Bahrami³, Mohammad Reza Khedri³

1. Department of Basic Sciences and Pathobiology, Microbiology section, Faculty of Veterinary Medicine, Razi University, Kermanshah, Iran

2. Veterinary Student, Pathobiology & Basic Sciences, Faculty of Veterinary Medicine, Razi University, Kermanshah, Iran

3. Veterinary Student, Clinical Sciences, Faculty of Veterinary Medicine, Razi University, Kermanshah, Iran

Background: There are studies on the biological activities of tested plants. Generally, Astragalus (Gavan) species have been used in Kurdish folk medicine to treat various conditions such as diabetes, gastric ulcer, hemorrhoids, inflammation, healing wounds, diuretic, treatment of upper respiratory, tract infections, improve physical endurance, lower blood pressure, antioxidative, antiviral, anti-inflammatory, analgesic, hypotensive, sedative, cardiogenic, hepato protective to, immune resistance and stimulant properties. The aim of this study was to evaluate the antibacterial activities of endemic plants found in Kurdish Ethno medicine.

Methods: The aerial parts of the endemic plants rasha gavan (*B.M.V*) were collected from Mountains at Kermanshah Province and air-dried in the shade. solvent acetone, were used for extraction. The plant sample were soxhlet extracted at 60 °C for 12 h. The extract were then vacuum evaporated. The extracts were prepared in different dilutions (800,400, 200,100, 50 and 25mg/ml). After preparing various concentrations of extract the method agar diffusion and Disk diffusion, MIC and MBC of dilution series for *E.coli* (ATCC10536) and *S. aureus* (ATCC25923), were used in Mueller Hinton Broth. Tetracycline and Vancomycin as a positive control and 10% of dimethyl sulfoxide (DMSO) was used as a negative Control respectively.

Results: Disk diffusion test results showed, Average Diagonal zone of growth inhibition dilution to 100, 200,400 and 800mg/ml Extract, were respectively $7 \pm 0/57$, $7/33 \pm 0/57$, $8/66 \pm 0/57$, $10/66 \pm 0/57$ mm and Vancomycin $19 \pm 0/33$ mm for *S. aureus* and $7/66 \pm 0/57$, $10 \pm 0/32$ mm (400 & 800mg/ml) and Tetracycline $17 \pm 0/39$ mm for *E.Coli*. The results showed an increased concentration, the Diagonal zone of growth inhibition increased. The acetone extract was observed the greatest impact inhibition zone diameter $10/66 \pm 0/57$ mm on *S. aureus* and $10 \pm 0/32$ mm on *E. coli* strain. The results showed MIC value for *S. Aureus* and for *E. coli* 800 mg/ml.

Conclusion: The results achieved in this study showed that Rasha Gavan Acetonic extract have Antibacterial Activity and growth inhibitory impact on both *E. coli* and *S. aureus* (*S.aureus* is known as more sensitive to the extract than *E.Coli*).

Keywords: Acetonic Extract, Rasha Gavan, *E.Coli*, *S. aureus*, MIC, MBC



ICIA2018\Immunology of Infectious Diseases Bacteria & Fungi\Poster\8671
8671

Presentation of Like Melanoma Onychomycosis due to *Fusarium solani* Species Complex

M. Soltania, A.R. Khosravia, A. Sharifzadeha

Mycology Research Center, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran

Background: Onychomycosis is a fungal infection of the nail caused by molds, yeasts and dermatophytes.

Method: In this study, we report a case with unusual fingernail onychomycosis in a 31-year-old Iranian woman suspected having nail bed melanoma. Biopsy was performed from the nail bed and no any evidence of melanoma was reported by the pathologist.

Result: In direct microscopic examination (DME) and culture numerous hyaline septated hyphae and canoe-shaped conidia on conidiophores were observed, respectively. Diagnosis was confirmed by molecular DNA sequencing and *Fusarium solani species complex* was identified as etiological microorganism.

Keywords: Onychomycosis, Melanoma, Non-dermatophyte fungi, *Fusarium*, DNA sequencing, Itraconazole

ICIA2018\Immunology of Infectious Diseases Bacteria & Fungi\Poster\9932

9932

Detection of IS903, IS26 and ISEcp1 Elements in CTX-M-Producing *Escherichia coli* and *Klebsiella pneumoniae* isolates from Leukemic Patients in Iran

Mahdaneh Roshani¹, Hossein Goudarzi¹, Ali Hashemi¹, Abdollah Ardebili², Soroosh Erfanimanesh³,
Aghil Bahramian¹

1. Department of Microbiology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Department of Microbiology, Faculty of Medicine, Golestan University of Medical Sciences, Gorgan, Iran

3. Department of Microbiology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

Background: The ability of Extended Spectrum Beta-Lactamases (ESBLs) production is one of the main mechanisms for the emergence of antibiotic resistance in *E. coli* and *K. pneumoniae*. The aim of this study was to evaluate the occurrence of IS903, IS26 and ISEcp1 insertion elements among the CTX-M-producing *E. coli* and *K. pneumoniae* isolates from leukemic patients in Tehran.

Methods: Eighty *E. coli* and *K. pneumoniae* isolates were recovered from patients admitted in hospitals in Tehran. Antibiotic susceptibility tests were performed by Kirby-Bauer disc diffusion and broth microdilution methods. Detection of ESBL producers was evaluated by phenotypic confirmatory test. The presence of IS903, IS26 and ISEcp1 insertion elements in CTX-M-positive *E. coli* and *K. pneumoniae* isolates were investigated by PCR-sequencing methods.

Results: The rate resistance of 80 *E. coli* and *K. pneumoniae* isolates against the 9 antibiotics was as follows: 100% to ampicillin, 15% to amikacin, 51% to ciprofloxacin, 30% to gentamicin, 58% to ceftriaxone, 10% to imipenem, 63% to cefotaxime, 51% to levofloxacin and 55% to ceftazidime. Using phenotypic confirmatory test, 51 (63.75%) isolates were ESBL producers. The prevalence of CTX-M-1, CTX-M-2, CTX-M-9, CTX-M-8 and CTX-M-25 genes was 87.5%, 13.75%, 23.75%, 10 and 0%, respectively. IS903, IS26 and ISEcp1 elements were detected in 93.75%, 71.25% and 100% of isolates, respectively.

Conclusion: This study indicates that the occurrence of antibiotic resistance, IS and CTX-M-producing *E. coli* and *K. pneumoniae* isolates could be a major concern and highlights the need of infection control measures.

Keywords: Leukemic patients, *Escherichia coli*, *Klebsiella pneumoniae*, Extended-spectrum beta-lactamases, Insertion Sequence

ICIA2018\Immunology of Infectious Diseases Bacteria & Fungi\Poster\9934
9934

The First Report of Extended-Spectrum β -Lactamase (ESBL) Genes in an *Escherichia coli* Isolate from a One-Month-Old Infant with Acute Lymphoblastic Leukemia (ALL) in Iran

Mahdaneh Roshani¹, Hossein Goudarzi¹, Ali Hashemi¹

Department of Microbiology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background Bloodstream infection is one of the most life-threatening complications and remains the frequent cause of treatment failure in children with acute lymphoblastic leukemia. In this study, we describe the isolation of an extended-spectrum β -lactamase (ESBL) producing *Escherichia coli*.

Methods: Antimicrobial susceptibility tests were also determined on the isolate by the Kirby-Bauer disk diffusion method. In addition, ESBL production was examined for the isolate using the Combination Disk Diffusion Test (CDDT). PCR was used to screen the presence of CTX-M-1, CTX-M-2, CTX-M-8, CTX-M-9, CTX-M-25, ISEcp1, IS₂₆, and IS₉₀₃ genes in this isolate.

Results: We found an ESBL-producing *E. coli* in a 1 month old infant with blood cancer that carried CTX-M-1 group enzymes.

Conclusion: Our finding emphasizes the need to have more precise screening methods to identify the causative infectious agent at early stage of infection to choose the appropriate treatment in these severely immunocompromised patients.

Keywords: Bloodstream infection, Extended-spectrum β -lactamase, *Escherichia coli*, Acute lymphoblastic leukemia



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Immunology of Infectious Diseases Viruses

Poster Presentation

ICIA2018\Immunology of Infectious Diseases Viruses\poster\5307
5307

Evaluation of hepatitis B viruses serologic markers in thalassemia and dialysis patients of Mazandaran province

Sara Mohamadnezhad¹, Nour Amirmozafari², Mahdi Asmar³, Mohtaram Nasrollahi⁴

1. Faculty of Medical Sciences, Sari Branch, Islamic Azad university, Sari, Iran

2. Department of Microbiology, Iran University of Medical Sciences, Tehran, Iran

3. Department of Microbiology, Pasteur Institute of Iran, Tehran, Iran

4. Department of Microbiology, Mazandaran University of Medical Sciences, Sari, Iran

Background: Nearly 300 million people worldwide are carriers of the hepatitis B virus (HBV). Thalassemia and dialysis patients are one of the most important occupational groups at risk for the virus. Despite vaccination, in many of them, antibody is reduced and while the exposure to infected materials, there is a possibility of infection to HBV. This study examines some of hepatitis B serologic markers among special patients of Mazandaran province.

Methods: 94 special patients were enrolled to our study. For all patients in the study, a questionnaire including demographic information, history of the dose of the vaccination for hepatitis B, blood transfusion and blood products and surgical history were completed. 5ml of venous blood was taken from each student and serum was separated and stored in -20°C. For all samples, ELISA test operated for anti-HBs and anti-HBc and HBsAg.

Results: Among 94 under studied thalassemia and dialysis patients, 55 people (58.5%) were male and 39 people (41.5%) were female. In these patients 57 people (60.6%) were dialysis and 37 people (39.4%) were thalassemia, in which 4 people (7%) of dialysis patients and 8 people (21.6%) of thalassemia were infected by HBV. 12 people (22.8%) of samples were positive by the presence of HBsAg, 54 people (57.4%) of samples were positive by the presence of Anti-HBs and 13 people (13.8%) of samples were positive by the presence of anti-HBc.

Conclusion: About 12.8% of understudied patients were HBV positive and also 42.6% of them were sensitive to HBV. Therefore, screening of them for HBV antibody is recommended before any blood products usage. Also, in order to adequate protection of the patients, the serologic studies is necessary during blood production usage.

Keywords: Hepatitis B Viruses, Serologic makers, Vaccinations, Thalassemia and Dialysis.

ICIA2018\Immunology of Infectious Diseases Viruses\poster\6315
6315

Increased of PDGF- β gene expression is associated with ATLL development in HTLV-1 carriers

Atabati Hadi ^{1,2}, Rafatpanah Houshang ^{1,2*}, Allahyari Abualqasem³, Davoud Ghaffarian^{1,2}

1. Department of Immunology, School of Medicine, Mashhad University of Medical Sciences(MUMS)

2. HTLV-1 Center and related diseases, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

3. Department of Hematology, Ghaem Hospital ,Mashhad, School of Medicine, University of Medical, Sciences(MUMS)

Background: Human T lymphotropic virus (HTLV-1) is a human retrovirus which is associated with adult T cell leukaemia (ATLL). ATLL is a malignancy lymphoproliferative disease which infects CD4+ T cells. It is not clear why the majority of HTLV-1 infected individuals remain healthy carriers and minority develop ATLL. It has been shown that angiogenesis is increased in ATL patients and the plasma levels of angiogenesis factors such as VEGF are upregulated in the patients. In the present study, we examined the mRNA expression of another angiogenesis factor, PDGF (platelet derived growth factor) and its receptors in patients with ATL and HTLV-1 healthy carriers (HACs).

Materials and Methods: Peripheral blood mononuclear cells (PBMCs) were isolated from ATLL patients and asymptomatic carriers by using ficoll-hypopaque density centrifugation. RNA was extracted and cDNA was synthesised. A real-time PCR TaqMan method was designed and optimized for evaluation of PDGF, PDGF-R and Tax gene expression. HTLV-1 proviral load was also quantified in patients with ATLL and carriers.

Results: The mRNA expression of Tax was significantly higher in ATLL patients than HACs ($p=0.01$). The HTLV-1 proviral load was higher in ATLL patients compared to HTLV-1 carriers ($p=$ There was significant increases in PDGF gene expression in HACs than ATLL patients ($p=0.002$). The expression of PDGF was higher in HTLV-1 carries compared with ATLL patients and significant differences was observed between two groups ($p<0.001$). There was no significant differences in PDGF-R gene expression between ATL patients and HACs ($p>0.05$).

Conclusion: High expression of PDGF-R in HTLV-1 carriers might block the PDGF signalling pathway and prevent ATLL development.

Keywords: HTLV-1, PDGF β , PDGF β -R

ICIA2018\Immunology of Infectious Diseases Viruses\poster\6328
6328

Dissection of plasmacytoid dendritic cell's responses to influenza virus based network analysis

Fatemeh Hadi^{1,2}, Rasoul Godini¹, Hossein Fallahi¹

1. Department of Biology, Faculty of Sciences, Razi University, Kermanshah, Iran

2. Department of Stem Cells and Developmental Biology, Brain and Cognitive Center, Royan Institute for Stem Cell Biology and Technology, ACECR, Tehran, Iran

Background: Plasmacytoid dendritic cells are responsible for defense against viral infection in respiratory organs. These cells secrete massive amount of interferon- α through innate immune responses. Flu, as a life threatening respiratory disease through the world, is caused by influenza virus. It has been known that plasmacytoid dendritic cells are responsible for protecting lung against flue. In here, using network analysis, we dissected transcriptome profile of plasmacytoid dendritic cells affected by influenza virus to reveal underlying molecular mechanisms.

Methods: In order to do this study, we used freely available microarray dataset numbered GSE68849. Differentially expressed genes were determined using GEO2R tool of NCBI. Protein-protein interactions were constructed by validated data from STRING database. Annotations for gene ontology and disease were obtained from DAVID database. The networks were visualized and analyzed by Cytoscape and Gephi software programs. Protein complexes were determined by ClusterONE plug-in of Cytoscape. Network centrality factors including degree and betweenness were used to detect hub genes.

Results: 1127 differentially expressed genes were detected, 605 and 522 down- and up-regulated, respectively. Using medium confidence of STRING database, a network of protein-protein interaction with 707 nodes and 2552 edge was constructed. Overlapping neighborhood expansion algorithm detected 15 modules. Centrality factors analysis showed 33 hub genes including JUN, HSP90AA1, SYK, MAPK3 and IL8. Annotation of the genes showed defense response to virus, innate immune response, and type I interferon signaling pathway are from processes which the genes are involved in.

Conclusion: These results reveal principal information about molecular mechanisms involved in response to influenza virus that can help to development a better treatment of this disease.

Keywords: Immune response, Influenza, Network, Protein-protein interactions

ICIA2018\Immunology of Infectious Diseases Viruses\poster\6333
6333

Relative Frequency of Cytomegalovirus (CMV) in Tissue Samples of Women with Breast Cancer in Sanandaj, Iran

Maryam karimi^{1,2}, Seyede Zeinab Hosseini^{3*}, Bahram Nikkhoo⁴, Farid Soleimani Mohammadi⁵

1. M.SC Microbiology, School of Medicine, Kurdistan University of Medical Sciences, Sanandaj, Iran

2. Student Research Committee, Kurdistan University of Medical Sciences, Sanandaj, Iran

3. virology Department, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

4. MD Pathologist, Kurdistan University of Medical Sciences, Sanandaj, Iran

5. Department of Virology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

Background: Despite the researchers' efforts, the cause and development of breast cancer is still incompletely understood. Currently, in some reports, human cytomegalovirus has been referred as a risk factor for breast cancer. This study aimed to determine relative frequency of cytomegalovirus in tissue samples of women with breast cancer in Sanandaj county.

Methods: To determine the relative frequency of the human cytomegalovirus (CMV), 50 formalin-fixed tissues of breast cancer, which all were invasive ductal carcinoma, were studied using the nested-polymerase chain reaction.

Results: In 26 cases of breast cancer tissues (26/50), human cytomegalovirus was detected. Seventeen cases of breast cancer tissues were in a moderately differentiated stage, and nine cases had poor-differentiated stage tissues that were positive for viral DNA. At older ages (>45 years) the rate of human cytomegalovirus DNA was higher, but no significant association was seen ($p=0.16$). In general, due to the high prevalence of the DNA of human cytomegalovirus (58%).

Conclusion: in this study it is assumed that human cytomegalovirus (HCMV) has a contributing role in breast cancer; although more study is required to clearly define its part in this type of cancer.

Keywords: Human Cytomegalovirus (HCMV), Breast Cancer, Nested PCR, ductal carcinoma

ICIA2018\Immunology of Infectious Diseases Viruses\poster\7614
7614

Evaluation of camel serum immunoglobulin on inhibition of hepatitis C virus in cell culture

Fariba ghorbanzadeh^{1,2}, Jalil Tavakkol Afshar^{1,2}, Amin Reza Nikpoor^{1,2}

1. Department of Immunology, Faculty of Medicine, Mashhad University of Medical Sciences (MUMS), Mashhad, Iran

2. Immunogenetic and Cell Culture Department, Immunology Research Center, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

Background: Hepatitis C virus (HCV) represents a major public health problem, affecting 3% of the world's population. Although the development of new therapeutic approaches to combat HCV infection has taken place, no efficient vaccines exist yet. Therefore this study aimed to evaluate the role of immunoglobulins extracted from camel serum, the first in cell culture, then in future research in animal models and in clinical phase done as a pilot for the treatment of patients with HCV infection. Camels are special animals in antibody production as they possess a unique type of IgG antibody in their serum which lack light chains as well as the heavy chain constant domain "CH1", so-called heavy chain antibodies or nanobodies.

Methods: First immunoglobulins were extracted from camel serum with ammonium sulfate. Then three types of experiments were performed on HuH7.5 cell. In the first process, the HCV pre-infected cells were treated with the immunoglobulins and in the second process HCV was incubated with the purified immunoglobulins and then mixed with cell at the end of the process the immunoglobulins were incubated with the cells and then exposed to HCV. Eventually with Real time-PCR method reduction or no reduction in viral titers were measured.

Results: The results indicate that there is a direct interaction between HCV and camel IgGs, which led to inhibition of HCV binding to the cells. Camel IgGs were capable of inhibiting the intracellular HCV replication at concentrations of 18.5 mg/ml. But IgGs could not inhibit entry the HCV into the cell.

Conclusion: Camel serum immunoglobulins isolated from camel serum could inhibit the HCV infectivity. Immunoglobulins inhibit the HCV infectivity through inhibiting proliferation and binding the virus to the cell but immunoglobulins could not prevent the HCV from entering the cell.

Keywords: Hepatitis C Virus, Camel Antibody, Infectivity

ICIA2018\Immunology of Infectious Diseases Viruses\poster\7771
7771

The effects of curcumin on gene expression and serum levels of Toll like receptor -2 &4 (TLR-2 and TLR-4) in patients with myelopathy associated with HTLV-1 (HAM/TSP) and virus carriers

Marjan Zamani¹, Dr. Houshang Rafatpanah², Dr. Reza Boostani³

1. Department of Allergy and Immunology, faculty of medicine, Mashhad University of Medical Sciences, Mashhad, Iran

2. Department of Allergy and Immunology, Inflammation and inflammatory Diseases Research Centre, faculty of medicine, Mashhad University of Medical Sciences, Mashhad, Iran

3. Department of Neurology, Ghaem Hospital, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

Background: TLRs are innate immune receptors that are able to respond to pathogens and derived molecules of a host. Curcumin has anti-inflammatory and anti-oxidant effect which can be used for the treatment inflammatory diseases. In the present study the anti-inflammatory effects of curcumin in HAM/TSP patients by evaluation of TLR-2 and TLR-4 which have pivotal role in inflammatory response were examined.

Materials and Methods: The population studied included ٢١ HAM/TSP patients before and after treatment with curcumin, 21 asymptomatic carriers, and 21 healthy controls. Peripheral blood mononuclear cells were separated by using ficoll and cDNA was synthesized. The genes expression and serum protein levels of TLR-2 and TLR-4 by real time PCR and ELISA was evaluated.

Results: The expression of TLR-2 was increased in treated group compared with untreated group ($p < 0.05$), however no significant differences in TLR-2 serum levels was observed between two groups ($p > 0.05$). Furthermore, the expression of TLR-4 was increased after treatment with curcumin compared with untreated group ($p < 0.05$). No significant differences in TLR-4 serum levels was observed between two groups. The mRNA expression of TLR-4 was higher in asymptomatic carriers compared to HAM/TSP patients ($p < 0.05$), however no significant differences in the serum levels of TLR-4 was observed between HAM/TSP patients and asymptomatic carriers ($p > 0.05$). There were significant difference in TLR-2 and TLR-4 gene and protein expression between healthy controls and HAM/TSP, and healthy controls and carriers ($p < 0.05$).

Conclusion: In conclusion the results of the present study might suggest that viral factors interference with curcumin in order to maintain the balance between the virus and inflammatory immune response. It seems that the expression of viral factors might also contribute in the difference between TLR-2 and TLR-4 gene expression between two groups. Further studied are needed to clarify this issue.

Keywords: HTLV-1, HAM/TSP, TLR-2, TLR-4.

ICIA2018\Immunology of Infectious Diseases Viruses\poster\7782
7782

**Evaluation of relationship between HBs Ab titer with birth weight and mode
Of delivery in 3-5 years old children**

Motahareh Ehsani¹, Mehdi Shahbazi¹, Mousa Mohammadnia afrouzi¹, Kowsar Smaeilnejad, Mojgan Bagherzade¹

Department of Immunology, School of Medicine, Babol University of Medical Sciences, Babol, I.R.Iran

Background: Vaccination of hepatitis B infection has been known as a crucial step in prevention of HBV. Both environmental and genetic factors affect immune response to vaccination. Environmental factors in children include birth weight, mother age, birth season, place of residence, Feeding status and mode of birth. In this study the relationship between HBs Ab titer with birth weight and mode of delivery was assessed.

Methods: Peripheral blood samples were collected from 100 children between the ages of 3-5. Sera were tested for anti-HBs antibody by enzyme linked immunosorbent assay (ELISA).

Results: Findings demonstrated that there is no significant difference (P value > 0.05) between HBs-antibody titer in caesarean-born children and vaginally-born ones and also in children with low birth weight (< 2500) and high birth weight (> 2500), this means that, both birth weight and mode of delivery couldn't affect on HBs-antibody titer in children with the age of 3-5.

Conclusion: Vaccination has been considered as an attempt in inducing immunity and protection against HBV infection. According to these findings, vaccine immune responses and Anti-HBs antibody titer in children are not affected by weight and mode of delivery, However there is not any assured conclusion and more researches is needed in this field.

Keywords: Vaccination, HBs Ab, Mode of delivery, Birth weight

ICIA2018\Immunology of Infectious Diseases Viruses\poster\7901
7901

Sequencing of HTLV1 p12 region in HAM/TSP, ATL patients and HTLV1 carriers in Mashhad, Iran

MehranMoharreri¹, SeyedAliMirhosseini¹, SeyedRohollahArab¹, BarataliMashkani², SeyedAbdolrahimRezaee³, RezaFaridHosseini⁴, HoushangRafatpanah¹

1. *Inflammation and Inflammatory Diseases Research Center, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran*

2. *Biochemistry and Nutrition Research Center, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran*

3. *Immunology Research Center, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran*

4. *Allergy Research Center, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran*

Background: HTLV1-virus is associated with two important diseases ATL & HAM/TSP while most of the infected cases remain as asymptomatic carriers (ACs). Considering Mashhad is one of the HTLV-1 endemic areas in the world, here, we report genomic variability of p12 region of positive HTLV-1 patients. Our aim in this study was increase our understanding about similarities and differences of p12 genome in 3 groups of individuals with distinctive sign and symptoms.

Methods: Blood samples were collected from 15 individuals including 5 HTLV-1 asymptomatic carriers, 5 HTLV-1 associated with HAM/TSP and 5 patients with adult T-cell leukemia lymphoma (ATLL). DNA was extracted from the peripheral blood mononuclear cells (PBMCs) then PCR was done in p12 genes. Finally, PCR products were cloned and sequenced.

Results: P12 nucleotides and amino acids in 15 HTLV1 infected patients who participated in our study were strikingly similar. According to molecular analytic considerations P34L mutation in P12 gene of Mashhad HTLV1 infected patients does exist.

Conclusion: The results presented in this study suggest that some p12 natural mutations may be associated to HAM/TSP development and to the proviral load of HTLV-1-infected individual. These results indicate that genetic variations among HTLV-1 strains are very low. It is probable that these genetic variations, result of geographical differences among origins of HTLV-1 strains. We will hope that this valuable genomic data facilitates vaccine design against HTLV1. Moreover, there are a lot of controversial debates about function of p12 protein, so sequencing of p12 region could help researchers to explain whether different p12 genomic make up have specific effect on manifestation disorders in HTLV-1 infected people.

Keywords: HTLV-1, HAM/TSP, ATL, P12, DNA sequencing

ICIA2018\Immunology of Infectious Diseases Viruses\poster\7913
7913

Cytotoxic and inflammatory markers determined the outcome of HTLV-1 infection

Houshang Rafatpanah¹, Asadollah Mohammadi¹, Reza Boostani²

1. Immunology Research centre, Division of Inflammation and Inflammatory Diseases, Mashhad University of Medical sciences, Mashhad, Iran

2. Department of Neurology, Mashhad University of Medical sciences, Mashhad, Iran

Background: Human T-lymphotropic virus type I (HTLV-1) is a type C retrovirus which is associated with two main types of diseases known; adult T-cell leukemia/lymphoma (ATLL) and HTLV-1 associated myelopathy/tropical spastic paraparesis (HAM/TSP). We have previously showed that north east of Iran, Mashhad and Sabzevar, have been recognized as an endemic region of HTLV-1. It is not fully known why a small portion of HTLV-1-infected individuals develop HAM/TSP and majority remains lifelong asymptomatic carriers. The different outcome of HTLV-1 infection may be explained by the existence of viral agents, genetic background or even viral factors. High titers of HTLV-I antibody, high proviral load and high frequency of HTLV-1 specific CTL in HAM/TSP patients distinguish them from HTLV-I carriers. Among these factors, the CTL response is known to be a key factor in HTLV-I infection. Furthermore proinflammatory markers are also involved in the outcome of HTLV-1 infection.

Methods: In the present study, we examined the role a panel of cytotoxic markers including levels of mRNA and protein to clarify their roles in HAM/TSP patients and HTLV-1 carriers. Furthermore, we also investigated the role of inflammatory markers such as IL-6, Toll like receptors (TLR) and high mobility group box1 (HMGB1) and viral markers such as proviral load in studied groups.

Results & Conclusion: The results of the study showed that cytotoxic response is failed in HAM/TSP patients. Additionally, increased inflammatory markers and viral load is shown to be associated with HAM.

Keywords: HTLV-1, HAM/TSP, Cytotoxic markers, inflammatory markers

ICIA2018\Immunology of Infectious Diseases Viruses\poster\8051
8051

Evaluation of anti-HBS titer of vaccinated students of Guilan University of medical science (2013-2014)

Sara Rasouli¹, Masumeh Avari²

1. Department of Microbiology, College of Science, Rasht Branch, Islamic Azad University, Rasht, Iran

2. Department of Microbiology, College of Science, Rasht Branch, Islamic Azad University, Rasht, Iran

Background: Hepatitis B is a viral disease with high prevalence. Unfortunately, immunologic response to the vaccines is not sometimes perfect and it is necessary to examine the immune response.

Methods: In a cross-sectional study of 181 students that have been vaccinated against hepatitis B, blood samples were obtained for assessing antibody titer of HBV by ELISA. After collecting demographic data at the University of Medical Sciences, Guilan in 2014, data were analyzed by SPSS software and presented with chi-square test.

Results: Among 181 participants completed the three stages of vaccination against HBV, 64/7% had adequate response against hepatitis. There was no significant correlation between the immunity response and disease history, gender and other variables except on duration of vaccination ($p < 0.001$).

Conclusion: According to our findings health education and evaluation of immune responses against hepatitis in medical students is necessary. Also, it should be considered revaccination for persons with low titers of antibody.

Keywords: Hepatitis B, vaccination, antibodies

ICIA2018\Immunology of Infectious Diseases Viruses\poster\8098
8098

A comparison of Phylogenetic of Human Immunodeficiency Virus Type 1(HIV-1) from Khorasan Razavi

Faeze Fakour.k¹, Dr. Abdolrahim Rezaee ¹

1. Department of Mashhad University of Medical Sciences

Background: HIV is the virus with high genetic diversity, this genetic diversity because of the mutations of this virus that is produced due to the rapid proliferation and characterization of recombinant of reverse transcriptase. HIV-1 divided into four phylogenetic groups based on gag gene sequences, which are M, N, O and P groups. M Group is pandemic factor and divided to 9 subtypes (A to K) and 61 circulating recombinant forms (CRF). The HIV-1 infection in our country has been increased significantly in recent years.

Methods: Risk of HIV infection of the patients in this project confirmed by using ELISA and immunoblot. For this purpose, first samples were collect and then send them to the laboratory of immunology, by Sciences inflammation and inflammatory diseases Research Center in Mashhad University of Medical, genetic material (DNA) was extracted in accordance with international standard method. Then with NESTED -PCR method of gene sequences areas p17 of gag and c2-v5 env gene was amplified by using specific Primers and then was sent to Denazist Asia company for sequencing. The nucleotide sequences of gag and env genes reviewed and edited in MacroGen Korea company.

Results: The data obtained from sequencing was arranged by MEGA6 software and phylogenetic trees were drawn with maximum-likelihood. 80% of patients who entered the study were male and 20% were female. 65% of them were in the age group 20-39 years. 65% of them were infected through intravenous drug use and 19% through sex. In terms of co-infections with other viruses, 45% co-infection were observed with hepatitis B and 45% co-infection with hepatitis C, and 10% co-infection with HTLV1.

Conclusion: Samples formed a significant cluster with subtype CRF35-AD from Afghanistan, subtype A1 from Pakistan, subtype F from South America, subtype A and C and of East Africa and subtype B of Eastern Europe.

Keywords: HIV, Phylogeny, Subtype, Iran

ICIA2018\Immunology of Infectious Diseases Viruses\poster\8171
8171

Altered expression of Human Leukocyte Antigen G (HLA-G) in HIV+ drug naïve patients

Versa OmraniNava¹, Saeid Abediankenari¹, Jamshid Yazdani²

1. Department of Immunology, Immunogenetics research center, School of Medicine University of Mazandaran medical Science

2. Department of Biostatistics School of Health, University of Mazandaran Medical Science

Background: HLA-G is an immunomodulatory protein that alteration in its expression has been reported in several pathologic conditions. For as much as the results of HLA-G expression in HIV+ patients have been in consistent and some reports suggest the effects of antiretroviral Therapy (ART) on HLA-G, we investigate HLA-G expression and its clinical relevance with disease status in drug naïve patients.

Methods: We enrolled a total of 20 HIV+ drug naïve patients referring to the Behavior Disease Consultation Center of Mazandaran and 20 sex-Age matched healthy volunteers in this study. Blood mononuclear cells were isolated by ficoll method. After RNA isolation and cDNA synthesis, Real time PCR was carried out to assess gene expression in both groups. CD4+ cells count were determined in cases using flowcytometry. Data were analyzed by SPSS and REST softwares.

Results: The mean age of case and control groups were 34.6 (CI 95% 30.83-38.37) and 33.9 (CI 95% 30.66-37.14), respectively. Median CD4 count in patient was 515 (CI 95% 395-606). HLA-G expression was statistically different between groups and patients showed lower expression. There was no relation between disease status, CD4 Count and HLA-G expression in patients.

Conclusion: Our findings suggest that HLA-G expression is dysregulated in HIV+ patients. Lower expression in patients may influence Antigen recognition process. Further follow up are needed to clarify interactions between HIV and non-classical HLA molecules.

Keywords: HLA-G, HIV, CD4, gene expression

ICIA2018\Immunology of Infectious Diseases Viruses\poster\8173
8173

Relationship between Up-regulation of Interleukin-10 (IL-10), Regeneration and Tolerance Factor (RTF) and CD4+ cell count in HIV+ patients

Versa Omraninava¹, Saeid Abediankenari¹, Jamshid Yazdani²

1. Department of Immunology, Immunogenetics research center, School of medicine University of Mazandaran medical science

2. Department of Biostatistics, school of Health, University of Mazandaran medical science

Background: Regeneration and Tolerance Factor (RTF), is a potential Th2 response marker with two forms of membrane bound and soluble. RTF can induce IL-10 expression and block tumor cell lysis by NK cells, thus it has been considered as an immune suppressor. Considering contrary in literatures regarding Th1-Th2 response pattern in HIV infection and effects of antiviral therapy on mentioned molecules, we determined IL-10 and RTF expression in HIV+ Patients before treatment and possible association with CD4 levels.

Methods: We enrolled 40 subjects in this study, 20 of whom were HIV+ patients without treatment history and 20 were healthy individuals. RNA extraction and cDNA synthesis were done on isolated blood mononuclear cells of all participants. We used Real time PCR to assess gene expression of RTF. Plasma IL-10 levels were measured by ELISA test. SPSS was used to analyze data.

Results: In case group, 14 male and 6 female subjects (mean age, 34.6) and in control group 15 male and 5 female (mean age, 33.9) were studied. RTF and IL-10 were increased in HIV+ patients in comparison to healthy subjects. A negative correlation between plasma IL-10 and CD4 cell count were observed in patients ($r: -0.04$).

Conclusion: Increased expression of Th2 associated markers in HIV+ and its negative correlation with patient immune status may explain immune evasion mechanisms employed by HIV. Potential therapeutic objectives of mentioned molecules needs to be investigated.

Keywords: HLA-G, RTF, IL-10, HIV

ICIA2018\Immunology of Infectious Diseases Viruses\poster\8204
8204

Hepatitis C virus viremia among first time Iranian blood donors

Fahimeh Ranjbar Kermani¹, Sedigheh Amini Kafi Abad¹, Kamran Mousavi Hosseini¹, Zohreh Sharifi¹,
Mohammad Ali Mansournia², Mahtab Maghsudlu¹

1. Blood Transfusion Research Center, High Institute for Research and Education in Transfusion Medicine, Tehran, Iran

2. Department of Epidemiology and Biostatistics, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

Background: Hepatitis C virus (HCV) viremia is described as persistent HCV RNA among HCV exposed individuals. HCV viremia rate is defined as the proportion of HCV Ab positive and HCV RNA positive individuals to total HCV Ab positive individuals and provides the likelihood of HCV viremia infection in a given population. Understanding the HCV viremia rate is important to increase knowledge about HCV epidemiology. The aim of this study was to evaluate the HCV viremia rate and demographic parameters and risk factors correlates among Iranian first time blood donors.

Methods: In this cross sectional study, serologically confirmed HCV positive blood donors who came back to the Iranian blood transfusion centers over the country for counseling from November 2015 to May 2017 were included and interviewed for HCV risk factors. HCV RNA RT-PCR was carried out for subjects by an in-house qualitative assay according to manufacturer's instruction. Logistic regression was performed for data analysis and STATA software version 13 was used for statistical analysis.

Results: Out of 239 subjects, 226 (94.5%) were first time donors and HCV viremia was detected in 155 (68.6%, 95% CI 62.4% - 74.8%) of first time donors. No statistically difference was found in demographic parameters and risk factors between HCV Ab positive/ HCV RNA positive and HCV Ab positive/ HCV RNA negative first time blood donors.

Conclusion: The results of this study show the likelihood of occurrence of active HCV infection among first time Iranian blood donors. The majority of first time Iranian blood donors are in active phase of HCV infection. The viremic status was not associated with demographic parameters and risk factors. More effective donor selection is required to improve blood safety and follow-up studies on viremic first time blood donors are recommended to clarify impact of factors on the occurrence of HCV viremia.

Keywords: Blood donors, Hepatitis C, Viremia

ICIA2018\Immunology of Infectious Diseases Viruses\poster\8212
8212

Evaluation of coverage and HBsAb titer in the serum of Dental students

AbbasAzadmehr¹, ParvizZoulghadri², NiloufarSadat Miri¹, Aboulfazl Rahmani¹, Mohammad Sadegh Falahatkar²,
FatemehHajmanochehri²

1. Department of Immunology, Babol University of Medical Sciences, Babol, Iran

2. Department of Periodontics, Dental School, Qazvin University of Medical Sciences, Qazvin, Iran

Background: Hepatitis B is an infection disease that affects the liver and caused by the hepatitis B virus. Health workers such as dental students are at the higher risks of blood-transmitted infections. Up to now, the hepatitis B virus (HBV) vaccination was the useful and effective way to prevent HBV infection which requires antibody titers estimation following the vaccination to be sure about its efficiency.

Methods: The study included 33 clinical and postgraduate students (18 male and 15 female). The serum samples were collected and tested hepatitis B surface antigen (HBsAg) by enzyme-linked immunosorbent assay (ELISA). The B hepatitis vaccination coverage and antibody titer were reported. The data were subjected to chi-square and Mann-whitney U tests.

Results: 92% of all studies received the three doses of the vaccine. Of all 33 participants, 29 individuals (87.9%) vaccinates against hepatitis B infection. 28 students (84.8%) showed positive antibody titer and 5 ones (15.2%) had negative antibody titer. No significant differences were noted among males and females regarding antibody titer (mean 410IU/MI and 362.2 IU/mL for males and females).

Conclusion: Hepatitis vaccination coverage and positive antibody titer among the clinical and post graduate dental students were appropriate, however; regarding higher risks of blood-transmitted disease among them; it should be tried to increase the vaccination coverage more.

Keywords: B Hepatitis, Vaccination, Antibody titer, ELISA

ICIA2018\Immunology of Infectious Diseases Viruses\poster\8235
8235

Hepatitis B Virus DNA is absent in Lebanese Blood Donors Who Are Positive for Both Anti-HBc and Anti-HBs Antibodies

Ghada Kchour,¹ Mohammad Fayyad Kazan,^{2,*} Antoine Haddad,³ Bassam Badran,¹, Mahdi Tarhini^{4,*}

1. Laboratory of Cancer Biology and Molecular Immunology, Faculty of Sciences I, Lebanese University, Hadath, Lebanon

2. Institute de Biology et de Medicine Moleculaires, Universite Libre de Bruxelles, 6041 Gosselies, Belgium

3. Department of Clinical Pathology and Blood Banking, Sacrecoeur Hospital, Lebanese University, Lebanon

4. Faculty of Health Sciences V, Lebanese University, and director of Clinical Pathology and Blood Banking Department, Raee Hospital, Saida, Lebanon

Background: A major risk during blood transfusion is the transmission of infectious agents such as hepatitis B virus (HBV). Screening for anti-HBc antibodies has long been used to test HBV infection and thus, blood safety. However, the development of advanced tools enabling HBV-DNA detection rendered the diagnostic benefit of anti-HBc test uncertain.

Methods: Sera of 7200 blood donors were first screened for HBs antigen (HBsAg) and anti-HBc antibody. Samples that were found to be HBsAg-negative but anti-HBc-positive were further tested for anti-HBs and HBV-DNA.

Results: Of the 7200 tested samples, 7143 (99.2%) were HBsAg-negative, while the remaining 57 (0.8%) samples were HBsAg-positive. Among the HBsAg-negative samples, 490 (6.8%) were anti-HBc-positive. Of the anti-HBc positive samples, 397 (81%) were anti-HBs positive, while the remaining 93 (19%) samples were anti-HBs-negative. Interestingly, all of the anti-HBc positive samples, which also were positive for anti-HBs, exhibited negative HBV-DNA results. On the other hand, the anti-HBc positive samples, which were negative for anti-HBs, showed both negative and positive HBV-DNA results.

Conclusions: Blood samples which are positive for both anti-HBc and anti-HBs are characterized by negative HBV load.

Keywords: Hepatitis B Virus (HBV), Anti-HBc, Anti-HBs, Blood Safety

ICIA2018\Immunology of Infectious Diseases Viruses\poster\8378
8378

Hepatitis B Core Antibody Immunoglobulin M in Blood Donors with a History of Hepatitis B Virus Infection

Helia Azimi^{1, 2}, Maryam Vaezjalali^{1, 3}

1. Infectious Disease Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Department of Microbiology, Faculty of Biological Sciences, Shahid Beheshti University, Tehran, Iran

3. Department of Microbiology, Faculty of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Hepatitis B is still an issue after blood transfusion. A reason could be the window period of hepatitis B infection in blood donors. In countries such as Iran, enzyme-linked immunosorbent assay (ELISA) is the only test to detect hepatitis B surface antigen in blood donors. This procedure may miss the window period of hepatitis B infected donors. The current study aimed to look for hepatitis B core antibody immunoglobulin M in blood samples of Iranian donors with a history of hepatitis B virus infection to detect infection window period.

Methods: Eighty serum samples with hepatitis B core antibody were collected from 1000 healthy blood donors, forty of them had been positive for hepatitis B virus DNA in authors' previous study and were diagnosed as occult hepatitis B infection. All 80 samples were tested for hepatitis B core immunoglobulin M.

Results: One thousand blood samples were collected from 64 (6.4%) female and 936 (93.6%) male subjects. None of the blood samples contained hepatitis B core immunoglobulin M. The study found no significant differences between male and female subjects in term of HBcAb positivity.

Conclusions: Hepatitis B core antibody immunoglobulin M positivity is different in healthy blood donors of different countries according to the prevalence of chronic hepatitis B and its vaccination. Based on the current study findings, all positive samples of hepatitis B core antibody in Iranian blood banks should be considered as candidates for occult hepatitis B not just the window period infected samples.

Keywords: Hepatitis B Virus, Blood Donors, Viral DNA, Immunoglobulin M

ICIA2018\Immunology of Infectious Diseases Viruses\poster\8410
8410

Deceased serum samples have inhibitor components that distract viral RNA extraction procedure

H. Jafarinejad¹, R. Yarmohamadi¹, A. Heravian¹, A. Radmanesh^{1*}, S. Bayanolhagh^{1, 2*}

1. The Legal Medicine Research Center, Legal Medicine Organization, Iran, Tehran

2. Iranian Research Center for HIV/AIDS, Iranian Institute for Reduction of high risk Behaviour, Tehran University of Medical sciences, Tehran, Iran

Background: The blood viral RNA minikit is intended for molecular biology applications. This procedure intend for the diagnosis, treatment or research of a disease. Many of corpse's blood samples are hemolysis and have many protein that release in blood when dead and may have inhibitory role in RNA extraction procedure and decrease the efficiency of RNA production. The main of this study is how we can extract viral RNA from this cases?

Methods: During the procedure for purification of RNA from serum samples, first samples diluted in 1:10 concentration, then the solution are lysed using highly denaturing conditions that immediately inactivate RNases, allowing the isolation of intact RNA. The negative (not diluted) and positive (viral sample) control used in this study. After homogenization of the lysate by a brief centrifugation through spin column, ethanol is added to adjust binding conditions and the sample is applied to the spin column. RNA is bound to the silica membrane during a brief centrifugation step. Contaminants are washed away and total RNA is eluted in 60µl or more of RNase-free water for direct use in any downstream application.

Results: The samples that diluted in 1:10 concentration had viral RNA as positive control and detected in Real Time PCR method ($P < 0.05$) and in compared with samples that not diluted, had no viral RNA ($P > 0.05$).

Conclusion: It seems the corpse serum samples have inhibitor component that cover the spin column filter and decrease the yield of RNA product.

Key word: Deceased donors, Inhibitor components, Viral

ICIA2018\Immunology of Infectious Diseases Viruses\poster\8489
8489

Investigation of serum level of IL-35 in HIV patients and its association with co-infection with human Pegi virus (GBV-C) and Hepatitis-C Virus

Alireza Najafi¹, Mohammad Jafari², Mostafa Haji Mollahoseini³, Farah Bokharaei Salim^{4,5}, Maryam keykhaee⁶

1. Immunology Research Center, Iran University of Medical Sciences, Tehran, Iran

2. Cellular and Molecular Research Center, Gerash University of Medical Sciences, Gerash, Iran

3. Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

4. HIV Laboratory of National Center, Deputy of Health, Iran University of Medical Sciences, Tehran, Iran

5. Department of Virology, Iran University of Medical Sciences, Tehran, Iran

6. Department of pharmaceutical biomaterials, faculty of pharmacy, Tehran university of Medical sciences, Tehran , Iran

Background: Interleukin-35 (IL-35), a newly identified member of IL-12 family, functions as a crucial immunosuppressive factor in immune-mediated diseases, and the predominant mechanism of suppression is its ability to suppress T cell proliferation and effector functions. Although several previous studies investigated the role of IL-35 in infectious diseases such as hepatitis B virus and Influenza, however there are limited data available regarding the role of IL-35 in the immunopathogenesis of HIV. The aim of this study was to determinate the immune activation status in HIV patients with GBV-C and/or HCV coinfection through measuring IL-35 plasma levels.

Methods: A case-control study was carried out on 50 HIV infected patients (10 HIV mono-infected, 22 HIV/HCV co-infected, 7 HIV/GBV-C co-infected, and 11 HIV/HCV/GBV-C co-infected) and 30 matched healthy control. Plasmas were analyzed for HCV using serologic test and GBV-C by reverse transcriptase polymerase chain reaction (RT-PCR) and the level of IL-35 was measured by ELISA assay.

Results: The level of plasma IL-35 in all patients with HIV infection was more than healthy control (P: 0.011). In addition, HIV/HCV co-infected subgroup had an elevated IL-35 level in comparison to healthy controls (P: 0.008). Furthermore, the concentration of IL-35 was significantly higher in HIV/HCV/GBV-C co-infected patients compared with healthy controls (P: 0.0229).

Conclusion: These findings suggest that IL-35 may be a role in immunopathogenesis of HIV/HCV and HIV/HCV/GBV-C co-infection. Our results provide a basis for further investigation in the role of IL-35 in HIV infection.

Keywords: Interleukin (IL)-35, HIV, GBV-C AND HCV, co-infection

ICIA2018\Immunology of Infectious Diseases Viruses\poster\8591
8591

The Frequency of Parainfluenza virus infection in Acute Respiratory infections

Morteza Sagharjoghi Farahani ¹, Arash Kalantari ², Babak Khodadadi ^{*3}, setare Mamishi ⁴

1. Young researchers and Elite Club, Islamshahr Branch, Islamic Azad University, Islamshahr, Iran

2. Assistant professor of Tehran University of medical sciences. Imam Khomeini hospital complex vali-e-asr hospital Department of Pediatrics, Tehran, Iran

3. Young researchers and Elite Club, Khorramabad Branch, Islamic Azad University, Khorramabad, Iran

4. Department of Pediatrics, Division of Infectious Disease. Markaz Tebbi Hospital Tehran University of Medicine Sciences

Background: Acute respiratory infections are one of the most common causes of morbidity in children. Parainfluenza viruses are important viral pathogens causing upper and lower respiratory infections in children. There are four type of Parainfluenza. The most common cause of laryngotracheobronchitis is parainfluenza type 1. The parainfluenza virus is transmitted via arousal and direct contact between children. The aim of this study was to determine the prevalence of parainfluenza virus infection among children aged less than 5 years referred to Markaz Tebbi Hospital.

Methods: Our diagnostic method for evaluating an outbreak of parainfluenza virus infection is to investigate the nasopharyngeal secretion of patients with immunofluorescence antibody. Also, the most common age group involved and various clinical symptoms are recorded.

Results: In our study, the frequency of respiratory tract infection due to parainfluenza virus was 26%. The age group of 25 to 36 months was the most commonly involved age group, the most common symptom was rhinorrhea(96%.) and the most common clinical sign was fever (68%).

Conclusions: This study demonstrate that the prevalence of parainfluenza virus and the most commonly occurring age is similar to other studies. Therefore, in this age group of children, the clinical symptoms of rhinorrhea and fever should lead us to parainfluenza virus infection and effective prevention strategies for respiratory virus infection should be done.

Keywords: parainfluenza virus, immunofluorescence antibody, respiratory infections.

ICIA2018\Immunology of Infectious Diseases Viruses\poster\8609
8609

Association between Interleukin-18 gene Polymorphism (rs5744249 A/C) and Susceptibility to Chronic Hepatitis B Virus Infection

Haleh Sarrafnia¹, Seyed Reza Mohebbi¹, Shabnam Kazemian¹, Mahsa Saeedi Niasar¹, Afsaneh Sharifian¹, Hamid Asadzadeh Aghdaei², Mohammad Reza Zali¹

1. Gastroenterology and Liver Diseases Research Center, Research Institute for Gastroenterology and
2. Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran
3. Basic and Molecular Epidemiology of Gastrointestinal Disorders Research Center, Research In
4. stitute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Infection with hepatitis B is one of the most notable causes of chronic liver disease. The outcome of hepatitis B virus (HBV) infection is mainly influenced by host genetic background such as single nucleotide polymorphisms (SNPs). Recent studies have indicated that gene polymorphisms in cytokines are associated with the development of chronic HBV infection and progression of the infection. Interleukin-18 is a proinflammatory cytokine that plays a significant role in immune responses against viruses. In this study, we aimed to explore whether interleukin-18 (IL-18) gene single nucleotide polymorphism (rs5744249) is associated with the outcome of hepatitis B virus (HBV) chronic infection or not.

Methods: In this case-control study, blood samples were collected from 100 chronic HBV infected patients and 100 healthy controls. Selection criteria for patient group were anti-HBc antibody (anti-HBcAb) and HBsAg positivity for more than 6 months and the control group consisted of anti-HBcAb and HBsAg negative individuals. After genomic DNA extraction, the SNP was genotyped by the PCR and restriction fragment length polymorphism (RFLP) method and then we used SPSS for statistical analysis.

Results: The results showed rs5744294 genotype frequencies of AA=81.7%, AC=15.8% and CC=2.5% in chronic patients and AA=75.8%, AC=22.5% and CC=1.7% in the control group. Results revealed that no difference was observed in the frequencies of neither genotype (P value=0.396) nor allele (P value=0.888) frequencies between two groups.

Conclusion: No correlation was detected between interleukin 18 gene Polymorphism rs5744249 A/C with hepatitis B virus infection or disease progression and that it could not be regarded as a host genetic factor associated with the hepatitis B virus infection. Other IL-18 SNPs are possibly more notable than the one studied.

Keywords: Interleukin-18, Hepatitis B, Single nucleotide polymorphism, rs5744249

ICIA2018\Immunology of Infectious Diseases Viruses\poster\8648
8648

Seroepidemiology of Varicella Zoster Virus among nursery students in Langroud Nursing Faculty

Reza Jafari-Shakib¹, Ashkan Rasouli², Mojtaba Farahbakhsh³, Mahtab Gholamzadeh², Zahra Mohtasham Amiri⁴

1. Department of Immunology, Faculty of Medicine, Guilan University of Medical Sciences, Rasht

2. Faculty of Paramedicine, Guilan University of Medical Sciences, Rasht

3. Department of Medical Biotechnology, Faculty of Paramedicine, Guilan University of Medical Sciences, Rasht

4. Department of Community and Preventive Medicine, Guilan University of Medical Sciences, Rasht

Background: This study was done to investigate anti-VZV (varicella zoster virus) antibody among nursery students as a group of health care workers who can transmit the infection to coworkers or patients.

Methods: The anti-VZV IgG was measured in all nursery students in Langroud Nursing Faculty who wanted to participate in this cross-sectional study. The antibody was measured in 101 students by ELISA method.

Results: Seventy-three (72.3%) students had anti-VZV IgG, 27 students (26.7%) were seronegative and one students have borderline IgG.

Conclusion: There is a significant percentage of nursery student seronegative and they are susceptible to VZV infection. They can get infection from other staffs and patients in hospital and also transmit infection to others especially immunocompromised patients. Therefore, it is recommended to screen anti-VZV antibody in nursery students and vaccination seronegative ones.

Keywords: Varicella zoster virus, Nursery students, Seroepidemiology, Langroud

ICIA2018\Immunology of Infectious Diseases Viruses\poster\8649
8649

Beclin1 inhibition promotes autophagy and decreases daclatasvir-induced apoptosis in Huh7 cells

Somaieh Nasereslami ^{1*}, Taravat Bamdad ¹, Asghar Abdoli ²

1. Department of Virology, School of Medical Sciences, University of Tarbiat Modares, Tehran, Iran

2. Department of Virology, Pasture Institute, Tehran, Iran

Background: Beclin1 is a well-known key regulator of autophagy, which is also a haploinsufficient tumor suppressor. Current studies revealed that down-regulation or monoallelic deletions of beclin1 were frequently found in various cancers. The purpose of this study was to investigate the effects of beclin1 inhibition on autophagy and daclatasvir-induced apoptosis of pancreatic cancer cells.

Methods: Beclin1 expression was inhibited by siRNA transduction and gene expression was determined by Real-time PCR and Western blot. The effects of beclin1 inhibition on autophagy and daclatasvir-induced apoptosis of Huh7 cells were analyzed through LC3 expression, cell viability, cell cycle and apoptosis using Western blot.

Results: We observed that beclin1 silence promoted microtubule-associated protein 1 light chain 3-II (LC3-II) protein formation and increased punctate fluorescent signals in Huh7 cells transfected with green fluorescent protein (GFP)-tagged LC3. Beclin1 inhibition showed a greater suppressive effect on daclatasvir-induced apoptosis of Huh7 cells.

Conclusion: Our data suggest that beclin1 silence not only up-adjusted autophagy process, but also played an important role in the regulation of apoptosis. Beclin1 inhibition could inhibit apoptosis signaling induced by daclatasvir in Huh7 cells.

Keywords: Daclatasvir, Huh7, Beclin1, LC3, Apoptosis

ICIA2018\Immunology of Infectious Diseases Viruses\poster\8654
8654

Evaluation of anti-CMV antibody level in women of reproductive age

Shayesteh Z¹, Yazdani Neyshabouri Sh², Beikzadeh L^{3*}, Vazeh H⁴, Farahmand M⁵

1. Msc of medical Immunology, school of allied medical sciences, karaj, iran
2. PhD of virology Islamic Azad University-Tehran Medical Branch Department of Microbiology
3. Msc of medical virology, school of allied medical sciences, karaj, iran (correspond)
4. Msc of medical virology, Golestan University of Medical Sciences, Department of Microbiology
5. Department of virology, school of public health, Tehran university of medical sciences, Tehran, Iran

Background: One of the herpes family with the potency to life-extensive latency within the host is cytomegalovirus (CMV) that have double-stranded DNA genome. Alike all herpes viruses, HCMV undertake latency and reactivation within the host and has been appeared to pollute a broad cells in vivo. CMV inborn contaminations might be symptomless or may prompt hearing debilitation, mental hindrance, cerebral paralysis, and neurodevelopmental disabilities. The frequency of inborn CMV contamination is around 1%-7% of births. Anticipating the risk of CMV vertical transmission and planning preventive measures are facilitated by knowing these characteristics.

Methods: Among of 360 women in age of fertility attending health center chosen randomly from all women attending for test during the study period. Peripheral blood (Δ ml) was aspirated from them and CMV serology test was done using a standard ELISA kit to determine CMV IgM and IgG antibodies.

Results: The total number of participants in this study was 360. The evaluation of IgG concentration is indicated. According to the cut-off determined by the kit, which is 16.5, 280 specimens are positive and 80 cases are IgG negative. A Fisher's comparison test was used to compare IgG between two age groups. The results showed that the observed difference between the two age groups was not statistically significant ($p = 0.79$).

Conclusion: According to the findings, it can be seen that, there is not anyone in participants who is in the acute phase of diseases. Also, comparing IgG positive cases in two age groups >30 and <30 years old, there was no significant difference in age variable. Overall, 77% of cases were IgG positive, indicating a chronic phase of the disease in participants.

Key words: Reproductive age, antibody level, CMV



ICIA2018\Immunology of Infectious Diseases Viruses\poster\8697
8697

Hepatitis C virus associated mixed cryoglobulinemia

Ramin Farajpour¹, Shiva Farajpour²

1. Department of Neuroscience Research Center, Faculty of Medicine, Tabriz University of Medical Sciences
2. Department of Ophthalmology, Faculty of Medicine, Tabriz University of Medical Sciences

Background: Mixed cryoglobulinemia (MC) is common in patients with chronic hepatitis C virus (HCV) infection. However, clinically significant vasculitis occurs only in a minority of these patients.

Methods & results: We hereby report a driver man with chronic HCV infection and MC presenting with polyarthralgia and skin purpura of the lower limbs. He later developed clinical remission and anti-viral therapy resulted in good clinical response.

Conclusion: The classical presentation of MC is a triad of cutaneous vasculitis, peripheral neuropathy and arthralgia. Anti-viral therapy is the standard treatment and good response depends on viral clearance.

Keywords: Hepatitis C, Mixed Cryoglobulinemia, Anti-viral therapy

ICIA2018\Immunology of Infectious Diseases Viruses\poster\9757
9757

Polyclonal antibody preparation against bovine ephemeral fever virus

Shokoofeh Almasi¹, Dr. Mehran Bakhshesh ^{*1}, Dr. Mohammad Mehdi Ranjbar¹

1. Department of viral animal disease research and diagnosis, Razi vaccine and serum research institute, Agricultural Research, Education and Extension Organization (AREEO), Karaj, Iran

Background: Considering the numerous applications of polyclonal antibodies in research and clinical fields, the development of methods for the preparation of polyclonal antibodies against pathogens is very important. In this research, a method for producing polyclonal antibodies against the whole viral particle of the *Rhabdovirus*, bovine ephemeral fever virus (BEFV), is introduced.

Methods: Six rabbits of Dutch breed were selected and divided into experimental and control groups. Each rabbit from the experimental group received at least 10^6 TCID₅₀ of the whole viral particles intravenously without adjuvant while the control group was injected with the cell culture supernatant free from the virus. The injections were administered at 0, 1 and 4 weeks intervals, and two weeks after the last injection the rabbits were bled and their sera were separated by centrifugation. The sera were tested by virus neutralization test (VNT), calculating the titer of antibodies to the live virus.

Results: This schedule resulted in a high titer of polyclonal antibodies to BEFV particles. Based on Karber formula, the rabbits from experimental group produced 288 to 290 titer of neutralizing antibodies as no titer of antibodies was detected in control group.

Conclusion: Bovine ephemeral virus has recently been isolated in Iran and studies are now being undertaken on characterization, epidemiology and particularly manufacturing a potent vaccine for prophylaxis against the virus. Therefore, the hyperimmune serum as a biological product of this research will definitely be exploited in the experiments required for these studies as well as possible commercial applications in the future.

Keywords: Bovine ephemeral fever, hyperimmune serum, rabbit

ICIA2018\Immunology of Infectious Diseases Viruses\poster\9803
9803

Interferon-inducible protein interferes with HCV replication through the autolysosomal degradation of NS5A

Somaieh Nasereslami ^{1*}, Taravat Bamdad ¹, Asghar Abdoli ²

1. Department of Virology, School of Medical Sciences, University of Tarbiat Modares, Tehran, Iran

2. Department of Virology, Pasture Institute, Tehran, Iran

Background: SCOTIN overexpression inhibits HCV replication and infectious virion production in cells infected with cell culture-derived HCV. HCV nonstructural 5A (NS5A) protein, which is a critical factor for HCV RNA replication, interacts with the IFN- β -inducible protein SCOTIN, which transports NS5A to autophagosomes for degradation. Furthermore, the suppressive effect of SCOTIN on HCV replication is impaired in both ATG7-silenced cells and cells treated with autophagy or lysosomal inhibitors. SCOTIN does not affect the overall flow of autophagy, however, it is a substrate for autophagic degradation. The physical association between the transmembrane/proline-rich domain (TPMRD) of SCOTIN and Domain-II of NS5A is essential for autophagosomal trafficking and NS5A degradation. Altogether, our findings suggest that IFN- β -induced SCOTIN recruits the HCV NS5A protein to autophagosomes for degradation, thereby restricting HCV replication.

Methods: In the present study, we propose that SCOTIN is one of the ISGs that interferes with HCV replication. IFNs type I exhibit most of their antiviral activities via hundreds of ISGs, which involve various proteins, including enzymes, transcription factors, heat-shock proteins and apoptotic proteins. SCOTIN expression was induced in Huh-neo cells treated with IFN- β , and the suppressive effect of IFN- β on HCV replication was reduced when SCOTIN expression was silenced. These results indicate that SCOTIN acts as one of the ISGs that contribute to the antiviral activity of interferon- β that is directed against HCV.

Results: IFN- β -inducible protein SCOTIN restricts HCV replication and SCOTIN promotes autophagosomal degradation of NS5A.

Conclusion: In this report, we demonstrate that an IFN- β -inducible host factor, SCOTIN, contributes to suppressing HCV replication via the autolysosomal degradation of the HCV NS5A protein. SCOTIN is an ER transmembrane protein that physically associates with NS5A and is also a substrate for autophagy-mediated degradation. Similar to the role of p62 autophagy receptor in the autophagy-mediated clearance of intracellular viruses and bacteria, SCOTIN recruits NS5A to LC3-II-containing autophagosomal compartments through physical interactions. These physical interactions and the colocalization of SCOTIN and NS5A in autophagosomes are critical for controlling the autophagy-mediated proteolytic degradation of the NS5A viral protein and HCV replication.

Keywords: IFN- β , HCV, NS5A, LC3, SCOTIN.

ICIA2018\Immunology of Infectious Diseases Viruses\poster\9809

9809

HTLV-1 infected asymptomatic carriers overexpress the cell-mediated cytotoxicity genes

Asadollah Mohammadi¹, Bahare Fazeli², Reza Boostani³, Houshang Rafatpanah²

1. Cellular & Molecular Research Center, Faculty of Medicine, Kurdistan University of Medical Sciences, Sanandaj, Iran

2. Inflammation and Inflammatory Diseases Research Center, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

3. Department of Neurology, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

Background: HTLV-1-associated myelopathy/tropical spastic paraparesis (HAM/TSP) is a chronic viral neuroinflammatory disease. Apoptosis plays an important role in the control of viral infection, this process occasionally makes it possible to distinguish asymptomatic carriers (ACs) from HAM/TSP patients.

Methods: The mRNA expression of Fas, FasL, TRAIL, perforin, granzyme A, granzyme B, and granulysin in the peripheral blood mononuclear cells (PBMCs) of 21 HAM/TSP patients, 21 ACs, and 21 healthy controls (HCs) were quantified. The Fas, FasL, TRAIL, granulysin serum levels, and the granulysin secretion level were evaluated and their interaction with the HTLV-1 proviral load (PVL), Tax and HBZ mRNA expression were also analysed.

Results: Fas, FasL, TRAIL, perforin, and granzyme B were significantly overexpressed in ACs whereas granulysin secretion significantly decreased in ACs and HAM/TSP patients. HAM/TSP patients have a higher PVL, Tax, and HBZ expression than ACs. Also, negative correlations between the mRNA expression of the Fas, FasL, TRAIL, perforin, granzyme B, and granulysin with HBZ levels were observed.

Conclusions: We found that ACs overexpress the genes which are involves in cell-mediated cytotoxicity. We conclude that successful suppression of the HTLV-1 PVL, Tax and HBZ expression, and control of disease progression is associated with the strong cytotoxic activity in the PBMCs of ACs.

Keywords: HTLV-1, Neuroinflammatory disease, HAM/TSP, Apoptosis.

ICIA2018\Immunology of Infectious Diseases Viruses\poster\9821
9821

Frequency of influenza-A-H1N1 in patients with Acute Pneumonia admitted to Loghman Hakim Hospital

Helya Sadat Mortazavi ¹; Abdolkarim Pajoumand ²; Seyed Hamidreza Monavari ²; Davood Javanmard ²; Latif Gachkar ³; Meinoosh Shabani ^{3*}

1. Department of Medical Virology, School of Medicine, Iran University of Medical Sciences, Tehran, Iran

2. Department of Clinical Toxicology, Loghman Hakim Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran

3. Infectious Disease and Tropical Medicine Research Center, Shahid Beheshti University of Medical Sciences and Health Services, Tehran, Iran

Background: Here we scrutinized the incidence of Influenza-A-H1N1-related pneumonia in community-acquired pneumonia (CAP) and hospital-acquired pneumonia (HAP) at Loghman Hakim hospital, Tehran, Iran.

Methods: In this prospective study from November 22, 2016, to June 21, 2017, patients suspected of seasonal Influenza and confirmed CAP-HAP included. Samples tested using rapid antigen tests and quantitative Real-time PCR assay. P-value < 0.05 was considered significant.

Results: A total of 29 suspected cases of seasonal influenza had obtained either CAP or HAP. Of whom 51.7% were male, and the mean age was 60.07 ± 20.94 years. The most common findings were fever and chills, cough and phlegm (15 cases, 29.41%). The highest number of admission occurred in February (48.3%). Mean days of hospital stay was 11.31 ± 17.59 days and the average days of treatment with antibiotics were 10.78 ± 17.13 days. 6.9% were positive in QReal-Time PCR assay (> 30 copies/ml). The highest prescribed antibiotic combination was ceftriaxone + Azithromycin, and only 31.3% of patients received empiric oseltamivir therapy. Also, 27.6% of patients were drug users. Consolidation (44.8%) was the most predominant radiological finding. We found a significant correlation in patient's systolic blood pressure, viral load, white blood cell count, BUN, CURB65 and radiological findings.

Conclusion: This study highlights the importance of the empiric influenza treatment before receiving the results of molecular tests in suspected cases. Moreover, precautions, including influenza and pneumococcal vaccination, should be emphasized by health care workers.

Keywords: Influenza A Virus, H1N1 Subtype, Quantitative Real-Time Polymerase Chain Reaction, Pneumonia

ICIA2018\Immunology of Infectious Diseases Viruses\poster\9917
9917

Association between Chronic Hepatitis B Virus Infection and Polymorphisms of rs9277535 in HLA-DP in the Southern Khorasan population (East of Iran)

Hossein Javdani^{1,2}, Abdolghader Taene^{1,2}, Ali Farahi^{1,2}

1. Student Research Committee, Birjand University of Medical Sciences, Birjand, Iran

2. Cellular and Molecular Research Center, Birjand University of Medical Sciences, Birjand, Iran

Background: Chronic hepatitis B virus (HBV) infection is a major global health concern. It is assessed that two billion individuals worldwide are tainted with hepatitis B infection and approximately 350 to 400 million people are chronic carriers of HBV. The clinical results of HBV disease extend from asymptomatic disease to constant hepatitis, liver cirrhosis and hepatocellular carcinoma. The genes encoding HLA are the most polymorphic in the human genome, probably as a way to be able to respond to all capacity overseas antigens. In the present study, we aim to explore the association of the HLA-DP polymorphisms rs9277535 with HBV susceptibility in the Southern Khorasan population.

Methods: In our study, we used 60 patients with chronic HBV infections as a group of patients (HBsAg positive test results for at least 6 months) and 120 patients who have previously recovered from HBV infection, spontaneously (HBc-positive antibody / HBsAg negative) as a control group. The amplified PCR products were purified using PCR Purification Kit. To determined SNPs of HLA-DP (rs9277535), the purified PCR products were sequenced using the Applied Biosystems 3730/3730XL DNA analyzer sequencing.

Results: For the rs9277535 polymorphism, the patients carrying A allele or AT genotype had a higher risk of chronic HBV infection than the other patients carrying G allele or GT genotype. ($P<0/005$).

Conclusion: In conclusion, this study demonstrated that HLA-DP rs9277535 polymorphisms are strongly associated with HBV susceptibility in the Southern Khorasan population.

Keywords: Chronic hepatitis B, Immune response, Polymorphism, Human leukocyte antigens DP (HLA-DP)

ICIA2018\Immunology of Infectious Diseases Viruses\poster\9921
9921

Relation between IFN- γ polymorphism and the outcome of hepatitis B infection in an Iranian population

Ali farahi^{1,2}, Abdolghader Taene Taene^{1,2}, Hossein Javdani Javdani^{1,2}

1. Student Research Committee, Birjand University of Medical Sciences, Birjand, Iran

2. Cellular and Molecular Research Center, Birjand University of Medical Sciences, Birjand, Iran

Background: The chronic HBV infection is considered as a multifactorial issue, which can prompt to a vast variety of clinical outcomes. The host's immune system is the most crucial factor in viral persistence. IFN- γ is one of the immune factors that produce helpful substances essential for the protection against viral infections. The current study aimed to investigate the relation between a SNPs (rs2430561) of the IFN- γ first intron and the outcome of hepatitis B infection in an Iranian population.

Methods: Blood samples of 60 chronically HBV infected patients (cases) and 60 healthy subjects with the history of HBV infection (controls) were collected. Genomic DNA was extracted by salting-out method and DNA analysis genotype identification was performed by ARMs-PCR.

Results: The results indicated that the SNP frequency had a significant difference between the case and control groups. As the frequency of allele A was 73.3% and 46.7% in cases and controls, respectively.

Conclusion: This study demonstrate a significant relationship between the IFN- γ (+874T/A, rs2430561) polymorphism and chronic HBV infection susceptibility in a sample of Iranian population. The result of current study suggests that carriers of mutant allele A at position +874 are more susceptible to chronic HBV infection and carriers of wild-type allele T at position +874 are predisposed to recover from HBV infection.

Keywords: IFN- γ polymorphism, hepatitis B infection, Iranian population

ICIA2018\Immunology of Infectious Diseases Viruses\poster\9972
9972

Evaluation of HIV Viral Load and CD4 count distribution among HIV infected patients in co-infection with HCV and Human pegivirus (GBV-C)

Mohammad Jafari¹, Alireza Najafi², Mostafa Haji Mollahoseini³, Farah Bokharaei Salim^{4,5}, Maryam keykhaee⁶, Roghayeh babaei⁴

1. Cellular and Molecular Research Center, Gerash University of Medical Sciences, Gerash, Iran
- Immunology Research Center, Immunology and Infectious Disease Institute, IRAN University of Medical Sciences
2. Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran
3. HIV Laboratory of National Center, Deputy of Health, Iran University of Medical Sciences, Tehran, Iran
4. Department of Virology, Iran University of Medical Sciences, Tehran, Iran
5. Department of pharmaceutical biomaterials, faculty of pharmacy, Tehran University of Medical sciences, Tehran, Iran

Background: Systemic immune activation is critical to the pathogenesis of HIV disease, and is accentuated in HIV/HCV co-infected patients. On the other hand, harmful or beneficial effect of GBV-C on HIV progression is controversial. In this study, we examine immune pathogenesis of HIV co-infection through HIV viral load detection and CD4 percentage measurement in different infectious profile.

Methods: A case-control study was carried out on 50 HIV infected patients (10 HIV mono-infected, 22 HIV/HCV co-infected, 7 HIV/GBV-C co-infected, and 11 HIV/HCV/GBV-C co-infected). Plasmas were analyzed for HCV using serologic test and GBV-C by reverse transcriptase polymerase chain reaction (RT-PCR). The determination of the level of HIV-1 viral load was tested by an Artus HIV-1 RG RT-PCR Kit in accordance with the manufacturer's instructions. Whole blood analyzed for the percentage of CD4+ and CD8+ T-cells by CD3/CD4+ and CD3/CD8+ double staining method and flowcytometric analysis on lymphocyte population gate.

Results: HIV viral load in 36% of HIV/HCV/GBV-C sub group was detectable. In addition, patients with HIV/HCV co-infection had 13% HIV viremia. In monoinfected patients 10% of them had detectable viraemia. Surprisingly, The HIV virus in sample of HIV/GBV-C co-infected patients was undetectable. Furthermore, the average of CD4 percentage in HIV/HCV/GBV-C, HIV/HCV, HIV mono-infected, and HIV/GBV-C co-infected was 16.5%, 18%, 23, 4%, and 18% respectively.

Conclusion: These findings suggest that co-infection complicates the clinical progress of HIV in infected patients, and may also harmfully affect on treatment of HIV infection.

Keywords: HIV co-infection, HIV viral load, CD4 count.

ICIA2018\Immunology of Infectious Diseases Viruses\poster\11073
11073

Rotavirus infection induces nitric oxide production by Caco-2 cells

Behnoush Baradaran¹, Sara Soudi¹, Hoorieh Soleimanjahi¹

1. Department of Immunology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

Background: Intestinal epithelial cells are the first barrier against intestinal pathogens. Rotavirus is the leading cause of severe gastroenteritis in infants and young children that infects villus epithelial cells of small intestine. After virus entry to the cells, signaling pathways are activated, resulting in the induction of proinflammatory cytokine production and increase in oxidative stress. In this research the effect of rotavirus on nitric oxide production by Caco-2 cells were measured.

Methods: To this purpose, Caco-2 cells were cultured at 3×10^5 /ml in each well of 12 well plates. Rotaviruses were added at 10^4 , 10^3 , 10^2 virus/ml to the cells in triplicate. No virus treated group were considered as control. 24 and 48 hours post virus treatment, supernatant of the cells were collected and amount of the NO production were measured by griess method.

Results: According to obtained results, 10^4 , 10^3 , 10^2 and no virus treated groups produced the mean amount of 20.47, 18.5, 18.5 and 12.6 nM of nitric oxide after 24 hours of infection, respectively. This amount were increased significantly to 30.0 and 28.2 nM in 10^4 and 10^3 virus treated groups, but no significant differences were seen in both 10^2 virus treated and no treated groups.

Conclusion: Nitric oxide production by epithelial cells may insert both protective effect against infection and negative effect on host cells by induction of excessive oxidative stress. The result showed that nitric oxide production by Caco-2 cells are dependent to both primary infection dose and duration of pathogen-host interaction.

Keywords: Rotavirus, Caco-2 cells, nitric oxide



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Immunology of Mycobacterium Tuberculosis

Poster Presentation

ICIA2018\ Immunology of Mycobacterium Tuberculosis\poster\7969
7969

Evaluation of Host Interaction and *Mycobacterium Tuberculosis* in Outbreak of Tuberculosis Disease with Evaluation of MMP-3 and MMP-9 Genes Expression in Host and ESAT-6 in Microbe in Patient of TB

Shadi Abbasnia¹, Kiarash Ghazvini², Saeed Amel Jamehdar², Narges Valizadeh¹, S.A.R Rezaee¹

1. Inflammation and Inflammatory Disease Research Center, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

2. Department of Microbiology & Virology, Mashhad University of Medical Sciences, Mashhad, Iran

Background: Tuberculosis (TB) caused by infection of *Mycobacterium tuberculosis* (*Mtb*). Different manifestations of TB result from interactions between the bacteria and host defense. A family of host molecules which may complicate in TB manifestation by extracellular matrix degradation is matrix metalloproteinases (MMPs). In this study, the interaction of host and *Mtb* was evaluated by assessing MMP-9 and MMP-3 gene expression in host and ESAT-6 in *Mtb*.

Method: Lung lavage fluid from 30 patients with clinical symptoms of tuberculosis but negative direct and culture *Mtb* tests, was taken by a pulmonologist. Out of 30 patients, 14 were TB⁺ and 16 TB⁻. After extracting the RNA and synthesis of cDNA, expression of MMP-3 and MMP-9 in host and ESAT-6 in *Mtb* were evaluated by TaqMan Real Time PCR.

Results: The findings show that MMP-9 gene expression in TB⁺ and TB⁻ were (2.56±0.68) and (1.13±0.35), respectively, which the higher expression (2.26 times) in TB⁺ was significant (p=0.05). Gene expression of MMP-3 in TB⁺ was (0.22±0.09) and in TB⁻ was (0.64±0.23) which the difference was not statistically significant.

ESAT-6 gene expression in patients with TB⁺ was (0.28±0.091). In addition, although, there was not any correlation between MMP-3 or 9 with ESAT-6, there were significant correlations between MMP-9 and WBC (R=+0.61, P=0.02), BMI (R=-0.59, P=0.02) and albumin (R=+0.68, P=0.007), and also between MMP-3 and INOS (R=+0.94, P≤0.001), *Mtb*-Ag8 (R=+0.92, P≤0.001) and T-bet (R=+0.84, P=0.002), WBC (R=-0.60, P=0.04), PMN (R=-0.86, P=0.001) and lymphocytes (R=+0.80, P=0.003).

Conclusion: The results showed that increased expression of MMP-9 in TB⁺ could be complicated in lung injuries in TB, which is a good target for therapy. Moreover, ESAT-6 seems to have no effect on lung damage.

Keywords: *Mycobacterium tuberculosis*, ESAT-6, MMP-3, MMP-9, Taqman Real-Time PCR



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Immunology of Rheumatic Diseases

Poster Presentation

ICIA2018\Immunology of Rheumatic Diseases\poster\5326
5326

The correlation between vitamin D plasma levels with Foxp3 gene expression in Rheumatoid arthritis patients

Parisa Zafari¹, Nasrin Iranshai², Mahdi Taghadosi²

1. Department of immunology, school of medicine, Mazandaran University of Medical Sciences

2. Department of immunology, school of medicine, Kermanshah University of Medical Science

Background: Rheumatoid arthritis (RA) is an autoimmune disease with an inflammatory nature that affects the small joints of the hands and feet. It seems that the function of Regulatory T cells (Tregs), the main cells in controlling and inhibiting of inflammation, have been compromised in RA. Foxp3 expression as major transcription factor of these cells depends on variety of environmental and internal factors. Vitamin D is one of the environmental factors that in addition to its regulatory role in the immune system, it is also important to induce Foxp3 expression. Therefore, the aim of this study was to investigate the effect of plasma levels of vitamin D on the expression of Foxp3 gene in patients with RA.

Methods: 20 untreated RA patients and 40 healthy subjects participated in this study. Plasma levels of vitamin D were measured by competitive ELISA method. Foxp3 transcription factor gene expression was also measured using real-time PCR technique.

Results: As expected, Foxp3 expression was lower in RA patients than healthy subjects but there was not a significant correlation between vitamin D plasma levels and Foxp3 gene expression.

Conclusion: Vitamin D deficiency cannot be considered as an environmental factor affecting RA pathogenesis unlike undeniable effect of this vitamin in pathogenesis of other autoimmune disease such as multiple sclerosis.

Keywords: Foxp3, rheumatoid arthritis, vitamin D

ICIA2018\Immunology of Rheumatic Diseases\poster\6313
6313

Analysis of the gene expression of Helios and the methylation status of Foxp3 TSDR sequence in the newly diagnosed Rheumatoid Arthritis patients

Mahdi Taghadosi¹, Parisa Zafari², Nasrin Iranshahi¹

1. Department of immunology, school of medicine, Kermanshah University of Medical Science

2. Department of immunology, school of medicine, Mazandaran University of Medical Sciences

Background: The control of auto-reactive cells may be compromised in Rheumatoid Arthritis (RA). Regulatory T cells (Tregs), key controller of immune system, may have impaired function in RA disease. Foxp3 is a master regulator of Treg cells which its expression is under tight control of epigenetic mechanisms. By analysis of epigenetic modulation of Foxp3 Treg-specific demethylated region (TSDR) and Helios gene expression, the present study aims to evaluate Treg cells alterations in RA disease.

Methods: Twenty newly diagnosed RA patients and 41 healthy subjects participated in this study. Foxp3 and Helios expression was measured using real-time PCR technique. Methylation level of TSDR was analyzed by bisulfite treatment and Quantitative methylation specific PCR (Q-MSP) method.

Results: The gene expression of Foxp3 and demethylation rate of Foxp3 TSDR sequence was decreased in RA patients compared with healthy subjects ($P < 0.001$ and $P = 0.006$ respectively). The Helios expression was significantly higher in RA patients compared to healthy controls ($P = 0.048$). Positive correlation between Foxp3 expression with Helios expression and demethylation rate of TSDR was detected ($P = 0.016$ and $P = 0.010$ respectively).

Conclusion: Our results suggest that epigenetic alterations may have an important role in the pathogenesis of RA.

Keywords: Rheumatoid arthritis, Foxp3, Helios, epigenetic, TSDR

ICIA2018\Immunology of Rheumatic Diseases\poster\7351
7351

Determination of IL-17A & IL-17F gene polymorphisms and serum levels of Anti-CCP in patients with Rheumatoid arthritis

M.Khojasteh¹, R.Mansori¹, A.Dehghan¹, M.Mahmoodi², M.Vakili¹, A.Shahmohammadi¹, Mazarzadeh¹

1. Department of Immunology, International campus of Yazd, Iran

2. Department of Rheumatology, Medical university of Tehran, Iran

Background: Rheumatoid arthritis (RA) is a chronic, systemic inflammatory autoimmune disorder that is symmetric in small and sometimes large joints. This disease affects about 1% of the world's population. Studies have shown RA is a multifactorial disease that is the result of the interaction of genetic and environmental factors. Genetic factors have a special role in severity disease. On the other hand the disease process triggers a cascade of inflammatory cytokines leading to activation of autoreactive inflammatory cells, which ultimately leads to the destruction of bone and cartilage. IL17 is the main pro-inflammatory cytokine that secrets from Th17 and expresses in synovium of joint fluid in RA patients. IL17A and IL17F (two main family members of IL17) induce expression of cytokines and other inflammatory processes in bone and involved in rheumatoid arthritis. Therefore, in this study, single nucleotide polymorphism rs2275913 of gene IL17A and single nucleotide polymorphism rs763780 of IL17F gene in patients and controls were examined and compared.

Methods: In this study, 100 patients with RA and 92 healthy controls were selected for IL17A and 116 patients with RA and 107 healthy individuals were selected for IL17F. DNA was extracted from peripheral blood mononuclear cells by DNA extraction kit. Primers were designed by PRIMER 3 software and then the appropriate restriction enzymes were selected. Allele frequencies and genotypes of these polymorphisms were analyzed using PCR-RFLP technique. After statistical analysis, a significant association between rs2275913 single nucleotide polymorphism with the risk of disease was observed (P- Value:0.0001,OR:2.97,CI:95%). But there was no significant association between rs763780 and risk of disease.(P-Value: 0.201)

Results: Also in this study, the relationship between the genotypes mentioned above and Anti-ccp antibodies was investigated. Anti-ccp is a specific and offensive diagnostic marker for rheumatoid arthritis that can predicts disease, there was no significant correlation between the genotypes and positive Anti-ccp.

Conclusion: Based on this study, rs2275913 polymorphism of IL17A gene may be related to the risk of rheumatoid arthritis in Iranian patients.

Keywords: rheumatoid arthritis, IL 17, single nucleotide polymorphism

ICIA2018\Immunology of Rheumatic Diseases\poster\7487
7487

T helper subsets and regulatory T cell in different groups of Rheumatoid Arthritis patients

Saeid Taghiloo¹, Abolghasem Ajami^{1,2*}, Mohsen Tehrani^{1,2}, Arezou Abbasi¹, Reza Alizadeh Navaei², Mohsen Akhiani³, Alireza salami⁴

1. Department of Immunology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran
2. Molecular and Cell-Biology Research Center, Mazandaran University of Medical Sciences, Sari, Iran
3. Alborz Hospital, Alborz University of Medical Sciences, Karaj, Iran
4. Department of Biotechnology, School of Advanced Technologies in Medicine, Tehran University of Medical Sciences, Tehran, Iran

Background: Rheumatoid arthritis (RA) is one of the most common autoimmune diseases in the world. Recent studies have indicated the imbalance of Th subsets and Treg activity in development, progressing and remission of disease. In this study, we investigate the expression gene of four major transcription factors T-bet (Th1), GATA (Th2), RORc (Th17) and Foxp3 (T reg) in peripheral blood of different group of RA patients.

Methods: In a case-control study relative gene expression of transcription factor of Th subsets and T reg were investigated in patients and controls. Twenty newly diagnosed patients, 20 patients under treatment, 20 patients in remission state, 20 patients with osteoarthritis and 20 age and sex matched healthy controls were selected. In this study diagnosis and classification of patients was done according to American college of Rheumatology criteria. two ml of peripheral blood collected from all patients, RNA was extracted and cDNA was synthesized. The expression of transcription factors were evaluated by Real-time PCR, GAPDH gene was considered as housekeeping gene.

Results: The relative expression of T-bet in RA patients were significantly higher than healthy controls ($P = 0.002$). Also, the relative expression of Foxp3 in RA patients was significantly lower than healthy controls ($P < 0.0001$). There was no significant difference in the relative expression of GATA3 and RORc in the RA patients, osteoarthritis and healthy controls ($P = 0.3$, $P = 0.2$, respectively).

Conclusion: The results of this study indicated that Th1 and Treg cells are important in RA, nevertheless the role of Th17 cells seem to be of little importance in these patients. It appears that Th2 cells do not interfere with the development of RA.

Keywords: Rheumatoid Arthritis, Osteoarthritis, T-bet, GATA3, FOXP3, RORc

ICIA2018\Immunology of Rheumatic Diseases\poster\7516
7516

Changes in Th17 cells frequency and function after Nanocurcumin therapy in ankylosing spondylitis patients

Atiye Mellati^{1, 2}, Sanam Dolati^{1, 2}, Mehdi Yousefi^{1, 2}

1. Stem Cell and Regenerative Medicine Institute, Tabriz University of Medical Sciences, Tabriz, Iran.

2. Department of Immunology, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Ankylosing spondylitis (AS) is a type of inflammatory disease that commonly contains spine and sacroiliac joint. It was described by chronic inflammation, bone reformation and back pain. Though there is no certain cure for AS some anti-inflammatory drugs can improve clinical symptoms and slow down the progression. Th17 play a key role in pathogenesis of inflammatory diseases. For this reason we tried to investigate anti-inflammatory effects of nanocurcumin on Th17 in AS patients.

Methods: Tewnty four AS patients were enrolled and divided into two groups of 12 patients in which one group received nanocurcumin and another received placebo as control group. After 16 week treatment, PBMCs were isolated from all 24 patients' blood. Then mRNA expression of Th17 associated cytokines (IL17 and IL23) and transcription factor (ROR γ t) in pre and post-treatment patients were assessed using Real-time PCR and CYBR green method. Secretion of Th17 associated cytokines was assessed using ELISA method and finally Th17 frequency was evaluated by flow cytometry technique.

Results: Nanocurcumin resulted in significant decrease in ROR γ t mRNA expression (P value=0.0001). IL17 mRNA expression decreased (P value=0.011) while IL23 mRNA didn't change compared with placebo group. IL17 secretion showed significant decrease compared with placebo and post treatment (P value=0.029) but no significant decrease in IL23 was observed. Furthermore results of flowcytometry technique showed that there is no decrease in Th17 frequency in nano-curcumin receiving group.

Conclusion: The results demonstrated that nanocurcumin may provide a promising treatment of AS and development of inflammatory cytokines suppression.

Keyword: Ankylosing spondylitis, Nanocurcumin, T helper17, Inflammation

ICIA2018\Immunology of Rheumatic Diseases\poster\7859
7859

Genepolymorphisms and serum levels of IL-10 in behcet's disease

Zohreh Babaloo¹, Babak Afkari ², Shahriar Alipour³, Ebrahim Sakhinia ⁴

1. Immunology Department, Medicine Faculty, Tabriz University of Medical Sciences

2. Immunology Department, Medicine Faculty, Tabriz University of Medical Sciences

3. Genetic Department, Medicine Faculty, Tabriz University of Medical Sciences

4. Genetic Department, Medicine Faculty, Tabriz University of Medical Sciences

Background: Interleukin 10 (IL-10) is a cytokine with potent anti-inflammatory properties that plays a fundamental role in restricting host immune response to pathogens, by means of that is a crucial importance for chronic inflammatory disease studies. Therefore, the goal of this study was to measure the correlation of the IL-10 gene polymorphisms with the susceptibility of Behçet's disease compared with the control group in the Azeri population and to determine the expression of this gene in the two groups. Also, real-time PCR was performed for evaluate the IL-10 mRNA expression of the associated polymorphisms.

Methods: In this study, blood samples from 47 (1 missed) patients and 58 (3 missed) healthy control were taken and then mononuclear cells isolated with ficoll protocol. The DNA and RNA were subsequently extracted and were then examined for -592A/C (rs1800872) of IL-10 gene single nucleotide polymorphism (SNP) using RFLP-PCR. Allele and genotype distributions were evaluated among groups using chi-square or Fisher's test. Then, the extracted RNA was converted to cDNA using the RT-PCR method, afterward the expression of IL-10 was evaluated by Real-time PCR. Serum levels of IL-10 were measured using Enzyme-linked immunosorbent assay (ELISA).

Results: Rate of the rs1800872 A allele was statistically lower in the control group compared with BD patients ($p = 0.0315$ and $OR = 1.90$ (1.05–3.42)). Also, as we expected, the expression level of the IL-10 gene was significantly decreased in the patient group compared to the control.

Conclusions: Our study showed that the rs1800872 A allele of the IL-10 gene may contribute to the genetic susceptibility of BD by regulating the expression of IL-10. Also as we expected, the expression level of this gene was significantly decreased in the patient group compared to the control group.

Keywords: rs1800872 SNP, Behçet's disease, IL-10

ICIA2018\Immunology of Rheumatic Diseases\poster\7984
7984

The effect of progesterone on MMP7 & MMP13 expression in a model of experimental systemic sclerosis

Elahe Izadi¹, Fatemeh Vafashoar¹, Hadi Poormoghim², Mohamad Ali Assarehzadegan¹, Jalil Kuhpayehzadeh³
Nazanin Mojtavavi^{1*}

1. Immunology department, Iran University of medical sciences, Tehran/ Iran

2. Scleroderma Study group, Firuzgar Hospital, Iran University of medical sciences, Tehran/Iran

3. Community Medicine and Public health research center, Iran University of medical sciences, Tehran/Iran

Background: Gender medicine is a new era of science which focuses on the impact of sex hormones and gender on normal physiology, pathobiology and clinical features of diseases. In our previous study we showed that supra physiological dose of progesterone exacerbate the lung fibrosis in a mouse model of scleroderma. Matrix metalloproteinases are a group of enzymes which play a role in tissue remodeling and fibrosis. Whereas the abnormal expression of MMP2 and MMP9 are indicated in the pathogenesis of systemic sclerosis, fewer studies are done on MMP13 and MMP7 which are expressed by epithelial cells, fibroblasts and macrophages of the lungs. They are involved in the pathogenesis of COPD, IPF and different lung diseases therefore we aimed to investigate the effect of progesterone on the expression of these two enzymes in lungs of mouse model of scleroderma.

Method: Female mice received progesterone for 28 and 21 days in addition to 28 days bleomycin. On day 29 mice were sacrificed and the expressions of these two enzymes in lungs were analyzed by real time PCR.

Result: We found that bleomycin significantly downregulated the expression of MMP7 and MMP13 and co-administration of progesterone and bleomycin declined the expression of these enzymes but not in a significant manner.

Conclusion: While progesterone cannot reduce the expression of MMP7 and MMP13 in a mouse model of lung fibrosis further investigations are required to find modulators of MMP-7 and MMP-13 in this disorder.

Keywords: Systemic sclerosis, Progesterone, Fibrosis, Bleomycin, MMP7, MMP13

ICIA2018\Immunology of Rheumatic Diseases\poster\8090
8090

The correlation between rheumatoid factor (RF) and anti cyclic citrullinated peptides (Anti-CCP) with gene expression of FoxP3 in rheumatoid arthritis patients

Nasrin Iranshahi¹, Seyed Mojtaba Amiri², Parisa Zafari^{3,4}, Mahdi Taghaddosi^{1*}

1. Department of Immunology, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

2. Department of Epidemiology and Biostatistics, School of Public Health, Kermanshah University of Medical Sciences, Kermanshah, Iran

3. Department of Immunology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran
Student Research Committee, Mazandaran University of Medical Sciences, Sari, Iran

4. Assistant professor in Department of Immunology, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

Background: Rheumatoid arthritis (RA) is a chronic autoimmune disorder that affects 1-2% of people worldwide. Inflammation is an important factor in the pathogenesis of RA. Anti- Cyclic Citrullinated Peptide (Anti-CCP) antibody and Rheumatoid Factor (RF) are autoantibodies that promotes inflammatory reactions and have a crucial role in RA pathogenesis. Treg cells are necessary for the maintenance of immune homeostasis and prevention of autoimmunity. FoxP3 is essential transcription factor for development of these regulatory cells. In this study, we surveyed the effects of FoxP3 gene expression in peripheral blood on plasma levels of Anti-CCP and RF.

Methods: Peripheral blood samples were collected from 47 patients and 44 healthy subjects. Then plasma levels of Anti-CCP were evaluated using ELISA method. Also RF was detected with latex agglutination test and gene expression of FoxP3 analyzed by real time PCR.

Results: The levels of Anti-CCP ($P < 0.001$) and RF ($P < 0.001$) were significantly higher in the patients in comparison with healthy subjects. Also significant reverse correlation between RF and Anti-CCP with gene expression of FoxP3 have been shown in our study ($r = -0.630$, $r = -0.584$) respectively. The sensitivity and specificity of Anti-CCP and RF was (89.1%, 86.95%) and (91.3%, 91.1%), respectively for the diagnosis of RA.

Conclusion: Our data illustrated that FoxP3 gene expression have reverse significant correlation with plasma concentration of anti-CCP and RF.

Keywords: Rheumatoid arthritis, FoxP3, RF, Anti-CCP

ICIA2018\Immunology of Rheumatic Diseases\poster\8091
 8091

Importance of immunological biomarkers in assessment of disease activity and progression in rheumatoid arthritis

Nasrin Iranshahi¹, shirin Assar², Parisa Zafari^{3,4}, Seyed Mojtaba Amiri⁵, Adel Fekri¹, Mahdi Taghaddosi*

1. MSc student in department of Immunology, Kermanshah University of Medical Sciences, Kermanshah, Iran

2. Assistance professor of Rheumatology in Clinical Research Development Center, Imam Reza Hospital, Kermanshah University of Medical Sciences, Kermanshah, Iran

3. Departmen of Immunology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran
 Student Research Committee, Mazandaran University of Medical Sciences, Sari, Iran

4. MSc student in department of Biostatistics and Epidemiology, school of health, Kermanshah University of Medical Sciences, Kermanshah, Iran

5*. Assistance professor in department of Immunology, Kermanshah University of Medical Sciences, Kermanshah, Iran

Background: Rheumatoid arthritis (RA) is a chronic autoimmune disorder that affect 1-2% of people worldwide. Disease Activity Score (DAS-28) is a clinical parameter of RA progression. This factor is measured according to joints tenderness and swelling. Many elements could cause inflammation in synovial microenvironment including pro-inflammatory cytokines (IL-1, IL-6 and TNF- α) and other mediators produced during immune response. Neopterin is one of the metabolites that is produced following catabolism of guanosine triphosphate (GTP). The high concentration of neopterin is indicator of a more severe inflammation. Other disease mediators including Anti-CCP and RF are autoantibodies which are indispensable in the diagnosis of RA.

Methods: Peripheral blood samples were collected from 47 patients and 44 healthy subjects. Then plasma levels of neopterin and Anti-CCP were evaluated using ELISA method. Also RF positive patients were detected with latex agglutination test and ESR levels were obtained from patient's records.

Results: The levels of neopterin ($P < 0.038$), Anti-CCP ($P < 0.001$) and RF ($P < 0.001$) were significantly higher in the patients in comparison with healthy subjects. Also a significant correlation between RF and DAS-28 ($P = 0.001$) were shown in our study. The sensitivity and specificity of anti-CCP and RF were (89.1%, 86.95%) and (91.3%, 91.1%) respectively for the diagnosis of RA.

Conclusion: Our data showed that neither neopterin nor Anti-CCP has any significant correlation with DAS-28 while the correlation between RF and DAS-28 was statistically meaningful.

Keywords: Rheumatoid arthritis, DAS28-ESR, Neopterin, RF, Anti-CCP

ICIA2018\Immunology of Rheumatic Diseases\poster\8425
8425

Frequency of different autoantigens among ANA seropositive patients referred to Shafa medical laboratory of Birjand City, Iran

Arghavan Asghari¹, Neda Mahavar¹, Elham Atabati², Mohammad Fereidouni¹

1. *Asthma, Allergy & Immunology Research Center, School of Medicine, Birjand University of Medical Sciences, Birjand, Iran*

2. *Department of Rheumatology, Vali-Asr Hospital, Birjand University of Medical Sciences, Birjand, Iran*

Background: Prevalence of autoimmune disorders has increased dramatically during the last decades particularly in developing countries. Autoantibodies are responsible for autoimmune inflammation and are considered as the helpful markers for accurate diagnosis. It is shown that specific autoantibodies can lead to a specific disease, so it is important to figure out which autoantibodies are more prevalent in our population to have a proper diagnosis, management and follow up. The aim of this study was to evaluate the frequency of different autoantibodies in patients with positive anti-nuclear antibodies referred to the Shafa laboratory of Birjand city of Iran 2017.

Methods: Serum samples of patients with symptoms of autoimmune diseases were collected and presence of anti-nuclear antibody (ANA) as well as the profile of autoantibodies against at least 16 auto antigens were evaluated by immunofluorescence and immunoblotting test respectively.

Results: one hundred and fifty five (mean age: 43.58 ± 14.50 , range 13-83 years, M/F ratio: 0.11) subjects with positive ANA test participated in this study. The most observed autoantibodies were anti SSA and anti-RO52 (42.22% ;42.22%). autoantibodies against gp120, mi2b, mi-2 and pm-scl 75 had the least frequency (0.6%;0.64%;0.64%), Respectively.

Conclusion: The results of current study showed that 90% of patients with positive ANA test were female and autoantibodies are the most frequent autoantibodies.

Keywords: Anti-nuclear antibody, Autoimmune disorders

ICIA2018\Immunology of Rheumatic Diseases\poster\8483
8483

Association of polymorphisms in ERAP1 and risk of ankylosing spondylitis in an Iranian population

Hamed Mohammadi^{1,2}, Maryam Hemmatzade^{1,2}, Farhad Babaie^{1,2}, Alireza Rezaeiemanesh³, Jafar Majidi^{1,2}, Behzad Baradaran^{1,2}

1. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

2. Department of Immunology, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

3. Department of Immunology, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

Background: Endoplasmic reticulum aminopeptidase 1 (ERAP1) gene variants reduce the risk of progression ankylosing spondylitis (AS) through a decrease in ERAP1 activity. We determined the role that ERAP1 gene plays in Iranian patients with ankylosing spondylitis (AS) in terms of disease susceptibility and disease severity.

Methods: We evaluated two variants (rs27044 and rs17482078) of ERAP1 gene in HLA-B27 positive individuals (160 ankylosing spondylitis and 160 controls). The genotypes of involved polymorphisms (rs27044 and rs17482078) in ERAP1 were identified by SSP-PCR.

Results: There was statistically association between ERAP1 rs27044 polymorphism and risk of AS and the carriers with rs27044 CG genotype have an enhanced risk for AS (CG versus GG, OR = 1.60, 95% CI = 1.10–2.1, P = 0.006). However, we found no evidence for the association of 17482078 polymorphism in ERAP1 with AS risk.

Conclusion: Our results demonstrated that ERAP1 rs27044 polymorphism associates with the susceptibility of AS.

Keywords: Ankylosing spondylitis, ERAP1, Polymorphism, HLA-B27

ICIA2018\Immunology of Rheumatic Diseases\poster\8509
8509

Gene expression profile of pro-inflammatory and anti-inflammatory cytokines in Peripheral Blood Mononuclear Cells of Patients with Ankylosing Spondylitis

Farhad Babaie^{1,2}, Hamed Mohammadi^{1,2}, Sorayya Salimi^{1,2}, Alireza Rezaeiemanesh³, Behzad Baradaran^{1,2}, Zohreh Babaloo^{1,2}

1. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

2. Department of Immunology, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

3. Department of Immunology, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

Background: Ankylosing spondylitis (AS) belongs to a typical group of arthritides called seronegative spondyloarthritides. AS targets fundamentally the spine and sacroiliac joints and is characterized by enthesal inflammation, dactylitis and uveitis. The role of pro-inflammatory and anti-inflammatory cytokines in ankylosing spondylitis is still unrevealed. Our study aims to determine gene expression profile of pro-inflammatory and anti-inflammatory cytokines and their correlations with AS clinical manifestation patterns, including the age of symptom onset, Bath AS Disease Activity Index (BASDAI), Bath AS Functional Index (BASFI) and Bath AS Metrology Index (BASMI).

Methods: Peripheral blood mononuclear cells (PBMCs) were isolated from 40 patients with AS and 30 healthy controls, then total RNA of PBMCs was extracted, followed by cDNA synthesis from the mRNA of PBMCs. Quantitative Real time PCR was performed to evaluate mRNA expression of pro-inflammatory (INF- γ , IL-17F, IL-17A, TNF- α , IL-33, and IL-6) and anti-inflammatory cytokines (IL-10, IL-37 and TGF- β).

Results: A significant up-expression of pro-inflammatory (IL-17A, IL-17F, TNF- α , IL-33 and IL-6) and anti-inflammatory cytokines (IL-10, IL-37, and TGF- β) was observed in AS patients compared to the control group but, there was no significant difference in INF- γ expression between groups. There was no significant correlation between gene expression and AS clinical manifestation patterns.

Conclusion: These results demonstrated the critical role of pro-inflammatory and anti-inflammatory cytokines in the immunopathogenesis of AS. On the other hand, these cytokines may be responsible for inflammatory-related outcomes of AS.

ICIA2018\Immunology of Rheumatic Diseases\poster\8575
8575

Study of retinal damage following cyclosporine-A eating using electroretinogram technique

Samaneh Gholami¹, Agheel Tabarmolahasan¹, Mohammad Masoud Shoushtarian¹, Seyed Hamid Aghajanzadeh¹, Safdar Mehdizadeh Shermeh¹

1. Islamic Azad University of Babol.

Background: Cyclosporine A is prescribed as an immunosuppressive agent in many diseases. One of the side effects of cyclosporine is visual disturbances and retinal damage. Different methods are available to investigate the damage of retina, the best of which is the electroretinogram technique. In this study, the effects of cyclosporine on the retina of patients with rheumatoid arthritis were studied.

Methods: In this observational study, which was a cross-sectional descriptive-analytic study, 55 patients receiving cyclosporine due to rheumatoid arthritis between 2012-2014 were randomly selected and subjected to electroretinography and voltage then delay phase was determined and compared with the normal range. The results were analyzed by means of statistical analysis of variance. Significant level was set at 0.05.

Results: The average recorded voltage in ERG was 74.7 with a standard deviation of 29.79, which showed a statistically significant difference from normal value of 125 ($P=0.0001$). The mean recorded latency was 33.9 with a standard deviation of 3.58, which means no significant difference from normal value of 32.39 ($P=0.545$).

Conclusion: Based on the results, it is concluded that the use of cyclosporine significantly reduces the voltage recorded in the electroretinogram, but does not affect the latency phase. Therefore, patients taking this medicine should continue their treatment according to the ophthalmologist

Keywords: Electrotonography, Cyclosporine, Retina

ICIA2018\Immunology of Rheumatic Diseases\poster\8685
8685

TNF- α -induced protein 3 gene polymorphisms in Azari patients with Behcet's Disease

Saeede Ghasemi¹, Yasamin Pahlavan², Alireza Khabbazi³

1. Department of Biology, Faculty of Sciences, Sciences and Researches of East Azarbijan Branch, Islamic Azad University

2. Department of Molecular Medicine, School of Advanced Medical Sciences, University of Tabriz Medical Sciences

3. Department of Romathology, School of Medicine, University of Tabriz Medical Sciences

Background: Behçet's Disease is an inflammatory and chronic autoimmune disease with an unknown cause, which is associated with oral ulcers, skin and genital lesions. The aim of our study was the investigating the association of polymorphisms of TNF- α -induced protein 3 (TNFAIP3) gene with susceptibility to Behcet's Disease (BD) in Azari population of Iran.

Methods: In this cross sectional study we considered the single-nucleotide polymorphisms rs9494885 and rs7753873 of TNFAIP3 in 50 Iranian Azary patients with BD and 50 healthy controls by polymerase chain reaction-restriction fragment length polymorphism.

Results: A significant difference was found for the rs7753873 polymorphism between the BD and control groups. The frequency was 6.6% in BD patients compared to 20% of controls ($p=0.04$). We found no significant differences between the BD and control groups regarding the distribution of the polymorphism frequencies.

Conclusion: The rs7753873 polymorphism of TNFAIP3 has negative association with BD in Iranian Azari population.

Keywords: TNF- α , polymorphisms, Behcet's Disease

ICIA2018\Immunology of Rheumatic Diseases\poster\9945
9945

Evaluation of CD4⁺/CD25⁺/high/CD127^{low}/- Regulatory T Cells in Rheumatoid Arthritis Patients

Samaneh Delavari¹, Ata Ghadiri², Mehri Ghafourian³, Elham Rajaei⁴, Afshin Amari⁵

1. Department of Immunology, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

2. Cellular and Molecular Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

3. Department of Immunology, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

4. Department of Internal Medicine, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

5. Department of Immunology, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

Background: Rheumatoid arthritis (RA) is the most common inflammatory rheumatic disease with an unknown etiology which is characterized by destruction of articular cartilage and bone loss. The hallmark of disease is defect in immune tolerance. Regulatory T cells (Treg) play a critical role in protection of peripheral tolerance. We investigated the percentage of CD4⁺/CD25⁺/high/CD127^{low}/- Treg cells in RA patients in comparison to healthy individuals.

Methods: The number of Treg CD4⁺/CD25⁺/high/CD127^{low}/- cells was assessed by multi-color flow cytometry. The clinical disease activity of RA patients was determined by disease activity score 28 (DAS-28). The correlation of DAS-28 and ESR with Treg cells in peripheral blood of RA patients was evaluated.

Results: The percentage of CD4⁺/CD25⁺/high/CD127^{low}/- Treg cells in RA patients was significantly decreased compared with healthy individuals. The percentage of CD4⁺/CD25⁺/high/CD127^{low}/- Treg cells negatively correlated with DAS-28 and ESR.

Conclusion: In our study, we conclude that defect of Treg cells play a key role in the pathogenesis of this disease. Further studies are necessary to clarify the role of Treg cells in clinical course of rheumatoid arthritis.

Keywords: Rheumatoid arthritis, Regulatory T cells, Flow cytometry

ICIA2018\Immunology of Rheumatic Diseases\poster\9984
9984

MicroRNAs as potential and novel biomarkers in adult Iranian patients with systemic lupus erythematosus

Majid Khoshmirsafa¹, Farhad Seif¹, Reza Falak¹, Mehdi Shkarabi^{*1}

1. Department of Immunology School of Medicine University of IRAN

Background: MicroRNAs (miRNAs) have emerged as effective immune-modulators of gene expression in immune responses. Previous miRNA studies have been showed some unique miRNA signatures that are associated with systemic lupus erythematosus (SLE), but their role as diagnostic and prognostic biomarkers of Lupus Nephritis (LN) has not been well surveyed.

Methods: In this ongoing study, we assessed miRNA expressions (miR-200a, b and c) in peripheral blood mononuclear cells (PBMCs) obtained from SLE patients with different kidney involvements using quantitative PCR (q-PCR).

Results: MiR-200c displayed significant increment in SLE patients (LN) in comparison with healthy controls (P value < 0.05).

Conclusion: The role of miRNAs in immunopathogenesis of rheumatic diseases was determined, it is possible that certain miRNAs may serve as therapeutic and diagnostic biomarkers for SLE and LN.

Keywords: Systemic lupus erythematosus, Lupus Nephritis, MicroRNA, q-PCR



Immunomodulation and Immunoregulation

Poster Presentation

14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Immunomodulation and Immunoregulation\Poster\8515

8515

Evaluation of Immunomodulatory Effect of Monolaurin

Elham Rahimi¹, Maryam Ekhtelat¹, Mojgan Naghitorabi¹, Ata Ghadiri²

1. Cellular and Molecular Research Center, Faculty of Pharmacy, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

2. Department of Immunology, Faculty of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

Background: Fatty acids, monoglycerides and some esters of fatty acids show antitumor and antimicrobial effect. In this study, we evaluated the monolaurin (1-glycerol-monolaurate; the esterified form of lauric acid) effect on macrophage activation.

Methods: For this purpose, monolaurin effect on peritoneal macrophages of mice (BALB/c) and LPS stimulated macrophages were examined in triplicate. MTT assay with different concentrations (10, 50, 100, 200, 400 and 800 µg/ml) of monolaurin was done for evaluation of macrophages viability and the level of nitric oxide in supernatant of culture was measured by nitric oxide (NO) assay kite. Data analysis was performed by the SPSS software using ANOVA.

Results: MTT assay showed that monolaurin in amount of 100 µg/ml and greater significantly reduced viability of macrophages (stimulation index). Also, the amount of NO production in stimulated macrophages in concentrations of more than 100 µg/ml of monolaurin, significantly was lower compared to the LPS control ($p < 0.05$).

Conclusion: Regarding immune-modulatory effects of monolaurin, it can be used to help cure disorders of immunity and inflammation. Of course, in vivo evaluation are necessary to confirm it.

Keywords: Immunomodulatory, Monolaurin, Macrophage, MTT assay, Nitric oxide

ICIA2018\Immunomodulation and Immunoregulation\Poster\8536

8536

Effect of low molecular weight garlic proteins on the T regulatory cell induction in co-culture of PBMCs and colorectal cancer cell linesMona Amani¹, Elahe Safari¹, Samaneh khorami¹*1. Immunology Department, School of Medicine, Iran University of Medical Sciences*

Background: Cancer is one of the non-infectious diseases with high mortality all over the world. In this regard, colorectal cancer is the third most common cancer worldwide. There is different therapeutic methods such as chemotherapy, radiotherapy and surgery using to treat the cancer. Between this methods perhaps we can bringing to account the herbal medicine to use as supplement in prevention and/or control the cancer. Among various types of herbal medicine it can mention to garlic as the eldest plant with medicinal properties that its medicinal usage was very common from ancient years. Garlic active ingredients have different biological effect. They can improve immune system to defend against different microorganism and also cancer. It can also use as treatment of cardiovascular disease. Thus in this study we were decided to extract the low molecular weight proteins of garlic and then treat the co-culture of PBMCs and colorectal cancer cell lines SW48 and SW837 with the garlic protein extract to examine the immunomodulatory effect of this protein in the tumor microenviroment as the most important factor that affect tumor growth and expansion.

Methods: After extraction from garlic cloves, protein fractions were separated with G-75 gel filtration chromatography. To check the purity of separation, Sodium Dodecyl Sulfate Poly Acrylamide Gel Electrophoresis(SDS-PAGE) were done. To define the protein identity, MALDI-TOF spectrometry were done. In the cell culture phase, PBMCs and cell lines alone and in co-culture were treated with desire protein and PBMCs proliferation was assayed by using fluorescent dye, CFSE. Also these cell culture supernatants were collected to evaluate the secretion of inflammatory and inhibitory cytokines in this milieu. Finally, to measure the inhibition rate of inhibitory cell induction like Treg and MDSC in co-culture medium in the presence of desire protein, PBMCs were cultured with cell lines and then stained with specific antibodies against Treg and MDSC surface antigens and the results were assayed with FACS.

Results: We identify that this protein extraction was a lectin binding protein with molecular weight of 11-16 kDa that exhibit immunomodulatory effect. In proliferation assay, this protein was able to stimulate PBMCs alone and in co-culture with tumor cell lines. In cytokine assay we observed that PBMC treatment with protein extraction caused reduction in TGF- β and Galectin-3 secretion; in opposite IL-6 and IFN- γ secretion level was upregulated and it has no significant effect on IL-10 secretion in comparison with negative control which had no treatment. Also we investigated that it could inhibit immune suppressor cell, Treg and MDSC induction in the co-culture milieu.

Conclusion: By stimulating PBMCs proliferation, inhibiting suppressor cell induction and upregulating inflammatory cytokine and reversely reducing inhibiting cytokines; garlic protein extract may use as immunomodulatory supplement in cancer treatment. Also in vivo study should be done.

Keywords: Immunomodulation , Tumor , garlic.

ICIA2018\Immunomodulation and Immunoregulation\Poster\8718
8718

Comparison of cytokine profile between patients with end-stage renal diseases and healthy individuals

Farshid Fathi¹, Hossien Motedayyen², Noushin Lotfi¹, Shohreh Mohammadi³, Hossien Hafezi⁴, Erfan Sadeghi⁵

1. Department of Immunology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

2. Department of Laboratory Medicine, Kashan University of Medical Sciences, Kashan, Iran

3. Khorshid Hospital, Isfahan University of Medical Sciences, Isfahan, Iran

4. Department of Dermatology, Isfahan University of Medical Sciences, Isfahan, Iran

5. Department of Biostatistics and Epidemiology, School of Public Health, Isfahan University of Medical Sciences, Isfahan, Iran

Background: Cytokines have a pivotal role in regulating the function of the cells from both the innate and adaptive immune system. Imbalance and dysregulation in cytokine profile produced by different immune cells can influence the risk of developing autoimmune and allergic disorders. We, therefore, focused to determine whether there is a difference in cytokine profile between renal patients who suffered from some autoimmune disorders and healthy subjects.

Methods: Heparinized whole blood (5 ml) was obtained from patients with end-stage renal failure and healthy subjects. Plasma was isolated from whole blood and the level of interferon-gamma (IFN- γ), tumor necrosis factor-alpha (TNF- α), interleukin-4 (IL-4), interleukin-17A (IL-17A) and interleukin-10 (IL-10) was measured by an Enzyme-linked immunosorbent assay (ELISA) kit.

Results: Our results indicated that the level of IL-17A and TNF- α was significantly increased in renal patients with autoimmune disorders compared to healthy individuals ($P < 0.05$). However, there was no significant difference in the level of IFN- γ and IL-10 between renal patients suffered from autoimmune disorders and healthy subjects. Moreover, a significant reduction in IL-4 level was observed in renal patients compared to healthy subjects ($P < 0.001$).

Conclusion: Regarding the fact that cytokines play an important role in regulating the immune responses which lead to the development of autoimmune disorders, dysregulation of cytokine production from different immune cells in patients with autoimmune diseases may be contributed to the immunopathogenesis of renal failures.

Keywords: Renal patients, Autoimmune diseases, cytokines



Immunoparasitology

Poster Presentation

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Immunoparasitology\Poster\8336
8336

The influence of fetal calf serum and hemin in growth and infectivity of *Leishmania major* in BALB/c mice

Sima Habibzadeh ^{1,2}, Sima Rafati¹

1. Immunotherapy and Leishmania Vaccine Research Dept., Pasteur Institute of Iran, Tehran, Iran

2. Islamic Azad University, Pharmaceutical Sciences Branch, Tehran, Iran

Background: Leishmaniasis is a common disease between humans and animals in tropical and subtropical regions. The causative agent is *Leishmania* which exists in two forms of promastigote and amastigote. Like other cells require nutrients to grow. Various culture media especially in liquid form designed for parasite cultivation. Most of these media contain either Fetal Calf serum (FCS) or blood as essential ingredient which is highly problematic. We have designed liquid media without FCS. Although is not expensive but the addition of hemin is obligatory for *in vivo* infectivity of *L. major* in BALB/c mice. This molecule as the prosthetic group of hemoproteins, has important roles in regulating vital cell functions and transformation of promastigote to amastigote.

Methods: In the present study, *L. major* parasites were cultured in two different media, LBM (LB-BHI-M199) and LBM with hemin (LBM H+) in comparison with M199 FCS 5% (M199 5%) as control positive group. After several passages, stationary-phase promastigotes were injected to the left footpad of mice. *In vivo* infectivity of parasites were evaluated at six weeks post infection, using footpad swelling and quantitative real-time PCR.

Results: There is no swelling in the footpad of mice infected with parasite growing within LBM without hemin. In addition, the level of swelling was similar in animals infected with parasite growing in LBM H+ media and M199 5%. Furthermore, by real-time PCR, the lowest parasite was detected in group infected by parasite in LBM without hemin in compare with LBM+ hemin and M199 with 5%FCS.

Conclusion: *Leishmania* like other trypanosomatids needs an external source of heme such as hemoglobin, blood, serum, hemin and hematin for their growth. This is due to lack of two essential enzymes for heme biosynthesis pathway. In this study we examined the effect of hemin on the infectivity of *L. major* parasite in BALB/c mice and prove it as an essential substance.

Keywords: *L. major*, Heme, Infectivity, BALB/c

ICIA2018\Immunoparasitology\Poster\8600
8600

Prevalence Cutaneous Leishmaniasis in the Western Cities Areas of Kermanshah Province, West Iran, During 2010- 2015

Mehrdad Pooyanmehr¹, Mohammad Reza Khedri², Erfan Bahrami², Amir Amiri Paryan^{3*}

1. Department of Basic Sciences and Pathobiology, Microbiology section, Faculty of Veterinary Medicine, Razi University, Kermanshah, Iran

2. Veterinary Student, Clinical Sciences, Faculty of Veterinary Medicine, Razi University, Kermanshah, Iran

3. Veterinary Student, Pathobiology & Basic Sciences, Faculty of Veterinary Medicine, Razi University, Kermanshah, Iran

Background: Leishmaniosis is one of the most important tropical diseases and zoonotic infection. Natural transmission of *Leishmania* parasites is carried out by biting of female sandflies genus *Phlebotomus* from reservoir hosts to human (caused by obligate intracellular protozoa of the genus *Leishmania*). There are three forms of leishmaniosis: Cutaneous leishmaniasis (CL, rural and urban), visceral leishmaniosis (VL) and muco cutaneous leishmaniosis (MCL). Cutaneous leishmaniosis only causes ulcers on skin. In past ten years, prevalence and incidence of leishmaniosis remarkably increased in many regions of Iran.

Methods: The data were taken from internal reports of Health Centers western cities areas of kermanshah province (Ravansar, Javanrood, Pavehe and Salasebabajany) in the period six years (2010-2015). Thirty-five patients with CL were included in this study. The diagnosis of CL was based on clinical, Para Clinical findings and microbiologic procedures on chronic (Non-Healing) skin lesions. Laboratories in Health Centers four city confirmed of the diagnosis for detecting *Leishmania* parasites in infected tissue (through light-microscopic examination of stained specimens, culture techniques and molecular methods).

Results: Results indicated that All Thirty-five patients' referent were who diagnosed as CL. The clinical presentation of CL was typical in 31 patients, Ravansar 13 (0.013%), Javanrood 8 (0.008%), Pavehe 1 (0.001%) and Salasebabajany 9 (0.009%) per100.000 persons, by clinical para clinical findings and pathologic documentation in 4 cities medical centers registries patients at six years period respectively.

Conclusion: CL is a very contagious and important infectious disease worldwide. The diagnosis of CL may be difficult, especially in atypical presentations and Non-endemic areas. Thus, CL should be kept in mind, especially in many regions of Iran.

Keywords: Zoonotic, cutaneous, leishmaniosis, western cities areas, Kermanshah

ICIA2018\Immunoparasitology\Poster\9850

9850

The Effect of *Leishmania Major* Lipophosphoglycan 3 on Human T Lymphocytes In Vitro

Maryam Hosseini¹, Mostafa Haji-Fatahaliha², Sanaz Rasolzadeh¹, Mehdi Yousefi¹

1. Department of Immunology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

2. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Leishmaniasis is a vector-borne disease threatens 350 million people in 98 countries over the world. No effective drug or vaccination strategy has been designed against leishmaniasis so far. Lipophosphoglycan 3 (LPG3) as one of the class II LPG genes from HSP90 family involves in assembly and synthesis of lipophosphoglycan (LPG) and other phosphoglycan residues implicate in parasite virulence. The aim of this study was to investigate the ability of recombinant LPG3 (rLPG3) to induce human T-cell lineages in vitro.

Methods: Peripheral blood mononuclear cells were obtained from heparinized blood samples of 10 healthy volunteers. Then CD4⁺ naïve T-cells were separated using magnetic-activated cell sorting technique and their purity was evaluated by flow cytometry. Isolated T-cells were treated with different concentrations of recombinant *Leishmania major* LPG3. After 48 h incubation, the activity of T-cells was investigated by assessing hallmark cytokines of T helper (Th) 1, 2 and 17 lineages using real-time PCR and ELISA techniques.

Results: The purity of isolated CD4⁺ T-cells was about 90%. rLPG3 could induce IFN- γ mRNA expression in moderate and high doses but no significantly effect on IL-4 and IL-17A expression was shown in treated T cells. Moreover, similar observations were achieved in cytokine secretion and showed stimulatory effect of rLPG3 in IFN- γ secretion especially in high concentration ($p < 0.05$).

Conclusion: Our results demonstrated that LPG3 could stimulate Th1 responses effectively but further researches are needed to investigate its anticipated adjuvant usage in *Leishmania* vaccination.

Keywords: Leishmaniasis, LPG3, T cell



Immunopharmacology and Medicinal Plants

Poster Presentation

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Immunopharmacology and Medicinal Plants\poster\5301

5301

The hydro alcoholic extract of *Hypericum perforatum* attenuates the inflammatory reactions in acetic acid induced ulcerative colitis

Siavash Mashhour¹, Seyyed Meysam Abtahi Froushani²

1. Graduate of Veterinary Medicine, Department of Microbiology, Faculty of Veterinary Medicine, University of Urmia

2. Department of Microbiology, Faculty of Veterinary Medicine, University of Urmia

Background: This study was done to investigate the beneficial effects of the hydro alcoholic extract of *Hypericum perforatum* to control the inflammation in animal models of ulcerative colitis.

Methods: Luminal instillation of acetic acid was used to induce ulcerative colitis in the male Wistar rats. Rats in the treatment groups received herbal extract (150 mg/kg PO) or Prednisolone (4 mg/kg PO) daily for 10 consecutive days. When the animals were sacrificed, the disease activity index, the levels of malondialdehyde, myeloperoxidase, nitric oxide, total protein and the concentration of TNF α IL1 β and IL6 were determined in the colonic homogenized tissues specimens.

Results: Both therapeutic regimes with herbal extract and Prednisolone could ameliorate the clinical scores and the mortality rate of ulcerative colitis in a comparable manner. The levels of TNF α , IL1 β and myeloperoxidase activity were regressed in the gut tissues of Prednisolone treated rats more than ulcerative colitis rats treated with herbal extract. Nevertheless, *H. perforatum* could suppress the levels of IL6, nitric oxide and malondialdehyde more significantly than Prednisolone.

Conclusion: It seems that the hydro alcoholic extract of *Hypericum perforatum* may be used as a natural source to control the inflammation in the ulcerative colitis.

Keyword: *Hypericum perforatum*, ulcerative colitis, Acetic acid, inflammation.

ICIA2018\Immunopharmacology and Medicinal Plants\poster\6363

6363

Effects of *Viola tricolor* flower hydroethanolic extract on lung inflammation in a mouse model of chronic asthma

Elham Harati, Alireza Razavi²., Maryam Bahrami¹, Mohammad Kamalinejad³., Maryam Mohammadian¹., Hamid Reza Sadeghipour¹.

1. Department of Physiology, Faculty of Medicine, Tehran University of Medical Sciences, Tehran, Iran

2. Department of Pathobiology, Division of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

3. Department of Pharmacognosy, School of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Asthma is a chronic inflammatory disease of the lungs driven by T cell activation. *Viola tricolor* L. as a traditional medical herb could suppress activated T lymphocytes and has been used empirically for asthma remedy. In the present study, we investigated the anti-inflammatory effect and underlying mechanism of *Viola tricolor* on asthma characteristics induced by ovalbumin (OVA) in mice.

Methods: Sixty BALB/c mice were randomly divided into six groups: normal control, OVA control, OVA mice treated with *Viola tricolor* (50, 100 and 200 mg/kg) and dexamethasone (3 mg/kg). All mice except normal controls were sensitized and challenged with OVA. Asthmatic mice were treated orally in the last 7 days of OVA challenge. The total and differential leukocyte counts, Interleukin (IL)-4 and interferon (IFN)- γ levels in bronchoalveolar lavage fluid (BALF) were determined. H&E staining for lung inflammation was performed.

Results: *Viola tricolor* treatment at 200 mg/kg significantly decreased IL-4 level but did not considerably affect the IFN- γ level. Therefore, it effectively reduced asthma characteristics including infiltration of leukocytes particularly eosinophil and peribronchial inflammation as compared to dexamethasone. However, *Viola tricolor* at 100 mg/kg had the most prominent inhibitory effect on the IL-4 level and also markedly increased IFN- γ level. As result, it prevented further reduction of inflammatory parameters in this group compared to the *Viola tricolor*-treated group at 200 mg/kg. It seems that a significant increase in IFN- γ level at 100 mg/kg could prevent the beneficial effects of reducing IL-4 in the asthma treatment.

Conclusion: Our study demonstrated that *Viola tricolor* has anti-inflammatory effects via inhibition of Th2 cytokine production and validated its empirical usage in traditional medicine.

Keywords: Asthma; Inflammation; Murine model; *Viola tricolor* L.

ICIA2018\Immunopharmacology and Medicinal Plants\poster\7338
7338

Improving Doxorubicin Chemotherapy Efficacy by Using Citrus Unshiu Extract That Reduces Inflammatory Cytokines-Induced Cachexia in a Murine Colon Cancer Model

Sahar Tahaghoghi Hajghorbani ^{1*}, Alireza Rafiei ¹, Mohamad Ali Ibrahim Zade ², Javad Akhtari³

1. Department of Immunology, Medical School, Mazandaran University of Medical Sciences, Sari, Iran

2. Pharmaceutical Sciences Research Center, School of Pharmacy, Mazandaran University of Medical Sciences, Sari, Iran

3. Mazandaran university of Medical Sciences, Nanobiomedicine, Sari, Iran

Background: Several medical trials on the treatment of cancer cachexia have been performed using synthetic drugs that can stimulate appetite and suppress systemic inflammation. But the long-term use of drugs induces adverse effects. Therefore, finding novel agents that efficiently prevent or treat cancer-induced cachexia with minimal adverse effects is very important to improve the cancer chemotherapy efficacy and survival time. In the present study, Citrus Unshiu (a seedless and easily peeled citrus fruit) Peel (CUP) extract was evaluated as a safe and potent agent along with chemotherapy.

Methods: Water CUP extract was prepared. The phytochemical profile of CUP was analyzed using the Hitachi HPLC system. Prophylactic and therapeutic extract was fed to male BALB/c mice alone or together with doxorubicin chemotherapy. Inflammatory cytokines (IL-1 β , TNF- α and IL-6) and NO production were evaluated using enzyme-linked immunosorbent assay (ELISA) and Griess reagent, respectively. Body weight, tumor size, food intake and survival rate were assessed in mice

Results: HPLC analyses revealed the high levels of Naringin and Hesperidin content in the CUP extract. We found that daily oral administration of CUP to male BALB/c mice bearing C26 reduced the losses in body weight and serum inflammatory cytokines level that eventually result in cancer-induced cachexia symptoms. Also the tumors in mice who received extract grew more slowly and the survival rates were greater than mice in the other groups

Conclusion: These findings collectively suggest that WCUP reduces systemic inflammation in tumor-bearing mice and suppresses the production of pro-cachectic cytokines followed by the prevention of weight loss and tumor growth.

Keywords: Immunotherapy, Cachexia, Colon Cancer, Unshiu Extract

ICIA2018\Immunopharmacology and Medicinal Plants\poster\7384
7384

Effect of D-panthenol on the adjuvant induced rheumatoid arthritis (RA) in Wistar rats.

Navid Mijani, Norouz DelireZh, Seyyed Meysam Abtahi Froushani

Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Background: The anti-inflammatory benefits of D-panthenol have been documented earlier. This study was done to evaluate the effect of D-panthenol on the rat model of rheumatoid arthritis (RA).

Methods: RA was induced by injection of complete Freund's adjuvant into the footpad of Wistar rats. Then, rats were allocated in 3 groups: treated with D-panthenol (500 mg/kg-orally), treated with Prednisolone (10 mg/kg-orally) and un-treated group. All treatments were initiated at day 5 after induction. The change in the dorso-plantar diameter of hands and legs of each rat were recorded every other day until 23 days after induction.

Results: The edema and swelling of the soles of the feet of RA rats received D-panthenol or prednisolone were significantly decreased in a similar manner. The serum levels of nitric oxide and myeloperoxidase concurrent with proliferation of spleen lymphocytes were significantly lessened in treatment groups compared to control rats. The level of decrease in the level of serum nitric oxide was higher in the D-panthenol group compared to the prednisolone group.

Conclusion: D-panthenol may be considered as a useful drug to control RA disease.

Keywords: D-panthenol, Rheumatoid arthritis, Immunity responses

ICIA2018\Immunopharmacology and Medicinal Plants\poster\7386
7386

Dexpanthenol has beneficial effects on the acetic-acid induced ulcerative colitis

Seyede Mehrnaz Mostatabzadeh, Seyyed Meysam Abtahi Froushani

Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Background: The anti-inflammatory benefits of Dexpanthenol have been documented earlier. Here, we investigated the protective potential of Dexpanthenol in ameliorating animal model ulcerative colitis (UC).

Methods: UC was induced in the male Wistar rats by luminal instillation of acetic acid. Rats in the treatment groups treated with Dexpanthenol (100 mg/kg PO) or Prednisolone (2 mg/kg PO) daily for 10 consecutive days. At the end, the animals were sacrificed and the disease activity index, the levels of myeloperoxidase, malondialdehyde, nitric oxide and total protein were evaluated in the colonic homogenized tissue specimens.

Results: The data indicated that both therapies with Dexpanthenol and Prednisolone could suppress the clinical scores and the mortality rate of ulcerative colitis in a comparable manner. The levels myeloperoxidase activity was lessened in the guts of Prednisolone treated rats more than Dexpanthenol groups. Nonetheless, Dexpanthenol downregulated the levels of nitric oxide and malondialdehyde more significantly than Prednisolone.

Conclusion: These data suggest that the Dexpanthenol may be used as an appropriate medication to ameliorate the signs of ulcerative colitis.

Keywords: Dexpanthenol, Ulcerative colitis, Anti-inflammatory.

ICIA2018\Immunopharmacology and Medicinal Plants\poster\7405
7405

Comparison of Immunomodulatory Effects of Some *Allium* species from Iran on Cytokine Generation

Hosseinpur Z¹, Radjabian T^{1,2}, Ghazanfari T¹, Naghizadeh MM¹

1. Immunoregulation Research Center, Shahed University, Tehran, Iran

2. Department of Biology, Faculty of Basic Sciences, Shahed University, Tehran, Iran

Background: The literature review has demonstrated that extract of *Allium sativum* (garlic) modulates the cytokines release patterns in lymphocytes. But there are unveiled data about immunomodulatory potentials of the other *Allium* species. So, in the present research, the effects of extracts from some wild grown *Allium* species of Iran on IFN- γ , IL-4 and IL-17 production were evaluated.

Methods: The bulbs of some *Allium* species were collected from their natural localities and the aqueous extracts of them were prepared. Lymphocytes were isolated from BALB/c mice and were incubated with various concentrations of the bulb extracts. Finally, the levels of IFN- γ , IL-4 and IL-17 were measured.

Results: Bulb extracts of *A. sativum* had stimulatory effects on production of IFN- γ and IL-17, so the extracts could probably switch immune responses towards Th1 and Th17 cells pattern. Bulb extracts of *A. asarense* increased secretion values of IL-4 from lymphocytes. Whilst, bulb extracts of *A. sativum* decreased the amounts of IL-4, especially in the higher concentrations. Therefore, *A. asarense* and *A. sativum* may be significantly acted as a stimulator and inhibitor of Th2 cells, respectively, in some concentrations. On the other hand, bulb extract of *A. jesdianum* significantly decreased the amounts of IL-4 and IL-17, especially in the higher concentrations. Hence it seems that this species reduced the response of Th2 and Th17 cells.

Conclusion: In general, bulb extracts of the *Allium* species had immunomodulatory effects on the lymphocyte cytokines levels, which are involved in immune related-diseases, and these species may be considered for treatment studies.

Keywords: *Allium* species, Bulb extract, Cytokines, IFN- γ , IL-4, IL-17

ICIA2018\Immunopharmacology and Medicinal Plants\poster\7407

7407

Assessment of the Immunomodulatory Effects of Some *Allium* species on TNF- α and NO Production

Hosseinpur Z¹, Radjabian T^{1,2}, Ghazanfari T¹, Nasiri Z¹

1. Immunoregulation Research Center, Shahed University, Tehran, Iran

2. Department of Biology, Faculty of Basic Sciences, Shahed University, Tehran, Iran

Background: The effects of *Allium sativum* (garlic) extract, as a medicinal plant, have been demonstrated on variety of immune cells and their secretory mediators associated with various immune-related diseases. According to the efficiency of garlic bulb extracts in animal studies and unveiled data about the other *Allium* species on immune responder cells, the effects of aqueous extracts from the bulbs of some wild grown *Allium* species from Iran on TNF- α and NO generation in BALB/c mice macrophages were evaluated.

Methods: The bulbs of some *Allium* species were collected from their natural localities and the aqueous extracts of them were prepared. Peritoneal macrophages were isolated from BALB/c mice and were incubated with different concentrations of the bulb extracts. Then, the amounts of TNF- α and NO production in macrophages were measured.

Results: In non-pathogenic conditions of this research, NO generation in macrophages was increased after incubation with bulb extracts of *A. elburzense* and *A. Asarensis*, but their stimulatory effects was not significant. The effects of different concentrations of the bulb extracts of the examined *Allium* species were not significant on TNF- α generation, except for 0.05 mg/ml ($P < 0.001$) bulb extract of *A. sativum*. In spite of the fact that the immunomodulatory effects of the bulb extracts on TNF- α and NO production were not significant, but the identical patterns were observed for the stimulatory effects of the bulb extracts of *A. elburzense* and *A. Asarensis* on the production of these factors in the pretreated macrophages. Also, similar patterns were obtained in reduction of NO production by macrophages after treatment with the bulb extracts of *A. sativum* and *A. Iranicum*.

Conclusion: In general, bulb extracts of the *Allium* species had modulatory effects on the studied mediator factors, which are involved in immune related-diseases, and these species may be considered for treatment studies.

Keywords: *Allium* species, Bulb extract, Macrophage, NO, TNF- α

ICIA2018\Immunopharmacology and Medicinal Plants\poster\7451
7451

Synergistic benefits of thymol and nicotine in alleviating rat model of rheumatoid arthritis (RA)

Sara Golbahari, Seyyed Meysam Abtahi Froushani

Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Background: One logical approach to enhancing rheumatoid arthritis (RA) treatment result is the combination of available medications to provide more favorable outcomes. Immunomodulatory benefits of thymol and nicotine were documented. Here, we investigated synergistic effects of combination therapy by suboptimal doses of atorvastatin and ATRA in experimental autoimmune encephalomyelitis (EAE), an animal model of MS.

Methods: RA was induced by injection of complete Freund's adjuvant into the footpad of Wistar rats. Then, animals were allocated in 4 groups: treated with nicotine (2.5 mg/kg-orally), treated with thymol (100mg/kg-orally), combined therapy (one half doses of each medication) and untreated group. All therapies were initiated at day 5 after induction and continued throughout the study until the day 23 when animals were sacrificed.

Results: Combined in EAE mice with half doses of thymol and nicotine lead to the more favorable clinical outcome than treatment with optimal doses of either medication alone. All medications caused a significant decrease in the serum levels of CRP, myeloperoxidase, splenocytes proliferation index and IL-17 and IFN- γ secretion after ex vivo stimulation of splenocytes. Without any advantage in anti-proliferative effect, combined therapy regressed secretion of pro-inflammatory IL-17 cytokine and nitric oxide in splenocytes more than either medication alone.

Conclusion: Combined thymol and nicotine have synergistic benefits and this approach may be a useful strategy to control RA.

Keywords: Rheumatoid arthritis, thymol, Nicotine, Combination therapy.

ICIA2018\Immunopharmacology and Medicinal Plants\poster\7463

7463

Hydroalcoholic extract of Cinnamon has anti-inflammatory benefits on the acetic-acid induced ulcerative colitis

Mojtaba Salamatian¹, Vahid Mohammadi¹, Seyyed Meysam Abtahi Froushani²

1. Department of internal medicine and clinical pathology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

2. Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Background: Recent evidence has proposed the anti-inflammatory benefits of Cinnamon. Here, we evaluated the protective potential of hydroalcoholic extract of Cinnamon in alleviating ulcerative colitis (UC) induced in rats by acetic acid.

Methods: Luminal instillation of acetic acid was applied to induce UC in the male Wistar rats. Rats in the treatment groups received hydroalcoholic extract of Cinnamon (400 mg/kg PO) or Prednisolone (2 mg/kg PO) daily for 10 consecutive days. At the end, the rats were sacrificed and the disease activity index, the levels of myeloperoxidase, nitric oxide and total protein were evaluated in the colonic homogenized tissue specimens.

Results: The results exhibited that both therapies with hydroalcoholic extract of Cinnamon and Prednisolone could suppress the clinical scores and the mortality rate of ulcerative colitis in a comparable manner. The levels myeloperoxidase activity was decreased in the colonic homogenate of Prednisolone treated rats more than Cinnamon group. Nevertheless, hydroalcoholic extract of Cinnamon regressed the levels of nitric oxide in gut tissues more prominent than Prednisolone.

Conclusion: Hydroalcoholic extract of Cinnamon may be used as an appropriate medication to reduce the inflammatory mediators in UC.

Keywords: Hydroalcoholic extract of Cinnamon, Ulcerative colitis, Anti-inflammatory.

ICIA2018\Immunopharmacology and Medicinal Plants\poster\7497
7497

Immunostimulatory effects of hydroalcoholic extract of Mallow in NMRI mice immunized by Sheep Red Blood Cells.

Zahra Zeinaly, Seyyed Meysam Abtahi Froushani, Hadi Esmaili Govarchin Galee

Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Background: Mallow (*Malva sylvestris*) is a medicinal herb used in folklore medicine for treatment of some infection and inflammation. Here, we investigated the effects of hydroalcoholic extract of Mallow on the immune responses of in NMRI mice Immunized by Sheep Red Blood Cells (sRBC).

Methods: 14 male NMRI mice were divided into two groups and were immunized with the sRBC antigen, twice, with one-week interval. Treated mice received the hydroalcoholic extract of Mallow (200 mg/Kg- orally, every day) for 3 weeks from the beginning of the survey. Finally, the levels of anti-sRBC antibody and the specific cellular immune responses were assumed by microhemagglutination test and footpad thickness, respectively. Furthermore, neutrophils were checked for phagocytosis and respiratory burst.

Results: Hydroalcoholic extract of Mallow significantly potentiated specific cellular immunity and concurrently potentiated specific humoral immunity in mice after challenge with sRBCs. Furthermore, hydroalcoholic extract of Mallow significantly increased the respiratory burst in neutrophil population. Nonetheless, neutral red uptake by neutrophils didn't show any significant change between groups.

Conclusion: It seems that Hydroalcoholic extract of Mallow can be applied as the immunostimulatory agent.

Keywords: Mallow (*Malva sylvestris*), Immunostimulator, NMRI mice.

ICIA2018\Immunopharmacology and Medicinal Plants\poster\7510

7510

Synergistic antitumor effect of NVP-BEZ235 and CAPE on MDA-MB-231 breast cancer cells

Samira Torki¹, Amin soltani¹, Hedayatollah Shirzad², Nafiseh Esmail³, Mahdi Ghatreh Samani¹

1. Cellular and Molecular Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran

2. Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran

3. Department of Immunology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

Background: Triple negative breast cancer (TNBC) is the most lethal and aggressive kind of breast cancer. Studies with TNBC cells suggest that tumor environmental cytokines such as tumor growth factor $\beta 1$ (TGF- $\beta 1$) have important roles in tumors fate.

Methods: In the present study, we aimed to investigate the effect of phosphatidyl inositol 3-kinase/AKT/mammalian target of rapamycin (PI3K/Akt/mTOR) signaling pathway dual inhibitor, NVP- BEZ235 and Caffeic acid phenyl ester (CAPE) on TNBC cell lines (MDA-MB-231), stimulated with TGF- $\beta 1$ for 14 days in vitro.

Results: We found that TGF- $\beta 1$ as a local tumor environmental cytokine plays important role in progression and invasiveness of TNBC cells. NVP-BEZ235 inhibited the enhanced cell viability and CXCR4 expression induced by TGF- $\beta 1$. In addition, the combined treatment of TNBC cell lines with CAPE and NVP-BEZ235 synergistically inhibited cell growth and reduced CXCR4 expression. Also, treatment of MDA-MB-231 cells with CAPE and NVP-BEZ235 led to decreasing the expression levels of p-FOXO3a in a time-dependent manner.

Conclusion: Overall, these results suggest that, tumor metastasis and progression in TNBC cells can be effectively reduced through the concurrent use of NVP-BEZ235 and CAPE. This could be of particular interest in assessing the effects of this therapy in the reduction of tumor metastasis and progression in other tumor types.

Keywords: Caffeic acid phenyl ester; NVP-BEZ235; Triple negative breast cancers; TGF $\beta 1$

ICIA2018\Immunopharmacology and Medicinal Plants\poster\7570
7570

Effect of cinnamon and turmeric aqueous extracts on serum interleukin-17F level of high-fructose fed rats

Maryam Sohrabi¹, Mahdi Alahgholi-Hajibehzad^{2,3}, Zahra Gholami Mahmoodian¹, Sayed Ali Hosseini Siyar⁴, Alireza Zamani^{2,3*}

1. Department of Anatomy, School of Medicine, Hamadan University of Medical Sciences. Hamadan, Iran

2. Department of Immunology, School of Medicine, Hamadan University of Medical Sciences. Hamadan, Iran

3. Molecular Immunology Research Group, Research Center for Molecular Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

4. Department of Animal Science, Agriculture Faculty, Bu-Ali Sina University, Hamadan, Iran

Background: Studies indicate overweight and obesity induce a chronic low-grade inflammation. Inflammation predisposes obese people to some chronic diseases and cancers. This study was conducted to evaluate the effect of two weight lowering and anti-inflammatory agents, cinnamon and turmeric, on the serum level of interleukin-17 (IL-17) as a pro-inflammatory cytokine.

Methods: Sixty-four rats were designated in eight groups. All groups except control, cinnamon and turmeric groups received fructose solution. The control group received normal diet. The other groups fed with normal diet plus: cinnamon, turmeric, fructose with cinnamon and fructose with turmeric. The serum level of IL-17F was detected by enzyme-linked immunosorbent assay (ELISA).

Results: High fructose consumption led to increase the body weight gain and IL-17. While, feeding with cinnamon and turmeric caused to decline body weight gain but surprisingly increased IL-17F.

Conclusion: Although, some studies showed cinnamon and turmeric supplementation decrease IL-17 under the standard diet but in the presence of high fructose diet and overweighting their effects were reversed.

Keywords: Interleukin-17, Cinnamon, Turmeric, Fructose, Inflammation, Overweight

ICIA2018\Immunopharmacology and Medicinal Plants\poster\7583
7583

Citrus Unshiu Extract Decreases Inflammatory Cytokines and MDA Production In LPS-Stimulated Macrophage Cell line and C-26 Tumor-Bearing Mice

Sahar Tahaghoghi Hajghorbani ^{1*}, Alireza Rafiei ¹, Mohamad Ali Ibrahim zade ², Zahra Hoseini Khah³

1. Department of Immunology, Medical School, Mazandaran University of Medical Sciences, Sari, Iran

2. Pharmaceutical Sciences Research Center, School of Pharmacy, Mazandaran University of Medical Sciences, Sari, Iran

3. Department of Molecular Cell-Biology, Research Center, Faculty of Medicin, Mazandaran University of Medical Sciences, Sari, Iran

Background: Citrus unshiu (Marcov) is an easy-peeling citrus fruit, which has been used as a traditional Chinese medicine to treat common cold, relieve exhaustion, and cancer. Hesperidin, naringin and nobiletin are representative components of citrus unshiu peel. However, its effects on inflammation remain unknown. In this study, we investigated the in vitro and in vivo effects of citrus unshiu peel (CUP) on the production of pro-inflammatory mediators. In addition, the effects of daily oral administration of CUP on tumor growth in mice bearing tumor was evaluated.

Methods: Water extract of citrus unshiu peel (WCUP) was prepared. Cytotoxicity was analyzed using MTT assay to find out the effective dose of extract. The phytochemical profile of WCUP was analyzed using the Hitachi HPLC system. Murine macrophage RAW 264.7 cell lines were pretreated with CUP and stimulated with LPS. Inflammatory cytokines (IL-1 β , TNF- α and IL-6) and NO production was evaluated using enzyme-linked immunosorbent assay (ELISA) and Griess reagent, respectively. Body weight, tumor size, food intake and survival rate were assessed in male BALB/c mice.

Results: HPLC analyses revealed that the naringin content in the CUP extract was 22% (2, tR: 13.87 min). We found that CUP represses LPS-induced MDA, as well as TNF- α , IL-1 β and IL-6 production. In addition we observed that daily oral administration of CUP to male BALB/c mice bearing C26 reduced the losses in body weight and serum inflammatory cytokines levels compared with saline treatment. Tumor growth was inhibited in 45% of the mice in this group.

Conclusion: WCUP has the in vitro and in vivo anti-inflammatory effects and it can be used as a nutritional supplement for the management of cancer patients with severe weight loss.

Keywords: Immunotherapy, TNF- α , Colon Cancer, Unshiu Extract

ICIA2018\Immunopharmacology and Medicinal Plants\poster\7597
7597

Modulatory effect tretinoin on histopathology of pancreas of streptozotocin-induced diabetes in C57BL/6 mice

Farin Malekifard¹, Nowruz Delirezh², Rahim Hobbenaghi³, Hassan Malekinejad⁴

1. Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

2. Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

3. Department of Pathobiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

4. Department of Pharmacology, Urmia University of Medical Sciences, Urmia, Iran

Background: It has been shown that some drugs such as tretinoin have immunomodulatory and anti-inflammatory activity, which might represent a potential preventive therapy for autoimmune diseases. The present study was undertaken to examine the protective effect of tretinoin on histopathology of pancreas in diabetic mice.

Methods: Diabetes was induced by multiple low-dose of streptozotocin (MLDS) injection (40 mg/kg/day for 5 consecutive days) in male C57BL/6 mice. Mice were considered diabetic when their fasting blood glucose levels were >250 mg/dl. Subsequently, the mice were allocated to three therapeutic groups (n=7 per group) (normal control group, MLDS group and treatment group). In Treatment group, mice were treated with tretinoin (20 mg/kg/day i.p.) for 21 days. Animals were euthanized on day 22 and pancreases were isolated and stained with hematoxylin & eosin (H&E) and Gomori aldehyde fuchsin (GAF) for histological analyses (the number of islets and β cells, diameter of islets) of pancreas.

Results: The number of islets and β cells and diameter of islets were decreased in diabetic rats compared to control animals ($P<0.05$). Tretinoin treatment in streptozotocin-induced diabetic mice increased the mean diameter of islets and the number of islets and beta cells compared with the diabetic group ($p<0.05$).

Conclusion: Tretinoin improved pancreas tissue during destruction of the pancreatic beta-cells in streptozotocin-induced type 1 diabetes in mice.

Keywords Type 1 diabetes, Tretinoin, Streptozotocin, Pancreas

ICIA2018\Immunopharmacology and Medicinal Plants\poster\7625
7625

Immunomodulatory effect of *Cinnamomum aromaticum* on dendritic cell

Abbas Azadmehr^{1,2}, Niloufar Sadat Miri¹, Mojtaba Shahnazi³, Robabe Latifi³, Sahar Mosalla³, Reza Hajiaghachee⁴

1. Department of Immunology, Babol University of Medical Sciences, Babol, Iran

2. Department of Immunology, Qazvin University of Medical sciences, Qazvin, Iran

3. Department of Parasitology, Qazvin University of Medical Sciences, Qazvin, Iran

4. Medicinal Plants Research Center, Institute of Medicinal Plants, ACECR, Karaj, Iran

Background: Extracts from medicinal plants have been reported to possess good immunomodulation. This study was conducted to assess the immunomodulatory effect of *Cinnamomum aromaticum* extract on dendritic cells (DC).

Methods: For this study, DCs were isolated from the spleen of Balb/c mice by using monoclonal antibody and magnetic beads (MACS method). 10^5 cell/well were cultured with 10, 50 and 100 $\mu\text{g/ml}$ concentration of *Cinnamomum aromaticum* extract. The expression of maturity markers of DCs (CD40, CD86, MHCII) were measured by flow cytometry method.

Result: In this present study, the result indicated that the extract of *Cinnamomum aromaticum* significantly ($P < 0.05$) decreased all maturity markers dose-dependently (10, 50 and 100 $\mu\text{g/ml}$) in DC cells via the immunomodulatory effect.

Conclusion: These results suggest that *Cinnamomum aromaticum* extract has potential immunomodulatory effect by inhibition of maturity markers on APCs. So, this medicinal plant can be used as an immunomodulator agent in inflammatory disease. However future studies are necessary.

Keywords: *Cinnamomum aromaticum*, Antigen presenting cells, Immunomodulation

ICIA2018\Immunopharmacology and Medicinal Plants\poster\7626
7626

Immunosuppression properties of *Cinnamomum camphora* extract on dendritic cells maturation markers

Niloufar Sadat Miri¹, Abbas Azadmehr^{1,2}, Mojtaba Shahnazi³, Sahar Mosalla³, Robabe Latifi³, Reza Hajiaghvaei⁴

1. Department of Immunology, Babol University of Medical Sciences, Babol, Iran

2. Department of Immunology, Qazvin University of Medical sciences, Qazvin, Iran

3. Department of Parasitology, Qazvin University of Medical Sciences, Qazvin, Iran

4. Medicinal Plants Research Center, Institute of Medicinal Plants, ACECR, Karaj, Iran

Background: In these studies we examined the impact of *Cinnamomum camphora* extract on dendritic cells. Here, we investigated the effect of *Cinnamomum camphora* extract on maturation and function of DCs.

Methods: DCs were collected from Balb/c mice spleen undergoing MACS method. Then, the cells were cultured and treated with 10, 50 and 100 µg/ml concentration of *Cinnamomum camphora* extract. The expression of CD40, CD86 and MHCII were evaluated by flow cytometry method.

Result: The result of our study demonstrated that this medicinal plant have immunomodulatory effect by significantly inhibition of CD40 and MHCII of DCs markers dose-dependently ($P < 0.05$) in all concentration (10-100 µg/ml). In addition, the extract of *Cinnamomum camphora* only in the concentration of 100 µg/ml significantly ($P < 0.05$) decreased the expression of CD86 marker.

Conclusion: Our findings showed that *Cinnamomum camphora* extract has immunosuppression property. However, further investigations to identify effective compounds in these extract and their effect on immune response are necessary.

Keywords: *Cinnamomum camphora*, Antigen presenting cells, Immunosuppression

ICIA2018\Immunopharmacology and Medicinal Plants\poster\7650

7650

Apoptosis and necrosis of a human colon cancer cell line after treatment by the extract of *Scrophularia megalantha*

Narges Amirjamshidi¹, Hesam Babaei Khameneh¹, Azin Aghamajidi¹, Abbas Azadmehr¹, Ebrahim Zabihi Neyshabouri², Reza Hajiaghache³, Amrollah Mostafazadeh²

1. Department of Immunology, Babol University of medical sciences, Babol, Iran

2. Cellular and Molecular Biology Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, Iran

3. Medicinal Plants Research Center, Institute of Medicinal Plants, ACECR, Karaj, Iran

Background: Medicinal plants are often used as cost-effective and safe anticancer agents. In this line, there is some evidence in literature indicating that *Scrophularia* species are used in traditional medicines to cure many diseases such as eczema, ulcers, cancer, etc. The objective of the present study was to investigate the apoptosis-inducing potential of the *S. megalantha* extract on a human colon cancer cell line; HCT-116.

Methods: HCT-116 cells were cultured with various concentrations of *S. megalantha* extract for 48 h. MTT assay was done to determine cytotoxicity and GI50 index. The cell death pattern was determined by staining with Annexin V-FITC (fluorescein isothiocyanate) and PI (propidium iodide).

Results: The GI50 of the *S. megalantha* extract was found for the colon cancer cells. Flow cytometry analysis showed that the dominant pattern of cell death induced by the *S. megalantha* extract was necrosis, whereas apoptosis was detected with higher doses of this extract.

Conclusion: Our experiments indicated that the extract of *S. megalantha* induced both apoptosis and necrosis to inhibit the proliferation of colon cancer cells. This extract would be a useful agent for improving the treatment of colon cancer. However, further experiments are necessary.

Keywords: Colon cancer; *Scrophularia megalantha*; Apoptosis; Herbal extract

ICIA2018\Immunopharmacology and Medicinal Plants\poster\7733
7733

Effect of dimethyl fumarate on *in vitro* differentiation of naïve CD4⁺ T cells to regulatory T cells

Haideh Namdari¹, Maryam Izad², Zahra Amirghofran^{1, 3*}

1. Department of Immunology, Medical School, Shiraz University of Medical Sciences, Shiraz

2. Department of Immunology, Faculty of Medicine, Tehran University of Medical Sciences, Tehran

3. Autoimmune Diseases Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Over-activated Th17 cells have been associated with the pathogenesis of autoimmune disorders. Regulatory T cells (Tregs) by controlling these cells can dampen autoimmunity and protect body against tissue injury. We investigated the effects of dimethyl fumarate (DMF), a fumaric acid ester isolated from *fumaria officinalis*, on cell proliferation and differentiation of Tregs and the Th17 response.

Methods: The effect of DMF on cell proliferation and the viability was determined by BrdU and propidium iodide (PI) assay. CD4⁺ naïve T cells were isolated from mouse spleen and after stimulation with anti-CD3/anti-CD28 antibodies, cultured in the presence of Treg-inducing cytokines and DMF. We performed flow cytometry for assessment of Tregs development and real-time PCR for gene expressions of specific transcription factors and cytokines.

Results: Cell proliferation analysis showed the inhibitory effects of DMF at various concentrations. Non-cytotoxic concentrations determined by PI assay were used for the next experiments. In differentiation study, naïve CD4⁺ T cells were treated with TGF- β and IL-2 cytokines to differentiate toward Treg cells. The rate of Treg development was approximately 47%. Treatment of cells with DMF reduced the rate of naïve CD4⁺ T cells differentiation to Tregs to less than 18%. Real-time PCR showed that treatment of mice splenocytes with DMF significantly decreased Foxp3 (0.32 ± 0.11 RFC, $p < 0.01$), IL-10 (0.3 ± 0.15 RFC, $p < 0.01$) and IL-17 (< 0.2 RFC, $p < 0.001$) gene expressions. The expressions of TGF- β and ROR γ t also decreased but the levels compared to the control were not significant.

Conclusion: Decreased Tregs generation and IL-10 expression as well as reduced IL-17 cytokine by DMF, suggest the suppressive effect of this compound on Tregs and Th17 cells. Further studies for determination of the effect of DMF on the Treg/Th17 balance at cellular level are recommended.

Key words: DMF; immunomodulation; CD4⁺ T cells; regulatory T cells

ICIA2018\Immunopharmacology and Medicinal Plants\poster\7860

7860

Anti-inflammatory effects of *Lavandula angustifolia* in rat model of myocardial infarction

Nahid aboutaleb¹, Faramarz Souri², Yaser Azizi¹

1. Physiology Research Center, Iran University of Medical Sciences, Tehran, Iran

2. Physiology Department, Faculty of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahwaz, Iran

Background: Studies have demonstrated that acute inflammatory response after myocardial infarction confers irreversible loss of a large number of cardiomyocytes, myocardial necrosis and subsequent heart failure. In the present study, we focused on the anti-inflammatory effects of Lavender in myocardial ischemia/reperfusion injury and its potential mechanisms.

Methods: Male Wistar rat weighing 250-300 g were randomly assigned into three major groups: Sham (n=10), control IR-vehicle (n=10), two cohorts treated with different doses of Lavender oil from *Lavandula angustifolia* +IR (L1/100mg/kg and L2/200mg/kg, n=10 per group). Lavender was administrated interaperitoneally in rats instantly after left anterior descending coronary artery (LAD) ligation for 20 min. At 24 h after reperfusion, rats were sacrificed under deep anesthesia and the heart was stored at -80 °C prior to inflammation assessment. The expression of pro-inflammatory and anti-inflammatory cytokines was evaluated by the immunohistochemical assay.

Results: LAD ligation remarkably increased the expression of pro-inflammatory cytokines (TNF α and IL1 β) and decreased the expression of anti-inflammatory cytokines (IL10) 24 h after reperfusion. After administration of different doses of Lavender in myocardial ischemia/reperfusion injury rats, the expression of TNF α and IL1 β were significantly decreased compared with control IR-vehicle in a dose dependent manner. Moreover, Lavender post-treatment markedly increased the expression of IL10 relative to control IR-vehicle cohort in a dose dependent mechanism.

Conclusion: In summary, our result showed that Lavender post-treatment could ameliorate myocardial injury after MI by targeting inflammatory cytokines. These findings establish a fundamental foundation for future drug design to treat MI.

Keywords: Lavender, Inflammatory Cytokines, Ischemia reperfusion

ICIA2018\Immunopharmacology and Medicinal Plants\poster\7869
7869

Evaluation of the anticancer effects of aqueous extract of persimmon seed on K562 cell Line (Chronic Myeloid Leukemia)

Mohammad Khojasteh¹, Elham Ahmadi², Rozhin Jamali¹

School of Veterinary Medicine, Islamic Azad University, Sanandaj Branch, Sanandaj, Iran

Department of Pathobiology, School of Veterinary Medicine, Islamic Azad University, Sanandaj Branch, Sanandaj, Iran

Background: Chronic myeloid leukemia (CML) is a myeloproliferative disease originating from a constitutively active tyrosine kinase termed Bcr-Abl, which is expressed by an oncogene resulting from a reciprocal translocation between chromosome 9 and chromosome 2. Plant-derived compounds and their semisynthetic serve as a major source of pharmaceuticals for human diseases. It is estimated that approximately 25% of prescriptions handled contain a plant-derived natural product and 74% of the 119 most important drugs currently contain ingredients from plants used in traditional medicine. Persimmon is used in traditional medicine against several human ailments, including cancer. The aim of this study was the investigation of the effect of aqueous extract of persimmon seed on K562 cell line.

Methods: First, the chronic myeloid leukemia K562 cell line was cultured. Concentrations of 25, 50, 100, 200, 400 and 800 µg/ml of aqueous extract of persimmon seed were prepared and the cells were treated for 24, 48 and 72 hours. Next, the inhibitory effects of this extract on K562 cells proliferation was measured by MTT assay. Hoechst (33342) staining and DNA electrophoresis was used to check apoptosis. Data analyses were performed using SPSS 23 software.

Results: In this study, anticancer effect of aqueous extract of persimmon seed on K562 leukemia cell line was confirmed. The results also showed that the aqueous extract of persimmon seed, compared with the control group, had cytotoxic activity that is highest in 800 µg/ml at 72 hours. Statistical analysis indicated that this extract inhibited the growth of K562 cells ($P < 0.05$).

Conclusion: The findings of the present study showed that the aqueous extract of persimmon seed possessed inhibitory effect on K562 cell line. In order to find the underlying mechanism of this activity, further research should be carried out.

Keywords: Leukemia, Apoptosis, Persimmon seed

ICIA2018\Immunopharmacology and Medicinal Plants\poster\7897
7897

Dopamine can instruct M2 anti-inflammatory macrophages

Hanieh Nikkhah, Seyyed Meysam Abtahi Froushani

Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Background: Previous documents indicated that dopamine possesses a spectrum of the direct and indirect role in modulating the immune system. This survey was designed to evaluate the effects of dopamine on the polarization of macrophages.

Methods: The resident macrophages were collected from the peritoneal cavity of Wistar rats by injecting 20 ml of ice-cold PBS and treated with dopamine for 24h in the concentration of 5×10^{-7} M (maximum physiologic concentration).

Results: The flow cytometry results demonstrated that the dopamine treatment significantly increased the levels of CD68+/CD206+ cells (M2 macrophage marker), when macrophages were cultured with dopamine, compared to macrophages alone ($46.9\% \pm 2.1$ vs. $23.7\% \pm 2.98$, $p < 0.001$). The mRNA level of NF- κ B p65 in macrophages pulsed with dopamine was down-regulated compared macrophages without treatments. Albeit, MTT reduction assay indicated that dopamine treatment couldn't any significant changes in the vitality of macrophages.

Conclusion: Collectively, dopamine treatment of macrophages instructs M2 anti-inflammatory macrophages. These findings might offer new insight into the potential mechanisms that underlie the immunomodulatory effects of dopamine.

Keywords: Macrophages, Dopamine, anti-inflammatory.

ICIA2018\Immunopharmacology and Medicinal Plants\poster\7907
7907

Zerumbone may diminish the expression of TGF- β produced by inflammatory fibroblasts

Zahra Radaee¹, Alireza Zamani^{1,2}, Razieh Amini^{2,3}

1. Immunology Department, Faculty of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

2. Research Center of Molecular Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

3. Student Research Committee, Hamadan University of Medical Sciences, Hamadan, Iran

Background: Inflammatory cytokines, TGF- β , play a pivotal role in controlling tumor progression. TGF- β belongs to the family of regulatory cytokines that interfere with the dual use of a wide range of cells that contribute to physiological and pathological processes such as fetal development, tumorigenicity, immune response, inflammation, and cancer. Fibroblasts play an important role in progression of human cancers. Tumor necrosis factor- α (TNF- α) is a proinflammatory cytokine that act as a major role in regulation of the inflammatory process in tumor growth. Zerumbone, the first sesquiterpene isolated from ginger, has been observed to show antitumor activities in multiple cancer cells. The current study was done to explore the role of TNF- α and zerumbone on TGF- β expression.

Methods: The cytotoxicity of zerumbone in fibroblast was determined using MTT assays. The influence of TNF- α on expression of TGF- β was measured using the SYBR green based QPCR and enzyme-linked immunosorbent assays. We also examined the effects of zerumbone on expression of TGF- β .

Results: We could show a significant increase of the expression of TGF- β in fibroblast cells after stimulation by TNF- α . On the other hand, we found that treatment of TNF- α stimulated fibroblasts by zerumbone decreased TGF- β expression.

Conclusion: Our study had suggested that the expression of TGF- β produced by inflammatory fibroblasts can be regulated by zerumbone and as a results, It can be suggested that treatment with the Zerumbone may be an ideal choice for immunological treatment of cancer.

Keywords: TGF- β , TNF- α , Fibroblast, Zerumbone.

ICIA2018\Immunopharmacology and Medicinal Plants\poster\7942

7942

Effects of *Achillea millefolium* Extract on Pancreas Tissue in Diabetic Rats

Mahsa Safari¹, Mohammad Shahbazzadeh²

Department of biochemistry, Faculty of management, Payam-e-Noor University, Tehran, Tehran, Iran

Background: Diabetes is the third leading cause of death in word. Hyperglycemia and oxidative stress have the main factors involved in pathogenesis of diabetes. On the other hand, the antioxidant system is the first defense mechanism of body against oxidative stress. *Achillea millefolium* possesses hypoglycemic and antioxidant effects. This study surveyed the effects of different doses of *Achillea millefolium* Hydro- alcoholic extract (50,100,150 mg/kg) on histological changes of Langerhans islets and serum insulin, nitric oxide and glucose levels.

Methods: A total of 45 male Wistar rats were divided into 8 groups (control, diabetic with STZ, treatment with *Achillea millefolium* (50,100,150 mg/kg) and diabetic treated with *Achillea millefolium*) (50,100,150 mg/kg). Data were analyzed by one-way ANOVA, and P value <0.05 was considered significant.

Results: *Achillea millefolium* extract (100 and 150 mg/kg) significantly decreased serum glucose level ($p<0.01$) and improved the diameter of islets ($p<0.05$) in diabetic rats treated with *Achillea millefolium* extract, compared with the diabetic group. Moreover, at dose of 150 mg/kg, the extract improved serum insulin ($p<0.01$), decreased nitric oxide ($p<0.01$) and increased the weight ($p<0.01$) and number of islets of diabetic rats ($p<0.05$). Histopathological studies also confirmed these changes.

Conclusion: *Achillea millefolium* can improve insulin secretion and serum glucose levels in an animal model of STZ induced diabetes, possibly by reducing nitric oxide production and preventing pancreatic tissue oxidative damage.

Keywords: *Achillea millefolium*, Diabetes, Pancreas, Hyperglycemias, Rats

ICIA2018\Immunopharmacology and Medicinal Plants\poster\8097
8097

Anti-angiogenic activity of β -D-mannuronic acid (M2000) as a novel NSAID with immunosuppressive properties under *in vitro* and *in vivo* models

Mohsen Rastegari Pouyani^{1,2}, Ali Mostafaie³, Kamran Mansouri³, Shahabeddin Mortazavi Jahromi^{1, 4},
Hamid Reza Mohammadi Motlagh³, Abbas Mirshafiey¹

1. Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran
2. Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran
3. Medical Biology Research Center, Kermanshah University of Medical Sciences, Kermanshah, Iran
4. Department of Cellular and Molecular Biology, Kish International Campus, University of Tehran, Kish, Iran

Background: Angiogenesis, the process by which new capillaries are formed from pre-existing ones, plays a pivotal role in the pathogenesis of several disorders including cancer and chronic inflammatory disorders. The β -D-mannuronic acid (M2000), patented (DEU;102016113018.4), is a novel NSAID with immunosuppressive effects and is a matrix metalloproteinase (MMP) inhibitor. Since MMPs play an important role during angiogenesis and due to the existence of various molecular links between angiogenesis and inflammation, we hypothesized that M2000 might also have anti-angiogenic properties.

Methods: Cytotoxic and anti-proliferative effects of M2000 on human umbilical vein endothelial cells (HUVECs) were examined by trypan blue and MTT assays, respectively. For *in vitro* angiogenesis assay we designed a three-dimensional (collagen-cytodex) model. Finally, we used the chick chorioallantoic membrane (CAM) model for *in vivo* assessment of M2000 effects on angiogenesis.

Results: Based on our cytotoxicity assay, endothelial cells showed a high tolerability towards increasing amounts of M2000. According to the anti-proliferative assay M2000 had little or no anti-proliferative effect on HUVECs. Our *in vitro* anti-angiogenesis model was indicative of weak anti-angiogenic effects of M2000. In the *in vivo* model, however, M2000 showed a much more potent anti-angiogenic activity in a dose-dependent manner.

Conclusion: M2000 could be considered as an anti-angiogenic molecule which more likely exerts its activity mainly via indirect effects on endothelial cells and its anti-inflammatory effects may partly be attributable to its anti-angiogenic activity. Therefore, it might be recommended as a candidate for prevention and treatment of cancer, chronic inflammatory diseases and other angiogenesis-related disorders.

Key words: Angiogenesis, M2000, Cancer, Collagen-cytodex, Inflammation, CAM

ICIA2018\Immunopharmacology and Medicinal Plants\poster\8132

8132

Hydroalcoholic extract of *Lavandula officinalis* (lavender) suppresses the phagocytosis and respiratory burst of neutrophils

Negin Mohammadpoor¹, Seyyed Meysam Abtahi Froushani¹, Bahman Mansouri Miotlagh¹, Sahar Etemadi¹

1. Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Background: *Lavandula officinalis* (lavender) is a kind of medicinal plants. It is under clinical research for its anti-inflammatory benefits. This study was performed to investigate the effects of lavender on the neutrophils isolated from orally treated mice with hydroalcoholic extract of Lavender.

Methods: 20 male NMRI mice were allocated into two equal groups: Treatment and Control groups. The animal in treatment groups orally received with hydroalcoholic extract of lavender for 4 constitutive weeks (400 mg/Kg-Daily). Control mice received PBS at the same volume. At the end, peripheral blood neutrophils were isolated by dextran and used for *ex vivo* assays.

Results: Neutral red uptake and MTT reduction assay test showed that Lavender couldn't change the vitality of neutrophils. Lavender significantly decreased the respiratory burst and phagocytosis potential in neutrophil population after challenge with opsonized yeast.

Conclusion: It seems that Lavender can be used as an anti-inflammatory agent in immunopathologic conditions induced by neutrophils.

Keywords: Neutrophils, Lavender, Inflammation.

ICIA2018\Immunopharmacology and Medicinal Plants\poster\8231
8231

Isolation of Ellagic acid from pomegranate peel and study of it's anti-inflammatory effect in mouse

Elham Abbasi

Kermanshah University of Medical Sciences School of Medicine

Background: Inflammation is a protective response to eliminate the primary stimuli, causing cell injury or necrosis and tissue damages from the tissues of reference. Previous studies have shown that there is a correlation between inflammation and heart disease, cancer and chronic inflammatory diseases. On the other hand, pomegranate has been proved to encompass various notable polyphenols in its peel, including punicalagin, hydrolysable tannin which could be hydrolyzed into ellagic acid. Ellagic acid has long been considered a phenolic compound with anti-inflammatory and anti-cancer effects compound to be considered. So, the present study aimed to purify ellagic acid from pomegranate husk and analyze the anti-inflammatory effects of ellagic acid on inflammation model induced by LPS in mouse.

Methods: After grounding pomegranate peel into powder, dried powder was extracted with aqueous-alcoholic solution. In order to purify ellagic acid, recrystallization by methanol was undertaken. The purified product was then assayed and confirmed using FTIR and C-NMR. To assess the anti-inflammatory activity of purified ellagic acid mouse samples were classified into test and control groups before induction by LPS. Test groups were treated with ellagic acid in doses of 10, 50 and 100 milligrams per kilogram of mouse body weight. Finally, the level of TNF- α cytokine in serum and peritoneal fluid in control and test groups was determined by Eliza kit.

Results: The results from purification of ellagic acid showed high purity and yield of 2% from pomegranate peel. Test groups treated with three doses of purified ellagic acid exhibited a significantly decreased TNF- α level in serum and peritoneal fluid comparing to control groups.

Conclusion: Purified ellagic acid with high purity and efficiency reduced the level of TNF- α cytokine. In this study it was shown that a significant impact on the TNF level is observed. The results presented here could be investigated by further studies and various other methods.

Key words: inflammation, Ellagic acid, tumor necrosis factor, lipopolysaccharide

ICIA2018\Immunopharmacology and Medicinal Plants\poster\8318

8318

The effect of Iranian propolis on pro- inflammatory cytokines IL-1B, IL-6 and TNF- α

Mina Motallebnejad¹, Hamid Reza Nouri², Fariba Asgharpour³

1. Department of Oral and Maxillofacial, Oral Health Center, Health Research Institute, Babol University of Medical Sciences, Babol, I.R. Iran

2. Cellular and Molecular Biology Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, I.R. Iran

3. Dental Materials Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, I.R. Iran

Background: Propolis is a resinous substance collected by bees from different plant sources and its compounds are divers by location and their characteristics depend on the type of plants in each region. Propolis is rich in flavonoids and antioxidants and indicate a broad spectrum of biological activities, such as antioxidant, antimicrobial, anticancer and anti-inflammatory properties. In this study investigated the pro-inflammatory effects of Iranian propolis on cytokine production (IL-6, IL-1 β and TNF- α) in murine macrophage cell line (RAW 264.7).

Methods: Propolis was collected in the beehive located in the Polur of Mazandaran province (PEEP) in the spring of 2016. RAW264.7 cells were treated for 12 h with or without various concentrations of PEEP and LPS (1 pg/ml). After that, total RNA was isolated cDNA synthesized and, then cytokines expression was evaluated by quantitative real-time PCR.

Results: IL-1 β significantly was decreased in presence of LPS plus PEEP in concentrations of 0.15 μ g/mL and 1.5 μ g/mL and PEEP (15 μ g/mL) significantly inhibited IL-6 mRNA expression, but PEEP was not effective in of TNF- α production.

Conclusion: These results suggest that PEEP has inhibitory effects on macrophage activity by reducing expression of IL-1 β and IL-6, and may have further-reaching implications for the pharmaceutical industry.

Keywords: Propolis, Pro-inflammatory, Cytokines, Murine macrophage

ICIA2018\Immunopharmacology and Medicinal Plants\poster\8497
8497

The Effect of Methanol Extract of *Echinops Lasiolepis* on TNF- α Production in LPS-activated J774 A.1 Mouse Macrophages

Fateme Sadat Dashti, Hossein Hadinedoushan, Maryam Asadi

Department of Immunology, Reproductive Immunology Research Center, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

Background: Plants as medicines have always played a vital role in human life. Tumor necrosis factor alpha (TNF)- α is one of the macrophage-derived inflammatory cytokine with pleotropic effects in the inflammation process. Some studies have been demonstrated that some of the *Echinops* species have anti-inflammatory activity. In fact, *Echinops lasiolepis* is introduced as one of the native plants of Yazd. Thus, the present study intended to assess the inflammatory activity of *Echinops lasiolepis* on TNF- α secretion in J774 A.1 mouse macrophages.

Methods: At first, methanol extract was prepared by maceration. 105 cells/ well were seeded in 96-well plate in triplicate and were treated with different concentrations of extract and 100 ng/ml Lipopolysaccharides. MTT cytotoxicity assay was used to determine the cell viability. Concentrations of extract with cell viability of more than 90% were used to evaluate the level of TNF- α in the macrophage culture using enzyme-linked immunosorbent assay.

Results: Viability of cells at different extract concentrations of 0.1, 1, 10, 50, 100 and 200 μ g/ml were 91.68, 95.27, 94.2, 90.8, 85.38 and 71.38, respectively. Therefore, cells treated with 50 μ g/ml and lower concentrates of extracts showed more than 90% of viability and their supernatants were used for TNF- α assay. The study results revealed that all concentrations of extract reduced the production of TNF- α .

Conclusion: Our findings showed that methanol extract of *Echinops lasiolepis* may have anti-inflammatory activity via reducing TNF- α production.

Keywords : Tumor necrosis factor , J774A.1 cell line , *Echinops Lasiolepis*

ICIA2018\Immunopharmacology and Medicinal Plants\poster\8578
8578

Influence of Acemannan and Glucomannan as an immunomodulator on E7d vaccine

Mahboubeh Nikookalam¹, Zeinab Salehian¹, Sahar Mirzaee¹, Mehdi Mahdavi², Mohammad Hossein Yazdi^{3,4}

1. Department of Biotechnology, Faculty of Advanced Sciences & Technology, Pharmaceutical Sciences Branch, Islamic Azad University, Tehran-Iran (IAUPS)

2. Department of Immunology, Pasteur Institute of Iran, Tehran, Iran

3. Department of Pharmaceutical Biotechnology and Biotechnology Research Center, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran

4. Evidence-Based Medicine Group, Pharmaceutical Sciences Research Center, Tehran University of Medical Sciences, Tehran, Iran

Background : Acemannan and glucomannan are mannose containing polysaccharides in Aloe Vera mucilage with immune enhancement capabilities were studied for its impact on the immunization profile of E7d Vaccine. These polysaccharides absorbed after oral feeding of mice and remained unchanged in the blood stream. Subsequently it can act as an immunomodulator which is induced both Th1 and Th2 immune response.

Materials and methods: 60C57BL/6 mice were divided into six groups as follow: Group 1, E7d-Alum-Gel; Group 2, E7d-Alum; Group 3, E7d-Montanid -Gel; Group 4, E7d-Montanid; Group 5 and Group 6, Control-Gel and PBS respectively. The oral feeding of Aloe vera gel with 0.05% polysaccharide was done for a period of 30 days. The mice were immunized with E7d vaccine three times on day 0, 14 and 28th. Then tumor was implanted subcutaneously and 2 weeks later, the serum samples of the mice were collected to evaluate the cytokines and antibodies levels by ELISA assay.

Result: analysis of humoral immune responses represented a noteworthy increase of total antibody level in E7d-MONTANID-Ge lgroup in comparisonto E7d-montanid group. Result of survival rate was best in E7d –ALUM-GEL and CONTROL GEL groups.

Conclusion: Aloevera polysaccharides as the potential of enhancing the immune system and also has antitumor activity and may have the capacity to use along with chemotherapy drugs. This effect was possibly derived from inducing IL-2, TNF- α , IFN- γ and other cytokines which are producedin the body and also improves the immunity response activity

Keyword: Acemannan, Glucomannan, E7d vaccine, Immunomodulator

ICIA2018\Immunopharmacology and Medicinal Plants\poster\8608
8608

The Study of Immunomodulatory Effect of Hydroalcoholic Extract Hypericum Scabrum on Wound Healing in Kurdish Ethnomedicine

Amir Amiri Paryan^{1*}, Mehrdad Pooyanmehr², Mohammad Reza khedri³, Erfan Bahrami³

1. *Veterinary Student, Pathobiology & Basic Sciences, Faculty of Veterinary Medicine, Razi University, Kermanshah, Iran*

2. *Department of Basic Sciences and Pathobiology, Microbiology section, Faculty of Veterinary Medicine, Razi University, Kermanshah, Iran*

3. *Veterinary Student, Clinical Sciences, Faculty of Veterinary Medicine, Razi University, Kermanshah, Iran*

Background: Several plants are used in traditional medicine for treatment of human diseases such as wound. Wound is the result of damages to the skin and other soft tissues. In traditional medicine, it has been reported that are species of *Hypericum Scabrum* (HS) has healing effect on peptic ulcer, depression and wound. The aim of this study is to appraise the effect of hydroalcoholic extract of HS on wound healing.

Methods: In this study 24 male wistar rats weighting about 200±25gr. Animals were anesthetized by mixture of ketamine and xylazine in ratio of 3 to 1. A wound was created on the back of rats by incision an area of 2×2 cm. The animals were divided into four equal groups, first group the wound treated and dressed with Phenytoin 1%, second group cream of hydroalcoholic extract of HS 10%, third group received Eucerin and fourth group was left untreated (control). 14 days after incision, wound healing process was evaluated microscopically and macroscopically.

Results: The results showed wound healing process and re-epithelialization in group 2 were more quickly than other groups and compared to group 3 and 4 there was significant difference. Treatment with HS reduced lymphocytes, macrophages and total white cell count compared with other treatments. Probably, effective compounds in the HS extract such as flavonoids, in addition to stimulating macrophages, natural killer cells and fibroblastic factors, angiogenesis and the production of interferon γ , can repair damaged tissue.

Conclusion: In this study, cream of hydroalcoholic extract of HS is effective in wound healing and accelerated the treatment. HS may have reduced inflammation.

Keywords: Wound healing, Macrophage, *Hypericum scabrum*, Inflammation, Treatment

ICIA2018\Immunopharmacology and Medicinal Plants\poster\8612

8612

Umbelliprenin cytotoxicity increased in combination with arsenic trioxide in MT-2 leukemia/lymphoma cells

Zahra Sadat Delbari¹, Houshang Rafatpanah¹, Fatemeh B. Rassouli², Mehrdad Iranshahi³

1. Inflammation and Inflammatory Diseases Research Center, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran
2. Cell and Molecular Biotechnology Research Group, Institute of Biotechnology, Ferdowsi University of Mashhad, Mashhad, Iran
3. Department of Pharmacognosy and Biotechnology, Biotechnology Research Center, Faculty of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

Background: Adult T-cell leukemia/lymphoma (ATLL) is a peripheral T-cell lymphoma caused by human T-cell leukemia/lymphoma virus type1 (HTLV-1). Iran, especially Khorasan province, is known as one of the endemic regions for HTLV-1. Despite advances in treatment of ATLL, the average survival rate of this malignancy is low. Umbelliprenin is a sesquiterpene coumarin with broad-spectrum anticancer and antiproliferative activities. To introduce more effective anticancer strategies against ATLL, here we investigate whether Umbelliprenin could enhance the efficacy of arsenic trioxide, a chemical drug prescribed for ATLL.

Methods: In this regard, at first MT-2 cells, an ATLL cell line, were treated with increasing concentrations of Umbelliprenin (25, 30, 40 and 50 µg/ml) and arsenic trioxide (2, 4, 8 and 16 µM) for 24, 48 and 72 h, and the IC50 value of drugs was determined. Then, cells were treated with combination of Umbelliprenin (25 µg/ml) and arsenic trioxide (2 µM) for 72 h. Finally, viability of cells was measured by alamarBlue[®] test, and apoptosis was determined by PI staining and flow cytometry.

Results: Obtained results indicated that 25 µg/ml Umbelliprenin increased the toxicity of 2 µM arsenic trioxide, and therefore, had synergic effects with this anticancer drug.

Conclusion: In conclusion, current findings suggest that combinatorial use of Umbelliprenin and arsenic trioxide can be helpful in ATLL treatment.

Keywords: Umbelliprenin, Adult T-cell Leukemia/Lymphoma, Cytotoxicity, Arsenic trioxide.

ICIA2018\Immunopharmacology and Medicinal Plants\poster\8626
8626

Effects of sesquiterpene lactone parthenolide on the expression of NF- κ B in human lymphoma cells

Hamide Kouhpeikar¹, Mohammad Hadi Sadeghian¹, Houshang Rafatpanah², Fatemeh B. Rassouli³,
Mehrdad Iranshahi⁴

1. Cancer Molecular Pathology Research Center, Department of Hematology and Blood Bank, Faculty of Medicine, Mashhad University of Medical Sciences, Iran
2. Inflammation and Inflammatory Diseases Research Center, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran
3. Cell and Molecular Biotechnology Research Group, Institute of Biotechnology, Ferdowsi University of Mashhad, Mashhad, Iran
4. Department of Pharmacognosy and Biotechnology, Biotechnology Research Center, Faculty of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

Background: Parthenolide (PTL), a sesquiterpene lactone (SL) originally purified from the shoots of feverfew (*Tanacetum parthenium*), has shown potent anticancer and anti-inflammatory activities. It is currently being tested in cancer clinical trials. Adult T-cell leukemia/lymphoma (ATLL) is an aggressive malignancy of mature activated T cells caused by human T-cell lymphotropic virus type I (HTLV-1). Iran, especially Khorasan province, is known as one of the endemic regions for HTLV-1. Despite advances in treatment of ATLL, the average survival rate of this malignancy is low. The virus carries the Tax oncogene which constitutively activates the NF- κ B pathway.

Methods: In present study, we investigated parthenolide effects on the expression NF- κ B (REL-A) in human lymphoma cells. In this regard, MT2 cells were treated with 1 μ g/ml parthenolide for 72 h, while cells treated with 0.2% DMSO (used as parthenolide solvent) were considered as relevant control. Then, the total cellular RNA was extracted and treated with DNase I. In the following, cDNAs were synthesized and their fidelity was confirmed by PCR using GAPDH primers. Real-time RT-PCR was conducted using Taq man prob and specific primers for NF- κ B (REL-A).

Results: Results of current study revealed that parthenolide (in non-toxic concentration) significantly ($p < 0.05$) down regulated the expression of NF- κ B (REL-A) in MT2 cells.

Conclusion: Accordingly, this natural sesquiterpene lactone could be considered as an effective agent to attenuate malignant properties of human lymphoma cells in future *in vivo* studies.

Key words: parthenolide, Adult T-cell leukemia/lymphoma, HTLV-1, NF- κ B

ICIA2018\Immunopharmacology and Medicinal Plants\poster\9753
9753

A Comparative Study on the Effects of *Allium asarense* and *Allium sativum* Extracts on Lymphocytes

Hatami H¹, Radjabian T^{1,2}, Hosseinpour Z¹, Ghazanfari T¹, Naghizadeh MM¹

1. Immunoregulation Research Center, Shahed University, Tehran, Iran

2. Department of Biology, Faculty of Basic Sciences, Shahed University, Tehran, Iran

Background: Garlic (*Allium sativum*) is used as an effective herbal medicine for the treatment of the autoimmune diseases and cancers. Although there are many studies on two *Allium* species (*Allium sativum* and *Allium cepa*), there is no scientific report on the other *Allium* species. The aim of this study was to compare the *in vitro* effects of aqueous bulb extracts of wild *A. asarense* and cultivated *A. sativum* on lymphocytes.

Methods: Lymphocytes were isolated from BALB/c mice and were incubated with various concentrations of the aqueous bulb extracts of the studied *Allium* species. Finally, the cytokines concentration and lymphocyte viability were evaluated and compared.

Results: The results did not show significant increases in the viability of lymphocytes after incubation with *A. asarense* bulb extract, however significant increases were observed in the viability of lymphocytes after treatment with lower concentration (≤ 0.05 mg/ml) of *A. sativum* extract. Although no significant changes were found in IL-17 and IL-4 concentration after treatment with two *Allium* extracts, IFN- γ concentration was decreased by *A. asarense* extract and was increased by *A. sativum* extract.

Conclusion: It suggests that similar to *A. sativum*, the extract of *A. asarense* may be regulate the IFN- γ levels.

Keywords: *Allium* species, IFN- γ , IL-4, IL-17

ICIA2018\Immunopharmacology and Medicinal Plants\poster\9754
9754

Evaluation of *Allium asarense* and *Allium sativum* Aqueous Extracts on Macrophages

Hatami H¹, Hosseinpour Z¹, Radjabian T^{1,2}, Ghazanfari T¹

1. Immunoregulation Research Center, Shahed University, Tehran, Iran

2. Department of Biology, Faculty of Basic Sciences, Shahed University, Tehran, Iran

Background: It has been showed that some of *Allium* species have immunomodulatory effects on human immune system. Recent studies on *A. cepa* and *A. sativum* showed that their extract have therapeutic effects including anti-cancer, anti-diabetes, anti-oxidant activities. In this research we performed a comparative study about the effects of bulb extracts from wild *A. asarense* and cultivated *A. sativum* on macrophages responses. Methods: Peritoneal macrophages were isolated from BALB/c mice and were incubated with various concentrations of the aqueous bulb extracts of the studied *Allium* species. Then, the amounts of TNF- α and NO generation and MTT in macrophages were measured.

Results: Our results showed that the bulb extracts of both *Allium* species had positive effects on macrophages viability, but the significant stimulatory effects of *A. asarense* bulb extract were observed at the lower concentrations (≤ 0.01 mg/ml). Also significant increases were found in TNF- α concentration after treatment with 0.1, 0.05 and 0.001 mg/ml of the *A. asarense* bulb extract and 0.05 mg/ml *A. sativum* bulb extract. No significant changes were observed in nitric oxide levels after treatment with the extracts.

Conclusion: Generally, it seems that in addition to *A. sativum*, *A. asarense* could be used as an effective medicinal plant for immunoregulation.

Keywords: *Allium sativum*, *Allium asarense*, Nitric Oxide, TNF- α

ICIA2018\Immunopharmacology and Medicinal Plants\poster\9765
9765

Parthenolide cytotoxicity increased in combination with Arsenic trioxide in MT2 lymphoma cells

Hamide Kouhpeikar¹, Mohammad Hadi Sadeghian¹, Houshang Rafatpanah², Fatemeh B. Rassouli³, Mehrdad Iranshahi⁴

1. Cancer Molecular Pathology Research Center, Department of Hematology and Blood Bank, Faculty of Medicine, Mashhad University of Medical Sciences, Iran

2. Inflammation and Inflammatory Diseases Research Center, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

3. Cell and Molecular Biotechnology Research Group, Institute of Biotechnology, Ferdowsi University of Mashhad, Mashhad, Iran

4. Department of Pharmacognosy and Biotechnology, Biotechnology Research Center, Faculty of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

Background: Adult T cell leukemia-lymphoma (ATLL) is an aggressive lymphoid neoplasm that occurs only in patients with human T-lymphotropic virus, type I (HTLV-1) viral infection. Iran, especially Khorasan province, is known as one of the endemic area for HTLV-1. Despite advances in treatment of ATLL, long term recovery of this malignancy has been failed. Parthenolide, a naturally occurring sesquiterpene lactone derived from fever few (*Tanacetum parthenium*), exhibits exceptional anti-cancer and anti-inflammatory properties, making it a prominent candidate for further studies and drug development. To introduce more effective anticancer strategies against ATLL, here we investigate whether parthenolide could enhance the efficacy of arsenic trioxide, a chemical drug prescribed for ATLL

Methods: In this regard, at first MT-2 cells, an ATLL cell line, were treated with increasing concentrations of parthenolide (1.25, 2.5 and 5 µg/ml) and Arsenic (1, 2, 4 and 8 µM) for 24, 48 and 72 h, and the IC50 value of drugs was determined. Then, cells were treated with combination of parthenolide (1 and 2 µg/ml) and Arsenic (1 and 2 µM) for 72 h. Finally, viability of cells was measured by alamar blue test

Results: Obtained results indicated that 1 µg/ml parthenolide increased the toxicity of Arsenic 2 µM, and therefore, had synergic effects with this anticancer drug

Conclusion: In conclusion, Current findings suggest that combinatorial use of parthenolide and arsenic trioxide could be considered for designation of novel chemotherapy regimes

Key words: parthenolide, Arsenic trioxide, Adult T-cell leukemia/lymphoma, HTLV-1

ICIA2018\Immunopharmacology and Medicinal Plants\poster\9798
9798

Anti- inflammatory effect of *Beta vulgaris* leaf aqueous extract in rat

Azad Bahrampour¹, Roonak Azizbeigi²

1. Faculty of Veterinary Medicine, Sanandaj Branch, Islamic Azad University, Sanandaj, Iran

2. Department of Physiology, Faculty of Veterinary Medicine, Sanandaj Branch, Islamic Azad University, Sanandaj, Iran

Background: Inflammation is a local response of tissue to injury or infection. It has been indicated that there is a protective response which is involving leukocytes, blood vessels, and inflammatory mediators. Inflammation which is unregulated mechanism can be result in excessive free radical activity and tissue destruction. *Beta vulgaris*, belonging to Chenopodiaceae, is used in traditional medicine to treat a wide variety of diseases. *B. vulgaris* leaves contain many possesses different potential included antioxidants, anti-rheumatic, and anti-cancer activity. Therefore, this study was conducted to evaluate the effect of *B. vulgaris* leaf aqueous extract on the inflammation induced by xylene.

Methods: A number of 25 adult male Wistar rats were divided randomly into 5 groups (n=5) Control (C), xylene (S), and different dose of (100, 200, and 300 mg/kg;ip) which is ,contain B1, B2, and B3 groups ,respectively. *B. vulgaris* leaf aqueous extract administrated 30 min before xylene. The volume injection of xylene was (0.03 ml/kg;sc). After treatment, the animal's leukogram and neutrophil/lymphocyte ratio (NLR) were calculated. Also, the inflammation was measured according to the weight difference of a plug with 7 mm of thickness in treated and untreated ears.

Results: The results indicated that the inflammation was significantly reduced in the B3 group compared to S group ($P<0.01$). The leukocytes count decreased significantly in extract-treated groups compared to the xylene group ($P<0.05$). NLR in the S group was significantly higher than B2 and B3 groups ($P<0.05$). There was no significant difference between B1 and S groups in the NLR ($P>0.05$).

Conclusion: Our finding suggested that the *B. vulgaris* leaf aqueous extract can control xylene induced inflammation. It seems that more researches are needed to investigate the role of this plant in the inflammatory processes.

Keywords: *Beta vulgaris*, Inflammation, Leukocytes, rat

ICIA2018\Immunopharmacology and Medicinal Plants\poster\9888
9888

Medicinal plant *Ganoderma Lucidum* reduced l-arginine induced nitric oxide release by macrophages

Sajjad Ahmadi Renani¹, Kazem Ahmadi²

1. Tehran University of Medical Science, Tehran, I.R.Iran

2. Department of Immunology, Faculty of Medicine, Baqiyatallah University of Medical Science, Tehran, I.R.Iran

Background: Inflammation is an unpleasant process which occurs in response to a wide range of agents such as infection and immune disorder. In this regard, over production of some pro-inflammatory mediators such as Nitric Oxide (NO) could lead to chronic diseases of the inflammatory origin. *Ganoderma lucidum* (*G. lucidum*) is a medicinal plant that in healthy individual boosts the immune system. The aim of this study was looking on the effect of *G.Lucidum* on NO release as an immune mediator by activated macrophages treated with L-Arginine.

Methods: In the in vivo study, mice were orally treated with *G.Lucidum* ranging from 2.5µg to 80µg/day/mouse in drinking water for 30 days. Peritoneal Macrophages of *G.Lucidum* treated and non-treated mice were extracted, counted and the 1×10^6 suspension cells were cultured in 24 wells plate in the presence or absence of 1 mM L-arginine and appropriate LPS in 2 ml volume phenol red and L-arginine free RPMI 1640 medium. Cells were incubated at 37°C and 5% CO₂ in air for 24h. Supernatants were collected, centrifuged and the amount of NO₂ as an indicator of NO production was measured by green,s method.

Results: The results showed that L-Arginine treated macrophages were produced more NO comparing to control groups $p < 0.001$. Meanwhile *G.Lucidum* reduced L-arginine induced NO release in a dose dependent manner.

Conclusion: Previous studies indicate that NO is partly involved in inflammatory disease, and the *G.Lucidum* works as an anti-inflammatory factor. Therefore, the reduction of NO release by *G.Lucidum* treated macrophages show the possibility of using *G.Lucidum* treatment as a safe option in inflammatory disease.

Keywords: *G.Lucidum*, Inflammation, Macrophages, Nitric Oxide.

ICIA2018\Immunopharmacology and Medicinal Plants\poster\9897
9897

Anticancer activity of Nanoemulsion formulation of Medical leech protein extraction on human breast cancer cells MCF-7

Sara Mohamadi Saani¹, Amir Shakouri², Jalal Abdolalizadeh³, Saeed Zeinali Heris¹

1. Faculty of Chemical and Petroleum Engineering of Tabriz University

2. Drug Applied Research Center of Tabriz medical science

3. Paramedical Faculty, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Nanoemulsion serves as an attractive vehicle for the delivery of drugs. In medicine, Nanoemulsion has glowed a rapidly growing notice as it potentials to solve an amount of issues associated with conventional therapeutic agent. Over the past several decades, significant progress has been made in the improvement and application of engineered nanoparticles to treat cancer more effectively. For example, therapeutic agents have been combined with nanoparticles engineered with optimal sizes, shapes, and surface properties to increase their solubility, prolong their circulation half-life, develop their bio distribution, and reduce their immunogenicity. Nowadays nanoemulsion is using to increase penetrate tissue and release drug at the target cells with linking target antibody on surface of Nanoemulsion.

Methods: High energy techniques are used for the preparation of nanoemulsions. In high energy methods, great disruptive forces are provided by the usage of mechanical devices such as ultrasonicators, and high pressure homogenizers which yields small sized droplets. This method is widely used for the producing nanoemulsions. High pressure homogenizer is used to yield nanoemulsions. Sonication can be widely used to formulate nanoemulsions on small scale. In addition, we use dynamic light scattering (DLS) to monitor the formation of the nanoemulsions complexes. For demonstrate the anti-cancer effect of encapsulation of medical leech protein extract in nanoemulsions, human breast adenocarcinoma cell line (MCF-7) was obtained from the American Type Cell Collection (ATCC). Cells were cultured at an initial inoculum cell concentration of 10^4 cells/cm² in 15 ml Roswell Park Memorial Institute medium (RPMI) with 10% FBS (v/v) strep in Corning® 75 cm² cell culture flask. The cultured cells were incubated at 37°C in 5 % CO₂ humidified atmosphere. Nanoemulsions treatment on cell capability of MCF-7 cells were specified using a colorimetric technique, which is an LDH assay on a 96-well plate in RPMI with 10% of FBS per well at cell density 1×10^4 cell/wells incubated for 24 h, 48 h (37 °C, 5% CO₂ and 95% humidity). DAPI staining is used for exhibit the apoptosis of the cells that treated with nanoemulsions.

Results: DLS showed that the size of nanoemulsions droplets is less than 100 nm and PdI is about 0.130. LDH assay demonstrated that MCF-7 Cancer Cells were sensitive to nanoemulsions that encapsulate with medical leech protein extract and it can be destroyed over 60% of cancer cells agent's normal cell line (HUVEC). In DAPI staining DNA of cancer cells that treated with nanoemulsions were degraded.

Conclusion: Nanoemulsions technology is useful way to treat cancer cells with targeting drug like this project with no side effect and Cytotoxicity in human body.

Keywords: Nanoemulsions; Breast Cancer; Medical leech; MCF-7 Cell line

ICIA2018\Immunopharmacology and Medicinal Plants\poster\9910

9910

The effect of Ginger powder on IL2, TNF α and IL1 β cytokines gene expression levels in Patients with Active Rheumatoid Arthritis

Aryaeian Naheed¹, Mahmoudi Mahdi², Shahram Farhad², Poursani Shiva², Jamshidi Fatemeh², Tavakoli Hajar³

1. Department of Nutrition, School of Public Health, Iran University of Medical Sciences, Tehran, Iran

2. Rheumatology Research Center, Tehran University of Medical Sciences, Tehran, Iran

3. Departments of Cellular and Molecular Nutrition, School of Nutritional Sciences and Dietetics, Tehran University of Medical Sciences, Tehran, Iran

Background: Rheumatoid arthritis (RA) is a chronic inflammatory disease that affects the joints and consequently leads to the destruction of cartilage and bone lesions. Ginger is a native plant to India, which traditionally has been used in treatment of osteoarthritis, joint and muscle pain, neurological diseases, and inflammation of gums, tooth pain, asthma, stroke, diabetes, and constipation. This study aimed to determine the effect of ginger on some immunological and inflammatory factors in patients with rheumatoid arthritis.

Methods: In this study, 66 patients with active rheumatoid arthritis were enrolled. Patients were randomly divided into two groups that were treated using ginger and placebo (wheat flour) respectively. To determine the effect of confounding factors on the findings of the study, questionnaires for nutrient intake, physical activity, medication, and determination of body mass index were filled for each participant, at the beginning and end of the study. Serum CRP and mRNA levels of IL-1 β , IL-2, and TNF- α were measured by ELISA and Quantitative Real-Time PCR respectively.

Results: Results of the study showed ginger supplementation caused a significant decrease in CRP ($P=0.05$), and IL-1 β ($P=0.02$). TNF α mRNA level in ginger group compared to placebo group decreased, although the difference was not significant between the two groups ($P=0.09$). Ginger had no effects on IL2 expression.

Conclusions: The study showed that ginger reduces inflammatory markers, CRP, and IL-1 β in patients with active RA and it seems that ginger can improve the inflammation in the patients.

Keywords: Ginger, Active rheumatoid arthritis, Inflammatory markers

ICIA2018\Immunopharmacology and Medicinal Plants\poster\9911
9911

The therapeutic effects evaluation of *Boswellia serrata* extract in experimental autoimmune encephalomyelitis in mice C57BL / 6

Abdolghader Taene ¹, Gholamreza Anani Sarab ²

1. Student Research Committee, Birjand University of Medical Sciences, Birjand, Iran

2. Cellular and Molecular Research Center, Birjand University of Medical Sciences, Birjand, Iran

Background: Multiple sclerosis (MS) and its animal model, experimental autoimmune encephalomyelitis (EAE), are inflammatory autoimmune diseases of the central nervous system. *Boswellia serrata* is widely used as an anti-inflammatory, anti-arthritic, anti-proliferative, anti-microbial and analgesic effect remedy in traditional and herbal medicine. In this study, we investigated the effect of methanolic extracts from *Boswellia serrata* on EAE and on the serum cytokine level and Total Antioxidant Capacity (TAC) in C57BL/6 mice.

Methods: EAE is induced in C57BL/6 mice by immunization with an emulsion of MOG35-55 in complete Freund's adjuvant (CFA), followed by administration of pertussis toxin (PTX) in PBS, first on the day of immunization and then again the following day. The mice were administered with PBS in the control groups and *Boswellia serrata* extract (200 mg/kg, daily) in the treatment group. Serum levels of IL17A and IL23 were measured using enzyme-linked immunosorbent assay (ELISA). Histopathologic assessments were performed by hematoxylin and eosin (H and E) and luxol fast blue (LFB) staining. TAC of samples were measured by using the determination of ferric reducing ability power (FRAP).

Results: Treatment of mice with daily oral 200 mg/kg *Boswellia serrata* significantly reduced the clinical symptoms and cerebral infiltration of inflammatory cells in C57BL/6 mice with EAE. Also treatment prevented weight loss and increased serum levels of IL17A, IL23 in immunized mice with MOG35-55. TAC production was significantly elevated in *Boswellia serrata* treated mice (P<0/005).

Conclusion: These results showed that treatment with *Boswellia serrata* may attenuate disease severity, inflammatory responses in EAE-induced mice and may be potentially useful for the treatment of Multiple Sclerosis (MS).

Keywords: experimental autoimmune encephalomyelitis, *Boswellia serrata*, Total Antioxidant Capacity, multiple sclerosis

ICIA2018\Immunopharmacology and Medicinal Plants\poster\9915
9915

The anti-inflammatory effects of *Matricaria chamomilla* hydro-alcoholic and aqueous extracts

Zahra Asadi¹, Mojtaba Bagheri¹, Tooba Ghazanfari²

1. Department of Biochemistry, Institute of Biochemistry and Biophysics, University of Tehran

2. Immunoregulation research center, University of Shahed

Background: *Matricaria chamomilla* (MC) is one of the most useful medicinal herbs in Iran, with anti-inflammatory, anti-oxidant and anti-bacterial properties in traditional medicine. Essential oil and phenolic compounds (e.g., coumarins, flavones, flavonols, and flavanones) are two major groups of the MC compounds. Although the anti-inflammatory activities of some of these compounds have been investigated, the effects of the MC extracts on the immune cells inflammatory parameters have not fully been investigated so far. We, therefore, planned to investigate the effects of the MC hydro-alcoholic (70%) and aqueous extracts on the cell viability and the amount of nitric oxide (NO) production in mouse macrophages *in vitro*.

Methods: The mouse peritoneal macrophages were separated and after preparation they were incubated with various concentration of the MC hydro-alcoholic and aqueous extracts (25, 50, 100, 200, 400, 800 and 1600 µg/ml) in the presence of lipopolysaccharide (LPS) (10 µg/ml) for 20 hrs. Extract without cells was taken as the negative control. Then, the macrophages cell viability was measured by using the MTT assay. The amount of the NO produced by the macrophages was measured by Griess test using the cell culture supernatant.

Results: Both hydro-alcoholic and aqueous extracts of the MC significantly reduced the amount of NO in all concentration except 1600 µg /ml. The MC hydro-alcoholic exposure significantly reduced the macrophages cell viability in the high concentration; i.e., 800 and 1600 µg/ml. The macrophage cell viability exposed to the MC aqueous extract increased for the concentration of 100, 400 and 800 µg/ml.

Conclusion: The MC hydro-alcoholic and aqueous extracts have strong anti-inflammatory property that can result in the reduction of the NO production in macrophages. Also, the MC hydro-alcoholic extract can kill activated macrophages. It seems that the MC aqueous extract exerted it anti-inflammatory effect by inducing the alternative macrophage activation.

Keywords: MC hydro-alcoholic extract, MC aqueous extract, Macrophage, Inflammation.

ICIA2018\Immunopharmacology and Medicinal Plants\poster\9959
9959

Anti-inflammatory effects of aqueous extract of *Eryngium campestre* on Ethylene Glycol-Induced calcium oxalate kidney stone in rats

Saman Seyedabadi¹, Mohammad Sadegh Naghizadeh², Sajjad Esmaeili², Mehran Falahpour², Gholamreza Anani Sarab³

1. Department of medical biotechnology, faculty of medicine, Birjand University of Medical Sciences, Birjand, Iran

2. Student Research Committee, Birjand University of Medical Sciences, Birjand, Iran

3. Infectious Disease Research Center, Birjand University of Medical Sciences, Birjand, Iran

Background: Kidney stone is one of the most prevalent diseases known to man and is the third most common disorder of the urinary tract. It is a common chronic disorder that affects 10-15% of the general population around the world. This study aims to evaluate the anti-inflammatory effect of *E. campestre* using the aqueous extracts obtained from aerial parts on Ethylene Glycol (EG)-induced calcium oxalate (CaOx) kidney stone in rats.

Methods: 64 male Wistar rats randomly divided into 8 groups including: negative control: standard diet and drinking water; received 1 ml of normal saline orally each day, kidney stone control: standard diet and 0.1 % of EG in drinking water and during the study; received 1 ml of normal saline orally each day, 3 prophylactic groups with 100, 200, 400 doses: standard diet and 0.1 % of EG in drinking water during the study; received 100, 200, 400 mg/kg extract daily, respectively and 3 treatment groups with 100, 200, 400 doses: standard diet and 0.1 % of EG in drinking water during the study, and from the day 15 until the end of the study; received the extract at the doses of 100, 200, 400 mg/kg daily, respectively. After the day 30, rats anaesthetized and killed. Blood collected and the kidneys excised immediately. Slides from each one's kidneys prepared and stained with Hematoxylin & Eosin method, also levels of IL-1 β and IL-6 determined in rat's serum by rat competitive ELISA kit.

Results: *E. campestre* reduced both cytokines IL-1 β and IL-6, showing a significant reduction in IL-1 β and IL-6 in all prophylactic groups. Also, IL-1 β and IL-6 reduced significantly in the treatment groups in 400 and 200 dosage, respectively (p-value <0.5).

Conclusion: This study suggests a potential therapeutic application of *E. campestre* extract for prophylaxis and treatment of the inflammation caused by CaOx kidney stone.

Keywords: kidney stone, cytokines, *E. campestre*, inflammation

ICIA2018\Immunopharmacology and Medicinal Plants\poster\10019

10019

Cytotoxic effects of combined copper oxide nanoparticles produced by walnut shells using and hydroalcoholic extract of *Aloe vera* against K562 cell line

Ardeshtir abbasi¹, Nasim Rahmani Kukia ², Seyyed Meysam Abtahi Froushani ²

1. Department of Immunology, Tarbiat Modares University of Medical Sciences, Tehran, Iran

2. Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Background: In recent studies, anticancer effects of *Aloe vera* and copper oxide nanoparticles (CuO NPs) have been documented. Here, we tested whether combining of the hydroalcoholic extract of *A. vera* and green CuO NPs produced by using eco-friendly and non-toxic walnut shells could provide synergistic cytotoxicity against K562 cell lines and peripheral blood mononuclear cells (PBMC).

Methods: first, green CuO nanoparticles were synthesized and characterized. And, hydroalcoholic extract of *A. vera* was produced. The K562 cells or PBMCs (1×10⁵ cells/100 µl/well) were incubated for 24h with serial dilution of *Aloe vera* extract (0, 20, 40, 60, 80, 160 and 320 mg/ml) or serial dilution of green CuO NPs (0, 50, 100, 200 and 400 µM). After incubation, the survivability of present cells was determined by MTT methods. In another experiment, The K562 or PBMC cells were treated with a combination of *Aloe vera* extract and Green CuO NPs at minimal cytotoxic concentrations and the inhibitory present was calculated.

Results: The hydroalcoholic extract of *A. vera* and green CuO NPs had cytotoxic effects against K562 cell-line in a dose-dependent manner. Unlike *A. vera*, the marginal safety of green CuO NPs is low because the IC₅₀ value of the green CuO NPs against K562 cell line had no significant difference comparing the IC₅₀ value of the Green CuO NPs against PBMCs. Moreover, combined treatment with minimal cytotoxic doses provided synergistic benefits and lead to more cytotoxic effect against K562 cell-line than their individual treatment of K562 without any additive cytotoxic effect against PBMCs.

Conclusion: This combination provides more favorable cytotoxicity against K562 cell-line without any additive cytotoxicity against PBMCs.

Keywords: K562, Green synthesis, walnut shells, Copper oxide nanoparticle, *Aleo vera*.

ICIA2018\Immunopharmacology and Medicinal Plants\poster\10036
10036

Effects of Umbelliprenin on the expression of NF- κ B in human leukemia/lymphoma cells

Zahra Sadat Delbari¹, Houshang Rafatpanah¹, Fatemeh B. Rassouli², Mehrdad Iranshahi³

1. *Inflammation and Inflammatory Diseases Research Center, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran*

2. *Cell and Molecular Biotechnology Research Group, Institute of Biotechnology, Ferdowsi University of Mashhad, Mashhad, Iran*

3. *Department of Pharmacognosy and Biotechnology, Biotechnology Research Center, Faculty of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran*

Background: Adult T-cell leukemia/lymphoma (ATLL) is a peripheral T-cell lymphoma caused by human T-cell leukemia/lymphoma virus type1 (HTLV-1). The virus carries the Tax oncogene. Tax is a potent oncogene that deregulates cellular gene expression by persistently activating signaling pathways such as NF- κ B. Iran, especially Khorasan province, is known as one of the endemic regions for HTLV-1. Despite advances in treatment of ATLL, the average survival rate of this malignancy is low. In this regard, introducing novel anticancer agents *in vitro* could help clinicians to design more effective therapeutic regimes for ATLL. Umbelliprenin (UMB) is a natural coumarin extracted from dried roots of *Ferula szwitsiana* with broad-spectrum anticancer effects.

Methods: In present study, we investigated Umbelliprenin effects on the expression NF- κ B (REL-A) in human lymphoma cells. In this regard, MT-2 cells were treated with 25 μ g/ml Umbelliprenin for 72 h, while cells treated with 0.2% DMSO were considered as relevant control. Then, the total cellular RNA was extracted and treated with DNase I. In the following, cDNAs were synthesized and their fidelity was confirmed by PCR using GAPDH primers. Real-time PCR was conducted using Taq man prob and specific primers for NF- κ B (REL-A).

Results: Results of current study revealed that Umbelliprenin (in non-toxic concentration) significantly ($p < 0.0001$) down regulated the expression of NF- κ B (REL-A) in MT-2 cells.

Conclusion: Regarding the role of NF- κ B in the proliferation of cancer cells and chemoresistance, the results of this study, introduces this natural coumarin as an effective factor in designing novel approaches against ATLL.

Keywords: Umbelliprenin, Adult T-cell leukemia/lymphoma, HTLV-1, NF- κ B



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Innate Immunity and Inflammation

Poster Presentation

ICIA2018\Innate Immunity and Inflammation\Poster\5260

5260

Effects of acetyl salicylic acid (ASA) and its derivatives on inflammation in diabetesShabnam Ghazanfari Sarabi¹, Mehran Habibi Rezaei^{1,2}*1. School of Biology, College of Science, University of Tehran, Tehran, Iran**2. Nano-Biomedicine Center of Excellence, Nanoscience and Nanotechnology Research Center, University of Tehran, Tehran, Iran*

Background: Glycation- the leading cause of Diabetes- is a non-enzymatic interaction between carbohydrates and proteins forming advance glycated end products (AGEs). AGEs and RAGEs (receptor for AGEs) interaction cause the formation of pro-inflammatory molecules and free radical such as reactive oxygen species (ROS) which promote the pathology of diabetes. Gene expression by NF- κ B and activator protein-1 (AP-1) transcription factors originate from ROS release. NF- κ B not only directly responds to the oxidative stress but also induces several inflammatory cytokines, which associate with insulin resistance. Oxidative stress can give rise to ROS formation. Hemoglobin (Hb) is one of the main targets of carbohydrates attack. Hb glycation leads to its structural modification such as heme degradation, the release of iron and free radical generation. The anti-glycation agents thought to be useful as therapeutics for prevention of diabetes inflammation. Aspirin (ASA) is among the first identified glycation inhibitors. However, ASA has its own side effects. In this regard, discovering a new glycation inhibitor without side effects has grown in importance.

Methods: In this study, glycation of 15 μ M bovine serum methemoglobin (MetHb), with 30mM fructose during 20 days of incubation was done. One and half mM ASA and two other rationally related compounds, including benzoic acid (BA) and p-nitrobenzoic acid (NBA) were added in order to analyze the formation of AGEs by determining their absorbance at 340nm as reported previously.

Results: Consequently, absorbance increased with time for all preparations except the native Hb. BA showed the maximum amount of AGEs production, whereas the absorbance of NBA-treated sample was not above ASA-treated sample.

Conclusion: We reported the more preventive effect of NBA against ASA on AGEs formation consequently decreasing heme-loss and ROS production during Hb glycation. No clinically human side-effects have been reported for NBA.

Keywords: Diabetes Inflammation, Free radicals, NF- κ B, Aspirin

ICIA2018\Innate Immunity and Inflammation\Poster\6365

6365

Lack of Association Between rs17568 Polymorphism in OX40 Gene and Myocardial Infarction

Abdolreza Sotoodeh Jahromi¹, Saeideh Erfanian¹, Mehrnoosh Maalagh¹, Mohammad Shojaei¹

1. Research Center for Non-communicable Diseases, Immunology Department, Jahrom University of Medical Sciences, Jahrom, Iran

Background: Tumor necrosis factor (TNF) is one of the inflammatory cytokines which has an important role in inflammation and migration of other inflammatory cells to the atherosclerotic plaques. OX40 is a member of the TNF superfamily receptor protein. OX40 and OX40 ligand are co-stimulators for T-cells and can increase inflammatory response in atherosclerotic plaques. The aim of this study was to determine the association of rs17568 polymorphism in OX40 gene with premature myocardial infarction.

Methods: This case control study was done on 100 patients with premature acute myocardial infarction (AMI) and a similar number of sex, age and some other cardiovascular risk factor matched healthy people. The OX40 rs17568 polymorphism was genotyped, using PCR-RFLP method.

Results: Allele frequency of rs17568 SNP was lower non-significantly in Premature AMI, compared to healthy subjects (49% vs. 51%). The analysis of rs17568 (A/G) polymorphism showed an odds ratio of 1.127 (95% CI: 0.635-1.999; P= 0.686) for the GG genotype and 5.761 (95% CI: 1.200-27.655; P= 0.029) for the AG genotype, compared to the AA genotype.

Conclusion: The results of this study indicate that the rs17568 SNP of OX40 gene is not associated with premature AMI in the evaluated population.

Keywords: rs17568 Polymorphism, OX40 Gene, Myocardial Infarction

ICIA2018\Innate Immunity and Inflammation\Poster\7378
7378

An Investigation into the Relation of High Sensitive C Reactive Protein (hs-CRP) with Cardiac Diastolic Function in Primary Hypertension

Reza Gheitasi¹, Samiramis Qavam², Mohammad Reza Hafezi Ahmadi*

1. Department of Immunology, School of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

2. Student Research Committee, Hamadan University of Medical Sciences, Hamadan, Iran

3. Department of Cardiology, Ilam University of Medical Sciences, Ilam, Iran

4. Department of Pathology, Ilam University of Medical Sciences, Ilam, Iran. B. Biotechnology and Medical plants research Center, Ilam University of Medical Sciences, Ilam, Iran

Background: Hs-CRP is an acute phase serum protein the synthesis of which is increased during tissue damage, inflammatory and infectious conditions by hepatocytes. High rate plasma levels of CRP indicates the risk of cardiovascular diseases. Diastolic dysfunction is one of the most basic cardiovascular complications in patients with hypertension, which may be diagnosed by screening hs-CRP.

Methods: This case-control study was conducted on 71 patients, diagnosed with primary HTN and between 20-55 years of age who visited the specialized cardiac clinic in the city of Ilam. Through a detailed history, physical examination and laboratory measures, those patients with effective factors in their serum levels of hs-CRP (kidney, liver and inflammatory diseases; infectious diseases; peripheral vascular or cerebrovascular disease; connective tissue disease; malignant tumor; trauma; drugs such as statins and aspirin) and diastolic function (e.g. ischemic cardiac disease; cardiomyopathy and pericardial disease; arrhythmias and valvular disease) were excluded from the study. Serum Hs-CRP were measured through Nephelometry method. Based on tissue Doppler echocardiographic findings, the patients were divided into two groups: with and without diastolic dysfunction. The serum levels of hs-CRP in these patients were also compared.

Results: 31% of the patients were male, and 69% were female with an average age of 45.42 ± 7.37 . 54 patients (76.1%) had diastolic dysfunction. The results revealed that a close relationship existed between hs-CRP in the serum and diastolic dysfunction ($P < 0.05$).

Conclusion: Based on the findings a close relationship existed between hs-CRP in the serum and diastolic dysfunction. The higher the rate of hs-CRP, the more likely the emergence of diastolic dysfunction. Therefore, inflammatory factors especially hs-CRP can be used as a diagnostic and prognostic marker for diastolic dysfunctions.

Keywords: hs-CRP, Biomarker, Diastolic dysfunction, Ilam

ICIA2018\Innate Immunity and Inflammation\Poster\7419

7419

Correlation of Serum hs-CRP and Lipid Profile with Coronary Artery Disease

Fatemeh Khaki Khatibi¹

1. Drug Applied Research Center and Department of Clinical Biochemistry, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Coronary Artery Disease (CAD) is one of the major causes of morbidity and mortality in the world. Atherosclerosis is a cause of CAD. Atherosclerosis leads to the stenosis of the vessels and there are several causes. These factors include: inflammation, lipids and their peroxidation, and other risk factors. The aim of the present study is to evaluate serum levels of hs-CRP and lipid profile and the correlation of these parameters with extent of CAD. Measuring these parameters, can help prevent the progression of the disease.

Methods: In this study, 200 individuals including 160 CAD subjects as patient group and 40 age and sex matched healthy persons as control group were chosen. Patients were angiographically divided into four groups: patients with normal angiography (Non-CAD, n=40), one coronary artery occlusion (n=40), two coronary artery occlusion (n=40) and three coronary artery occlusion (n=40). Control group was selected from healthy persons with no history of other diseases. Hs-CRP serum level was measured by immunoturbidimetric method. Serum lipid levels were measured by standard methods.

Results: There was no significant difference between two groups in terms of age and sex, but a significant difference existed among familial history ($p=0.03$). Serum level of hs-CRP was significantly higher in the patient group than the control group ($p=0.01$). Also Serum levels of cholesterol, TG, HDL and LDL were significantly higher than the control group ($p=0.001$). Furthermore, serum concentration of hs-CRP in double-vessel disease was significantly higher as compare with control and no-vessel groups ($p=0.003$ and $p=0.05$, respectively). A significant correlation was found between some lipid parameters and hs-CRP.

Conclusion: Serum levels of hs-CRP and lipid parameters were increased in CAD patients compared to normal controls. These biochemical parameters may have roles in the progression of atherosclerosis. Timely measurement and control of these parameters in normal range may be helpful in preventing CAD and its development. It can also be used to diagnose and control the treatment of vascular disease.

Keywords: hs-CRP, lipid profile, coronary artery disease, Atherosclerosis

ICIA2018\Innate Immunity and Inflammation\Poster\7431
7431

The associations between serum levels of vitamin D and TGF- β in coronary artery disease patients

Nadereh Naderi^{1,2}, Mahsa Rahimzadeh³, Fereshteh Rasa³, Ebrahim Eftekhari³, Elahe Mansoori³

1. Department of Immunology, Faculty of Medicine, Hormozgan University of Medical Sciences, Bandar Abbas, Iran

2. Food health research center, Hormozgan University of Medical Sciences, Bandar Abbas, Iran

3. Department of Biochemistry, Faculty of Medicine, Hormozgan University of Medical Sciences, Bandar Abbas, Iran

Background: The protective role of vitamin D in atherosclerosis and coronary heart disease is well established but the mechanism has not been fully elucidated. The aim of this study was to investigate the associations between vitamin D status and regulatory T cells (Tregs) inhibitory cytokine, TGF- β 1, in patients suffering from coronary artery disease (CAD).

Methods: Vitamin D, and TGF- β 1 were measured in 81 patients. The patients were divided into single (n= 20), double (n=20) and triple (n=20) vessel disease groups and compared to no vessel disease (n=21).

Results: Vitamin D and TGF- β 1 concentrations in controls (32.4 ± 15.2 , 667.7 ± 427.6 respectively) were significantly higher than CAD patients (18.1 ± 9.8 , 360.4 ± 354.1 respectively, $p < 0.05$). TGF- β 1 was significantly higher in double vessel disease patients (591.9 ± 465.7) compared to those with triple vessel disease (173.1 ± 163.3 , $p < 0.05$). Vitamin D, TGF- β 1 revealed a negative correlation ($r = -0.36$, $r = -0.46$ and $r = -0.024$ respectively) with severity of CAD ($p < 0.05$). TGF- β 1 in vitamin D deficient patients was significantly lower compared to normal serum vitamin D patients (326.6 ± 351.7 pg/mL vs. 754.5 ± 560 pg/mL, $p = 0.036$ respectively).

Conclusion: We found *significant association* between TGF- β 1 and vitamin D deficiency in CAD patients. It seems that the *anti-atherosclerotic* effect of vitamin D is attributed, *at least partly*, to the up-regulation of anti-inflammatory cytokines, TGF- β 1.

Key words: TGF- β , coronary artery disease, vitamin D

ICIA2018\Innate Immunity and Inflammation\Poster\7438
7438

Role of monocyte chemotactic protein-1 in irritable bowel syndrome

Arezoo Gowhari Shabgah^{1,2}, Mojgan Mohammadi^{3*}, Mohammad Mahdi Hayatbakhsh⁴, Saeid Pishgoee⁵, Jalil Tavakol Afshari¹, Hadi Zeidabadi⁵

1. Immunology Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

2. Student research committee, Mashhad University of Medical Sciences, Mashhad, Iran

3. Allergy Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

4. Gastroenterology and Hepatology Research Center, Institute of Basic and Clinical Physiology sciences, Kerman University of Medical Sciences, Kerman, Iran

5. Physiology Research Center, Institute of Neuropharmacology, Kerman University of Medical Sciences, Kerman, Iran.

Background: Irritable bowel syndrome (IBS) is one of the most common functional gastrointestinal disorders worldwide, with the prevalence ranging from 1.1% to 29.2% in the general population diagnosed by the Rome III criteria. IBS causes digestive problems like cramping, abdominal pain, and bloating; and diarrhea (IBS-D), constipation (IBS-C), or both (IBS-A: alternating). Little is known about the immunopathogenesis of these three subtypes of IBS. Monocyte chemotactic protein-1 (MCP-1) is a cytokine and belong to chemokine family. MCP-1 causes infiltration of antigen presenting cells and memory T cells to the sites of inflammation. We aimed to evaluate the serum levels of MCP-1 in patients with IBS and compare it with healthy controls.

Methods: The serum concentration of MCP-1 was assessed by enzyme linked immunosorbent assay (ELISA) in patients with IBS (n=96) and healthy controls (n=44).

Results: Our results showed a significant decrease in serum levels of MCP-1 in patients with IBS as compared to controls (p= 0.001).

Conclusion: As all patients who participated in our study had taken medications for IBS, significant decline in the MCP-1 serum levels in comparison with controls might be due to the conventional therapy. Further investigations on the new case of IBS patients to evaluate the effect of MCP-1 in the immunopathogenesis of IBS is highly recommended.

Keywords: Irritable bowel syndrome, monocyte chemotactic protein-1, inflammation.

ICIA2018\Innate Immunity and Inflammation\Poster\7454
7454

Stimulation of cholinergic anti-inflammatory system by nicotine ameliorates animal model of rheumatoid arthritis (RA)

Sara Golbahari, Seyyed Meysam Abtahi Froushani

Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Background: It has been revealed that the signaling by $\alpha 7$ nicotinic acetylcholine receptor can promote the anti-inflammatory condition. This study was designed to evaluate therapeutic benefit of nicotine on the animal model of rheumatoid arthritis (RA).

Methods: Intra-footpad injection of complete Freund's adjuvant was used to induce RA in Wistar rats. Then, rats were allocated in 3 groups: treated with nicotine (2.5 mg/kg-orally), treated with Prednisolone (2 mg/kg-orally) and un-treated group. All therapies were started at day 5 post-induction. The changes in the dorso-plantar diameter of hands and legs of each rat were monitored every day until the day 23 post-intrafootpad injection of complete Freund's adjuvant.

Results: Nicotine treatment after the occurrence of clinical symptoms significantly reduced the edema and swelling of the soles of the feet of RA rats in a similar manner with prednisolone. The serum levels of myeloperoxidase and nitric oxide concurrent with proliferation of spleen lymphocytes were significantly decreased in treatment groups compared to control rats. Albeit, the level of decrease in the splenocytes proliferation index and serum nitric oxide level was somewhat higher in the prednisolone group compared to the nicotine group.

Conclusion: This pharmacological approach may be as a useful strategy to control RA.

Keywords: Nicotine, Rheumatoid arthritis, Wistar rat.

ICIA2018\Innate Immunity and Inflammation\Poster\7467

7467

Comparison of serum and tissue antioxidant changes in rat model of ulcerative colitis treated with hydroalcoholic extract of Cinnamon

Mojtaba Salamatian¹, Vahid Mohammadi¹, Seyyed Meysam Abtahi Froushani²

1. Department of internal medicine and clinical pathology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

2. Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Background: Cinnamon is a spice obtained from the inner bark of the genus *Cinnamomum*. It has been used for a long time in folk medicine. Here, we investigated and compared serum and tissue antioxidant changes in rat model of ulcerative colitis (UC) treated with hydroalcoholic Cinnamon extract.

Methods: UC was induced in the male Wistar rats by luminal instillation of acetic acid. Animals in the treatment group treated with hydroalcoholic extract of Cinnamon (400 mg/kg PO-daily) for 10 consecutive days. Finally, the rats were sacrificed and the disease activity index, the levels of malondialdehyde (MDA) and total antioxidant capacity (TAC) were measured in the colonic homogenized tissue specimens and sera.

Results: Attained data indicated that hydroalcoholic extract of Cinnamon could reduce the clinical scores and the mortality rate of ulcerative colitis. The levels of TAC were significantly decreased and conversely the levels of MDA were increased in both sera and gut samples. Nevertheless, hydroalcoholic extract of Cinnamon, concurrent with a significant decrease in clinical scores, could regress the levels of MDA and conversely increased the levels of TAC in gut tissues more prominent than sera samples.

Conclusion: It seems that the changes in the antioxidant levels of the tissue samples is a more suitable criterion for the study of beneficial effects of a therapeutic agent compared to study of these change in sera.

Keywords: Hydroalcoholic extract of Cinnamon, Ulcerative colitis, Antioxidant.

ICIA2018\Innate Immunity and Inflammation\Poster\7470
7470

A lower mRNA expression level of TLR2, 3, 4 in PBMCs isolated from non-responder children to HBV vaccine

Kousar Smailneja ganji ^{1,2}, Mehdi Shahbazi ², Mousa Mohammadnia-Afrouzi ²,
Mojdeh Ghias Tabari³, Motahareh Ehsani², Majgan Bagherzadeh²

1. Student research committee, Babol University of medical sciences, Babol, Iran
2. Department of Immunology, School of Medicine, Babol University of Medical Sciences, Babol, Iran
3. Department of Biochemistry, School of Medicine, Babol University of Medical Sciences, Babol, Iran

Background: Toll like receptors (TLR) have an important role in the innate immune system and also able to recognize Hepatitis B virus that effect on immune response. Many host related factors such as the expression of TLRs, contribute to vaccine response. This study was designed to determine the expression levels of TLR2, 3, 4 genes as indicator of innate immunity response to the hepatitis B vaccine among 3-5 years old children.

Methods: The study was carried out among 60 children with age 3-5 years old who received HBV vaccine (43 responders, 17 non responders). Blood samples were collected, then the expression of TLR2, 3, 4 genes in peripheral blood mononuclear cells (PBMCs) were analyzed by Real Time PCR method.

Results: The results of this study indicated that the average expression of TLR2, 3, 4 genes in the responder group was higher than the non-responder group. Additionally, the expression levels of TLR 3, 4 genes were significantly higher in responders compared to non-responders (P-Value ≤ 0.01).

Conclusion: This study suggest that the expression of TLR2, 3, 4 could be considered as a marker of protection along with serological markers. Indeed, evaluation of the HBs antibody titer and the expression level of immune response genes together can also be used to determine efficiency of hepatitis B vaccine in responder and non-responder individuals.

Keywords: Hepatitis B vaccine, TLR2, 3, 4 genes and Real Time PCR

ICIA2018\Innate Immunity and Inflammation\Poster\7499

7499

The role of CCR1 and different monocyte subsets in progression of coronary atherosclerotic lesion

Saeid Taghiloo¹, Abolghasem Ajami^{1,2*}, Hossein Asgarian Omran^{1,3}, Hamideh Rahimi¹, Reza Alizadeh Navaei², Seyed Mohammad Mehdi Peyghambari⁴, Mahin Nikogoftar⁵

1. Department of Immunology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

2. Molecular and Cell-Biology Research Center, Mazandaran University of Medical Sciences, Sari, Iran

3. Immunogenetics Research Center, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

4. Department of Cardiology, School of Medicine, Tehran University of Medical Sciences, Sari, Iran

5. Department of Research Center of Iranian Blood Transfusion Organization Tehran, Iran

Background: The inflammation underlying atherosclerosis is strongly related to monocyte action. Circulating monocyte divide into subpopulation, according to CD14 and CD16 expression. In this study, we investigated the relationship between the severity of coronary artery disease (CAD) and the two distinct monocyte subsets and CCR1 receptor.

Methods: We enrolled 88 patients who underwent diagnostic CAG (coronary angiography). Patients were divided into four groups. Those without CAD (normal group), those with 50% stenosis in one vessel (MVD), those with single vessel stenosis (1VD) and those with 2 or 3 vessels stenosis (2+3 VD). In addition, the severity of CAD was evaluated by Gensini score. The two monocyte subsets: classical (CD14⁺CD16⁻) and non-classical (CD14⁺CD16⁺) were measured by flow cytometry. Mean fluorescence intensity (MFI) for CCR1 marker was measured and compared between monocyte subset in different group of patients.

Results: Circulating non-classical monocytes were more frequently observed in patients with 2+3 VD than normal, minimal and 1VD groups, but this increase is not statistically significant ($P > 0.05$). According to the results of the CCR1, both classical and non-classical monocytes were surface expressed, but a larger number of classical monocytes expressed this molecule. The number of non-classical monocytes that express the CCR1 molecule decreased with CAD, but not statistically significant.

Conclusions: This study showed that non-classical monocytes increased with increase CAD, and CCR1 is not associated with atherosclerosis, because of this molecule decreased at the level of non-classical monocytes.

Keywords: monocyte, CCR1, coronary artery disease, cardiovascular diseases

ICIA2018\Innate Immunity and Inflammation\Poster\7562
7562

Anti-APOH IgG antibody in diabetic patients with atherosclerosis

Mohammad Rafiee¹, Hossein Golmoghaddam¹, Shahdad Khosropanah², Mehrnoosh Doroudchi¹

1. *Memory T-cell Laboratory, Department of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran*

2. *Cardiovascular Research Center, Shiraz University of Medical Sciences, Shiraz, Iran*

Background: Self-proteins such as Apolipoprotein-H (APOH), Heat Shock Proteins, and Apolipoprotein-B100 may be involved in atherosclerosis. APOH is shown to be associated to vascular alterations in type-2 diabetes; however, the information regarding its antigenicity in these disorders is scarce. In this study, we assessed the anti-APOH-antibody levels in patients with atherosclerosis.

Methods: In total, 134 participants who underwent coronary artery angiography for diagnostic purposes were included in this study. Thirty-one subjects had normal/insignificant coronary artery disease (group 1). Twenty-eight subjects had diabetes type-2 but normal/insignificant coronary artery disease (group 2). Thirty-nine subjects had no diabetes but showed $\geq 50\%$ stenosis in one or more coronary arteries (group 3) and thirty-six subjects had diabetes type-2 and showed $\geq 50\%$ stenosis in one or more coronary arteries (group 4). Blood samples were obtained from all 134 individuals and IgG class anti-APOH-Ab was measured by a commercial ELISA in the sera.

Results: In total, 11 (8.2%) of subjects showed IgG anti-APOH-Ab in their sera. The frequency of anti-APOH Ab+ samples in group 4 was the highest of all (16.7%). In group 2, 7.1% and in group 3, 5.1% of subjects had anti-APOH-Ab in their sera. The least frequency of anti-APOH-Ab+ samples was found in group 1 in which only one subject was positive (3.2%). Statistical analysis between the 4 groups did not reach the significant level by Chi-square test ($p=0.17$) or by Kruskal-Wallis test ($p=0.18$). However, the diabetic patients were more likely to have IgG anti-APOH-Ab in their sera ($p=0.08$).

Conclusion: Our results suggest that anti-APOH-Ab is higher in type-2 diabetic patients and increases with the existence of artery stenosis. Therefore, not only APOH protein but also anti-APOH-Ab is produced in diabetic patients. The mechanism by which this self-antigen bypasses immune-tolerance is worth investigating in a larger group of patients.

Keywords: ApoH, Atherosclerosis, Diabetes type 2, IgG

ICIA2018\Innate Immunity and Inflammation\Poster\7591

7591

Serum levels of interleukin (IL)-33 in patients with ischemic heart disease

Morteza Jafarinia¹, Mohsen Khalili, Sina Vakili

Department of Immunology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

Department of Immunology, School of Medicine, Kerman University of Medical Sciences, Kerman, Iran

Department of Biochemistry, Shiraz University of Medical Sciences, School of Medicine, Shiraz, Iran

Background : Objective: It has been reported that cytokines play a crucial role in pathogenesis of different diseases such as cardiovascular diseases. The aim of this study was to evaluate the serum level of IL-33 in patients with ischemic heart disease (IHD) and also clarify its association with traditional risk factors of the disease.

Methods: A total of 400 patients with IHD as having acute myocardial infarction (AMI; n=100), unstable angina (UA; n=100) or stable angina (SA; n=100) and 100 sex- and age- matched healthy subjects as control group were enrolled to this cross-sectional, case controlled study. Serum samples were collected from all participants (for AMI patients at 3-5 days after events and for UA or SA at admission time) and tested for IL-33 by use of ELISA method.

Results: The mean serum concentration of IL-33 in AMI (103.33 ± 19.29 Pg/ml), SA (157.60 ± 33.43 Pg/ml) and UA (122.21 ± 22.26 Pg/ml) was significantly higher than that observed in healthy control (61.85 ± 7.67 Pg/ml). There was no significant difference between patient with or without certain traditional risk factor including hypertensive patients, dyslipidemic patients, smoker, obese patients and diabetic patients in the mean serum level of IL-33.

Conclusion: These results showed that the higher serum concentration of IL-33 were associated with IHD. The presence or absence of certain traditional risk factors of IHD did not influence the serum level of cytokines.

Keywords: ischemic heart disease, acute myocardial infarction, unstable angina, stable angina, interleukin (IL)-33



ICIA2018\Innate Immunity and Inflammation\Poster\7605
7605

Thymol ameliorates acetic-acid induced ulcerative colitis via selective COX-2 inhibition

Pourya Tahmasebi, Seyyed Meysam Abtahi, Nahideh Afzale Ahangaran

Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Background: Thymol, a natural monoterpene phenol derivative of Thymus, has anti-inflammatory properties. Here, we investigated the potential of Thymol in ameliorating animal model of ulcerative colitis (UC).

Methods: UC was induced in the male Wistar rats by luminal instillation of acetic acid. Rats in the treatment groups treated with Thymol (100 mg/kg PO) or Prednisolone (2 mg/kg PO) daily for 10 consecutive days. At the end, colon tissue specimens were used for Cyclooxygenase-2 expression by immunohistochemistry. Also, the levels of IL-1 β and IL-6 were checked in colonic homogenates.

Results: The data indicated that both therapies with Thymol and Prednisolone could regress the clinical scores and the mortality rate of ulcerative colitis in a comparable manner. The levels of COX-2 was significantly decreased in the guts of Thymol treated rats more than Prednisolone groups. Nevertheless, Thymol somewhat decreased the levels of IL-1 β and IL-6 more than Prednisolone.

Conclusion: Thymol can ameliorate acetic-acid induced ulcerative colitis via selective COX-2 inhibition compared to Prednisolone.

Keywords: Thymol, Ulcerative colitis, COX-2.

ICIA2018\Innate Immunity and Inflammation\Poster\7718
7718

Effect of aqueous extract and saponin fraction of *Tribulus terrestris* L. on the expression of *ICAM1* and *VCAM1* genes in the human endothelial cell lines in vitro

Zahra Fereydouni¹, Elahe Amirinezhad Fard¹, Ali Mostafaie¹

1. Department of immunology, Kermanshah University of Medical sciences, Kermanshah

Background: Atherosclerosis is a chronic inflammation during which low density cholesterol and lipid deposition on the wall of arteries are formed. *Tribulus terrestris* L. as a medicinal plant, due to its anti-inflammatory properties, has been used for treatment of various kinds of diseases such as atherosclerosis. These observations motivated us to investigate the effect of aqueous extract and saponin fraction of *T.terrestris* on the expression of Intracellular Adhesion Molecule-1 (*ICAM1*) and Vascular Cell Adhesion Molecule-1(*VCAM1*) genes in the Human Umbilical Vein Endothelial Cell (HUVEC) and Human Bone Marrow Endothelial Cell (HBMEC) lines in vitro during normal and LPS-induced conditions.

Methods: After the culture of HUVEC and HBMEC cell lines and induction with LPS, the cells were treated with aqueous extract and a saponin fraction of the *T.terrestris*. Then, the expression of *ICAM1* and *VCAM1* genes under normal and LPS-induced conditions was investigated using Real-Time PCR technique.

Results: The data obtained showed that the aqueous extract and saponin fraction of *T. terrestris* exerted a decreasing effect on mRNA expression of the investigated genes in the LPS-induced HUVEC and HBMEC lines.

Conclusion: As revealed by our study, anti-inflammatory activity of *T. terrestris* could be partly attributable to its potential to downregulate *ICAM1* and *VCAM1*. More reduction in the expression of above-mentioned genes by aqueous extract compared to saponin fraction may be due to the presence of other chemical constituents, such as flavonoids and alkaloids in the aqueous extract. In conclusion, *T.terrestris* could be considered as a candidate for the treatment of atherosclerosis.

Keywords: atherosclerosis, *ICAM-1*, *VCAM-1*, *Tribulus terrestris* L.

ICIA2018\Innate Immunity and Inflammation\Poster\7739
7739

Macrophage function can alter toward anti-inflammatory phenotype after oral administration of Hypiran

Somayeh Ahmadiashar, Mahsa Alem, Seyyed Meysam Abtahi Froushani

Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Background: *Hypericum perforatum* is a medicinal herb with anti-inflammatory properties as an arachidonate 5-lipoxygenase inhibitor and COX-1 inhibitor. Hypiran is a commercial hydro-alcoholic extract of this plant. The current research was conducted to investigate the effects of oral administration of Hypiran on the some functions of the peritoneal macrophages in NMRI mice.

Methods: The study population consisted of 10 NMRI mice randomly allocated into the treatment and control groups. Hypiran was orally administered to treatment group in daily doses of 110 mg/kg from the beginning of the study and continued for 1 week. The control mice received PBS at the same volume. The resident macrophages were isolated from the peritoneal cavity of mice by injecting 20 ml of ice-cold PBS and used for experiments.

Results: The NBT reduction assay test showed that oral administration of Hypiran couldn't induce any change in the potential of reactive oxygen production by of macrophages. Moreover, MTT reduction assay test showed that Hypiran can increase vitality and biological activity of macrophages isolated from Hypiran treated mice compared to macrophages isolated from untreated mice.

Conclusion: Despite increase in biological activity of macrophages, potential harmful reactive oxygen substances production by macrophages couldn't showed any significant changes. Therefore, oral administration of Hypiran can alter the phenotype of macrophages toward anti- inflammatory phenotype.

Keywords: Macrophages, Hypiran, *Hypericum perforatum*.

ICIA2018\Innate Immunity and Inflammation\Poster\7862

7862

Evaluation of interleukin-1 beta (IL-1 β), before and after curcumin treatment in HAM/TSP patients and HTLV-1 carriers

Zahra Soleimani¹, Houshang Rafatpanah^{2,*}, Reza Boostani³, Reza Farid Hosseini⁴

1. *Inflammation and Inflammatory Diseases Research Center, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran*

2. *Inflammation and Inflammatory Diseases Research Center, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran*

3. *Department of Neurology, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran*

4. *Allergy Research Center, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran*

Background: Human T-cell leukemia virus (HTLV-1) is the first human lymphotropic retrovirus which is associated with adult T cell leukemia (ATL) and HTLV-1-associated myelopathy/tropical spastic paraparesis (HAM/TSP), a chronic neuroinflammatory disease that lead to destruction of central nervous system. Some of the proinflammatory cytokines such as IL-1 β , IL-6, IL-8 and TNF- α play important roles in the diseases progression. Curcumin, a bioactive polyphenol from turmeric, has wide range of biological activities including anti-inflammatory, anti-oxidant and anti-tumor properties. Curcumin mediates its anti-inflammatory effects through the downregulation of inflammatory transcription factors, protein kinases, enzymes and cytokines such as IL-1, -2, -6, -8, -12 and TNF- α . Thus the aim of this study was to evaluate the serum levels of IL-1 β , before and after curcumin treatment in HAM/TSP patients. Furthermore we compared the levels of this cytokine between HAM/TSP patients and HTLV-1 asymptomatic carriers (ACs).

Methods: The Serum levels of IL-1 β was measured by ELISA before and after curcumin treatment in 21 HAM/TSP patients. Furthermore, the Serum levels of this cytokine was also measured in 21 ACs.

Results: The serum levels of IL-1 β was lower after treatment with curcumin in HAM/TSP patients. The serum levels of IL-1 β was significantly higher in HAM/TSP patients compared with ACs.

Conclusion: The results of the present study showed that curcumin reduces the levels of IL-1 β and could be useful as a supplement for treatment of HAM/TSP patients.

Keywords: HTLV-1, HAM/TSP, Curcumin, IL-1 β , Proinflammatory cytokines

ICIA2018\Innate Immunity and Inflammation\Poster\7902
7902

Dopamine modulates inflammatory functions of the monocytes.

Hanieh Nikkhah, Seyyed Meysam Abtahi Froushani, Habbib Dastmalchi Saei, Nahideh Afzale Ahangaran

Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Background: Dopamine, produced in many cell types like immune cells, has autocrine and paracrine effects. The current research was designed to evaluate the effects of dopamine on the inflammatory functions of the monocytes.

Methods: The peripheral blood cells were collected by a Ficoll-Hypaque density gradient from the blood of Wistar rats. Then, non-adherent plastic cells were removed after 2 h. Monocytes pulsed with dopamine for 24h within the concentration of 5×10^{-7} M (maximum physiologic concentration).

Results: The mRNA level of NF- κ B p65 in monocytes treated with dopamine was reduced compared macrophages without treatments. MTT assay showed that dopamine treatment does not significantly change the viability of monocytes. Nevertheless, the potential of nitric oxide and reactive oxygen species production of monocytes stimulated with opsonized yeast were regressed after dopamine exposure.

Conclusion: Dopamine treatment of monocytes can regress the inflammatory function of monocyte without altering any changes in the vitality of monocytes.

Keywords: Monocytes, Dopamine, anti-inflammatory.

ICIA2018\Innate Immunity and Inflammation\Poster\7903

7903

The effects of dopamine on the respiratory burst and nitric oxide production by macrophages.

Hanieh Nikkhah, Seyyed Meysam Abtahi Froushani

Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Background: Dopamine is a neurotransmitter in both central and peripheral nervous system. Nowadays, its role in the neural-immune communication has been revealed. The current research was set out to investigate the effects of dopamine on the inflammatory functions of the peritoneal macrophages.

Methods: The resident macrophages were isolated from the peritoneal cavity of Wistar rats by injecting 20 ml of ice-cold PBS and pulsed with the maximum physiologic concentration of dopamine (5×10^{-7} M) for 24 h.

Results: MTT reduction assay test showed that dopamine couldn't alter the vitality of macrophages. Phagocytosis of opsonized yeast also couldn't show any changes after dopamine treatment of macrophages. Nevertheless, concurrent with the killing activity of macrophages after challenge with opsonized yeast, the potential of nitric oxide and reactive oxygen production by these cells were significantly reduced after treatment with dopamine.

Conclusion: Dopamine exposure of macrophages can reduce the inflammatory potential of macrophages without altering any changes in the vitality of these cells.

Keywords: Macrophages, Dopamine, Respiratory burst, Nitric oxide.

ICIA2018\Innate Immunity and Inflammation\Poster\8109

8109

Curcumin can suppress experimental autoimmune encephalomyelitis via ameliorating oxidative stressLaleh Mavaddatiyan¹, Shiva Khezri¹, S.Meysam Abtahi frushani²

1. Department of Biology, Faculty of Sciences, Urmia University, Urmia, Iran

2. Department of Microbiology, Faculty of Veterinary, Urmia University, Urmia, Iran

Background: Despite the advances made in recent years in controlling multiple sclerosis, many patients respond less well to existing treatments. Previous studies have pointed to the role of oxidative stress in the pathogenesis of multiple sclerosis. The aim of this study was to investigate the effects of curcumin on reducing the effects of oxidative stress in the animal model of multiple sclerosis.

methods: In this experimental study, Autoimmune encephalomyelitis was induced by immunization of wistar rats (12 vertebrae) with homogenized spinal cord of Indian guinea pigs and Compound Freund's Adjuvant. Then, the animals were safely placed in two groups of 6 heifers. Six vertebrae were considered as healthy controls. Treatment with curcumin (10 mg / kg daily) from the onset of clinical symptoms in the treatment group (day 12). At the same time, the control group received only the drug solvent. Treatment continued until 30 days after immunization.

Results: Administration of curcumin in rats with multiple sclerosis significantly reduced nitric oxide levels and serum myeloperoxidase activity. Both malondialdehyde and uric acid levels were reduced at the same time in the serum of affected and treated animals. However, there was no significant change in total antioxidant capacity of the serum.

Conclusion: Adding curcumin to a multiple sclerosis regimen may be helpful in reducing the harmful effects of oxidative and direct anti-inflammatory effects.

Keywords: Multiple Sclerosis, Experimental Autoimmune Encephalomyelitis, Curcumin, Oxidative Stress

ICIA2018\Innate Immunity and Inflammation\Poster\8133

8133

Glycyrrhizin instructs M1 pro-inflammatory macrophages.

Sahar etemadi¹, Seyyed Meysam Abtahi Froushani¹, Hadi smaili Govarchin Ghalee¹, negin mohammadpoor¹

1. Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Background: Glycyrrhizin is under preliminary clinical research for its activity against common viruses, like hepatitis C. This study was designed to evaluate the effects of Glycyrrhizin on the polarization of macrophages.

Methods: The resident macrophages were isolated from the peritoneal cavity of Wistar rats by injecting 20 ml of ice-cold PBS and treated with dopamine for 24h in the concentration of 100 µg/mL.

Results: MTT reduction assay and neutral red uptake assay indicated that Glycyrrhizin couldn't alter the vitality of macrophages. Nevertheless, concurrent with the killing activity of macrophages after challenge with opsonized yeast, the potential of nitric oxide and reactive oxygen production by these cells were significantly up-regulated after treatment with Glycyrrhizin.

Conclusion: Glycyrrhizin exposure of macrophages potentiates the inflammatory potential of macrophages without altering any changes in the vitality of these cells. These findings might offer new insight into the potential mechanisms that underlie the immunostimulatory effects of Glycyrrhizin.

Keywords: Macrophages, Glycyrrhizin, Inflammation.

ICIA2018\Innate Immunity and Inflammation\Poster\8527
8527

Trimethylamine-N-oxide induce the expression of the TLR2 and NADPH oxidase in U937-derived macrophages

Zakaria Vahabzadeh¹, Abbas Ahmadi², Leila Farhadi², Mohammadraman Molodi³, Sara Shirahmadi¹

1. Department of Clinical Biochemistry, Faculty of Medicine, Kurdistan University of Medical Science

2. Department of Molecular Medicine and Genetics, Faculty of Medicine, Kurdistan University of Medical Science

3. Department of Medical Physiology and Pharmacology, Faculty of Medicine, Kurdistan University of Medical Science

Background: The exact mechanism by which gut microbiota-dependent trimethylamine-N-oxide (TMAO) induce inflammation and bring about atherosclerosis is not well understood. Engagement of toll-like receptors (TLRs) on macrophages and consequently its downstream NADPH oxidases (NOX) activates the signaling pathways that initiate pro-inflammatory cytokines as well as ROS production. The present study was designed to evaluate the expression of TLR2, TLR6 and NOX2 at mRNA levels in macrophages.

Methods: U937-derived macrophages were treated with different concentration (37.5, 75, 150 and 300 μ M) of TMAO for 24 h. The cells were also treated with tunicamycin, as a positive control for stress. RT-qPCR was used to evaluate the expression of TLR2, TLR6, and NOX2 at mRNA levels.

Results: Unlike tunicamycin, high dose of TMAO significantly increased TLR2 and NOX2 mRNA levels compared to the control cells ($P < 0.05$). Tunicamycin significantly increased the mRNA levels of TLR6 ($p = 0.010$).

Conclusion: Our results provide findings to support the contribution of TLR2 and NOX2 in proatherogenic mechanism of TMAO during the foam cell formation and abnormal activation of macrophages.

Keywords: Toll-like receptors, Atherosclerosis, Trimethylamine-N-Oxide, Macrophages

ICIA2018\Innate Immunity and Inflammation\Poster\8532

8532

The proinflammatory cytokine response of U937-Derived macrophages treated with Trimethylamine-N-oxide

Zakaria Vahabzadeh¹, Mohammad Abdi¹, Leila Farhadi², Amin Ghanivash¹, Javad Hosseini³

1. Department of Clinical Biochemistry, Faculty of Medicine, Kurdistan University of Medical Science

2. Department of Molecular Medicine and Genetics, Faculty of Medicine, Kurdistan University of Medical Science

3. Cellular and Molecular Research Center, Kurdistan University of Medical Sciences

Background: Atherosclerosis is known as an inflammatory disease. Trimethylamine N-oxide (TMAO) as a risk factor, has the potential to trigger or promotes the immune inflammatory reactions in atherosclerosis. The exact mechanism by which TMAO induce inflammation during atherosclerosis is not well understood. The present study was designed to evaluate the expression of IL-1 β , IL-6, and TNF- α at mRNA levels in response to treatment of macrophages with TMAO.

Methods: U937-Derived macrophages were treated with different concentration of TMAO for 24 h. The cells were also treated with tunicamycin as a positive control for stress. RT-qPCR was used to evaluate the expression of IL-1 β , IL-6, and TNF- α at mRNA levels.

Results: although TMAO induced the expression all IL-1 β , IL-6, and TNF- α at mRNA levels, only 300 μ M of TMAO significantly increased the expression of IL-1 β at mRNA level compared to control cells. Tunicamycin increased the mRNA levels of IL-6.

Conclusion: The results of this study provide documentation to support the possible contribution of IL-1 β as a proinflammatory cytokine, among the others, to involve in immune inflammatory reactions induced by TMAO during atherosclerosis.

Keywords: Cytokine, Atherosclerosis, Trimethylamine-N-Oxide, Tunicamycin, Macrophage

ICIA2018\Innate Immunity and Inflammation\Poster\9943
9943

Increased concentrations of chemokines CXCL10 in patients with ischemic heart disease

Mojde Kazemi¹, Abdollah Jafarzadeh^{1*}, Amin Safa³, Hamid Reza Rashidinejad⁴, Maryam Nemati¹,
Mohammad Mehdi Mohammadi¹

1. Department of Immunology, Medical School, Kerman University of Medical Sciences, Kerman, Iran

2. Department of Immunology, Medical School, Rafsanjan University of Medical Sciences, Rafsanjan, Iran

3. Physiology Research Center, Institute of Neuropharmacology, Kerman University of Medical Sciences, Kerman, Iran

Background: Recruitment of leukocytes is one of the earliest events in the pathogenesis of ischemic heart disease (IHD) and chemokines play an important role in the migration of these cells into the inflammation sites. The aim of this study was to evaluate the CXCL10 in patients with IHD.

Methods: A total of 300 patients with IHD as having acute myocardial infarction (AMI; n=100), stable angina (SA; n=100) or unstable angina (UA; n=100) and 100 healthy subjects as a control group were enrolled to study. Serum samples from all participants were tested for the CXCL10 levels by using ELISA.

Results: The mean serum concentrations of CXCL10 in AMI patients (395.97 ± 21.20 pg/mL), SA patients (405.48 ± 27.36 pg/mL) and UA patients (396.69 ± 22.79 pg/mL) were significantly higher than in the healthy group (179.38 ± 8.85 pg/mL, $P < 0.001$). Similarly, the mean serum levels of CXCL10 in total IHD patients (399.38 ± 13.77 pg/mL) were also significantly higher as compared with healthy subjects ($P < 0.001$).

Conclusion: These results showed that the higher levels of CXCL10 were associated with IHD. The serum levels of chemokines may influence by the certain traditional risk factors of IHD.

Key words: Acute myocardial infarction, Stable angina, Unstable angina, Chemokine, CXCL10

ICIA2018\Innate Immunity and Inflammation\Poster\9948
9948

Does HLA-DR influence susceptibility to pulmonary cystic fibrosis?

Esmail mortaz¹, alireza asef², Hossein Ali Ghaffaripour³, Ali Arabkhazali¹

1. Department of Immunology, Faculty of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Young Researchers Club, Rasht Branch, Islamic Azad University, Rasht, Iran

3. Pediatric Respiratory Diseases Research Center, NRITLD, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Cystic fibrosis (CF) is the most common lethal autosomal recessive disease in white Caucasians. It affects many organs including the lung, pancreas and liver. Whilst CF is a monogenic disease, several studies reveal a complex relationship between genotype and clinical phenotype. Objective of study: we examined the expression of HLA class II alleles among Iranian CF patients in relation to disease-related criteria namely infection with *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Candida albicans* and with the serum total IgE level.

Materials and Methods: Our study was conducted on 50 CF patients (27 male, 23 female) hospitalized in the Masih Daneshvari Hospital in Tehran and 50 healthy age- and gender-matched control subjects from March 2016 to April 2017. Individuals with a family background of CF were excluded. Clinical and demographic data including lung function and comorbidities such as diabetes and gallstones were evaluated. HLA-DRB1 subtypes were determined by Single Specific Primer Polymerase Chain Reaction (SSP-PCR).

Results: HLA-DRB1*09 and HLA-DRB1*10 were less frequent and HLA-DRB1*13 and HLA-DRB1*11 were the most frequent in CF patients compared with healthy controls although this did not reach significance. 16 CF patients had high serum IgE levels (430.25 ± 219.7 IU/ml) which was most closely associated with the HLA-DRB1*04 allele. Twenty seven CF patients were *P. aeruginosa* positive with the HLA-DRB1*04 being the most frequent allele. Thirty one CF patients were *candida albicans* colonized with HLA-DRB1*11 as the most frequent allele. The small numbers of patients with allergic bronchopulmonary aspergillosis (ABPA, n=3) or with diabetes (n=2) prevented further exploration of their data.

Discussion: Our data suggests that the DR4 and DR11 serotypes are the most prominent in the Iranian CF population and further research on should be conducted on DR4+ CF patients to understand the role in clinical phenotypes of CF.

Key words: HLA-DRB1, SSP-PCR, lung cystic fibrosis

ICIA2018\Innate Immunity and Inflammation\Poster\10009

10009

Nicotine and Caffeine Alter the Effects of the Pro-Inflammatory Mesenchymal Stem Cells on the Co-Cultured Neutrophils FunctionRahmani Kukia Nasim¹, Abbasi Ardeshir ^{2*}, Abtahi Froushani Seyyed Meysam³, Hashemi Seyyed Mahmoud⁴*1. Division of Immunology, Department of Microbiology, Veterinary Faculty, Urmia University, Urmia, Iran**2. Division of Immunology, Tarbiat Modares, University of Medical Sciences, Tehran, Iran**3. Department of Microbiology, Veterinary Faculty, Urmia University, Urmia, Iran**4. Department of Immunology, Shahid Beheshti University of Medical Sciences, Tehran, Iran*

Background: Mesenchymal stem cells (MSCs) express some of the nicotinic receptor subunits and adenosine receptors. Recent studies indicate that polarized MSC1 mostly elaborate pro-inflammatory mediators. The communication between pro-inflammatory MSCs with neutrophils has not been yet studied. Therefore, the main desire is to determine the role of nicotine or caffeine on MSC1s and its effects on neutrophils.

Methods: After isolation, characterization and differentiation of MSCs into osteocyte and adipocyte, they were pulsed with LPS (10 ng/ml) for 1 hour. MSCs were incubated with different concentrations of caffeine (0.1, 0.5 and 1 mM) and with different concentrations of nicotine (5, 25 and 50 μ l) separately for 48 hours. Then the medium was aspirated and cells were used for co-culture experiment with neutrophil and the behavior of neutrophils was studied.

Results: Attained data indicated that when pro-inflammatory MSCs treated with nicotine or caffeine, increased significantly the vitality of co-cultured neutrophils. The rate of the respiratory burst of neutrophils after co-culture MSC1s, didn't show any significant change after challenging with non-opsonized yeast in non-treated groups. Nicotine or caffeine treatment, at a high dose, could increase the respiratory burst of neutrophils after challenging with non-opsonized yeast. Interestingly, after challenging with opsonized yeast, the rate of the neutrophil respiratory burst was decreased after co-culturing with pro-inflammatory MSCs comparing the respiratory burst of neutrophil alone. Nicotine and/or caffeine treatment could reverse this reduction.

Conclusion: Generally, these findings provide a new insight into understanding the anti-inflammatory and immunomodulatory effects of nicotine and caffeine.

Keywords: Mesenchymal stem cells, Caffeine, Nicotine, Neutrophil

ICIA2018\Innate Immunity and Inflammation\Poster\11068
11068

Investigating level of hs-CRP in steatohepatitis patients

zivar zangeneh

Isfahan University of medical sciences, Isfahan, Iran

Background: Non-alcoholic fatty liver disease (NAFLD) is characterized by triglyceride accumulation in hepatocytes, which occurs without alcohol abuse. It is one of the most commonly encountered chronic liver diseases. The clinical spectrum of NAFLD ranges from asymptomatic steatosis to steatohepatitis, fibrosis and cirrhosis. Most NAFLD patients have asymptomatic simple steatosis without adverse sequelae. Recently, studies have reported that NAFLD is an independent risk factor of numerous diseases, including diabetes, cardiovascular disease, hypertension, kidney disease and colon cancer. Therefore, interest is growing with regard to NAFLD. C-reactive protein (CRP) is a biomarker of inflammation. Plasma CRP concentrations increase rapidly and dramatically (100-fold or more) in response to tissue injury or inflammation. High-sensitivity CRP (hs-CRP) is more precise than standard CRP when measuring baseline (ie, normal) concentrations and enables a measure of chronic inflammation. The aim of this study is to investigate relationship between CRP and fatty liver diseases.

Methods: 113 patient with steatohepatitis were determined based on the results of abdominal ultrasonography grade I and II. Hs-CRP was measured by kit and auto-analyzer and TG and cholesterol levels were determined. Statistical analysis was done by prism graph-pad.

Results: Seventy eight patients had hs-CRP level average of more than 3mg/dl with grade II. But others had an average level of less than 3mg/dl. Seventy eight patients had higher levels of TG and cholesterol.

Conclusion: Hs-CRP is suggested as a biomarker for diagnosis and also prognosis of steatohepatitis.

Keywords: hs-CRP, steatohepatitis



Monoclonal Antibody Diagnostic and Therapeutic

Poster Presentation

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\ Monoclonal Antibody Diagnostic and Therapeutic\ Poster\ 8583
8583

Expression and purification of Recombinant scFv antibodies against EpEX in *Escherichia coli* Rosetta™(DE3) Rosetta-Gami™, Origami™ and SHuffle™

Mohammadi, S.A ¹, Hashemi, A¹, keramati, M²

1. Pharmaceutical Biotechnology Department, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Pilot of nanobiotechnology Department, Pasteur institute of IRAN, Tehran, Iran

Background: Overexpression of some cell surface antigens in malignancies make them good targets for active and passive immunotherapy of cancers. EpCAM (epithelial cell adhesion molecule) is an antigen that over expressed in many types of epithelial cell cancers. Recently, several monoclonal antibodies have been developed to directly interfere with EpCAM extracellular domain (EpEX) antigen. Single-chain variable fragment (scFv) is a class of engineered antibodies generated by the fusion of the heavy (VH) and light chains (VL) of immunoglobulins through a short polypeptide linker. Now, scFvs have been developed as potential diagnostic and/or therapeutic agents. *Escherichia coli* (*E.coli*) is one of the most widely used hosts in recombinant proteins production.

Methods: This report describes the production and purification of a scFv specific for EpEX in periplasm of *E.coli* strains containing Rosetta™(DE3) Rosetta-Gami™, Origami™ and SHuffle™.

Results: The majority of expressed anti-EpEX-scFv protein was produced in soluble form. A Ni-NTA affinity column was used to purify the anti-EpEX scFv protein. The molecular weight of anti-EpEX-scFv protein was estimated to be approximately 30 kDa, as confirmed by SDS-PAGE and western blotting assay. The anti-EpEX-scFv showed near 95 % purity.

Conclusion: Our results revealed that *E.coli* is a well-established host system for the production of antibody fragments such as anti-EpEX-scFv. This novel targeting agent can be used in naked or conjugation forms in the treatment of the wide range of tumor cells with epithelial origin including colon cancer.

Key words: ScFv, EpEX, EpCAM, *E.Coli*



Mucosal and Gastrointestinal Immunology

Poster Presentation

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Mucosal and Gastrointestinal Immunology\Poster\6314
6314

The differences between bovine and human sIgA in binding to gut-associated bacteria

Milad Ameri¹, Vedad Alkhamis²

1. Department of Pathobiology, University of Bu Ali Sina, Hamedan

2. Department of Microbiology University of Shahid Chamran Ahvaz

Background: In our study, we tested the hypothesis that human and bovine sIgA have similar innate binding activity for bacteria

Methods: SIgAs, isolated from human and bovine milk, were incubated with a selection of commensal, pathogenic and probiotic bacteria from gastrointestinal tract (gut). Using flow cytometry, we measured numbers of bacteria binding SIgA and their level of SIgA binding

Results: The percentage of bacteria bound by human and bovine SIgA varied from 30 to 90% depending on bacterial species and strains, but was remarkably consistent between human and bovine SIgA. The level of SIgA binding per bacterial cell was lower for those bacteria that had a higher percentage of SIgA-bound bacteria, and higher for those bacteria that had lower percentage of SIgA-bound bacteria. Overall, human and bovine SIgA interacted with bacteria in a comparable way

Conclusion: The in vitro data have shown that milk-derived hSIgA and bSIgA have a similar propensity to bind a range of gut-associated bacteria. Further studies are required to understand why the level of SIgA binding varies between different bacteria. These data will contribute to longer term research about the potential benefits of bovine SIgA for human consumers

.Keywords: Secretory IgA, Bovine IgA, Human IgA, infection, milk

ICIA2018\Mucosal and Gastrointestinal Immunology\Poster\7459
7459

Association between *helicobacter pylori* infection in Subjects with Gastritis and serum levels of LL-37, MBL and M-Ficolin

Mehdi soltanzadeh Yamchi¹, Mohammad mirzakhani¹, Mehdi shahbazi¹, Mousa Mohammadnia Afrouzi¹, Mojgan Bagherzadeh¹, Javad Shokri Shirvani²

1. Department of Immunology, School of Medicine, Babol University of Medical Sciences, Babol, I.R. Iran

2. Department of Internal Medicine, School of Medicine, Babol University of Medical Sciences, Babol, I.R. Iran

Background: *Helicobacter pylori* (*H. pylori*) can stimulate immune responses and lead to release of pro-inflammatory factors and antimicrobial peptides such as LL-37, MBL and M-Ficolin. The aim of this study was to determine the level of changes in serum levels of the three mentioned factors in people with gastritis and their association with the presence or absence of *Helicobacter Pylori*.

Methods: Study populations were divided in two groups of 35 patients with *H. pylori* and 25 patients without *H. pylori* infection. Biopsy and blood samples were collected from each subject. *H. pylori* positivity was investigated regarding the serum level of anti-*H.pylori* IgG by enzyme-linked immunosorbent assay (ELISA) kit and its presence in the tissue was examined by histopathology observations and rapid urease test (RUT). LL-37, MBL and M-Ficolin serum levels were measured in each group using the standard ELISA kit.

Results: It was revealed that 58% of the subjects were infected with *H. pylori*. Subjects with MBL levels lower than 500 ng/ml in the sera were significantly infected with *H. pylori* and subjects with MBL levels higher than 1000 ng/ml often did not have *H. pylori* infection. LL-37 had an increased level while M-Ficolin showed no significant change in the presence of *H. pylori*.

Conclusion: Our findings indicate that MBL with a lower presence and LL-37 with a higher presence might be involved in *H. pylori* infection while M-ficolin seems to be less effective in the infection.

Keywords: *Helicobacter pylori*, LL-37, MBL, M-Ficolin

ICIA2018\Mucosal and Gastrointestinal Immunology\Poster\7514
7514

Gene expression and serum concentration of Thioredoxin in patients with celiac disease is more than healthy controls despite gluten free diet

Manizhe Faghih¹, Mohammad Rostami Nejad², Davar Amani¹, Elham Aghamohamadi², Amir Sadeghi¹, Mohammad Reza Zali²

1. Department of Immunology, School of medical science, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Gastroenterology and Liver Diseases Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Thioredoxin (TRX) as a ubiquitous redox-active protein is induced in several cells against oxidative stress, as well as increasing in the extracellular matrix in various infectious and autoimmune diseases. Additionally TRX is a highly specific activator of oxidized human extracellular TG2 in small intestinal biopsies, thus it can play a role in the pathogenesis of celiac disease. We investigated the gene expression and serum level of TRX in celiac disease patient's compared to healthy group.

Methods: In this study, duodenal biopsies were collected from 60 confirmed celiac disease patients under gluten-free (between 6 months and 2 years) diet and 60 healthy subjects as control group. RNA was extracted from tissue according to the protocol of the commercial kits, cDNA was synthesis, primer pairs were designed and then TRX gene expression was run by using Real-time PCR technique. Also the serum concentrations of TRX was determined by using ELISA method.

Results: Out of 60 CD patients and 60 controls, with mean age of 38 and in control group with an average age of 35. The result of this study was showed that gene expression and serum concentration of TRX in GFD-patients was more than healthy controls ($P < 0.4$ and < 0.001), respectively.

Conclusion: TRX plays an important role in the onset of tissue damage in celiac disease, and the result of this study confirmed that thioredoxin gene expression and serum level is higher in CD patients than healthy group. According to the results of this study, thioredoxin may interfere with the pathogenesis or process of celiac disease, and further studies are needed to make this clear.

Keywords: Celiac Disease, TRX (Thioredoxin), Gene Expression, serum level

ICIA2018\Mucosal and Gastrointestinal Immunology\Poster\8353
8353

Serum mucosa-associated epithelial chemokine (MEC/CCL28) in COPD: a specific marker for severity

Erfan babaei¹, Ali jalili², Rasoul Nasiri kalmazzi², Naseh Sigari³

1. Student Research Committee, Kurdistan University of Medical Sciences, Sanandaj, Iran

2. Cancer and Immunology Research Center, Kurdistan University of Medical Sciences, Sanandaj, Iran

3. Department of Internal Medicine Faculty of Medicine, Kurdistan University of Medical Sciences, Sanandaj, Iran

Background: Chronic obstructive pulmonary disease (COPD) is a complex inflammatory disease of the lung with systemic reflections. The disease is one of the major health problems around the world that affects 10 to 15 percent of the adult population of 40 and older, this disease is thought to be associated with local and systemic inflammation. However, the exact mechanism of inflammation in this disease has not been fully understood. The chemokine CCL28, known as mucosal epithelial chemokine, is shown to be produced by epithelial cells and chemoattracts inflammatory cells into mucosal tissues.

Methods: Blood sample were taken from 40 patients with COPD and 40 age matched normal individuals. Serum were collected and frozen at -80 °C until use. CCL28 was measured by ELISA as described by manufacturer. All the data were analyzed by SPSS 24.

Results: Our data show that (i) the levels of CCL28 almost equal in both genders, (ii) CCL28 levels are significantly higher in the serum of patients with COPD than the normal group (p value <0.05) and (iii) treated patients showed to have a lower levels of CCL28 than untreated patients. In addition, spirometric findings show that, if the disease is more severe, the concentration of ccl28 is higher.

Conclusion: For the first time we demonstrate that CCL28 Is elevated in serum of patients with COPD and this inflammatory marker could be used as biomarker for diagnosis of COPD.

Keywords: CCL28, COPD, Spirometry

ICIA2018\Mucosal and Gastrointestinal Immunology\Poster\8512
8512

Evaluating of the expression level of CXCL12 and CXCL13 in animal model of Ulcerative Colitis disease

Mohammad amin Boshagh¹, Poorya Foroutan¹, Ali Jalili², Shohreh Fakhari²

1. Student Research Committee, Kurdistan University of Medical Sciences, Sanandaj, Iran

2. Department of Immunology, Faculty of medicine, Kurdistan University of Medical Sciences, Sanandaj, Iran

Background: Ulcerative colitis (UC) and Crohn's disease (CD), two forms of inflammatory bowel disease (IBD) that are accompanied by the uncontrolled activation of effector immune cells in the mucosa. In recent years, experimental studies in rodents have led to a better understanding of the role played by these inflammatory mediators in the development and progression of colitis. Therefore, the aim of this study was to evaluate of the expression level of CXCL12 and CXCL13 in animal model of Ulcerative Colitis disease.

Methods: In this experimental study, 6 male Wistar rats were randomly divided into two groups: control and treated groups. In the treated group, colitis was induced by rectal injection of acetic acid. After 48hours, the animal was anesthetized and then the mucosa samples of its colon were separated in order to consider the histopathological indices and polymerase chain reaction.

Results: Based on PCR results the expression of these chemokine receptors in the treated group has changed in comparison with the control group. CXCL12 and CXCL13 up-regulated in treated group in comparison with the control group.

Conclusion: According to the information obtained from this study, the likely expression of these receptors in the animal model of the ulcerative colitis has changed. Given the fact that this study is part of my thesis, the final confirmation, and publication of exact information require more samples, as well as quantitative techniques such as Real-Time PCR, as well as verification methods such as Immunohistochemistry. Finally, in the current study, we intend to evaluate the expression of the whole CXCLs in this model.

Keywords: Inflammatory bowel disease, Ulcerative colitis, Chemokine, Acetic acid

ICIA2018\Mucosal and Gastrointestinal Immunology\Poster\9825
9825

The Effect of iFlora Probiotic on Hepatic Enzymes Serum Level NAFLD Patients

Zahra Asadi¹, Hossein Ghaedamini², Hamidreza mohajerani¹, Yasaman Alimohammadi¹

1. Islamic Azad University Arak, Arak, Iran

2. Kerman University of Medical Sciences, Kerman, Iran

Background: Today there is evidence of using probiotic products on the improvement of non-alcoholic fatty liver disease (NAFLD). This study was conducted to consider the effect of iFlora probiotics on serum levels of liver enzymes in patients with NAFLD.

Methods: This randomized, double-blind clinical trial was performed on 40 patients (24 women and 16 men) with NAFLD. The intervention group (n=20) used probiotic powder iFlora in dairy products for 8 weeks. The percentage change in body mass index, blood pressure, and serum levels of fasting glucose, triglyceride, total cholesterol (total, LDL, HDL), AST and ALT were collected at the beginning and end of this research. Statistical analysis was performed by SPSS software using Chi-square, independent t-test, covariance analysis and paired t-test.

Results: The results showed that using probiotic iFlora caused a reduction of 3.45% of serum alanine aminotransferase ($p < 0.002$), a reduction of 4.65% in aspartate aminotransferase ($p < 0.001$) and a reduction of 7.8% in total cholesterol ($p < 0.001$) in the intervention group compared with the control group.

Conclusion: Using of probiotic iFlora improves the levels of liver enzymes and total cholesterol in patients and can be effective in treating NAFLD.

Keywords: Probiotics, iFlora, liver enzymes, NAFLD

ICIA2018\Mucosal and Gastrointestinal Immunology\Poster\9928
9928

Comparison of LncRNA13 gene expression in patients with celiac disease

Naghmeh Zamani, Farzaneh Tafvizi, Mohammad Rostami Nejad, Mohammad Reza Zali

1. Basic Department of Biology, Parand Branch, Islamic Azad University, Parand, Iran
2. Basic and Molecular Epidemiology of Gastrointestinal Disorders Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, IR Iran
3. Gastroenterology and Liver Diseases Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Celiac disease has been known to be one of the health concerns with an increasing trend worldwide. This autoimmune disease is due to a reaction to gluten resulting in destroyed intestinal villi. Recent studies have shown that LncRNA13 affects the pathogenicity of the disease. This part of genome does not result in production of protein and were previously known as junk DNA. However, recent studies showed that this group of genes can affect the inflammation due to the disease. We aimed to evaluate the expression of LncRNA13 in a group of celiac patients.

Materials and Methods: In a case-control study, LncRNA13 was evaluated in a total of 50 celiac patients and 50 healthy controls. To evaluate the expression of target gene, qRT-PCR method was applied.

Results: No significant difference was detected between the LncRNA13 expression between the cases and controls ($P=0.27$). No such significant difference was detected between males and females, either ($P=0.99$). In the celiac patients with positive and negative IgA tTg, LncRNA13 expression was significantly different ($P=0.03$)

Conclusions: LncRNA13 expression is recommended to be evaluated in celiac patients to determine the prognosis and treatment of choice. Evaluation of LncRNA13 expression is recommended in other gastrointestinal tract diseases.

Keywords: LncRNA13, gene expression, celiac disease.

ICIA2018\Mucosal and Gastrointestinal Immunology\Poster\9973
9973

Investigation of Caspase 3 gene expression pattern in HepG2 cell line under Graphene Oxide exposure

Mahsa Saeedi Niasar^{1&2}, Seyed Reza Mohebbi³, Mehrangiz Dezfoulan², Mohammad Reza Zali³

1. Basic and Molecular Epidemiology of Gastrointestinal Disorders Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Department of Biology, Faculty of Science, Karaj Branch of Islamic Azad University, Karaj, Iran

3. Gastroenterology and Liver Diseases Research Center, Research Institute for Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Graphene based nanomaterials have been widely used as carrier for drug delivery, scaffold for tissue engineering purposes and clinical diagnosis during recent years due to their unique and attractive physical and chemical properties. However, evaluating the safety of these materials is necessary and a matter of great concern. The present thesis aimed to evaluate the possible cytotoxicity of graphene oxide nanomaterial on the HepG2 cell line.

Methods: Different concentrations of graphene oxide including 1, 5, 10, 20, 50, 100, 200, 400 µg/ml have been utilized to treat the HepG2 cells as sample treated cells and compared to untreated cells. Cytotoxicity of these groups was determined by XTT assay. In addition, Caspase 3 gene expression was evaluated on different time 4 and 24 hours after cell exposure to graphene oxide using real time PCR technique.

Results: The results showed decreasing levels of cell viability in higher doses of graphene oxide 100µg/ml and more (P value=0.004). Inductions of apoptosis pathway and increasing of caspase 3 mRNA levels were also observed expression of caspase 3 was increased (RQ=3.921).

Conclusion: Based on the present results, graphene oxide at the low concentrations is a safe nanomaterial for cells with hepatocyte origin nevertheless its probable risk of graphene oxide in high concentrations should be considered prior to biomedical applications.

Keywords: Graphene oxide, cytotoxicity, Apoptosis, Cell viability



14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

Nanoimmunology

Poster Presentation

ICIA2018\Nanoimmunology\Poster\5282
5282

Nanoencapsulation of modified protective antigen in PLA-PEG nanoparticles

Seyed Masih E'temad aubi¹, Hosein Honari¹, Hamed Bagheri², Mojtaba Nofeli³

1. Department of Biology, Imam Hosein University, Tehran, Iran

2. School of Technology, Tarbiat Modarres University, Tehran, Iran

3. Department of Human Vaccine and Serum, Razi Vaccine and Serum Research Institute, Karaj, Iran

Background: The *Bacillus anthracis* causes anthrax lethal disease. Traditional Anthrax vaccines require multiple injection doses due to their low stability to be fully effective. Polylactic acid (PLA) has been approved as a biodegradable and biocompatible polymer for drugs and vaccines delivery applications. The aim of this study is to encapsulate modified *Bacillus anthracis* protective antigen (mPA63) by PEG-PLA copolymers and characterizing the synthesized nanoparticles.

Methods: After expression of the PA63 in *E. coli*, the produced antigen was purified by affinity chromatography and the encapsulation in PLA-PEG copolymer was done using double emulsion solvent evaporation method. The morphological characteristics of produced nanoparticles were evaluated by scanning electron microscopy and particle size, polydispersity index and Zeta potential were measured using dynamic light scattering.

Results: The results of this study showed that the size of free protein-PLA-PEG nanoparticles and the protein-loaded nanoparticles were 162 nm and 233 nm, respectively and the polydispersity index for above mentioned particles were 0.034 and 0.340, respectively. Also, electron microscopic images showed spherical shape of nanoparticles. The Zeta potential of PLA-PEG nanoparticles was -32.2 and PLA-PEG-PA63 nanoparticles have an average peak at -10.8mV. The encapsulation efficiency of the PA63 recombinant antigen in nanoparticles was 34%.

Conclusion: From the result of this study it was showed that PLA-PEG copolymer has an appropriate ability to encapsulate the mPA63 and this system can be recommended as a candidate for an effective vaccine carrier in future investigation.

Keywords: Anthrax vaccine, Recombinant antigen, Nano-carrier, PEG-PLA copolymer.

ICIA2018\Nanoimmunology\Poster\7721

7721

Targeting P68 and STAT3 molecules through Nanoparticles loaded with the siRNA in 4T1 and CT26 cancerous cells lines

Vida Hashemi^{1,2,3}, Ali Masjedi^{2,3}, Afshin Nikkhoo^{2,3}, Narges Rostami^{2,3}, Shima Bastaki^{4,2}, Farhad Jadidi Niaragh^{2,3}

1. Department of basic science, Faculty of Medicine, Maragheh University of Medical Sciences, Maragheh, Iran

2. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

3. Department of Immunology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

4. Department of Cellular and Molecular biology, Faculty of Basic Sciences, Azarbaijan University, Tabriz, Iran

Background: The ribonucleic acid (RNA) helicase P68 can acts as a coactivator of STAT3 and upregulate STAT3 target genes which are involved in various tumorigenic processes such as cellular survival, proliferation, angiogenesis and invasion. STAT3 is constitutively activated in breast and colon cancers.

Methods: P68- and STAT3- specific siRNA loaded chitosan-lactate (ChLa) nanoparticles (NPs) was produced by ionic gelation technique to suppress the expression of P68 and STAT3 molecule on 4T1 breast and CT26 colon cancerous cells lines, in vitro. All of physicochemical characteristics, serum stability, siRNA loading, siRNA release pattern, and cellular uptake of NPs were investigated. The expression of target genes was evaluated by real-time PCR. Anti-proliferative effect of NPs was assessed by MTT assay. Wound healing assay was performed to assay anti-metastatic effects.

Results: P68 and STAT3-siRNA-loaded ChLa NPs could efficiently suppress the expression of P68 and STAT3. Furthermore, downregulation of target molecules led to apoptosis of cancer cells and was associated with reduced expression of angiogenic and metastatic molecules. Consistently, migration capacity of cancer cells was significantly suppressed after treatment.

Conclusion: Combinatorial suppression of P68 and STAT3 by siRNA-loaded NPs may be considered as a promising therapeutic approach for cancer therapy.

Key words: P68, STAT3, Nanoparticle, siRNA, Cancer Therapy.

ICIA2018\Nanoimmunology\Poster\7801
7801

Downregulation of PD-L1 and STAT3 in 4T1 breast and B16/F10 skin cancer cells through siRNA-loaded chitosan-lactate nanoparticles

Shima Bastaki^{1,2}, Narges Rostami^{2,3}, Afshin Nikkhoo^{2,3}, Ali Masjedi^{2,3}, Vida Hashemi^{7,2,3}, Fatemeh Atyabi^{5,6}, Ali Rastegari⁵, Masoomah Baghaei⁶, Farhad Jadidi Niaragh^{2,3,4}

1. Department of Cellular and Molecular Biology, Faculty of Basic Sciences, Azarbaijan Shahid Madani University, Tabriz, Iran

2. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

3. Department of Immunology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

4. Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

5. Department of Pharmaceutical Nanotechnology, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran 1714614411, Iran

6. Nanotechnology Research Center, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran

7. Department of Basic Science, Faculty of Medicine, Maragheh University of Medical Sciences, Maragheh, Iran

Background: The immunosuppressive factors in tumor microenvironment play major role in tumor progression. One of the most critical factors is the programmed death-ligand1 (PD-L1), whose expression on tumor cells has been reported. The activation of STAT3 signaling is significantly associated with expression of PD-L1 in tumor cells, and blocking STAT3 signaling inhibits the PD-1/PD-L1 axis.

Methods: PD-L1 and STAT3-siRNA were encapsulated into chitosan-lactate (ChLa) nanoparticles (NPs) via ionic gelation strategy and were employed to decrease the expression of PD-L1 and STAT3 on 4T1 and B16/F10 tumor cells, *in vitro*. All of physicochemical characteristics, serum stability, siRNA loading efficiency, siRNA release pattern, and cellular uptake of NPs were assessed. The expression of target genes was evaluated by real-time PCR. Anti-proliferative effect of nanoformulation was assessed by MTT assay. Wound healing assay was performed to assay anti-metastatic effects.

Results: Generated NPs had about 100 nm size with a polydispersive index below 0.3 and a zeta potential about 12. Synthesized NPs had high siRNA loading, high serum stability, and efficient cellular uptake. Moreover, NPs had low toxicity during 72 hr cell culture. In addition, siRNA-loaded NPs significantly suppressed expression of PD-L1 and STAT3 in cancer cells. Downregulation of these molecules was associated with reduction of anti-apoptotic proteins, angiogenic, and metastatic molecules. Suppression of PD-L1 and STAT3 also led to decreased migration of cancer cells.

Conclusion: PD-L1 and STAT3 siRNA-loaded ChLa NPs can be considered as a promising therapeutic tool for cancer therapy; however it requires further *in vivo* investigations.

Key words: PD-L1, STAT3, cancer immunotherapy, chitosan-lactate

ICIA2018\Nanoimmunology\Poster\7803

7803

Silencing S1PR1 and GP130 in 4T1 breast cancer cells and B16F10 melanoma cancer cells through siRNA-loaded trimethyl chitosan nanoparticles

Narges Rostami^{1,2}, Farhad Jadidi iaragh^{1,2,3}, Afshin Nikkhoo^{1,2}, Ali Masjedi^{1,2}, Fatemeh Atyabi^{4,5}, Ali Rastegari⁴, Masoomeh Baghaei⁵, Vida Hashemi^{6,1,2}, Shima Bastaki^{7,1}

1. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran.

2. Department of Immunology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

3. Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

4. Department of Pharmaceutical Nanotechnology, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran 1714614411, Iran

5. Nanotechnology Research Center, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran

6. Department of Basic Sciences, Faculty of Medicine, Maragheh University of Medical Sciences, Maragheh, Iran

7. Department of Cellular and Molecular Biology, Faculty of Basic Sciences, Azarbaijan Shahid Madani University, Tabriz, Iran

Background: There are several stimulating molecules in tumor microenvironment which enhance tumor growth. Among them, S1PR1 promotes cancer cell survival, invasion, metastasis and radio/chemo-resistance in various cancers via JAK/STAT3 and other signaling pathways. Glycoprotein130 is a shared subunit of IL-6 receptor which mediates tumor progression following ligation with IL-6 on tumor cells. IL-6 via its receptor, gp130, triggers STAT3 stimulation. There is an amplification loop between S1PR1 and IL-6 for stimulation of STAT3 signaling in tumor cells. Therefore combinatorial blockage of these molecules can be considered as an effective immunotherapeutic approach for cancer therapy.

Methods: S1PR1-siRNA and gp130-siRNA were encapsulated into trimethyl chitosan (TMC) nanoparticles (NPs) were generated through ionic gelation of TMC by hyaluronic acid and employed to suppress the expression of S1PR1 and gp130 molecules on 4T1 and B16F10 cancer cells, *in vitro*. All of physicochemical characteristics, serum stability, siRNA loading efficiency, siRNA release pattern, and cellular uptake of NPs were assessed. The expression of target genes was evaluated by real-time PCR. Anti-proliferative effect of nanoformulation was assessed by MTT assay. Wound healing assay was performed to assay anti-metastatic effects.

Results: SiRNA-loaded NPs had about 100 nm size with a polydispersive index below 0.3 and a zeta potential about 15. Generated NPs exhibited high siRNA loading capacity, high serum stability, and efficiently uptaken by cancer cells. The NPs demonstrated low toxic effect during 72 hr cell culture. In addition, S1PR1-siRNA and gp130-siRNA-loaded TMC NPs could efficiently suppress the expression of S1PR1 and gp130. Combination treatment of tumor cells also led to downregulation of anti-apoptotic proteins, angiogenic, and metastatic molecules. Suppression of S1PR1 and gp130 was also associated with decreased migration of cancer cells.

Conclusion: Combinatorial treatment of tumor cells with S1PR1-siRNA and gp130-siRNA-loaded TMC NPs may be considered as a promising therapeutic tool for cancer therapy.

Key words: S1PR1, gp130, cancer immunotherapy, Trimethyl chitosan

ICIA2018\Nanoimmunology\Poster\7871

7871

Generation of M918/HIV-1 Nef nanoparticles and its delivery into mammalian cellsBahareh Rostami¹, Azam bolhassani^{2*}, Shiva Irani ¹, Reza Ahangari Cohan³

1. Department of Biology, School of Basic Science, Science and Research Branch, Islamic Azad University, Tehran, Iran

2. Department of Hepatitis and AIDS, Pasteur Institute of Iran, Tehran, Iran

3. Pilot Nano-Biotechnology Department, Pasteur Institute of Iran, Tehran, Iran

Background: Among different HIV-1 proteins, Nef protein plays an important role in the down-regulation of CD4 and MHC class I as a virulence factor for AIDS pathogenesis. Therefore, Nef protein can be considered as a possible attractive target in development of therapeutic HIV vaccine. The most important limitation of protein-based vaccines is their low immunogenicity. Recently, a group of short, highly basic peptides known as cell penetrating peptides (CPPs) were used to carry polypeptides and proteins into cells. In this study, the production of HIV-1 Nef protein was performed in *E. coli* expression system for delivery into a mammalian cell line using a novel cell penetrating peptide, M918, in a non-covalent fashion.

Methods: The cloning of HIV-1 Nef was performed into the cloning site of pET23a expression vector. The *E. coli* Rossetta strain was applied for expression of the recombinant Nef using IPTG inducer. The Nef expression was detected by SDS-PAGE and purified using affinity chromatography. The morphological features of the M918/Nef and TurboFect/Nef complexes were studied by scanning electron microscopy (SEM) and SDS-PAGE. The efficiency of Nef transfection using M918 and TurboFect reagent was evaluated by SDS-PAGE and western blot analysis in HEK-293T cell line.

Results: Our results showed a clear band of ~ 27 kDa for Nef protein in SDS-PAGE and western blot analysis. SEM data confirmed the formation of discrete nanoparticles with a diameter of below 300 nm. The dominant band of ~ 27 kDa was detected in the transfected cells with Nef/M918 at molar ratio of 1:20 using the anti-HIV-1 Nef monoclonal antibody. The corresponding band was not detected in the un-transfected cells and transfected with Nef protein alone indicating that M918 could transfer Nef protein into the cells.

Conclusion: These data suggest that M918 peptide would indicate promising applications for improvement of HIV therapeutic vaccines.

Keywords: HIV, Nef antigen, Protein vaccine, M918

ICIA2018\Nanoimmunology\Poster\8034

8034

Combinatorial Suppression of Interleukin-6 and STAT3 in breast, colon and melanoma cancer cells through siRNA-loaded trimethyl chitosan nanoparticles

Ali Masjedi^{1,2}, Fatemeh Atyabi^{3,4}, Narges Rostami^{1,2}, Afshin Nikkhoo^{1,2}, Vida Hashemi^{1,2,5}, Shima Bastaki^{1,6}, Ali Rastegari³, Masoomeh Baghaei⁴, Farhad Jadidi Niaragh^{1,2,7}

1. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

2. Department of Immunology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

3. Department of Pharmaceutical Nanotechnology, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran 1714614411, Iran

4. Nanotechnology Research Centre, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran

5. Department of basic science, Faculty of Medicine, Maragheh University of Medical Sciences, Maragheh, Iran

6. Department of Cellular and Molecular Biology, Faculty of Basic Sciences, Azarbaijan Shahid Madani University, Tabriz, Iran

7. Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Chronic inflammation in the tumor microenvironment enhances resistance against chemo- and radiotherapy, besides promoting growth. Interleukin-6 (IL-6) secreted by tumor cells is one of the main cytokines in the tumor microenvironment and its overexpression has been reported in most types of tumors. The role of IL-6 and its associated signaling pathway (IL-6/JAK/STAT3) has been proven in cancer growth, progression, metastasis and inducing therapeutic resistance.

Methods: IL-6 and STAT3s directed small interfering RNA (siRNA) molecules encapsulated into trimethyl-chitosan nanoparticles(TMC-NP) via ionic gelation of TMC by Dextran and used to suppress the expression of target molecules on murine 4T1(breast), CT26 (colon) and B16F10(melanoma) cancer cells, in vitro.

Results: Generated NPs exhibited optimum physicochemical characteristics associated with high siRNA loading, high stability and efficient cellular uptake. NPs could markedly suppress expression of target molecules which was associated with apoptosis induction. Moreover, expression of angiogenesis and metastatic related molecules was reduced following treatment. Accordingly, migratory function of cancer cells was significantly decreased after suppression of target molecules.

Conclusion: Combinatory suppression of IL-6 and STAT3 using siRNA-loaded TMC-NPs may be considered as a promising therapeutic approach for cancer therapy.

Keywords: IL-6, STAT3, cancer immunotherapy, Trimethyl chitosan, siRNA

ICIA2018\Nanoimmunology\Poster\8487
8487

The Cytotoxic and Apoptotic Effects of Different Sizes of Copper Oxide Nanoparticles in B92 Cancer Cell Line

Rahmani Kukia Nasim¹, Abbasi Ardeshir^{2*}, Abtahi Froushani Seyyed Meysam³

1. Division of Immunology, Department of Microbiology, Veterinary Faculty, Urmia University, Urmia, Iran

2. Division of Immunology, Tarbiat Modares, University of Medical Sciences, Tehran, Iran

3. Assistant professor of Immunology, Department of Microbiology, Veterinary Faculty, Urmia University, Urmia

Background: Chemotherapeutic researches based on the use of copper oxide nanoparticles (CuO NPs) for new anticancer drugs with the potential of being less toxic and more cytotoxic against tumors have been developed. Therefore this study was done to investigate the concentration and size dependent of CuO NPs exposure in B₉₂ cancer cells.

Methods: CuO NPs were first green synthesized and characterized by typical scanning electron microscopy, the zeta sizer and X-ray diffractometer, then were cultured in 30 and 60 nm particle sizes at several concentrations (5, 10, and 20 µg/ml) with B₉₂ glial cancer cells for 24 hr. The apoptosis and cytotoxicity of cells were estimated by acridin orange/ propidium iodide staining and MTT assay, respectively. Also, the RT-PCR method was used to evaluate the expression of *P₅₃* and *Bax* expression.

Results: The images of synthesized CuO NPs confirmed the average sizes of 30 and 60 nm. The both size have cytotoxic effects, even with the lowest concentration, the cytotoxic effect accommodates the increasing percentage of cell apoptosis (32% at 30 nm size). Also, as the concentration increases, viability decreases and apoptosis increases. The IC₅₀ of CuO NPs with 30 nm size was (12.015±1.9) and for NPs with 60 nm was (9.17±3.3). However, the amount of IC₅₀ was decreased as the size of particles increase, but there was no significant change. Both mRNA level of *P₅₃* and *Bax* was upregulated in treated cells.

Conclusion: Our results suggest that an exposure to CuO NPs had significant cytotoxic effects when compared to unexposed ones in a way that the smaller size and higher concentration displayed the maximum cytotoxicity. However, it seems that augmentation may not have any effect on CuO NPs cytotoxicity *in vitro* state.

Keywords: B₉₂ cell line, CuO NPs, Cytotoxicity, Apoptosis, Glioma

ICIA2018\Nanoimmunology\Poster\10049
10049

New nanoformulation for artemisinin and evaluation of its effect on MCF7 cell line

Yasaman salimi, zuhair saraf hasan

Department of immunology, faculty of medical sciences, Tarbiat Modares University, Tehran, Iran

Background: One major challenge to relieve cancer burden is to develop highly effective drugs with specificity on cancers but little or no side effects on normal mammalian cells. Because conventional therapies based on chemotherapy, surgery and radiotherapy and current anti-cancer regiment are in many cases of limited efficacy. Therefore, herbal antioxidants, including artemisinin, were expanded. Artemisinin is a sesquiterpene lactone with a 1, 2, 4-tioxane ring system that extracted from the Chinese herb qinghaosu (*Artemisia annua* or annual wormwood). Despite its efficacy, artemisinin, has pharmacokinetic limitations. artemisinin has low solubility in water and biological fluids, poor bioavailability, and a short half-life in vivo (~2.5 h). To overcome these problems, we conjugated albumin with artemisinin and produced nano-artemisinin. The important thing is that the Nano substance must be harmless, protect the anti-cancer properties of artemisinin and also increase its solubility in water and its stability in the blood. In this study artemisinin had been conjugated with albumin.

Methods: After making the nano artemisinin, we used FTIR graph and scanning electerone microscopy (SEM) In order to confirm the combination of these two substances. Also the cytotoxicity properties of artemisinin and nano-artemisinin effect on mcf7 cell lines in Gradient of concentration within 24 hours, examined by mtt test.

Results: The FTIR confirmed that artimesinin and albumin bonded together. Particle size was reported 51.31 nm by SEM. The analysis of MTT tests indicated that the rate of killing of artemisinin and nano-artemisinin at 20µg dose was 77% and 86% respectively. Previous studies have been shown that this dosage almost has no killing effect on normal cells.

Conclusion: Due to the drug loading which is about 25% , Therefore, the amount of artemisinin used is a quarter of the amount of free artemisinin. This indicates that the killing ability was 4 times higher, with about no effect on normal cells.

Key words: artemisinin, MCF7 cell line, nanoformulation

ICIA2018\Nanoimmunology\Poster\9823

9823

The effects of coq10 nanoparticle on inflammation markers in streptozotocin-induced diabetic ratsAzad bahrampour¹, Roonak azizbeigi²*1. DVM Student, Faculty of Veterinary Medicine, Sanandaj Branch, Islamic Azad University, Sanandaj, Iran**2. Assistant of Physiology, Department of Physiology, Faculty of Veterinary Medicine, Sanandaj Branch, Islamic Azad University, Sanandaj, Iran*

Background: Coenzyme Q10 is an intracellular antioxidant that protects cell membrane phospholipids against oxidative damage caused by free radicals. Diabetes and its complications have been related to increased levels of free radicals and systemic proinflammatory cytokines and to an abnormal lipid profile. The aim of this study was to investigate the effects of CoQ10 nanoparticle supplementation on some cytokine levels in streptozotocin-induced diabetic rats.

Methods: In this study, 40 healthy adult male rats were used. The rats were divided into 5 groups. The animals in group 1 was fed standard rat pellets for 4 weeks. It was administered at 0.3 mL corn oil intraperitoneally daily for four weeks in group 2 animals. The animals in group 3 was injected intraperitoneally with (0.3 mmol/lit) CoQ10 nanoparticle daily for 4 weeks. Group 4 was made diabetic by subcutaneous injections of streptozotocin at dose of 40 mg/kg in 0.1 M citrate buffer (pH 4.5) single daily dose for two days and group 5 was made diabetic by subcutaneous injections of streptozotocin at dose of 40 mg/kg in 0.1 M citrate buffer (pH 4.5) single daily dose for two days and then was injected intraperitoneal with (0.3mmol/lit) CoQ10 nanoparticle daily for 4 weeks.

Result: At the end of the study, blood samples were taken from all animals. In these blood samples, IL-6 and TNF- α plasma levels were determined with ELISA using sandwich enzyme-linked immunosorbent method via commercial kits. In this study, IL-6 and TNF- α levels in diabetic group significantly increased ($P < 0.05$) in parallel with each other compared to control group levels. The same parameters were reduced ($P < 0.05$) by CoQ10 nanoparticle application in diabetic animals.

Conclusion: CoQ10 nanoparticle suppressed the increments in plasma pro-inflammatory cytokine levels and play important role in regulation of imbalance between inflammation markers in diabetes conditions.

Keywords: CoQ10 nanoparticle, cytokine, diabetes, rat



14th International Congress of Immunology and Allergys (ICIA2018)
26-28 April 2018, Tehran, I.R.Iran

Psychoneuroimmunology and Immunoendocrinology

Poster Presentation

ICIA2018\Psychoneuroimmunology and Immunoendocrinology\Poster\7761
7761

Relationship between serum levels of vitamin D and anti-Tpo in Hypothyroid patients in Urmia city

Asma khanzade¹, Yousef yousefzadeh², Roya khanahmadi³, Amir asghari⁴

1. Department of microbiology, Islamic azad university of urmia

2. Social security organization of urmia, iran

3. Faculty of medicine, Tabriz University of Medical Sciences, Tabriz, Iran

4. Department of Biology, Faculty of Sciences, Urmia University, Iran

Background: Vitamin D is nowadays regarded as a vitamin that is associated with several diseases. The aim of this study was to determine the relationship between serum vitamin D level and anti-TPO antibodies in hypothyroid patients. Thyroperoxidase (TPO) is an enzyme that is involved in the synthesis of the thyroid hormone. Vitamin D is a group of fat-soluble vitamins and a natural moderator of the immune system, then the serum levels of vitamin D and anti-TPO antibodies were then evaluated using ELISA (Monobain, USA).

Methods: In this study, 90 subjects with hypothyroidism and 60 healthy subjects were matched with age and sex, and blood samples were taken.

Results: from 90 patients, 66 had a vitamin D deficiency (> 30 ng) and had positive anti-TPO. The vit. D level in 32 patients was less than 30ng/ml (48.5% deficiency) and in 20 patients (30.3%) it was more than 30ng/ml. 14 patients had vit.D level under 10ng/ml (severe deficiency in 21.2%). while this was in the control group (1.75 ± 31.78) and the patients average control group was (27.31).

Conclusion: In people with vitamin D deficiency, the anti-TPO antibody level is high. The results showed that there is a significant relationship between vitamin D deficiency and anti-TPO antibodies and vitamin D deficiency effects hypothyroid. Further consideration is suggested in the other geographic regions.

Keywords: Vitamin D, Anti-TPO, Hypothyroid deficiency

ICIA2018\Psychoneuroimmunology and Immunoendocrinology\Poster\8291
8291

Diminished circulating levels of interleukin-10 in patients with Parkinson's disease

Javad Masoumi¹, Hossain Khorramdelazad¹, Mohammad Taghi Rezayati¹, Azam Bagherizadeh¹, Shekoofeh Moghadam¹, Alireza Vakilian², Abdollah Jafarzadeh³

1. Department of Immunology, Medical School, Rafsanjan University of Medical Sciences, Rafsanjan, Iran.

2. Department of Neurology, Medical School, Rafsanjan University of Medical Sciences, Rafsanjan, Iran.

3. Department of Immunology, Medical School, Kerman University of Medical Sciences, Kerman, Iran.

Background: Parkinson's disease (PD) is one of the most prevalent neurodegenerative disorders that is specified by the destruction of dopaminergic neurons. A number of genetic as well as environment parameters have been introduced as risk factors of PD. However, the neuro-inflammation is one of the most important characteristic of PD. The aim of this study was to investigate the levels of anti-inflammatory cytokine IL-10 in patients with PD.

Methods: The blood samples collected from 23 patients with PD and 24 age and gender-matched healthy subjects as a control group. The serum levels of IL-10 were measured by ELISA.

Results: The mean serum levels of IL-10 was 7.7 ± 3.89 pg/ml in PD group and 12.61 ± 9.9 in healthy subjects. The mean serum levels of IL-10 in PD patients was significantly lower than healthy group ($P < 0.03$). No significant differences were observed between men and women neither in PD patients nor in healthy control group.

Conclusion: These results showed lower levels of IL-10 in patients with PD that represent the may plays an important role in pathogenesis of PD. The levels of IL-10 did not influence by gender of participants.

Keywords: Parkinson's disease, Interleukin-10, Serum, Gender

ICIA2018\Psychoneuroimmunology and Immunoendocrinology\Poster\8480
8480

Evaluation of the effects of *Lactobacillus plantarum* on the acquired conditional learning of rats

Minoo Mohamadi, Maryam Tajabadi Ebrahimi*, Bahareh Pakpour

Department of Biology, Faculty of basic science, Islamic Azad University, Central Tehran branch, Tehran, Iran

Background: Investigations have shown that foodstuffs can affect the brain and nervous system hormones through vagus nerve. Accordingly, probiotics can be effective in regulating activity of different parts of the brain by improving the function of the brain.

Aim: The aim of this study was to evaluate the effect of *Lactobacillus plantarum*, an indigenous probiotics isolated from traditional Iranian dairy products, on acquired conditional learning of Wistar rats.

Methods: In this study, 80 male Wistar rats with 200-250 grams weight range were used. For induction of amnesia, intra peritoneal pre-test injection of morphine (1 mg/rat) was performed in rats, and 1×10^9 cfu/ml probiotic bacteria (*Lactobacillus plantarum*) were administered to rats in the treatment group for nine month. The step-through passive avoidance used to examine long-term memory in control group and probiotic-treated group.

Results: In the control group, morphine significantly decreased learning. In the probiotic-treated group receiving morphine, there was a significant improving in the acquired conditional learning when compared with control group.

Conclusion: The findings of this study represented the additive effect of *Lactobacillus plantarum*, an indigenous Iranian probiotic isolated from dairy products, in acquired conditional learning of rats received intraperitoneal morphine, indicating beneficial effects of oral probiotics on memory and the learning ability of rats.

Keywords: Amnesia, Morphine, *Lactobacillus plantarum*, Rats

ICIA2018\Psychoneuroimmunology and Immunoendocrinology\Poster\8572

8572

Circulating specific immunoglobulin G to *Helicobacter pylori* in patients with Parkinson's disease

Saeed Mirzaee¹, Hossain Khorramdelazad¹, Mohammad-Taghi Rezayati¹, Azam Bagherizadeh¹, Shekoofeh Moghadam¹, Alireza Vakilian², Abdollah Jafarzadeh³

1. Department of Immunology, Medical School, Rafsanjan University of Medical Sciences, Rafsanjan, Iran

2. Department of Neurology, Medical School, Rafsanjan University of Medical Sciences, Rafsanjan, Iran

3. Department of Immunology, Medical School, Kerman University of Medical Sciences, Kerman, Iran

Background: Parkinson's disease (PD) is a progressive neurodegenerative disease that is specified by damage of dopaminergic neurons. The neuro-inflammation is one of the main feature of PD and there are some reports regarding the association of infections with PD. The aim was to evaluate the seroprevalence of anti-H. Pylori IgG in patients with PD and healthy individuals from Rafsanjan, a city located in south-east of Iran.

Methods: The blood samples collected from two groups including patients with PD and age and gender-matched healthy subjects as a control group. The sera were tested for the anti-H. pylori IgG antibodies using ELISA method.

Results: The seroprevalence of anti-H. pylori antibodies in patients with PD (78.3%) was similar to that observed in healthy subjects (79.2%). There was no also significant difference between healthy control group and control group regarding the mean titer of anti-H. pylori antibodies (67.17 ± 8.74 U/ml vs 56.54 ± 9.38 U/ml; $P = 0.41$).

Conclusion: These results show that H. pylori seropositivity rate was similar in PD patients and control group. No association was also found between H. pylori and PD in investigated population.

Keywords: Parkinson's disease, *Helicobacter pylori*, Seroprevalence

ICIA2018\Psychoneuroimmunology and Immunoendocrinology\Poster\9807
9807

The Role of Cytokines in Autism Spectrum Disorder

Fatemeh Mohammadizadeh^{1,2}

1. Student Research Committee, Sirjan Faculty of Medical Sciences, Kerman University of Medical Sciences, Kerman. Iran

2. Department of Laboratory, Sirjan Faculty of Medical Sciences

Background: Autism is a neurodevelopmental disorder characterized by deficits in communication and social skills as well as repetitive and stereotypical behaviors. While much effort has focused on the identification of genes associated with autism, research emerging within the past two decades suggests that immune dysfunction is a viable risk factor contributing to the neurodevelopmental deficits observed in autism spectrum disorders (ASD). Currently, it is estimated that ASD affects 1 in 68 children in the United States, with the average age of diagnosis at 4 years.

Methods: A systematic search was performed to identify studies published in multiple databases (Cochrane, Embase, Pubmed and google Scholar) up to 2018, and recently published abstracts were also reviewed. Using the key words Autism, Cytokines in Autism, Autism Spectrum Disorder, Neuroimmunology and pathophysiology of autism.

Results: There is strong evidence that disruption of normal cytokine levels has a significant role as a risk factor for neurodevelopmental defect, such as autism. Results showed that elevated mid-gestational levels of inflammatory cytokines and chemokines are more highly associated with the ASD subphenotype that presents with intellectual disability (ID), compared with ASD without ID, developmental delay (DD) without ASD, and typically developing controls. These included higher levels of GM-CSF, TNF- α , IFN- γ , IL-1 α , IL-1 β , IL-4, and IL-6.

Conclusion: Maternal inflammation history and the pro-inflammatory/anti-inflammatory cytokine balance are thus particularly important avenues for further study, especially at the interface between mother and fetus.

Keywords: Autism, Cytokine, Immune dysfunction, autism spectrum disorders

ICIA2018\Psychoneuroimmunology and Immunoendocrinology\Poster\7476
7476

Dysregulation of peripheral Th17 cells in elderly patients with ischemic stroke

Farnaz Hajizadeh^{1,2}, Sanam Dolati^{1,2}, Mehdi Yousefi^{1,2}

1. Stem Cell and Regenerative Medicine Institute, Tabriz University of Medical Sciences, Tabriz, Iran

2. Department of Immunology, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Ischemic stroke causes brain injury with activation of an inflammatory response that can contribute to clinical impairment. The involvement of IL-17A and IL-17-secreting T cell subsets in IS patients has not been verified. In the present study, we studied Th17 frequency, cytokine secretion, expression of transcription factors related to Th17 which is compared between IS patients and control group.

Methods: 30 patients with IS and 30 persons as control group were enrolled in this study. Blood samples were collected from IS patients at 24 hours, 5 and 10 days after stroke, Th17 was detected among PBMCs using flowcytometric analysis, SYBR Green method of polymerase chain reaction (PCR) was employed for the analysis of ROR γ t mRNA levels, Quantification of IL-17 was carried out with the TaqMan Real-Time PCR. Secretion of IL-17, was evaluated in the supernatant acquired from cultured PBMCs using ELISA.

Results: The proportion of Th17 cells were significantly elevated in IS patients, ROR γ t mRNA expression levels were significantly increased in IS patients after 1 and 5 days compared to the controls 2.06 ± 72 and 1.77 ± 0.5 ($p < 0.0001$), IL-17A levels were increased in IS patients at 1, 5 and 10 days after stroke compared with the controls from 118 ± 35.9 to 171 ± 34 ($p = 0.0003$), 147 ± 24.84 ($p = 0.005$) and 146 ± 27.57 ($p = 0.005$), respectively.

Conclusion: These studies suggest that the increase in proportion of Th17 cells might contribute to the pathogenesis of IS.

Keywords: Ischemic Stroke, Th17, Cytokines, Inflammation



Reproductive Immunology

Poster Presentation

14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

ICIA2018\Reproductive Immunology\Poster\5289

5289

Relation of Gene's expression Profile of IL-17 and IL-23 in Cumulus cells and their Follicular Fluid Concentrations in Infertile Patients at the Risk of Ovarian Hyperstimulation Syndrome (OHSS)

Leila Motamedzadeh ^{1,3}, Hossein Hadinedoushan ², Ehsan Farashahi Yazd ³, Farzaneh Fesahat ³, Mohammad Mahdi Mohammadi ^{1,4*}

1. Department of Immunology & Allergy, Kerman University of Medical Sciences, Kerman, Iran

2. Department of Immunology, Reproductive Immunology Research Center, Shahid Sadoughi University of Medical Science, Yazd, Iran

3. Research and Clinical Center for Infertility, Shahid Sadoughi University of Medical Science, Yazd, Iran

4. Physiology Research Center, Kerman University of Medical Sciences, Kerman, Iran

Background: Ovarian hyperstimulation syndrome (OHSS) is one of major complications during assisted reproductive technology.

Methods: The cumulus cells of oocytes from 80 infertile women were divided into two groups of case (n=40) with an average age 29.6 ± 1.4 years and control (n=40) with an average age 29.2 ± 1.1 according to the patients' infertility etiologies. The control group was the infertile patients with male factor infertility etiologies while; infertile women at the risk of OHSS were defined as case group in this study. IL-17 and IL-23 concentrations in FF were determined. mRNA expression levels of IL-17 and IL-23 of samples were measured using real time PCR method in all subjects.

Results: The concentrations obtained from FF demonstrated the higher levels of IL-17 in case group ($P=0.04$), while, it showed no remarkable difference in IL-23 ($P=0.3$). The mRNA levels of IL-17 and IL-23 showed no significant differences between two groups. There was a positive significant correlation between FF concentrations of IL-23 and the oocytes maturation rates in case group ($P=0.01$).

Conclusion: Our finding showed the mRNA expression of IL-17 and IL-23 were similar in two groups, and IL-17 may be considered as risk factor for OHSS.

Keywords: OHSS, IL-23, IL-17

ICIA2018\Reproductive Immunology\Poster\7412

7412

TH17 and Treg frequencies in treatment of RPL women with immune abnormalities after IVIG therapyGolshan Sheikhsaria^{1,2}, Leili Aghebati-Maleki^{1,2,3}, Mehdi Yousefi^{1,2}

1. Stem Cell and Regenerative Medicine Institute, Tabriz University of Medical Sciences, Tabriz, Iran

2. Department of Immunology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

3. Immunology research center, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Recurrent pregnancy loss (RPL) is a complication in pregnancy and is defined as three or more recurrent miscarriage in over 30 years of old women. Previous studies have described that immunological factors have essential role in RPL. After 1980 using of IVIG therapy for abortion associated with antiphospholipid antibodies (APA) have considered. In this study, we aimed to evaluate of immune cell (TH17, Treg) and pregnancy outcome after IVIG therapy in RPL patients.

Methods: 44 blood samples were taken from RPL women with immune abnormalities after positive pregnancy test. Simultaneously 400 mg/kg of IVIG was administered intravenously every 4 weeks until 28–30 weeks of gestation. Control group was included of 12 RPL patient with abnormal cellular immune profile without IVIG therapy. The frequency of Th17 and Treg cells, the expression of related transcription factors of these cells and the serum levels of associated cytokines were evaluated after IVIG therapy by flow cytometry, real-time PCR and ELISA, respectively.

Results: We have seen significantly reduced frequency of Th17 cells from $3.94 \pm 1.12\%$ to $1.83 \pm 0.56\%$ and increased frequency of Treg from 3.55 ± 1.65 to 9.13 ± 1.23 in IVIG treated group. Significant decreased secretion of IL-17 and IL-23 and expression of ROR γ t and IL-17 mRNAs also significant increased secretion of IL10 and TGF- β and expression of Foxp3. Pregnancy outcome in IVIG treated group in Comparison to untreated group had increased 40.8 %.

Conclusion: These studies suggest that, IVIG therapy can be very helpful in establishing the balance between Th-17 and Treg in successful pregnancy. Also, the imbalance TH17/Treg in RPL women with immune abnormalities could be improve by decreased TH17 and increased Treg frequency following IVIG therapy.

Keyword: Intravenous immunoglobulin G, Recurrent pregnancy loss, TH17, Treg

ICIA2018\Reproductive Immunology\Poster\7416
7416

Effect of IVIG on exhausted T cell and exhausted regulatory T cells in women with recurrent miscarriage

Amin kamrani¹, Leili Aghebati Maleki², Mehdi Yousefi³,

1. a Stem Cell and Regenerative Medicine Institute, Tabriz University of Medical Sciences, Tabriz, Iran b Department of Immunology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

2. a Stem Cell and Regenerative Medicine Institute, Tabriz University of Medical Sciences, Ta

3. briz, Iran b Department of Immunology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran c Immunology research center, Tabriz University of Medical Sciences, Tabriz, Iran

a Stem Cell and Regenerative Medicine Institute, Tabriz University of Medical Sciences, Tabriz, Iran b Department of Immunology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran.

Background: Exhausted T cell, exhausted regulatory T cells and Treg cells have been suggested as novel main factors for recurrent miscarriage (RM). In current study, we studied the effect of IVIG on exhausted T cell and exhausted regulatory T cells and pregnancy outcome in women with RM.

Methods: 94 pregnant women with RM were participated in this study. Blood samples were taken. Then, IVIG (400 mg/kg) was administered for 44 patients, intravenously. 50 other RM patients were included as no IVIG interfering control group. Following the first administration, IVIG was given every 4 weeks through 32 weeks of pregnancy. Blood sample was collected after the last administration. Exhausted T cell, exhausted regulatory T cell and regulatory T cell population were assessed before and after treatment in the two groups.

Results: IVIG administration reduced the population and function of exhausted regulatory T cells and increased the population and function of regulatory T cell CD4 + CD25 + cells. Also IVIG treatment don't effect on population and function of Exhausted T cells. But, no significant difference in population and function of exhausted T cell, regulatory T cell and exhausted regulatory T cells was detected in the untreated groups before and after pregnancy. Pregnancy outcome in IVIG treated subjects was successful in 38 out of 44 RM women (86.3%). Nevertheless, pregnancy outcome was successful in 21 out of 50 untreated RM women (42%).

Conclusion: Administration of IVIG in RM women with cellular immune cells irregularities during pregnancy effects exhausted T cell and exhausted regulatory T cells and regulatory T cells in peripheral blood and improves Treg and decreases exhausted regulatory T cells responses. These studies suggest that, this immune modulatory influence of IVIG may be associated with effective pregnancy.

Keywords: IVIG; exhausted T cell; regulatory T cell; recurrent miscarriage

ICIA2018\Reproductive Immunology\Poster\7513
7513

Investigation of T cell subtypes in women with recurrent pregnancy loss and control group

Samaneh Abdolmohammadi Vahid ^{1,2}- Majid Ahmadi ^{1,2}-Mehdi Yousefi ^{1,2}

1. Stem Cell and Regenerative Medicine Institute, Tabriz University of Medical Sciences, Tabriz, Iran.
2. Department of Immunology, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran.

Background: Recurrent pregnancy loss (RPL) includes approximately 15% to 25% of all pregnancies and is determined as 3 pregnancy losses before the twentieth week of gestation. Immunologic abnormalities, such as elevated rate of Th1, Th17 and reduced number of Th2 and Treg, are among the etiologies which are mentioned for RPL. Here, we evaluated T cell subtypes frequency and related genes expression level in RPL women and control group.

Methods: Totally 40, 20 RPL and 20 healthy women enrolled in the study. Evaluation of Treg and Th17 frequency was done by flowcytometry and mRNA expression level of T-bet (Th1 transcription factor), GATA-3 (Th2 transcription factor), GITR (Treg associated factor) and IRF-4 (Th17 related marker), the genes responsible for T cells subpopulation function, by real time PCR.

Results: Treg frequency was significantly decreased in RPL women compared with healthy women, while an increase was observed in Th17 level. Comparison of expression of GATA-3 and GITR in RPL patients, showed a significant decrease when compared with healthy women (p value=0.0008 and 0.0001, respectively), while expression level of IRF-4 and T-bet were higher in RPL women (p value=0.0001 and 0.011, respectively).

Conclusion: Increased rate of proinflammatory parameters such as Th17 frequency, T-bet and IRF-4 expression and reduction of suppressive parameters, Treg and expression of GATA-3 and GITR, in RPL women, represent the undeniable role of immune system in successful pregnancy. Immunologic factors may be used as helpful prognostic biomarkers in high risk women.

Keywords: recurrent pregnancy loss, T regulatory, T helper17

ICIA2018\Reproductive Immunology\Poster\7593
7593

Modulatory effect of pioglitazone on pro-inflammatory cytokines (TNF- α and IL-1 β) levels in testes of streptozotocin-induced diabetic rats

Farin Malekifard¹, Ali Soleimanzadeh²

1. Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

2. Department of Theriogenology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

Background: Diabetes mellitus has adverse effects on the male sexual and reproductive functions. Oxidative stress is one of the major pathophysiological routes during diabetes. Pioglitazone is a high-affinity PPAR- γ agonist. Previous studies have demonstrated that pioglitazone is a potent inhibitor of inflammation and a potent antioxidant. Oxidative stress causes the activation of various transcription factors including the nuclear factor- κ B (NF- κ B). NF- κ B plays a crucial role in inflammation, immune function and expression of pro-inflammatory cytokines such as TNF- α , IL-1 β , and IL-6. IL-1 β , alone or in combination with TNF- α and IFN γ induces transcription of the inducible nitric oxide synthase (iNOS) gene.

The present study was undertaken to examine the protective effect of pioglitazone on the anti-inflammatory system in the testis of streptozotocin-induced diabetic rats.

Methods: Induction of experimental diabetes was done using a single intraperitoneal injection of streptozotocin (STZ) dissolved in citrate buffer (pH 4.5) at the dose of 65 mg/kg to overnight fasted rats. Only rats with blood glucose concentrations above 250 mg/dl were considered as diabetic. Animals were randomly divided into four groups of eight rats: control group, STZ-induced diabetic group and STZ-induced diabetic groups treated with low or high doses of pioglitazone of 1 or 10 (mg/kg/day, orally) for 5 weeks. Mice were euthanized on day 35 and pro-inflammatory cytokines (IL-1 β and TNF- α) levels in testicular tissue were performed by ELISA kits.

Results: Testicular levels of pro-inflammatory cytokines including TNF- α and IL-1 β were increased in diabetic rats compared to control animals ($P < 0.05$). IL-1 β and TNF- α levels significantly decreased only in the high (10 mg/kg) dose of pioglitazone group when compared to diabetic rats ($p < 0.05$).

Conclusion: Pioglitazone reduced testicular inflammation by decreasing pro-inflammatory cytokine levels.

Keywords: Pioglitazone, IL-1 β , TNF- α , Testicular damage

ICIA2018\Reproductive Immunology\Poster\7594
7594

Effect of pioglitazone, a ligands of peroxisome proliferator-activated receptor gamma (PPAR- γ), on expression of NF- κ B in testes of streptozotocin-induced diabetic rats

Farin Malekifard¹, Ali Soleimanzadeh², Navideh Mirakhani³

1. Department of Microbiology, Facutly of Veterinary Medicine, Urmia University, Urmia, Iran

2. Department of Theriogenology, Facutly of Veterinary Medicine, Urmia University, Urmia, Iran

3. Department of Pathobiology, Facutly of Veterinary Medicine, Urmia University, Urmia, Iran

Background: Diabetes mellitus caused testicular damage by increasing oxidative stress and inflammation. Pioglitazone is high-affinity PPAR- γ agonists with antioxidant properties and potential anti-inflammatory effects. Previous studies have suggested that the anti-inflammatory effect of pioglitazone is mediated by the inhibition of NF- κ B activation. It has been shown that Native or synthetic PPAR- γ ligands inhibit the production of several inflammatory mediators, nitric oxide synthase and NF- κ B transcription activity. Previous reports showed that diabetes is associated with up-regulations of iNOS and NF- κ B-p65 expressions with a concomitant increase in the level of NO in testicular tissues of diabetic rats. The purpose of this study was to investigate the preventive effects of pioglitazone on expression of NF- κ B of testicular tissues.

Methods: Induction of experimental diabetes was done using single intraperitoneal injection of streptozotocin dissolved in citrate buffer (pH 4.5) at the dose of 65 mg/kg to overnight fasted rats. Only rats with blood glucose concentrations above 250 mg/dl were considered as diabetic. Animals were randomly divided into four groups of eight rats: control group, STZ-induced diabetic group and STZ-induced diabetic groups treated with low or high doses of pioglitazone (Sigma) of 1 or 10(mg/kg/day, orally) for 5 weeks. Mice were euthanized on day 35 and expression of NF- κ B in testicular tissue was performed by western blot.

Results: Diabetic rats exhibited significant high level in the expression of NF- κ B-p65 in testicular tissues ($P < 0.05$). Administration of pioglitazone effectively inhibited expression of NF- κ B-p65($P < 0.05$).

Conclusion: Pioglitazone attenuated testicular damage in diabetic rats by decreasing expression of NF- κ B.

Keywords: Diabetes, Pioglitazone, NF- κ B, Testicular damage

ICIA2018\Reproductive Immunology\Poster\7595

7595

Effect of hydro-alcoholic garlic extract on serum level of testosterone of streptozotocin induced diabetes in C57BL/6 mice

Farin Malekifard¹, Ali Soleimanzadeh², Yaser jafarikhataylou³

1. Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran.

2. Department of Theriogenology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

3. Department of Microbiology, Faculty of Veterinary Medicine, Tabriz University, Tabriz, Iran

Background: Previous studies have shown that diabetes induces changes in reproductive functions especially on steroidogenesis, histology of testes, spermatogenesis, sperm quality and fertility both in diabetic men and experimental diabetic animals. Oxidative stress is one of the major pathophysiological routes during diabetes. It has been reported that hyperglycemia causes dysfunction of leydig cells thus these cells cannot secrete testosterone. The aim of present study was to examine the antioxidative effects of garlic extract in diabetic mice.

Methods: Diabetes was induced by multiple low-dose of streptozotocin injection (40 mg/kg/day for 5 consecutive days) in male C57BL/6 mice (15-20 gr body weight). After induction of diabetes, mice were divided into 6 groups: group 1 (normal control group); group 2 (diabetic control group); group 3 (treatment with garlic extract 200 mg/kg for 35 days) and group 4 (treatment with garlic extract 400 mg/kg for 35 days). Mice were euthanized on day 35 and serum testosterone levels were estimated by ELISA kit.

Results: In the present study, significant decrease in serum testosterone level was observed in diabetic control group ($P < 0.05$). Garlic treatment effectively increased ($p < 0.05$) serum level of testosterone

Conclusion: Garlic treatment protects against diabetes-induced testicular damage in diabetic mice by increasing serum testosterone level.

Keywords: Diabetes, Garlic Extract, Testosterone

ICIA2018\Veterinary Immunology\Oral\8689
8689

Soluble expression of IHNV immunogenic glycoprotein in the periplasm of Escherichia coli

Seyed Amir Hossein Jalali^{1,2}, Rezvan Mohammadi Nezhad³

1. Department of Natural Resources, Isfahan University of Technology, Isfahan 84156–83111, Iran

2. Research Institute for Biotechnology and Bioengineering, Isfahan University of Technology, Isfahan 84156–83111, Iran

3. Department of Biotechnology, College of Agriculture and Natural Resources, University of Tehran, Karaj, Iran

Background: Glycoprotein of the infectious hematopoietic necrosis virus (IHNV), is the only structural protein of the virus capable of inducing neutralizing antibodies in the fish host. Special attention has been paid for recombinant production of this vaccine candidate. However, due to the highly hydrophobic nature and the presence of several disulfide bonds, IHNV-G expression in Escherichia coli (E. coli) is associated within complete protein folding and aggregation. In the present study, we have explored the soluble expression of heterologous IHNV-G in the periplasm of E. coli using Pel B signal peptide.

Methods: The full coding sequence (CDs) of IHNV-G was amplified from infected tissues of rainbow trout (Oncorhynchus mykiss) using reverse transcription-polymerase chain reaction (RT-PCR). After cleavage with appropriate restriction enzymes, the PCR product was cloned in the bacterial plasmid pET22b containing Pel B signal sequence for periplasmic expression. Simultaneously, a truncated fragment of IHNV-G lacking N- and C-terminal hydrophobic parts was cloned in pET19b for cytoplasmic expression. Both constructs were used to transform the expression host E. coli Rosetta (DE3).

Results: Considerable amount of truncated protein was produced in E.coli cytoplasm after induction with IPTG, but predominantly in the insoluble fraction. However, targeting protein expression to the periplasm resulted in the production of soluble IHNV-G, as detected by both SDS-PAGE and western blotting analysis. The soluble IHNV-G was successfully purified in an optimized stabilizing buffer using affinity chromatography.

Conclusion: The expression procedure described in this study can provide sufficient amount of recombinant IHNV-G to immunize rainbow trout fish or to be used for producing specific antibodies for diagnostic or therapeutic purposes.

Keywords: Periplasmic expression, Glycoprotein, Recombinant vaccine, IHNV

ICIA2018\Veterinary Immunology\Oral\9978
9978

The lysozyme cross reactivity, pH-optimum activity and similarity between animal species

Mohammad Khosravi¹, Shokooh Salimi Moghadam²

1. Department of Pathobiology, Faculty of Veterinary Medicine, Shahid Chamran University of Ahvaz, Iran.
2. Student of Veterinary Medicine, Shahid Chamran University of Ahvaz, Iran.

Background: Lysozyme is distributed in serum, plasma, amniotic fluid, stool, saliva, tears, urine and other biological fluids. The most commercially available egg-white lysozyme and detection methods routinely used for creating standard curves and detection tools on other species; this experiment investigated sequences similarity, cross-reactivity and pH-optimum activity of the lysozyme in different animal species.

Methods: Polyclonal antibodies against chicken's lysozyme were obtained from the immunized rabbits and the IgGs were purified using ion exchange chromatography. An in-house ELISA was developed for detection of the lysozyme cross reaction between different animal species. The optimum pHs for the lyses of *Micrococcus* cell wall were detected using turbidimetric analysis. The lysozyme amino acid sequences were retrieved and analyzed using NCBI data base.

Results: The linearity of developed ELISA was evaluated in the range of 1mg–7μg. The coefficients of regression 0.98 confirmed that the method employing detection was linear for the determination of chicken lysozyme; however, the test has unacceptable results for the other species samples. The optimum pH of the buffer on turbidimetric method was determined as 5.6, 5, 5, 5, 5 and 5.6 in human, cattle, sheep, horse, dog and fish samples, respectively. The chicken's lysozyme amino acid sequence has the identity as 0.60, 0.61, 0.55, 0.50, 0.58, 0.54, 0.60, 0.59 and 0.61 with cattle, fish, sheep, horse, human, goat, buffalo, dog and donkey sequence, respectively.

Conclusion: It's suggested to use a species-specific lysozyme and detection tools for measurement of the lysozyme.

Keywords: Lysozyme, ELISA, Cross- reactivity

ICIA2018\Reproductive Immunology\Poster\7643
7643

Effect of pioglitazone, a ligands of peroxisome proliferator-activated receptor gamma (PPAR- γ), on expression of inflammatory marker (iNOS) in testes of streptozotocin-induced diabetic rats

Farin Malekifard¹, Ali Soleimanzadeh², Yaser jafarikhataylou³

1. Department of Microbiology, School of Veterinary Medicine, Urmia University, Urmia, Iran

2. Department of Theriogenology, School of Veterinary Medicine, Urmia University, Urmia, Iran

3. Department of Microbiology, School of Veterinary Medicine, Tabriz University, Tabriz, Iran

Background: Impaired homeostasis under diabetic conditions is connected with the increased production of free radicals and deficiency of antioxidant systems. Pioglitazone is potent inhibitor of inflammatory and potent antioxidants. Biological markers of inflammatory responses are elevated level of NF- κ B and the associated inflammatory mediators such as iNOS. iNOS catalyzes the production of large amount of NO from L-arginine, and importantly, NO inhibited insulin secretion. The purpose of this study was to investigate the preventive effects of pioglitazone on expression of inflammatory marker (iNOS) of testicular tissues.

Methods: Induction of experimental diabetes was done using single intraperitoneal injection of streptozotocin dissolved in citrate buffer (pH 4.5) at the dose of 65 mg/kg to overnight fasted rats. Only rats with blood glucose concentrations above 250 mg/dl were considered as diabetic. Animals were randomly divided into four groups of eight rats: control group, diabetic group and treated with low or high doses of pioglitazone of 1 or 10(mg/kg/day, orally) for 5 weeks. Mice were euthanized on day 35 and expression of iNOS in testicular tissue was performed by western blot.

Results: Expression of iNOS in testicular tissue was significantly increased in diabetic rats compared with normal control group. Administration of pioglitazone for five consecutive weeks markedly reduced expression of iNOS in testicular tissue.

Conclusion: Our data suggest that pioglitazone attenuated testicular damage in diabetic rats by decreasing expression of iNOS.

Keywords: Diabetes, Pioglitazone, iNOS, Testicular damage

ICIA2018\Reproductive Immunology\Poster\7644
7644

Effect of pioglitazone, a ligands of peroxisome proliferator-activated receptor gamma (PPAR- γ), on antioxidant status marker (SOD) levels in testes of streptozotocin-induced diabetic rats

Farin Malekifard¹, Ali Soleimanzadeh², Nowruz Delirezhi¹

1. Department of Microbiology, Urmia University, Urmia, Iran

2. Department of Theriogenology, Urmia University, Urmia, Iran

Background: Diabetes can cause sexual dysfunction in people including spermatogenic disorders. Enhanced oxidative stress and changes in antioxidant capacity have important roles in the pathogenesis of chronic diabetes mellitus. Previous studies have demonstrated that pioglitazone treatment could reduce superoxide radical generation in different tissue types. The aim of present study was to examine the antioxidative effect (SOD levels) of pioglitazone in diabetic rats.

Methods: Induction of experimental diabetes was done using single intraperitoneal injection of streptozotocin dissolved in citrate buffer (pH 4.5) at the dose of 65 mg/kg to overnight fasted rats. Only rats with blood glucose concentrations above 250 mg/dl were considered as diabetic. Animals were randomly divided into four groups of eight rats: control group, diabetic group and treated with low or high doses of pioglitazone of 1 or 10(mg/kg/day, orally) for 5 weeks. Mice were euthanized on day 35 and testicular SOD (Superoxide dismutase) activity was estimated following the method described by Kono. One unit of SOD is described as the amount of enzyme required to cause 50% inhibition of pyrogallol auto-oxidation at alkaline pH.

Results: Diabetic animals treated with pioglitazone showed the activity of anti-oxidative enzyme. Pioglitazone had enhanced the level of SOD. The improvement in the level of SOD indicated that pioglitazone could have either increased the proliferation of the antioxidant enzyme or reduced the formation of ROS.

Conclusion: Pioglitazone could reduce excessive production of ROS with a resulting decrease in the anti-oxidative defense.

Keywords: Diabetes, Pioglitazone, Superoxide dismutase, Testicular damage

ICIA2018\Reproductive Immunology\Poster\7884

7884

Sjögren's Syndrome Related Infertility: Case Report in IranFatemeh Rezayat¹, Sadegh Feyzollahzadeh², Abbas Rezaei¹*1. Department of Immunology, School of Medicine, Isfahan University of medical Science, Isfahan, Iran**2. Department of Immunology, School of Medicine, Urmia University of medical Science, Urmia, Iran*

Background: Infiltration in the glands. The etiology of Sjögren's syndrome has been remained unknown. The chronic immune system stimulation mediators may play an important role in the pathogenesis of this disorder. This includes various autoantibodies, especially anti-Ro/SS-A and anti-La/SS-B. These autoantibodies are involved to immunopathogenesis of infertility and recurrent spontaneous abortion. Moreover high levels of antinuclear antibodies in pregnant women may induce fetus rejection. Most individuals with Sjögren's syndrome have systemic symptoms, such as dry eyes, xerostomia (dry mouth) and parotid gland enlargement. Primary Sjögren's syndrome occurs in the absence of the other rheumatic disorders; however, secondary Sjögren's syndrome is associated with rheumatic disease, such as SLE, rheumatoid arthritis, or scleroderma.

Methods: A 42 years old woman with 10 years history of infertility referred to Dr. Rezaei's Immunological Laboratory. Routine and related immunological tests were performed and the presence of antinuclear antibodies (ANA) was detected. ANA profile test determined the reactivity of these autoantibodies with nuclear Ro/SS-A antigen. Other known causes of infertility and/or miscarriage were rule out by specific tests such as lupus anti-coagulant, Antiphospholipid antibodies, complement regulatory molecules (CD55, CD46 and CD59) and CD16/56.

Results: Moreover the infertility associated infections such as HIV, HCV, HTLV, CMV, Toxoplasma and rubella were negative. Symptoms of Sjögren's syndrome such as dry eyes with foreign body sensation, xerostomia and facial erythema confirmed.

Conclusion: According to these results this patient may be suffering from an early (Rheumatoid factor (RF) negative) primary Sjögren's syndrome and her infertility may be the cause of high levels anti-Ro/SS-A auto-antibodies.

Keywords: Sjögren's syndrome, Antinuclear antibody, anti-Ro/SS-A, Infertility.

ICIA2018\Reproductive Immunology\Poster\7915

7915

Effect of Cyclosporine on Th1 and Th2 Lymphocytes and Improvement of Pregnancy Outcome in Recurrent Pregnancy Loss (RPL)

Ramyar Azizi¹, Majid Ahmadi², Mehdi Yousefi³

1. Department of Immunology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

2. Stem Cell and Regenerative Medicine Institute, Tabriz University of Medical Sciences, Tabriz, Iran

3. Department of Immunology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

Background: Recurrent pregnancy loss (RPL) occurs in approximately 2-5% of clinically recognized pregnancies in reproductive-aged women. Cyclosporine is an immunosuppressant medication and decrease function and frequency of lymphocytes. In the present study, the effect of cyclosporine treatment on Th1/Th2 cell ratio and function is compared between cyclosporine treated and untreated RPL patients.

Methods: A total of 129 patients with a history of RPL were enrolled in this study. The frequency of Th1 and Th2 lymphocytes and the expression of related transcription factors were assessed by flow cytometry and real-time PCR respectively. 76 patients that have increased Th1/Th2 cell ratio selected for this study. 38 patients since the β -HCG test has been positive received oral CsA three times a day in a dose of 50mg for 6 months. Other 38 RPL patients were included as no CsA interfering control group. The dosage of CsA sustained between 40-80 ng/ml. flow cytometry and real-time PCR Tests were accomplished in both group before and after 32 weeks treatment.

Results: In the treated groups Th1 cells population and function down-regulated but for Th2 up-regulated also T-bet transcription factor down regulated and GATA3 transcription factor up regulated. Th1/Th2 cell ratio decreased in the treated group compared with untreated group after 32 weeks (p value = 0.0001). 31 cases from 38 patient have successful Pregnancy outcome in the treated group (81.5%) compared with 16 cases from 38 patients (42.1%) in the untreated RM women.

Conclusion: During pregnancy administration of CsA in RPL women with increased Th1/Th2 cell ratio, raises frequency and function of Th2 and decreases in Th1 as well as influences Th1/Th2 ratio in peripheral blood. These results indicate that CsA can be an effective treatment strategy for RPL women with immunological abnormality.

Keywords: Recurrent pregnancy loss, Cyclosporine, Th1, Th2

ICIA2018\Reproductive Immunology\Poster\7933

7933

Investigating the presence of autoantibodies against Glucose Regulated Protein78 (GRP78) in the sera of women with Recurrent Pregnancy Loss

Behrouz Ghahresani fard^{1, 3}, Mohammad Hossein karimi², Leila Bidar L³, Mohammad Ali-Hasanzadeh¹
Moslem Ahmadi¹

1. Department of Medical Immunology, School of Medicine, Shiraz university of Medical sciences, Shiraz, Iran

2. Organ transplant research center, Shiraz University of Medical Sciences, Shiraz, Iran

3. Infertility Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Recurrent pregnancy loss (RPL) is defined as 3 or more abortion prior to 20th week of pregnancy, and occurs in 15-20 % of all pregnancies. Glucose Regulated Protein78 (GRP78) is an endoplasmic reticulum(ER) protein which is expressed on the trophoblast cells. The purpose of this study was to evaluate the presence of anti-GRP78 antibodies in sera of women with RPL.

Methods: In a case-control study, 38 women with unexplained RPL as patients group and 42 healthy pregnant women with no history of abortion as control group who referred to the Infertility Center, Shiraz University of Medical Sciences were selected. In the present study, we evaluated the presence of anti-GRP78 antibody in the sera of RPL patients and healthy pregnant women using a Lab ELISA assay. Furthermore, western blot technique was used to confirm the expression of GRP78 by placental tissues.

Results: We detected the presence of anti-GRP78 antibody in both healthy and patient women by ELISA technique. However no significant difference regarding the presence of anti-GRP78 antibody levels between patients and healthy women was observed (P value= 0.1)

Conclusion: The results of this study did not shown a significant relationship between anti-GRP78 antibody level and susceptibility to unexplained RPL.

Keywords: Pregnancy, RPL, Placenta, GRP78

ICIA2018\Reproductive Immunology\Poster\7960

7960

Human amniotic epithelial cells (hAECs) regulates NK cells activation in women with recurrent spontaneous abortion (RSA)Fahimeh Khadem¹, Nafiseh Esmail¹, Abbas Rezaei¹, Faezeh Abbasi Rad¹, Behnaz khani²*1. Department of Immunology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran**2. Department of Obstetrics and Gynecology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran*

Background: Unexplained Recurrent Spontaneous Abortion (URSA) is the most common immunological complication during pregnancy. In recent years, it has been found that the cells such as human amnion epithelial cells (hAECs) have the potency to modulate immune responses in vitro and in vivo. However, the immunomodulatory effect of hAECs on NK cells in women with URSA is largely unknown.

Methods: Blood mononuclear cells (PBMCs) were obtained from 14 URSA patients and co-cultured with isolated hAECs. NK cells were identified using anti-CD56 and anti-CD3 monoclonal antibodies (mAb). The expression of the activating receptor CD69 and the degranulation marker CD107a on NK cells was detected using specific mAb and analyzed by flow cytometry.

Results: In this study, we demonstrated that CD69+ activating receptor expression on NK cells was significantly decreased by incubation with hAECs in a dose-dependent manner. Also, the degranulation marker CD107a was significantly downregulated on NK cells following incubation with hAEC.

Conclusion: Our results suggest hAECs have immune regulatory effects on activation and cytotoxicity of NK cells. Potential therapeutic application of hAECs for dysregulated NK immunity should be investigated in the future.

Keywords: Human amnion epithelial cells (hAECs), Unexplained recurrent spontaneous abortion (URSA), Immunomodulation, Natural killer cell.

ICIA2018\Reproductive Immunology\Poster\7993

7993

Altered frequencies of CD4⁺CD25⁺Foxp3⁺ and CD8⁺CD25⁺Foxp3⁺ regulatory T cells in pre-eclampsiaMaryam Zare¹, Mehrnoosh Doroudchi¹, Behrouz Gharezi Fard^{1,2}, Masha Hoseini¹

1. Department of Immunology, Shiraz University of Medical Sciences, Shiraz, Iran

2. Infertility Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Regulatory T cells are of utmost importance for tolerating the fetus during a normal pregnancy. In some pregnancy complications such as pre-eclampsia, the frequency of regulatory CD4⁺CD25⁺Foxp3⁺ regulatory T cells is altered, but there is no consistency regarding the results. Besides, little is known about the frequency of CD8⁺CD25⁺Foxp3⁺ Treg cells in pregnancy complications. We aimed to investigate the frequency of both CD4⁺ and CD8⁺ regulatory T cells in the peripheral blood of women afflicted by pre-eclampsia.

Methods: Ten non-pregnant, ten healthy pregnant, and ten preeclamptic women participated in this study. Four colors flow cytometry method was used to identify the frequency of the CD4⁺ and CD8⁺ regulatory T cells in the peripheral blood.

Results: Results indicated that the frequencies of CD4⁺CD25⁺Foxp3⁺ and CD8⁺CD25⁺Foxp3⁺ cells were significantly lower in preeclamptic women compared to healthy pregnant and non-pregnant ones ($p < 0.05$). A positive correlation was also observed between CD4⁺ and CD8⁺ regulatory T cells ($R = 0.532$, $p = 0.002$). Moreover, CD4⁺ regulatory T cells negatively correlated with systolic and diastolic blood pressures ($R = -0.760$ and -0.753 , respectively; p values < 0.001). CD8⁺ regulatory T cells also had negative correlations with systolic ($R = -0.503$, $p = 0.001$) and diastolic ($R = -0.590$, $p = 0.005$) blood pressures.

Conclusion: In conclusion, a reduction in the frequencies of both CD4⁺CD25⁺Foxp3⁺ and CD8⁺CD25⁺Foxp3⁺ regulatory T cells might be important in the pathogenesis of pre-eclampsia.

Keywords: CD4⁺CD25⁺Foxp3⁺ regulatory T cells, CD8⁺CD25⁺Foxp3⁺ regulatory T cells, Pre-eclampsia, Pregnancy

ICIA2018\Reproductive Immunology\Poster\8020

8020

The effects of maternal immune system deviation on development of the neonate's spleen

Mohammad Khosravi¹, Kaveh Khazaeel², Mahmoud Khaksary Mahabady², Mohammad Amin Behmanesh³, Shokooh Salimi Moghadam⁴

1. Department of Pathobiology, Faculty of Veterinary Medicine, Shahid Chamran University of Ahvaz, Iran
2. Department of Basic sciences, Faculty of Veterinary Medicine, Shahid Chamran University of Ahvaz, Iran
3. Department of histology, Faculty of medicine, University of Dezful, Dezful, Iran
4. Student of Veterinary Medicine, Shahid Chamran University of Ahvaz, Iran

Background: Various organs derived from fetal stem cell's development; these processes are highly dependent on the endometrial niches. Circulating hormones and cytokines are released mostly, by maternal and fetal immune system. This experiment evaluated the effects of maternal T lymphocytes deviation on development of the neonate's spleen.

Methods: Total of the 20 female rats divided in 4 groups (L, S, T and C). The L and S groups were treated using *Leishmania major* and *Salmonella typhimurium*, respectively. Tacrolimus (2mg/kg/day) orally administrated to group T in continuous 5 days. The group C, as control, treated using sterilized normal saline. The animals were mated after 3 times treatments at 2 week intervals. The newborns were scarified, according to animal's welfares, at one and two months of age. The total body and spleen weight of the neonates were determined; the prepared samples were analyzed using routine histological methods.

Results: The neonates of the group S showed significant reduction ($P \leq 0.05$) on the spleen's average weight at two months of age; also, the mentioned group had a significant elevation on the white pulp's number than the other groups. The neonates of the groups L and T showed no significantly difference on weight or pulps number, despite the higher levels than group C.

Conclusion: Imbalanced responses of the Th1<Th2, even occurrences before pregnancy, have significant effects on development of the neonate's spleen.

Keywords: Maternal, Neonate, Immune system, Spleen.

ICIA2018\Reproductive Immunology\Poster\8021

8021

Evaluation of the effects of maternal immune response deviations on neonate's immune responseMohammad Khosravi¹, Mahmoud Khaksary Mahabady², Kaveh Khazaeel², Shokooh Salimi Moghadam³

1. Department of Pathobiology, Faculty of Veterinary Medicine, Shahid Chamran University of Ahvaz, Iran

2. Department of Basic sciences, Faculty of Veterinary Medicine, Shahid Chamran University of Ahvaz, Iran

3. Student of Veterinary Medicine, Shahid Chamran University of Ahvaz, Iran

Background: Development of the fetal immune system originated from stem cells expansion. The maternal circulating hormones and cytokines, which are released mostly by immune system components, may be transferred to uterine cavity and alter the endometrial niches. This experiment evaluated the effects of maternal immune response deviations on neonate's immune responses.

Methods: Total of the 20 female rats divided in 4 groups (L, S, T and C). The L and S groups were treated using *Leishmania major* and *Salmonella typhimurium*, respectively. Tacrolimus (2mg/kg/day) orally administrated to group T in continuous 5 days. The group C, as control, treated using sterilized normal saline. The animals were mated after 3 times treatments at 2 week intervals. The newborns were treated by injection of sheep Red Blood Cells (SRBC) at 4 weeks of age. The neonate's humoral immune responses to maternal treated antigens and SRBC were evaluated at 5 weeks of age using micro-agglutination test.

Results: The neonates of S and L groups have detectable antibodies titer against *S. typhimurium* and *L. major* respectively. Also, the neonates of S and T groups showed significant reduction ($P \leq 0.05$) on the anti-SRBC antibodies titer.

Conclusion: Stimulation of the Th2 and suppression of the immune responses, even occurred before pregnancy, have significant effects on neonate's immune responses.

Keywords: Maternal, Neonate, Immune response, Rat.

ICIA2018\Reproductive Immunology\Poster\8258
8258

Comparison of the therapeutic effect of syngeneic, allogenic, and xenogeneic mesenchymal stem cells on the reduction of abortion rates in murine model

Fatemeh Rezaei, Seyed Mohammad Moazzeni

Immunology Department, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

Background: Introduction: mesenchymal stem cells (MSCs), due to their immunomodulatory functions, are an ideal candidate for the treatment of immune related diseases. Recurrent spontaneous abortion (RSA) is one of the most common complications of pregnancy which in many cases is related to the immune system disorders. Our previous study has shown that syngeneic MSCs therapy could reduce the abortion rate in abortion prone mice. In this study the therapeutic effect of syngeneic, allogenic, and xenogeneic MSCs were compared in a mouse model of RSA.

Methods: MSCs were isolated from adipose tissue of CBA/j, BALB/c mice and human. After characterization, MSCs were injected (IP) at day 4 of gestation to female CBA/J mice following their mating with DBA/2 male. In control group PBS was injected and CBA/J x BALB/c mating was also used as normal pregnancy control. On day 14.5 of pregnancy embryo resorption rate was determined.

Results: All syngeneic, allogenic, and xenogeneic MSCs therapy reduced the abortion rates compared to MSCs non-treated group, however the therapeutic potential of xenogeneic MSCs was less than syngeneic and allogenic MSCs in this model.

Conclusion: Intraperitoneal administration of MSCs from various source could significantly reduce the abortion rate in abortion prone mice, this results indicates that the immunogenicity of allogenic and xenogeneic MSCs is not a contradictory problem for their therapeutic effects on RSA.

Keywords: Recurrent spontaneous abortion, mesenchymal stem cells, syngeneic MSC, allogenic MSC, and xenogeneic MSC

ICIA2018\Reproductive Immunology\Poster\8324
8324

The effect of leukemia inhibitory factor (LIF) on gene expression of vascular endothelial growth factor A (VEGF-A) in trophoblast tumor Cell Line (JEG-3)

Khodakaram Jahanbin¹, Mehri Ghafourian ^{1*}, Mohammad Rashno¹, Afshin Amari¹, Iman Naamipouran¹,

1. Department of Immunology, Faculty of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

Background: Several studies have shown that leukemia inhibitory factor (LIF) one of the most important cytokines in the process of embryo implantation and pregnancy, but so far the role of this cytokine in angiogenesis process have not been fully investigated. Our goal in this study is to evaluate the effect of LIF on gene expression of vascular endothelial growth factor (VEGF) in choriocarcinoma cell line (JEG-3) as a trophoblast cells.

Methods: JEG-3 choriocarcinoma cells stimulated with different concentrations of LIF for 6, 12, 24, 48 and 72 hours. Expression of VEGF analyzed by real-time PCR.

Results: In stimulated cells, different concentrations of LIF caused the significant decrease of VEGF gene expression ($p < 0.05$) at 12, 24 and 48 hours but in 72 hours there was an increase in VEGF gene expression and at 6 hours at concentrations of 1 and 10 ng, there was also an increased expression of the gene, but the expression of gen reduced at 50 ng concentration.

Conclusion: Gene expression of VEGF in trophoblast cells treated with LIF decreased, indicating that this cytokine plays an important role in angiogenesis in placenta. Therefore, LIF may have effects in pregnancy-related disorders in related to angiogenesis.

Keywords: JEG-3, leukemia inhibitory factor, trophoblast, VEGF-A.

ICIA2018\Veterinary Immunology\Poster\5264
5264

Survey Epidemiological of Anthrax in Kermanshah Province, West Iran During 2010- 2015

Pooyanmehr Mehrdad¹, Momeni Mohamad^{2*}, Zangeneh Mohamad Mahdi ²

1. Department of Pathobiology, microbiology section, Faculty of Veterinary Medicine, Razi University,

2. Veterinary Student, Faculty of Veterinary Medicine, Razi University

Background and Aim: Zoonotic diseases constitute a public health problem throughout the world. Anthrax is a potentially fatal zoonotic disease and is an important zoonotic disease which is primarily a job-related infection caused by *Bacillus anthracis*. The diagnosis of cutaneous anthrax (CA) may be very difficult, particularly in atypical presentations and non-endemic regions. Aim this study, anthrax cases seen between 2010 –2015 (six years) in Kermanshah Province.

Materials and Methods: Medical records of infected in patients with CA (Clinical, Para clinical information and pathological diagnosis of disease CA such as microbiologic procedures) who had been operated in referral all Health Centers Kermanshah Province were collected in the period six years (2010 – 2015). Twenty-five patients with CA were included in this study.

Results: Results indicated that All patients' referent Twenty-five patients were who diagnosed CA in medical center registries, 2010(3case), 2011 (4case), 2012 (7case), 2013 (7case), 2014(3case) and 2015(1case) patient) at six years' period (0,025%), per 100,000 persons respectively. Each Twenty-five patients with a diagnosis of CA were followed up. All patients had a history of animal contact. The clinical presentation of CA was typical in all patients were initially misdiagnosed with insect bites or angioedema. Cultures from the lesions were positive for *Bacillus anthracis* in Twenty-five cases. Gram stain from the lesions revealed Gram-positive rods in 25 cases. Patients were diagnosed by clinical presentation and a history of contact with sick animals or contaminated animal products.

Conclusion: CA is a very contagious and important infectious disease worldwide. Early and accurate diagnosis dramatically affects the prognosis of the disease. The diagnosis of CA may be difficult, especially in atypical presentations and non-endemic areas. Thus, CA should be kept in mind, especially in these situations.

Keywords: *Bacillus anthracis*, Zoonotic diseases, Kermanshah Province

ICIA2018\Transplantation\Poster\5266

5266

Gene Expression patterns of Toll Like Receptor (TLR)-2, TLR-4 and (MYD88) in renal transplant patients developing allograft dysfunction; A cohort study

Morteza Hosseinzadeh¹, Pedram Ahmadpoor², Mir Saeed Yekaninejad³, Farshad Foroughi⁴, Mehri Barabadi⁵, SanazKeshavarz shahbaz⁵, Ehsan Soltaninejad⁵, Aliakbar Amirzargar⁶

1. Department of Immunology, School of Medicine, Ilam University of Medical Sciences, Ilam, Iran

2. Department of Nephrology, Shahid Labbafi-nejad Medical Center and Chronic Kidney Disease Research Center, University of Shahid Beheshti, Tehran, Iran

3. Department of Epidemiology and Biostatistics, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

4. Department of Immunology, School of Medicine, Qazvin University of Medical Sciences, Qazvin, Iran

5. Department of Immunology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

6. Molecular immunology research centre, Tehran university of medical sciences, Tehran, Iran

Background: Innate immune responses play an essential role in the development of (IRI) and subsequent impaired graft function. This cohort intends to determine the sequential dynamic changes in (TLR)-4, TLR-2, and (MYD88) mRNA expressions in (PBMCs) and biopsy samples from kidney allograft recipients in relation to graft function within the first year post-transplant.

Methods: This study enrolled 52 renal transplant patients, 27 with well graft function (WGF) and 25 with graft dysfunction (GD). We collected peripheral blood samples from participants, pre- and post-transplantation (days 2, 90 and 180) in order to analyze mRNA expression levels of the TLR-2, TLR-4, and MYD88 genes in relation to allograft function during a one-year follow up period. Expression patterns of these molecules were evaluated in samples obtained by either protocol biopsy (PB,n=24) or cause biopsy (n=6) at the sixth month post-transplantation.

Results: There were significantly higher mean dynamic changes of post-transplant TLR-2, TLR-4, and MYD88 mRNA expressions in GD patients compared to WGF patients ($P=0.001$). (ROC) curve analysis based on (GFR) index showed the (AUC) values for the genes: TLR-2 (0.89; $P<0.001$), TLR-4 (0.86; $P<0.001$), and MYD88 (0.75; $P=0.003$) in the third month after transplantation for a GD diagnosis. Similar analyses showed the following AUC values at six months post-transplantation: TLR-2 (0.73; $P=0.006$), TLR-4 (0.73; $P=0.007$), and MYD88 (0.70; $P=0.02$). Also, the calculated AUCs for the expressions of these genes in allograft biopsies were 0.94 (TLR-2), 0.95 (TLR-4), and 0.98 (MYD88) in the sixth month post-transplant based on the pathology report ($P<0.001$).

Conclusion: Sequential monitoring of the expression patterns of TLR-2, TLR-4, and MYD88 in PBMCs as well as in biopsy samples could be considered predictive biomarkers for early and late kidney allograft outcomes.

Keywords: Kidney transplantation, acute rejection, Toll like receptors, Allograft Outcome.

ICIA2018\Transplantation\Poster\5267

5267

Increased Expression of Toll-Like Receptors 2 and 4 markers on Peripheral Blood Mononuclear Cells (PBMC) in Renal Transplant Recipients that Develop Allograft Dysfunction

Morteza Hosseinzadeh¹, Mohsen Nafar², Pedram Ahmadpoor², Mir Saeed Yekaninejad³,
Mohammad Hossein Niknam^{1,4}, Aliakbar Amirzargar^{1,4}

1. Department of Immunology, School of Medicine, Ilam University of Medical Sciences, Ilam, Iran

2. Chronic Kidney Disease Research Center, Shahid Beheshti University of Medical Sciences

3. Department of Epidemiology and Biostatistics, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

4. Molecular Immunology Research Center, Tehran University of Medical Sciences, Tehran, Iran

Background: The incidence of ischemic reperfusion injury (IRI) in early phase post transplantation and activation of toll-like receptor (TLR-2) and TLR-4 remarkably impacts the outcome of a renal allograft. To investigate whether the expression of TLRs in peripheral blood mononuclear cells (PBMCs) can predict the clinical outcome of kidney allografts.

Methods: We obtained blood samples from 52 renal transplant patients before transplant, and 2, 90, and 180 days post-transplantation in order to analyze the surface expressions of TLR-2 and TLR-4 on peripheral blood monocytes. The expression patterns of TLR-2 and TLR-4 were compared between patients with graft dysfunction (GD) and those with well-functioning graft (WFG).

Results: Significantly different mean dynamic changes in surface expression of TLR-2, according to percentage of TLR-2+ cells, between (the GD and WFG) groups existed at most time-points before and after renal transplantation ($p=0.007$) with the exception of day 2 post-transplantation. We observed significantly higher mean fluorescence intensities of TLR-2 and TLR-4 on CD14+ cells in the GD group compared to the WFG group. This finding was particularly observed 180 days post-transplantation ($p=0.001$). Based on TLR-2 and TLR-4 protein expression for each step, multiple logistic regression and ROC curve analysis revealed that an increase in CD14+ TLR-2+ monocytes within the 90 days post-transplantation was associated with increased risk of GD at 180 and 365 days post-transplantation [odds ratio (OR)=1.27, $p=0.005$].

Conclusion: Sequential monitoring of TLR-2 and TLR-4 expression patterns in peripheral blood monocytes appear to be prognostic and predictive biomarkers for early and late kidney allograft outcomes.

Keywords: Allograft Function, Kidney Transplant, Toll-Like Receptors

ICIA2018\Vaccine\Poster\5271

5271

Predicted B- and T-cell epitopes analysis of the genes encoding the malaria vaccine candidate, *Plasmodium falciparum* Cell Traversal Protein for Ookinetes and SporozoitesSakineh Pirahmadi¹, Sedigheh Zakeri^{1*}, AkramAbouie Mehrizi¹, NavidDinparast Djadid¹*1. Malaria and Vector Research Group (MVRG), Biotechnology Research Center (BRC), Pasteur Institute of Iran, Tehran, Iran, Pasteur Avenue, P.O. Box 1316943551, Tehran, Iran*

Background: Malaria still remains one of the most prominent public health problems. In modern malaria elimination and eradication strategies, a great deal of attention has been paid to the development of novel intervention strategies such as an effective malaria vaccine. However, one of the major concerns in malaria vaccine development is the antigenic diversity of the vaccine candidate antigens, since antigenic diversity limits the efficacy of acquired protective immunity to malaria. Cell-traversal protein for ookinetes and sporozoites (CelTOS) is a promising malaria vaccine candidate. Therefore, in the present investigation the antigenic diversity of CelTOS in global isolates of *P. falciparum* was assessed and the predicted immunodominant regions in PfCelTOS in relation to polymorphisms were analyzed.

Methods: The extent of sequence diversity of the *pfceltos* were assessed among both natural *P. falciparum* isolates collected from endemic areas of Iran (n = 93) and available global *pfceltos* sequence data retrieved from Plasmodb database (n = 159). Furthermore, PfCelTOS structure analysis in relation to antigenicity was done by using *in silico* approaches.

Results: The present results revealed that low-level diversity were observed in *pfceltos* gene. Also, most of the predicted B- and T-cell epitopes were located in the conserved region of the gene.

Conclusion: The present analysis demonstrates that there is a limited antigenic diversity in global PfCelTOS and most of the detected polymorphisms are located on C-terminal region. Also, most of the predicted B- and T-cell epitopes were located in the conserved region of the gene, but the limited detected polymorphisms in these regions should be considered for evasion of *P. falciparum* from immune responses in CelTOS-based vaccine. This study is the first investigation on the antigenic diversity of PfCelTOS, and the obtained results could provide knowledge for a better design of PfCelTOS-based malaria vaccine.

Key words: *Plasmodium falciparum*, CelTOS, immunodominant epitope, vaccine

ICIA2018\Research and Development\Poster\5323

5323

Effects of Monochromatic Different Colored Lights on Melatonin Secretion from in vitro Peripheral Blood Mononuclear Cells, Polymorphonuclear Cells and Whole Blood Cultures

Mehrnoosh Zagheh,¹ Rostam Golmohammadi,² Mahdi Alahgholi Hajibehzad^{1,3}, Roya Najafi Vosough,⁴ Zohre Zaerieghane,¹ Alireza Zamani^{1,3,*}

1. Department of Immunology, School of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran.

2. Department of Occupational Hygiene, School of Health, Hamadan University of Medical Sciences, Hamadan, Iran

3. Molecular Immunology Research Group, Research Center for Molecular Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

4. Department of Biostatistics, School of Public Health, Hamadan University of Medical Sciences, Hamadan, Iran

Background: The melatonin is mainly produced by pineal gland and controls the circadian rhythms. Recent studies have shown melatonin is also secreted from peripheral blood mononuclear cells (PBMCs). Therefore, we investigated whether melatonin could secrete from polymorphonuclear cells (PMNs) and then measurement monochromatic different colored lights effect on in vitro production of melatonin by PBMCs, PMNs and whole blood cultures.

Methods: Blood samples were obtained from an apparently healthy volunteer. PBMCs and PMNs were isolated by ficoll-paque and dextran solutions respectively. PBMCs, PMNs and whole blood cultures were exposed to white, red, blue and green lights (100 lux) and to darkness in with/without of phytohemagglutinin (PHA, 50 µg/ml) and phorbol myristate acetate (PMA, 10 ng/ml) activation for 8 hours. The melatonin secretion in the supernatants was measured by Enzyme-Linked Immunosorbent Assay (ELISA). Data were analyzed by multi-factorial, Kruskal-Wallis and Mann-Whitney tests.

Results: Results indicated in addition to PBMCs, PMNs also produce melatonin. The level of melatonin in the supernatant of PBMCs, PMNs and whole blood cultures was increased due to PHA (134.3 ± 29.31 pg/ml) and PMA (141.1 ± 31.4 pg/ml) activation comparing with no activated (92.30 ± 22.5 pg/ml) ($P < 0.001$). The monochromatic different colored lights have no effect on the melatonin production by PBMCs, PMNs and whole blood cultures.

Conclusion: Conclusion: It concluded PBMCs, PMNs and whole blood cells produce melatonin in vitro and mitogens activation caused to higher production. Moreover, although white light suppressed melatonin secretion but there was not difference between various types of lights.

Keywords: Peripheral Blood Mononuclear Cells, Polymorphonuclear Cells, Whole Blood, Melatonin, Light.

ICIA2018\Vaccine\Poster\6335

6335

CpG ODN and MPL are more efficient adjuvants than naloxone for induction of Th1 immune response against *plasmodium vivax* recombinant TRAP in C57BL/6 miceSaeed Nazeri¹, Sedigheh Zakeri¹, Akram Abouie Mehrizi¹, Navid Dinparast Djadid¹*1. Malaria and Vector Research Group (MVRG), Biotechnology Research Center (BRC), Pasteur Institute of Iran, Tehran, Iran*

Background: *Plasmodium vivax* is the major cause of malaria outside Sub-Saharan Africa and it has been indicated that the elimination of *P. vivax* is often technically challenging. A key tool for the control, elimination and eradication of *P. vivax* is the development of a broadly protective vaccine. The thrombospondin-related adhesion protein (TRAP) is one of the major sporozoite antigens that plays an important role in the invasion of mosquito salivary glands and hepatocytes by sporozoites and it is a promising malaria vaccine candidate. The main aim was to investigate the role of antibodies and cellular immune responses induced by purified recombinant PvTRAP delivered in individual adjuvants, naloxone (NLX), CpG oligodeoxynucleotides ODN1826 (CpG ODN), and 3-O-decylated monophosphoryl lipid A (MPL), in immunized C57BL/6 mice.

Methods: The recombinant PvTRAP alone or combined with NLX, MPL or CpG adjuvants were applied for immunization of mice. IgG, IgG1, IgG2b, IgG2c, and IgG3 antibody responses as well as the IFN- γ and IL-4 cytokines were determined in post-immunized mouse plasma.

Results: The significant highest level of anti-rPvTRAP IgG, IgG2b and IgG2c were identified in the groups received rPvTRAP/MPL (mean OD₄₉₀ nm: 2.41, 1.50, 1.30, respectively) and rPvTRAP/CpG (mean OD₄₉₀ nm: 2.40, 1.44 and 1.27 respectively) in comparison to the group received rPvTRAP/NLX (mean OD₄₉₀ nm: 2.01, 1.40, 1.158, respectively) ($P < 0.05$, independent sample *t*-test). Also, mice receiving rPvTRAP/MPL (2164 pg/ml) and rPvTRAP/CpG (2029 pg/ml) induced significantly the higher levels of IFN- γ than mice receiving rPvTRAP/NLX (1615 pg/ml) ($P < 0.05$, independent sample *t*-test) and no detectable IL-4 production in all mouse groups.

Conclusion: The present result revealed that the administration of rPvTRAP with CPG and MPL induced higher Th1 immune response than NLX adjuvant in mice and had more potential to increase the level of anti-TRAP antibodies as well as the high production of INF- γ .

Keywords: *Plasmodium vivax*, TRAP, Adjuvant, Vaccine

ICIA2018\Veterinary Immunology\Poster\7353

7353

Evaluation of some of the factors associated with immune system following effect of magnetic field in rats

Amin Lafzi¹, Mansour Khakpur²

1. Dvm, Veterinary Medicine Clinician , Tabriz ,Iran

2. Department Of Pathobiology, Tabriz Branch, Islamic Azad University, Tabriz, Iran

Background: This study was designed to investigate the effects of electromagnetic fields (EMF) uniformly on the immune system of 12 Norwegian male Wistar rats were used as experimental model.

Methods: These rats were divided into three control, treatment one and treatment two groups. The first treatment group was subjected to electromagnetic field for four hours and fifteen days. The second group received this treatment for eight hours and fifteen days. At the end of the fifteenth day, blood samples from rats were taken and were prepared to be analyzed. Then, blood cells were counted and after conducting biochemical tests, the data were analyzed using SPSS computer application, one-way ANOVA and Tukey range tests.

Results: The results revealed that the level of WBC increased in the first experimental group compared to control group; yet, it decreased in the second experimental group. The level of total protein relatively decreased compared to control group; however, level of albumin decreased significantly and at the same time Gamma globulin increased.

Conclusion: Eventually, the increase in the number of lymphocytes and decrease in the number of neutrophils and monocytes are the most significant effects of electromagnetic field on immune system.

Keywords: Immune system; Magnetic field; Rats.

ICIA2018\Tolerance and Autoimmunity\Poster\7413

7413

Preserving Th17/Treg Balance to be Considerable in the Pathogenesis of Behçet's DiseaseRaminPourakbari^{1,2}, Leili Aghebati-Maleki^{1,2,3}, Mehdi Yousefi^{1,2*}*1. Stem Cell and Regenerative Medicine Institute, Tabriz University of Medical Sciences, Tabriz, Iran**2. Department of Immunology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran**3. Immunology research center, Tabriz University of Medical Sciences, Tabriz, Iran*

Background: Behçet's disease (BD) is a systemic inflammatory disorder with a various range of clinical appearances, containing mucocutaneous, ocular, vascular, gastrointestinal, musculoskeletal, and central nervous system contribution. The BD pathophysiology is remarkable as it shares several features with both autoimmune and inflammatory syndromes, perhaps prompted by infections or additional environmental factors. In this study, we aimed to assess Th17 and Treg cell frequency, cytokine secretion, and expression of transcription factors relevant to Th17 and Treg cells of patients with BD.

Methods: Blood samples from 47 patients with Behçet's disease and 58 healthy individuals were taken, and then PBMCs were isolated by ficoll separation method. The frequency of Th17 and Treg cells were evaluated by flow cytometry. Then, the expression of related transcription factors were assessed by real-time PCR. Also, the serum levels of associated cytokines were determined using ELISA Technique.

Results: The ratio of Treg frequency and levels of IL-10, FOXP3 expression were significantly decreased in BD patients. In contrast, the proportion of Th17 was significantly increased accompany with increase in the levels of IL-17, IL-23, and ROR γ t expression in patients with BD.

Conclusion: These studies suggest that, the increase in ratio of Th17 cells and decrease in Treg cells might collaborate to the pathogenesis of BD. Therefore, controlling balance between Tregs and Th17 could be useful for the treatment of the BD patients.

Keywords: Behçet's disease; Peripheral blood; Treg; Th17

ICIA2018\Transplantation\Poster\7433

7433

Association of Delayed Graft Function with serum levels of soluble Fibrinogen-like protein 2 in patients with kidney transplantation

Ali Salmani ¹, Majid Faridgozar ², Javad Absari ³, Hassan Nikoueinejad^{*4}, Hossein Akbari ⁵, Behzad Einollahi⁶

1. Medical Student; Student Research Committee, Kashan University of Medical Sciences, Kashan, Iran

2. Master Student; Department of Microbiology and Immunobiology, Kashan University of Medical Sciences, Kashan, Iran

3. Master Student; Department of Microbiology, School of Medicine, Iran University of Medical Sciences, Tehran, Iran.

4. Medical doctor & Immunologist; Nephrology and Urology Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran

5. Statistician; Trauma Research center, Department of Community medicine, Kashan University of medical sciences, Kashan, Iran

6. Nephrologist; Nephrology and Urology Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran

Background: Delayed graft function (DGF) is an inflammatory complication of kidney transplant which means needing dialysis throughout the first week after transplantation. Considering the effects of regulatory T cells (Tregs) on DGF, we aimed to ascertain the relationship between soluble Fibrinogen-like protein 2 (sFGL-2), as Tregs' secretory factor, and DGF after kidney transplantation.

Methods: This Nested case-control study was done in a 6-month period among 2 groups of each 58-member of kidney transplanted patients with and without DGF. sFGL-2 serum level in all blood samples was measured by Elisa.

Results: The serum level of sFGL-2 in recipients with DGF was significantly higher than that in those without DGF ($P < 0.001$). There was a strong meaningful correlation between the serum level of sFGL-2 and the risk of DGF ($P < 0.001$). We found no significant effects of different variables of rejection, HLA mismatch, type of medication, family relation between recipient and donor, age, and sex on such correlation.

Conclusion: Our study showed a significant relationship between sFGL2 and DGF. Therefore, plasma levels of sFGL2 may be used as a diagnostic tool to determinate the risk of DGF.

Key words: Kidney Transplantation, Delayed Graft Function (DGF), Soluble Fibrinogen-Like protein 2 (sFGL-2)

ICIA2018\Tolerance and Autoimmunity\Poster\7434
7434

The Increased Expression of Integrin $\beta 7$ Gene in Peripheral Blood Leukocytes of the Newly Diagnosed Rheumatoid Arthritis Patients

Mahdi Taghadosi¹, Adel Fekri¹, Parisa Zafari², Nasrin Iranshahi¹

1. Department of immunology, school of medicine, Kermanshah University of Medical Sciences

2. Department of immunology, school of medicine, Mazandaran University of Medical Science

Background: Integrins and chemokines play critical role in Lymphocytes and leukocytes migration and recirculation during autoimmune disorders. Normally gut-homing T cells express CCR9 and integrin $\beta 7$ in order to home back to lamina propria of intestine from peripheral blood. Our study was conducted to evaluate expression levels of the chemokine receptor CCR9 and the integrin $\beta 7$ in circulating leukocytes and lymphocytes from newly diagnosed Rheumatoid arthritis (RA) patients and compare these values with the same data obtained from healthy individuals without inflammatory disease.

Methods: Total RNA was extracted from peripheral blood samples immediately cDNA was synthesized and finally gene expression of CCR9 and $\beta 7$ evaluated by quantitative Real time PCR. The T-test and ANOVA were used in order to compare the differences between two groups.

Results: The $\beta 7$ gene expression showed significant difference between RA patients and healthy controls (P value= 0.007).

Conclusion: In our newly diagnosed patients who did not received any medication for RA the gene expression of integrin $\beta 7$ was higher than healthy controls. CCR9 gene expression didn't reach significant different between two groups. Our results bring up disposability that over expression $\beta 7$ gene may have a role in pathogenesis of RA.

Key words: Rheumatoid arthritis, CCR9, Integrin $\beta 7$, MALT

ICIA2018\Veterinary Immunology\Poster\7456
7456

Possible allergic reactions, blood factors and the immune system alterations in response to ingestion of oral sumac (*Rhuscoriaria L*) Compared to Levamisole in dogs.

Ali Vahid Dastjerdi¹, Mohsen Jafarian dehkordi¹, saam torkan¹

1. Department of Clinical Science, faculty of veterinary medicine, Islamic azad university of Shahrekord, Shahrekord, Iran

Background: Since before any research on drugs we should be aware of its allergic dose, and given the importance of hematology system that implies the changes in the body, in this study, we decided to investigate the hematologic effects of sumac plant and compare it with a chemical agent (levamisole). In future studies, we'll use its valuable properties especially antibacterial and may replace it by some drugs including levamisole.

Methods: In this study, 8 native breed dogs were used for ten days, the dogs were divided into four groups of two, a group of control, a group receiving levamisole, a group receiving sumac at a rate of 10 mg per kg, and one group receiving 5 mg per kg sumac, and after the completion of prescribed period of sumac plant powder and oral levamisole powder, blood samples were taken and then the relevant tests were carried out.

Results: Given that in the majority of allergies and poisonings, increased eosinophils and basophils as well as red blood cell lysis, and decreased hematocrit can be seen, and because no study has been done so far on allergic factors such as eosinophils and basophils, and also due to the dosage of sumac plant used in this study that there was no significant difference between treatment and control group (), you can ensure the absence of allergies and sensitivities to this plant.

Conclusion: According to the same hematologic results of sumac plant and levamisole, valuable properties of sumac plant especially its antimicrobial and immune boosting effects used in future research without worrying about allergic reactions in certain doses of this research and without hematologic side effects.

Keywords: allergy, dog, Hematology, levamisole, sumac

ICIA2018\Tolerance and Autoimmunity\Poster\7464

7464

The Effect of Vitamin D3 on Proliferation of CD4⁺ T Lymphocytes in Patients with Type 2 Diabetes Mellitus

ElhamMahabadi-Ashtiyani^{1,2}, Vida Sheikh³, Shiva Borzouei³, AlirezaZamani¹, IrajSalehi⁴,
Mahdi Alahgholi Hajibehzad^{1,*}

1. Department of Immunology, Hamadan University of Medical Sciences, Hamadan, Iran

2. Student Research Committee, Hamadan University of Medical Sciences, Hamadan, Iran

3. Department of Internal Medicine, Clinical Research Development Unit of Shahid Beheshti Hospital, School Of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

4. Neurophysiology Research Center, Hamadan University of Medical Sciences, Hamadan, Iran

Background: It has been demonstrated that T helper cell subsets have a fundamental role in pathogenesis of type 2 diabetes mellitus (T2DM). Active form of vitamin D (1, 25 dihydroxyvitamin D₃, VitD₃) has an immunomodulatory effects on immune system. The aim of this study was to investigate the proliferation level of CD4⁺ T cells and effect of VitD₃ on CD4⁺ T cells proliferation in T2DM compared to healthy controls (HC).

Method: Blood sample of 18 patients with T2DM (mean: 51.68 ± 8.62; female=13, male=5) and 21 healthy control (mean: 42.80 ± 1.19; female=12, male=9) were collected. Peripheral blood mononuclear cells (PBMCs) were isolated by Ficoll density gradient method. CFSE-labeled PBMCs were cultured and stimulated with PHA in presence/ absence of VitD₃ for 5 days. The proliferation of CD4⁺ T cells were analyzed using flow cytometry.

Result: The addition of VitD₃ to the culture of PBMCs significantly decreased the proliferation of CD4⁺ T cells in both T2DM and HC, respectively (p= 0.001 and p=0.0001). The % proliferation was significantly lower in the HC group compared with T2DM patients in cultures with VitD₃. There was no significant changes in the % proliferation of CD4⁺ T cells between T2DM and HCs in cultures without VitD₃.

Conclusion: The data suggest potential insight into the consideration of VitD₃ in the control of T cells proliferation in T2DM disorders.

Keywords: T2DM, Proliferation, Lymphocyte, Vitamin D

ICIA2018\Tolerance and Autoimmunity\Poster\7518
7518

Gene Expression of CD226 and Its Serum Levels in Patients with Multiple Sclerosis

Fereshteh Alsahebhosul^{1,2}, Nahid Eskandari¹, Shokofeh Rahimpourkoldeh¹, Vahid Shayeghanejad³, Mazdak Ganjalikhani Hakemi¹, Arezo Dabiri⁴

1. Immunology department, Faculty of Medicine, Isfahan University of medical sciences, Isfahan, Iran

2. Isfahan Research MS and Neuroimmunology Center (IRMN), Isfahan, Iran

3. Neurology Department, Faculty of Medicine, Isfahan University of medical sciences, Isfahan, Iran

4. Immunology department, Shahid Sadoughi University of Medical Sciences, International Campus, Yazd, Iran

Background: Multiple sclerosis (MS) is a chronic disabling inflammatory demyelinating disease that is caused by immune responses directed against myelin proteins and progressive axonal loss of the central nervous system (CNS). CD226 one of T cell co-stimulatory molecule is suggested as genetic risk factors for TH1 dependent autoimmune disease. CD226 co-stimulatory signals potentially promote Th1 differentiation. We hypothesized that the CD226 gene expression and serum level can increase in MS patients.

Methods: The type of study is case-control. The qRT-PCR (qualitative real-time polymerase chain reaction) and ELISA (Enzyme-Linked Immunosorbent Assay) methods were applied for determining CD226 gene expression and serum protein level, respectively.

Results: We found that the gene expression and mean serum protein level of CD226 no significant differences was found in the MS patient and control groups. In addition, there was no significant difference between age and the CD226 gene expression and its protein in the individual patients. In the patients group also no significant differences in the mean of CD226 gene expression and protein between men and women was observed, although the mean of CD226 gene expression in women was higher than CD226 gene expression in the men, but this difference was not significant.

Conclusion: the gene expression and mean serum protein level of CD226 had no significant differences in the studied MS patient and control groups. Taken together, these data showed that there is no pathological relation between CD226 molecule and multiple sclerosis. It seems that CD226 can not be considered as a diagnostic marker for MS disease.

Key words: Multiple sclerosis, CD226, Gene expression, PCR

ICIA2018\Vaccine\Poster\7631

7631

Cloning of Immunogenic Moiety of Fusion Protein of NDV in a Plant Expression Vector

NargesArkian Boroujeni¹, Atena Mozafari¹, Mohammad Javad Motamedi¹, Somayeh Behjat Khatouni¹, Mahyat Jafari¹, Ali Hatef Salmanian^{1*}

National Institute of Genetic Engineering and Biotechnology (NIGEB), Department of plant Biotechnology, Tehran, Iran

Background: Newcastle disease (ND) is a highly contagious respiratory disease in poultry that affects many domestic and wild species. Outbreaks of virulent ND have a great economic impact on poultry industry in developing countries. The surface of Newcastle disease virus (NDV) particles containing two important functional antigens: the fusion (F) and hemagglutinin-neuraminidase (HN) glycoproteins which are considered as the main targets for the immune system against NDV. There are many killed or live-attenuated vaccines to prevent infection by NDV but they have many limitations which lead to develop novel vaccines having flexibility in administration, storage, transportation and cost-effective, also one of the promising platforms is using the genetic engineering tools and production of recombinant proteins in plant based systems such as hairy roots.

Methods: In the current study, a chimeric construct containing two antigens (F and HN), with the codon preference of plant which has already designed was used for further immunological analysis. The fusion (F) fragment was amplified from the synthetic HN-F gene by PCR using specific primers and cloned in pTZ57R/T vector. Then, the authentic fragment was subcloned into a plant expression vectors (pBI121), under the control of general (CaMV35S) and the specific hairy root (NtREL1) promoters, respectively. The recombinant vectors (pBI121-CAMV35S-*f*, pBI121-NtREL1-*f*) were transformed to an *Agrobacterium* host (LBA4404) and finally introduced to tobacco plant.

Results: The authentic fragment of *f* (~ 909 bp) in pTZ57R/T, pBI121-CAMV35S and pBI121-NtREL1 analyzed and confirmed by PCR, digestion and sequencing. Presence of *f* gene in recombinant *Agrobacterium* host confirmed by PCR method.

Conclusion: The hairy roots have several advantages such as genotype and phenotype stability, fast and indefinite in vitro growth. Analysis of expression level and immunological response of the recombinant constructs will be evaluated in an animal model are in progress.

Keywords: Newcastle Disease Virus, Fusion protein, Hairy root, Recombinant edible vaccine

ICIA2018\Tolerance and Autoimmunity\Poster\7635

7635

Frequencies of Antiphospholipid Antibodies in Women with Recurrent Abortion in Northeast of Iran

Tayebeh Sohrabi¹, Zahra Badiei², Mohammad Reza Keramati², AzamMoradi Zarmehri², Mohsen Rokni³, Mohammad Hadi Sadeghian², Hamid Farhangi², Masoud Shahdoodian², Geshvad Nasiri Pirbazari¹

1. Department of Hematology & Blood Bank, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran.

2. MD, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran.

Department of Immunology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran.

Background: Antiphospholipid Syndrome (APS) is taken into account as one of the most important immunologic cause of recurrent abortion, which is mostly not diagnosed. APS is an autoimmune hypercoagulable disorder caused by antiphospholipid antibodies. The majority of women with antiphospholipid antibodies show complications such as recurrent fetal abortions even after treatment. Due to the lack of accurate and reliable statistical data and the necessity of further research in this field in Northeast of Iran, this study aimed to investigate the rate and relationship of these antibodies with recurrent abortions

Methods: This cross-sectional study was performed on 239 patients with recurrent abortion who were referred to the Sarvar Thalassemia and Hemophilia Center between 2012 and 2014 years. Venous blood samples were collected to determine the serum levels of protein C, protein S, and antiphospholipid syndrome panel; including anti-cardiolipin, anti-β2 glycoprotein I, and lupus anticoagulant.

Results: In 239 patients with a mean age of 30 ± 5.4 years, the average rate of abortions was 2.75 ± 1.04 times. Anti-β2 glycoprotein I was detected in 5% (12), anti-cardiolipin and lupus anticoagulant identified in 4.6% (11) and 3.3% (8) of patients, respectively. The levels of anti-β2 glycoprotein I and anti-cardiolipin showed a .(remarkable relationship with the frequency of abortion (P -value=0.048 and 0.004, respectively

Conclusion: The frequency of antiphospholipid antibodies among women with recurrent abortion was low in Northeast of Iran. The frequency of abortion was significantly linked to the levels of anti-cardiolipin and anti-β2 glycoprotein I antibodies. The presence of more than one antibody is considered as a risk factor for increasing the frequency of abortions.

Keywords: Antiphospholipid antibody, Anti-cardiolipin, Anti-β2 glycoprotein I, Recurrent abortion

ICIA2018\Vaccine\Poster\7651

7651

Molecular Analysis of Chimeric Haemagglutinin-Neuraminidase Protein (HN) of Newcastle Disease Virus in Hairy Roots of Tobacco Plant (*Nicotiana tabacum L.*)

Somayeh Behjat Khatouni¹, Atena Mozafari¹, Mohammad Javad Motamedi¹, Narges Arkian Boroujeni¹, Mahyat Jafari¹, Ali Hatef Salmanian^{1*}

National Institute of Genetic Engineering and Biotechnology (NIGEB), Department of Plant Biotechnology, Tehran, Iran

Background: Newcastle disease (ND) is an important avian infectious agent that causes large economic losses to the poultry industry in developing countries. The Newcastle Disease Virus has two envelope glycoproteins, the fusion (F) and hemagglutinin-neuraminidase (HN) protein which considered as the main antigenic targets of the immune system. The HN protein plays an important role in attachment of the virus to host cell membrane. Due to a presence of various epitopes within the HN protein, it could be considered as an appropriate target for the immune system. Developments in genetic engineering have led to establishing a plant-based platform for production and oral delivery of candidate vaccine. The main benefits of edible vaccines are the facility of administration, as well as high-level expression of immunogens in different organs such as seeds and hairy roots via ordinary *Agrobacterium*-mediated transformation.

Methods: In the present study, a chimeric construct composed of F-HN epitopes with plant codon preference was designed. The HN fragment was amplified by PCR using specific primers and cloned in pTZ57R/T cloning vector. The fragment (HN) was subcloned into two plant expression vectors, one harboring a root-specific promoter "NtREL1" (pBI121-NtREL1-HN) and another with the CaMV35S promoter (pBI121-CaMV-HN). The recombinant vectors were transformed to the LBA4404 strain of *A. tumefaciens* and finally introduced to tobacco (*Nicotianatabacum L.*) plant.

Results: The authentic fragment of *hn* (~1038 bp) in pTZ57R/T, pBI121-CaMV and pBI121-NtREL1 analyzed by PCR, digestion and sequencing. Presence of recombinant vector in *Agrobacterium* and regenerated plants were confirmed by PCR method.

Conclusion: Production of recombinant antigenic proteins in hairy root platforms has several advantages such as high growth rate, easy genetic manipulations, high levels expression, and the potential for large-scale production in bioreactor. The next step, the analysis of expression level and immunological response of the recombinant construct will be evaluated in an animal model.

Keywords: Hemagglutinin-Neuraminidase protein, Newcastle Disease Virus, Hairy roots, Edible candidate vaccine

ICIA2018\Vaccine\Poster\7664

7664

Development of two novel chimeric VK210/VK247CSP based antigens for vaccine design against *Plasmodium vivax* malaria

Samaneh H. Shabani^{1,2}, Sedigheh Zakeri^{1*}, Ali H. Salmanian³, Jafar Amani⁴, Akram A. Mehrizi¹, Georges Snounou⁵, Yousef Mortazavi², Navid D. D. jadid¹

1. Institute, Iran

2. Zanjan University of Medical Sciences, Iran

3. National Institute of Genetic Engineering and Biotechnology, Tehran, Iran

4. Baqiyatallah University of Medical Sciences, Iran

5. Pasteur institute, France

Background: *Plasmodium vivax* is the most widespread species of *Plasmodium* outside Africa and is predominant in countries where malaria is in pre-elimination or elimination phase. As *P. vivax* cannot be maintained in *in vitro*, a sporozoite-based vaccine is not a practical option, and most current efforts have been focused on developing a *vivax*-based subunit vaccine. The circumsporozoite protein (CSP) of the *P. vivax* is a major target of pre-erythrocyte vaccine. In this work, we designed and constructed two novel synthetic chimeric genes of *P. vivax* CSP by using computer modelling. Subsequently, the expression and purification of both chimeric recombinant proteins were carried out, and their biological and functional properties were evaluated.

Methods: The two constructs were designed based on reference sequences (Sal-1 and PNG) encompassing different repeats from the two major alleles, VK210 and VK247 and entire N- and C-terminal to compare their structure, antigenicity and their biological activities. DNA encoding the selected chimeras were synthetically constructed based on *E. coli* codons and then cloned and expressed. Characterization of both constructs were performed by western blot, ELISA, heparin and HepG2 binding assay as well as IFA.

Results: Design, expression and characterization of both constructs were performed successfully. Strain specific mouse monoclonal antibodies recognized the chimeric antigens in ELISA, indicating correct conformation and that the B-cell epitopes are accessible to the antibodies. Heparin and HepG2 binding assay showed dose depended manner.

Conclusion: Although both designed constructs had structurally different characteristics at the molecular level, the expression levels were satisfactory, and chimeras had a conformational structure with biological function that signified their potential use in the development of *vivax*-based vaccine.

Keywords: *Plasmodium vivax*, malaria, subunit vaccine, circumsporozoite protein

ICIA2018\Veterinary Immunology\Poster\7692

7692

Horse Toxoplasmosis Seroprevalance in three Northwest Provinces of IranGolbahar Hosseini¹, Reza Rezaie², Diba karimi³*1. Department of Basic Sciences, Faculty of Veterinary Medicine, University of Tabriz**2. Department of Basic Sciences, Faculty of Veterinary Medicine, University of Tabriz**3. Department of Basic Sciences, Faculty of Veterinary Medicine, University of Tabriz*

Background: *Toxoplasma gondii* is one of the most important Zoonotics in the world. Despite many studies on human toxoplasmosis and ruminants, there are few to be found in Iran.

Methods: Ten ml of blood samples were collected from jugular vein of 385 horses among 68000 population in three northwest provinces of Iran. The sera were centrifuged at 3000 rpm for 10 minutes. To determine the prevalence of age, livestock was divided into three age groups ≤ 2 , 10-2, and 10. Rh antigens of *Toxoplasma gondii* were from prestigious research centers (such as the Pasteur Research Institute or the Razi Vaccine and Parenting Institute) for use in Modified Agglutination Test (MAT) *Toxoplasma gondii* antibodies were evaluated using the modified (MOD) agglutination test (MAT) according to the described methodology of Tavalla et al. (2015). Sera were diluted in binary dilutions (dilution) using PBS containing 2-mercaptoethanol. Fifty μ l of each dilution in each well was planted into 96 plates of U-shaped Eliza. Then, 50 μ l of whole formalin-preserved *T. gondii* Tachyzoites taxicides will be added to each serum dilution. incubated at 37 ° C overnight.

Results: Fifteen of 385 horses were identified positive which included 9 stallion and 6 mare. Of the total, 5 were horses less than 5 years old and 10 horses were above 5 years old.

Conclusion: Studies on seroprevalence of the disease with various methods of serology in the human population, indicated the presence of *Toxoplasma gondii* with different prevalence among humans and Livestock. since Equine are considered as interface hosts of *Toxoplasma gondii* and there is no comprehensive information on seroprevalance of the disease in the population we aimed to do this.

Keywords: *Toxoplasma gondii*. Horse, Ardabil, East/West Azerbaijan

ICIA2018\Vaccine\Poster\7775

7775

A quaternary vaccine against Enterohaemorrhagic *Escherichia coli* (EHEC)

Emad kordbacheh¹, Shahram Nazarian¹, Javad Fathi¹, Amin Farhang¹

1. Imam Hossein University, Faculty of Science, Department of Biology, Tehran, Iran.

Background: Enterohaemorrhagic *Escherichia coli* (EHEC) are a subset of *E. coli* pathovars and a frequent agent of diarrhoeal disease which can contaminate by food and water sources. Especially O157:H7 are most important serovar of EHEC and carry potent infectious agent; so it is associated with many outbreak that influenced on global human health. EHEC can lead to haemolytic uremic syndrome (HUS). This development kidney disease (HUS) 5% leads to death despite treatment with antibiotic or rehydration. Type-III secretion system made this pathotype formidable and help bacteria to attach and transfer virulence protein into host cell. Bacteria first attach to host cell by flagella temporarily and HcpA play an important role in this stage, afterward Tir protein as a specific receptor make this adhesion permanent. Also EspA which is a part of needle apparatus help to translocate enterotoxins (like Stx) into host cell.

Methods: A chimeric protein inclusive of mentioned protein (HETS) was constructed and subcloned into PET32a Bl21(DE3) vector. After western blot which confirmed expressed protein, it purified by Ni-NTA column. Mice vaccinated with HETS protein and immune sera assessed for IgG. Finally neutralization and inhibition assays were performed versus O157.

Results: 1721bp DNA fragment encoding HETS cassette were successfully subcloned into pET32a vector. This sequence expressed by induced T7 promoter and associated 80kDa protein monitored on SDS-PAGE. Western blotting confirmed protein by using anti-His tag antibody. Immunological analyses showed production of a high titer of specific immunoglobulin in immunized animals. Adherence inhibition and neutralization assessments were shown immune sera are considerable functional in this regard.

Conclusion: development of this chimeric vaccine that focus on various virulence agent of only one bacteria can be literally functional and leads to broad vaccine coverage intention.

Keywords: Enterohaemorrhagic *E. coli*, chimeric protein, Immunization

ICIA2018\Tolerance and Autoimmunity\Poster\7781
7781

The Influence of PCSK9 on CD36 Expression on Monocytes in Patients with Hashimoto's Thyroiditis

M.S.Fazaeli¹, A.R.Khoshdel², M.R.Shafipoor³, M.Mahmoodi, M.R.Hajizade, M.A.Fahmidekar, M.Rohban

Department of Clinical Biochemistry Faculty of Medicine University of Medical Sciences

Background: Proprotein convertase subtilisin/kexin type 9 (PCSK9) is a serine protease and a secreted protein which increases cholesterol levels in plasma via inducing degradation of low-density lipoprotein receptor (LDLR). So PCSK9 serum levels correlate positively with the plasma LDL-C. Some studies have reported that PCSK9 levels increase in hypothyroidism. Cluster of differentiation 36 (CD36) is a member of a family of cell surface proteins in many cells. CD36 is known as fatty acid translocase (FAT) because it imports fatty acids inside cells. Therefore CD36 involves in regulation of lipid metabolism by facilitating cellular uptake of fatty acids and participating in triglyceride storage. It has been suggested that PCSK9 regulates CD36 in some tissues.

Methods: Data and serum levels of TSH, FT4 and PCSK9 and expression level of CD36 on monocytes from 20 new untreated patients with Hashimoto's thyroiditis in subclinical stages and 20 age- sex- and BMI-matched euthyroid controls were analyzed in a cross-sectional study. Quantification of CD36 expression on monocytes was done by flow cytometry. Then the relationships between these parameters were examined.

Results: PCSK9 was significantly higher in patients ($p = 0.001$). CD36 expression was lower in the patient group than in the control group, but this difference was not significant. There was no significant relationship between PCSK9 and CD36.

Conclusion: Currently, PCSK9 inhibitors are used to reduce blood cholesterol levels as drugs. If it will be proven that PCSK9 can reduce the level of CD36, taking these drugs may have unwanted side effects. This study showed that there is no relationship between PCSK9 and CD36 and PCSK9 antibodies can still be used as an effective way to treat hypercholesterolemia.

Keywords: Hashimoto's thyroiditis, Proprotein convertase subtilisin/kexin type 9, Cluster of differentiation 36, Thyroid stimulating hormone.

ICIA2018\Vaccine\Poster\7791

7791

Design, expression and purification of recombinant EtpA-LTB chimeric protein from Enterotoxigenic *Escherichia coli*

Moslem Kulivandi¹, Shahram Nazarian², Yousef Tarverdizadeh³

1. Department of Biology, Faculty of Science, Imam Hossein University, Tehran, Iran

2. Corresponding author: Department of Biology, Faculty of Science, Imam Hossein University, Tehran, Iran.
Tel: +98 21 77104934. E-mail address: kpnazari@ihu.ac.ir

3. Department of Biology, Faculty of Science, Imam Hossein University, Tehran, Iran

Background: Enterotoxigenic *Escherichia coli* (ETEC) is the most common bacterial causes of children's diarrhea. Because there are no effective vaccines against ETEC infection, investigating on vaccines against that consider necessary. In the classic paradigm of ETEC infection, these organisms adhere to the small intestinal mucosa via fimbrial colonization factor (CF) molecules, where they elaborate heatlabile enterotoxin and/or heat-stable enterotoxin. Recently novel antigens include adhesions EtpA and EaeH, and mucin enzymes are identified. Fusion of these novel antigens with classic antigens may help to producing effective candidate vaccines against ETEC infection. In this study we focused on expressing EtpA-LTB chimera as candidate immunogen for ETEC infection.

Methods: For optimum level expression of protein, the gene was synthesized with codon bias of *E. coli*. EtpA-LTB chimeric gene was subcloned in pET28 and expression of recombinant protein was induced by IPTG. The protein was purified with Ni-NTA column and confirmed by western blot.

Results: Percentage of codon having a frequency distribution of 91–100 in the native chimeric gene was 56%, which was significantly improved to 78% in the optimized gene sequence. Increasing CAI index to 81% means that appropriate codon is used for expression in *E. coli*. SDS-PAGE analysis showed the presence of a 44 KD recombinant chimeric protein. Purification of the proteins was achieved by Ni-NTA affinity chromatography. Western blot analysis by anti LT antibody, confirmed a single band of recombinant chimeric protein.

Conclusion: The chimeric protein can be considered as a candidate immunogen against ETEC infection.

Keywords: Enterotoxigenic *Escherichia coli*, LT toxin, Chimeric protein, Bioinformatic

ICIA2018\Stem Cells Based Immunotherapy\Poster\7824
7824

Exosomes from mesenchymal stem cells as a microRNA delivery system to mouse splenocytes

Zohreh Bolandi^{1,2}, Seyed Mahmoud Hashemi³, Sara Soudi⁴, Hossein Ghanbarian^{1,2*}

1. Department of Biotechnology, School of Advanced Technologies in Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Cellular and Molecular Biology Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

3. Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

4. Department of Immunology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

Background: MicroRNAs (miRNAs) are a type of small non-coding RNA that control gene expression at the posttranscriptional level. miRNAs have a critical role in many biological activities including cell proliferation, apoptosis and inflammation. Recent studies showed that miRNAs stably present in different extracellular biofluids (e.g., blood or urine) and vesicles (e.g., exosomes or microvesicles). Exosomes protect miRNAs from digestion and exosomal miRNAs can be taken up by cells and subsequently modulate their gene expression. Due to these properties, exosomes have attracted attention as an efficient cell-free vector for gene delivery. The aim of this study was to determine whether the exogenous miRNAs could be transferred to mouse splenocytes via adipose-derived mesenchymal stem cells (AD-MSCs) exosomes.

Methods: AD-MSCs were cultured in DMEM with 1% ITS (insulin-transferrin-Selenium) and culture media was harvested after 72hrs. Exosomes were then isolated and characterized by dynamic light scattering (DLS) and scanning electron microscopy (SEM). Exosomes were loaded with miRNA mimic by electroporation in 0.4cm electroporation cuvettes. Splenocytes were isolated from 8 weeks old female C57BL/6 and were cultured in RPMI-1640 plus 10% FBS in presence of the loaded exosomes. After 24 or 120 hrs, RNA was extracted and qPCR was performed.

Results: For transfer of miRNA to splenocytes, exosomes were isolated from AD-MSCs. SEM and DLS analysis revealed spherical structures with a diameter of 117nm. After exosomes electroporation, splenocytes were cultured with the loaded exosomes. RNA was extracted and qPCR determination showed that miRNA were overexpressed significantly in splenocytes.

Conclusion: Here, we showed that AD-MSCs exosomes can deliver exogenous miRNA to mouse splenocytes. Loaded Exosomes provides a new platform for genetic information transfer and gene therapy of immune cells or modulate immune responses including immune stimulation and tolerance induction.

Keywords: Splenocytes, Exosomes, miRNA.

ICIA2018\Vaccine\Poster\7881

7881

Determination of tetanus immunization in pregnant woman in Zanjan province, Northwest of Iran

SafaTahmasebi¹, Sousan Ghasemi², Maryam Erfanmanesh³, Abdolreza Esmaeilzadeh^{*4, 5}

1. MSc student of Immunology, Tehran University of Medical Sciences, Tehran, Iran

2. Medical Laboratory of Shaheed Beheshti Hospital, Zanjan University of Medical Sciences, Zanjan, Iran

3. Young Researchers Club, Zanjan Branch, Islamic Azad University, Zanjan, Iran

Department of Immunology,⁵CancerGeneTherapyResearch Center, Faculty of Medicine, Zanjan University of Medical Sciences, Zanjan, Iran

Background: Despite the fact that neonatal Tetanus has been eradicated in all over the world, it is still a significant health problem in some developing countries. It could be even resulted in death. This disease is a worrying problem but it could be easily prevented by vaccination of mother. Vaccinating every pregnant woman with at least one dose of TT would be an affordable way to protect against neonatal tetanus, and would be a step toward eliminating the deaths that continue to occur due to this preventable disease.

Methods: The purpose of this study was evaluation of immunity in 576 pregnant women in Zanjan province, Iran. These women's serums were separated. Then tetanus antibody was measured in their serums using Immunoaffinity Chromatography Assay (IACH).

Results: Results indicated that 95 percent of these pregnant women which had received vaccine had acquired immunity and the remained 5 percent of them had no immunization.

Conclusion: In conclusion, not only vaccination is important in women, but also effectiveness of vaccination is vital for successful immunization. It is proposed that more attempts should be done in order to gain affordable immunization.

Keywords: Tetanus antibody, vaccination, pregnant women, Immunoaffinity Chromatography Assay, Iran

ICIA2018\Vaccine\Poster\7898
7898

A novel cell penetrating peptide for efficient delivery of HPV16 E7 antigen *in vitro*

Sepideh Shahbazi¹, Azam Bolhassani^{1*}, Fatemeh Motevalli¹

Department of Hepatitis and AIDS, Pasteur Institute of Iran, Tehran, Iran

Background: The E7 protein has been suggested as a candidate for generating protein vaccines against human papillomavirus infections. The cell penetrating peptides has been suggested for delivery of proteins and peptides *in vitro* and *in vivo*. In this study, the efficiency of a amphipathic peptide, hPP10, for delivery of the fluorescent HPV16 E7 protein was evaluated in HEK-293T cells.

Methods: Cloning of the recombinant hPP10-E7-GFP DNA performed in the pET-28a vector. After confirmation of the recombinant clones using digestion and PCR. The expression of hPP10-E7-GFP protein was induced by IPTG and purified by affinity chromatography and confirmed using SDS-PAGE and western blotting. HEK-293T cells were transfected with hPP10-E7-GFP and E7-GFP/TurboFect. The efficiency of hPP10 CPP and TurboFect reagent was compared by fluorescent microscopy and flow cytometry at 3 and 24 hours after transfection.

Result: The recombinant hPP10-E7-GFP gene construct was confirmed by PCR and digestion. The results showed a ~ 1100 bp band in gel electrophoresis. The purified hPP10-E7-GFP protein with apparent molecular weight of ~ 50 KDa, was recognized by SDS-PAGE and Western blotting. Flow cytometry results showed that the transfection efficiency of E7-GFP protein was ~ 63.66% and ~ 32.95% for hPP10 CPP and TurboFect, respectively. The cells transfected by hPP10-E7-GFP and E7-GFP/TurboFect demonstrated the green regions with different distribution using fluorescent microscopy.

Conclusion: The expression of the hPP10-E7-GFP protein was performed in *E. coli* system. *In vitro* transfection results showed the transfection of hPP10 was significantly higher than TruboFect commercial reagent. Thus, hPP10 can be considered as a promising delivery system for therapeutic purposes in Future.

Keywords: HPV16, E7, CPP, hPP10

ICIA2018\Tolerance and Autoimmunity\Poster\7945

7945

Survey of Alpha-1 Antitrypsin Activity in Patients with Multiple Sclerosis

Giti Farsi¹, Alireza Khoshdel², Abbas Lotfi³, Amir Moghadam Ahmadi⁴, Mahmood Sheikhfatollahi⁵, Mohammadreza Hajizadeh⁶, Mehdi Mahmoodi⁶, Fatemeh Ayoobi⁷

1. Department of Clinical Biochemistry, Faculty of Medicine, Rafsanjan University of Medical Sciences, Rafsanjan, Iran

2. Department of Clinical Biochemistry, Faculty of Medicine, and Pistachio Safety Research Center, Rafsanjan University of Medical Sciences, Rafsanjan, Iran

3. Department of Clinical Biochemistry, Faculty of Medicine, Tarbiat modares University of Medical Sciences, Tehran, Iran

4. Neurologist, Rafsanjan University of Medical Sciences, Rafsanjan, Iran

5. Department of Epidemiology and Biostatistics, School of Medicine, Rafsanjan University of Medical Sciences, Rafsanjan, Iran

6. Molecular Medicine Research Center and Department of Clinical Biochemistry, Faculty of Medicine, Rafsanjan University of Medical Sciences, Rafsanjan, Iran

7. Physiology-Pharmacology Research Center, Rafsanjan University of Medical Sciences, Rafsanjan, Iran

Background: Alpha-1 Antitrypsin (AAT) is the most abundant serine-protease inhibitor in blood which impedes degradation of the tissues by its anti-protease activity. Any lack or imperfection of this protein remarkably intensifies the risk of various diseases including pulmonary diseases, hepatitis, cirrhosis, autoimmune diseases such as multiple sclerosis. Multiple sclerosis (MS) is an inflammatory, degenerative disorder with biological associations. Hypothetically, AAT may play an essential role in mentioned disorder. The most important objective of the present study is to determine the association between AAT activity and Multiple Sclerosis degree.

Materials and Methods: In this descriptive study, the activity of AAT was determined using sera, by Trypsin Inhibitory Capacity (TIC). Patients with MS were identified from 42 patients included in the specific diseases center. Also, 42 healthy volunteers (controls) were recruited. Data was analyzed by SPSS software.

Results: The average activity of AAT (TIC degree) in MS patients was significantly lower than controls (2.23 $\mu\text{mol}/\text{min}/\text{ml}$). In addition, TIC levels in MS patients in accord with age and sex was meaningfully lower than healthy individuals ($p < 0.001$). Moreover, there was a reverse correlation between disease duration and AAT activity ($r = -0.481$, $p = 0.001$).

Conclusion: Alpha-1 Antitrypsin activity in patients with MS was lower than the controls. Since alpha-1 antitrypsin is important in controlling inflammation process in response to environmental exposure (free radicals, smoking, etc.), such disorders may be due to abnormalities of this protein. Efforts should focus on early recognition and effective interventions for AAT activity in individuals with MS. Detection and changing of lifestyle as soon as possible continues decisive to the management off AAT deficiency.

Keywords: Alpha-1 Antitrypsin, Multiple sclerosis, Trypsin Inhibitory Capacity

ICIA2018\Vaccine\Poster\7972

7972

Formulation of polio vaccine deactivated by gamma irradiation and comparing its immunogenicity with an Iranian commercial vaccineMaryam mollaee alamooti¹, mehrdad ravanshad², farahnaz motamedi³

1. Department of virology, School of medical science, University of tarbiat modares

2. Department of virology, School of medical science, University of tarbiat modares

3. Research Institute of Beam, Research Institute of Nuclear science and technology

Background: poliomyelitis (polio) is caused by various poliovirus serotypes. A live vaccine that is weak commercially exists, in some cases as a result of vaccine-induced paralysis. Therefore, the production of low-cost inactivated vaccines is important and necessary to eradicate polio. In this study, the inactivated vaccine by gamma-ray radiation was compared with the common polio vaccine in the mouse model.

Methods: First, the virus titre of the oral polio vaccine was determined, then deactivated by the appropriate dose of gamma radiation and formulated to form a radio vaccine. BALB/c mice received two different formulations of the intramuscular vaccine in two week intervals. Two weeks after the final injection, blood samples were collected from the mice and the humoral and cellular immune responses were evaluated.

Results: Humoral immune responses using neutralizing anti-serum titers of mice vaccinated with Vaccine conventional, Radio Vaccine showed a significant difference in some dilutions. Also, the study of cellular immunity using MTT test and statistical analysis of the data by Anova showed that the induction of cellular immunity in the radio vaccine group was significantly increased.

Conclusion: The anti-serum titre of the inactivated vaccine by gamma irradiation showed that it could produce proper humoral immunity in mouse groups.

Key words: Polio, vaccine, gamma radiation, immune responses

ICIA2018\Tolerance and Autoimmunity\Poster\7990

7990

Effects of Gliadin Administration on Gene Expression Profile of Chemokines in Central Nervous System of Mice

Parisa Esmaeilie¹, Hamed Bashiri, Shohreh Fakhari^{1,2}, Maryam Moradi¹

1. Department of Immunology and Microbiology, Faculty of Medicine, Kurdistan University of Medical Sciences, Sanandaj, Iran

2. Cellular and Molecular Research Center, Kurdistan University of Medical Sciences, Sanandaj, Iran.

Background: Gluten is the most widely used food to trigger development of multiple autoimmune diseases and neurological disorders. Recently, an association between anti-gliadin serum antibodies and CNS autoimmune diseases such as MS has been reported. Concerning the involvement of chemokines synthesized in activated astrocytes and microglial cells in cellular infiltration and brain defense mechanisms, the present study was conducted to investigate chemokines gene expression in CNS of mice immunized with gliadin.

Methods: Healthy female 6-8 weeks old C57BL/6 mice were assigned into 3 groups (N=6). Mice in group 1 were injected subcutaneously (sc) with PBS (400µl). In group 2, mice were immunized sc with complete freundadjuvant (CFA;400µl). In group 3, mice were immunized sc with peptic-tryptic-gliadin (300µg) emulsified in CFA (400µl). Boosters containing the same amount of antigen were injected on days 7 and 14. On day 28, mice were sacrificed and brain and spinal cord tissue were removed. RT-PCR was used to evaluate the mRNA expression of MCP-1, CXCL2, and CXCL10 chemokines.

Results: mRNA expression for MCP-1 and CXCL2 were increased significantly ($p<0.05$) in group 3 in comparison to control groups (group 1 and 2). Conversely, the level of CXCL10 expression in group 3 in comparison to group 1 and 2 was significantly ($p<0.05$) up-regulated and down-regulated respectively.

Conclusion: The changes in gene expression profile of chemokines may be involved in gliadin induced CNS pathology.

Keywords: Gliadin, MCP-1, CXCL2, CXCL10

ICIA2018\Research and Development\Poster\8010

8010

Gene expression and levels of IL-6 and TNF α in PBMCs correlate with severity and functional class in patients with chronic heart failure

V. Eskandari¹, A. A. Amirzargar¹, M. J. Mahmoudi², Z. Rahnemoon³, F. Rahmani^{4,5}, S. Sadati¹, Z. Rahmati¹, F. Gorzin¹, M. Hedayat^{6,7}, N. Rezaei^{1,4,5}

1. Department of Immunology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

2.. Department of Cardiology, Amir Alam Hospital, Tehran University of Medical Sciences, Tehran, Iran

3. Cardiac Heart Center, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

4. Research Center for Immunodeficiencies, Pediatrics Center of Excellence, Children's Medical Center, Tehran University of Medical Sciences, Tehran, Iran

5. Network of Immunity in Infection, Malignancy and Autoimmunity (NIIMA), Universal Scientific Education and Research Network (USERN), Tehran, Iran

6. Division of Immunology, Boston Children's Hospital, Harvard Medical School, Boston, MA, USA

7. Network of Immunity in Infection, Malignancy and Autoimmunity (NIIMA), Universal Scientific Education and Research Network (USERN), Boston, MA, USA

Background: Evidence shows that proinflammatory cytokines are important determinants of assessment of severity and prognosis of chronic heart failure (CHF).

Methods: In this report, we used real-time PCR assay to compare relative gene expression of TNF α and IL-6 in PBMC from CHF patients with various heart diseases (n = 42, EF < 45%, NYHA I to IV) and matched healthy control subjects (n = 42). We also determined the TNF α and IL-6 concentrations of cell culture supernatant of PBMCs with ELISA.

Results: There was a significant negative correlation between gene expression of TNF α and LVEF (r = 0.4, p < 0.05). Patients with CHF had increased gene expression of TNF α and IL-6 in PBMCs (p < 0.05). They also had elevated the supernatant levels of these cytokines in cultured PBMCs (p < 0.001). Levels of TNF α and IL-6 were increased in ischemic heart disease compared to non-ischemic heart disease. There was a positive correlation between TNF α and IL-6 levels in CHF patients and severity of CHF in patients. Levels of these cytokines were higher in patients with NYHA III-IV than in NYHA I-II and normal subjects.

Conclusions: Results of this study indicate that peripheral expression of proinflammatory cytokines, TNF- α and IL-6, is important indicators of severity and prognosis in patients with chronic heart disease.

Keywords: Heart failure; Proinflammatory cytokines; Tumor necrosis factor alpha; Interleukin 6

ICIA2018\Vaccine\Poster\8044

8044

Development of a multi-epitope peptide vaccine inducing robust T cell responses against Brucellosis using immunoinformatics based approaches

Ahmad Karkhah^{1,2}, Mahdiye Saadi^{1,2}, Hamid Reza Nouri^{1*}

1. Cellular and Molecular Biology Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, Iran

2. Student Research Committee, Babol University of Medical Sciences, Babol, Iran

Background: Current investigations have demonstrated that a multi-epitope peptide vaccine targeting multiple antigens could be considered as an ideal approach for prevention and treatment of brucellosis. According to the latest findings, the most effective immunogenic antigens of brucella to induce immune responses are included Omp31, BP26, BLS, DnaK and L7-L12. Therefore, in the present study, an *in silico* approach was used to design a novel multi-epitope vaccine to elicit a desirable immune response against brucellosis.

Methods: First, five novel T-cell epitopes were selected from Omp31, BP26, BLS, DnaK and L7-L12 proteins using different servers. In addition, helper epitopes selected from Tetanus toxin fragment C (TTFrC) were applied to induce CD4⁺ helper T lymphocytes (HTLs) responses. Selected epitopes were fused together by GPGPG linkers to facilitate the immune processing and epitope presentation. Moreover, cholera toxin B (CTB) was linked to N terminal of vaccine construct as an adjuvant by using EAAAK linker. A multi-epitope vaccine was designed based on predicted epitopes which was 377 residues in length. Then, the physico-chemical properties, secondary and tertiary structures, stability, intrinsic protein disorder, solubility and allergenicity of this multi-epitope vaccine were assessed using immunoinformatics tools and servers.

Results: Based on obtained results, a soluble, and non-allergic protein with 40.597 KDa molecular weight was constructed. Expasy ProtParam classified this chimeric protein as a stable protein and also 89.8% residues of constructed vaccine were located in favored regions of the Ramachandran plot. Furthermore, this multi-epitope peptide vaccine was able to strongly induce T cell and B-cell mediated immune responses.

Conclusion: In conclusion, immunoinformatics analysis indicated that this multi-epitope peptide vaccine can be effectively expressed and potentially be used for prophylactic or therapeutic usages against brucellosis.

Keywords: Brucellosis, multi-epitope vaccine, Omp31, BP26 BLS, Dnak, L7-L12

ICIA2018\Tolerance and Autoimmunity\Poster\8131
8131

Association between gene expression of Interlukine-35 subunits with plasmaneopterin and disease activity in rheumatoid arthritis patients

Nasrin Iranshahi¹, shirin Assar², Seyed Mojtaba Amiri³, Parisa Zafari^{4,5}, Mahdi Taghaddosi*

1. Department of Immunology, Kermanshah University of Medical Sciences, Kermanshah, Iran

2. Assistance professor of Rheumatology in Clinical Research Development Center, Imam Reza Hospital, Kermanshah University of Medical Sciences, Kermanshah, Iran

3. Department of Biostatistics and Epidemiology, School of health, Kermanshah University of Medical Sciences, Kermanshah, Iran

4. PhD student in department of Immunology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

5. Student Research Committee, Mazandaran University of Medical Sciences, Sari, Iran

*Assistance professor in department of Immunology, Kermanshah University of Medical Sciences, Kermanshah, Iran

Background: Interleukin-35 (IL-35), is the member of IL-12 family that produced by T regulatory (Treg) cells. This cytokine has immunosuppressive properties which can inhibit overt inflammatory responses such as those associated with rheumatoid arthritis. This study aims to determine the correlation between gene expression of IL-35 subunits in peripheral blood with immunological parameters and disease activity in rheumatoid arthritis patients.

Methods: Peripheral blood samples were collected from 47 patients and 44 of healthy subjects. Plasma levels of Neopterin and Anti cyclic citrullinated peptide (Anti-CCP) have been evaluated using ELISA method. Gene expression of FoxP3, IL-35 subunits (IL-12A/EBI-3) analyzed by qPCR. Also RF detected by agglutination test and clinical variables were extracted by physical examination.

Results: In this study, there was no correlation between gene expression of EBI-3, IL-12A and FoxP3 with disease activity score-28 (DAS-28), ($P=0.234$), ($P=0.634$) and ($P=0.457$) respectively but the significant negative correlation was observed between the plasma levels of neopterin and EBI-3 gene expression ($P=0.004$).

Conclusion: With respect to negative correlation between EBI-3, the subunit of immunosuppressive cytokine IL-35 with plasma neopterin, the inhibitory effect of this cytokine on inflammatory process in RA patients can be concluded.

Keywords: Rheumatoid arthritis, Interlukine-35, Neopterin, DAS-28, FoxP3

ICIA2018\Tolerance and Autoimmunity\Poster\8137
8137

Cold Autoimmune Hemolytic Anemia due to High-grade non-Hodgkin's B cell Lymphoma with Weak Response to Rituximab and Chemotherapy Regimens

Behzad Nazel Khosroshahi,^{1,2} Mohammad Jafari,³ Hossein Vazini,⁴ Alireza Ahmadi⁵, Keivan Shams⁶, Mahdi Kholoujini⁷

1. Emam Reza Hospital, Hamadan University of Medical Sciences, Hamadan, Iran

2. Blood transfusion research center, High institute for research and Education in Transfusion Medicine, Tehran, Iran

3. Department of Pathology, School of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

4. Department of nursing, Hamedan branch, Islamic Azad University, Hamedan, Iran

5. Department of Hematology and Oncology, School of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

6. Department of Immunology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

7. Department of Mycology, School of Medicine, Tarbiat Modares University, Tehran, Iran

Background: Autoimmune hemolytic anemia (AIHA) is characterized by shortening of red blood cell (RBC) survival and the presence of autoantibodies directed against autologous RBCs. Approximately 20% of autoimmune hemolytic anemia cases are associated with cold-reactive antibody. About half of patients with AIHA have no underlying associated disease; these cases are termed primary or idiopathic. Secondary cases are associated with underlying diseases or with certain drugs. We report herein a rare case of cold autoimmune hemolytic anemia due to high-grade non-Hodgkin's lymphoma of B-cell type with weak response to rituximab and chemotherapy regimens. For treatment B cell lymphoma, Due to lack of treatment response, we used chemotherapy regimens including R- CHOP for the first time, and then Hyper CVAD, R- ICE and ESHAP were administered, respectively.

Methods: For treatment of autoimmune hemolytic anemia, we have used the corticosteroid, rituximab, plasmapheresis and blood transfusion and splenectomy.

Results: In spite of all attempts, the patient died of anemia and aggressive lymphoma nine months after diagnosis.

Conclusion: To our knowledge, this is a rare report from cold autoimmune hemolytic anemia in combination with high-grade non-Hodgkin's lymphoma of B-cell type that is refractory to conventional therapies.

Keywords: Cold autoimmune hemolytic anemia, High-grade, non-Hodgkin's lymphoma, Rituximab, Chemotherapy

ICIA2018\Tolerance and Autoimmunity\Poster\8160

8160

Investigation of Chosen Polymorphisms in Forkhead box P3 (FoxP3) Gene in Patients with Hashimoto's Thyroiditis.

Mahsa Eshkevar Vakili¹, Solmaz Khansalar¹, Danial Ghasemi¹, Farshid Mossallaie¹, Zahra Amirghofran^{1,2}, Kurosh Kalantar¹

1. Department of Immunology, School of medicine, Shiraz University of Medical Sciences, Shiraz, Iran

2. Autoimmune Disease Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

Background: Loss of immune tolerance to thyroid autoantigens leads to autoimmune thyroid diseases such as Hashimoto's thyroiditis (HT). A subset of T cells that promote immune tolerance to autoantigens, called regulatory T cell (Treg). Reduced number and function of Tregs result in abnormal responses against autoantigens and subsequently cause autoimmune diseases. These cells modulate the immune responses. Forkhead box P3 (FoxP3) is a master transcription factor of Treg cells which is pivotal for maintaining and function of these cells. In this study, we investigated the association of two polymorphisms of FoxP3 gene in Hashimoto's thyroiditis (HT).

Methods: The study was performed in the group consisting of 110 patients with Hashimoto's thyroiditis (HT) recruited from the Motahari Clinic which is affiliated to Shiraz university of medical science, (mean age: 38.1±01years) and 105 healthy subjects (mean age: 44.4±2.2years). DNA was extracted from the peripheral blood leukocytes using the salting-out method. The two single nucleotide polymorphisms (SNPs) including rs3761549 (-2383C/T) and rs376154 (-3279A/C) in the FoxP3 gene. Genotyping were done by polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP).

Results: About rs3761549 (-2383C/T), there wasn't a significant difference in frequency of alleles ($v_{\text{alue}} > 0.05$) and genotype ($v_{\text{alue}} > 0.05$) between patients and control. In regards to the rs3761548 (-3279A/C) the same results obtained and no difference observed respect to the allele and genotype among studied subjects.

Conclusion: We have not shown association between FoxP3 polymorphisms and susceptibility to Hashimoto disease.

Keywords: Autoimmune thyroid diseases, Regulatory T cell, Forkhead box P3 (FoxP3), Single nucleotide polymorphism (SNPs)

ICIA2018\Tolerance and Autoimmunity\Poster\8221

8221

The Association Circulating IL-17 Levels and IL-23 Receptor Gene Polymorphisms with Multiple Sclerosis

Seyed Ali ghaffarikhalfmohammadi^{1,2}, AbdollahJafarzadeh^{1,2}, Abdolkarim Sheikhi³

1. Department of Immunology, Medical School, kerman University of Medical Sciences, kerman, Iran

2. Neurology Research Center, Department of Neurology, Kerman University of Medical Sciences, Kerman, Iran

3. Department of Immunology, Medical School, Dezful University of Medical Sciences, Dezful, Iran

Background: IL-17/IL-23 axis plays a prominent role in the pathogenesis of a number of autoimmune diseases. The aim of this study was to investigate the IL-17 levels in MS patients and its association with gender, treatment, disease patterns and single nucleotide polymorphisms (SNP) in *IL-23R* gene, including rs11209026 and rs1004819.

Methods: The blood samples were collected from 135 MS patients and 135 healthy subjects. The patients have relapsing-remitting (RRMS; n=65), primary progressive (PPMS; n=19), secondary progressive (SPMS; n=35) or progressive relapsing (PRMS; n=14) patterns. The DNA analyzed for SNPs using PCR-RFLP and IL-17 levels measured by ELISA.

Results: The serum IL-17 levels in MS patients was significantly higher than in control group ($P<0.001$). The IL-17 levels were significantly higher in MS men as compared to women patients ($P<0.05$). Untreated patients had significantly higher IL-17 levels than healthy group and treated patients ($P<0.001$ and $P<0.01$, respectively). The IL-17 levels in treated patients with interferon- β (IFN- β), methylprednisolone (MP) or both IFN- β and MP were significantly lower than untreated MS patients ($P<0.05$, $P<0.02$ and $P<0.05$, respectively). Patients with RRMS and PRMS, had significantly higher IL-17 levels than healthy group ($P<0.005$ and $P<0.01$, respectively). No significant differences were observed between patients and controls regarding the genetic variations at rs11209026 and rs1004819. The levels of IL-17 did not influenced by genetic variations at investigated SNPs.

Conclusion: These results indicated higher levels of IL-17 in MS patients that may be influenced by disease patterns, treatment and gender. There was no any association between investigated SNPs and MS.

Keywords: Multiple sclerosis, IL-17, IL-23 receptor, Gene polymorphisms.

ICIA2018\Vaccine\Poster\8250

8250

Naloxone reinforced immune response in compared to HBS antigen formulated with Montanide ISA-720Mina Mirzaee¹, Setareh Haghighat ², Mehdi Mahdavi ^{3,*}*1. Department of Microbiology, Pharmaceutical Sciences Branch, Islamic Azad University, Tehran, IR Iran**2. Department of Microbiology, Pharmaceutical Sciences Branch, Islamic Azad University, Tehran, IR Iran**Department of Immunology, Pasteur Institute of Iran, Tehran, Iran*

Background: Utilization of immunostimulatory factors to promote the Immune system have captured the attention of the scientists in the field of vaccine and have resulted in improvement of vaccines potency. Hepatitis vaccine and its development is one of the ongoing projects of the country carried out by scientists and injected clinically, but it has a main problem being its deficiency to follow a positive response to all people. In this study, Naloxone as an immunopotentiator is used in an oil-based Montanide ISA-720 formulation of hepatitis vaccine to promote higher cellular immune responses.

Methods: The HBs Ag (5µg/dose) was formulated in Montanide ISA-720 adjuvant by hemojenizer and Naloxone was added to the Formulation with final concentration of 5 mg/kg and 10 mg/kg of experimental mice. Then, the prepared formulation was injected to the BALB/C mice three times with two week interval and the cytokine responses of IL-4, IFN-γ, TNF-α, IL-2, total IgG antibody and IgG1 and IgG2a isotypes were assessed via ELISA method.

Results: The results indicated that addition of Naloxone to the final Formulation was followed by an enhancement of IFN- γ, IL-2 responses against control groups (P<0.05) and versus commercial vaccine.

Conclusion: Naloxone could be used as immunopotentiator in mixture adjuvant for induction of Th1 immunologic platform

Keywords: HBS-Ag¹, vaccine², Adjuvant³, Naloxane⁴, Montanide, ISA720⁵

ICIA2018\Stem Cells Based Immunotherapy\Poster\8275

8275

Developing Rat Bone Marrow derived mast cells by the Splenic Cells Culture Supernatant of rat and mouse

Amani Saeede ¹, Shahrooz Rasoul ^{2,*}, Karimi Ali¹, Bakhtiari Zahra ¹, Asl iranifam Neda ¹, Mortaz · Esmail ^{3,4,5,*}

1. PhD student, Comparative Histology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

2. Associate Professor, Anatomy sciences, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

3. Clinical Tuberculosis and Epidemiology Research Center, National Research Institute of Tuberculosis and Lung Diseases (NRITLD), Shahid Beheshti University of Medical Sciences, Tehran, Iran

4. Department of Immunology, Faculty of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

5. Division of Pharmacology, Utrecht Institute for Pharmaceutical Sciences, Faculty of Science, Utrecht University, Utrecht, The Netherlands

Background: Mast cells are playing a crucial role in the pathogenesis of diverse diseases that include MC-driven disorders. The aim of the current study was to develop simple and cost-efficient method for differentiation and culture of rat mast cells from bone marrow by using rat and mouse spleen supernatant.

Methods: Bone marrow cells from male 10 to 15-weeks-old rats was obtained through femur bone and cultured for three weeks at complete cell culture medium. After that, the purity of cells was confirmed by flowcytometry, toluidine blue and immunohistochemistry (IHC).

Results: After 3 weeks continuous culturing rat bone marrow cells with spleen supernatant of rat and mice, we found high purity of cells in expression of specific lineage markers of cells. Results similar to each other be derived for both spleen supernatants of rat and mouse. Specific markers for differentiated bone marrow stromal stem cells. CD117, CD34 markers and tryptase were positive 80.1%, 76.89% and 87.9%, respectively, with rat splenic supernatant whereas were positive 85.4%, 83.07% and 82.1%, respectively for cells which cultured with mouse splenic supernatants. IHC result is confirmed by flowcytometry analysis. In this case with rat supernatant of spleen (89.1%) and with mouse splenocyte supernatant (91.8%) double positive for FCεRI and CD117 markers were obtained.

Conclusion: By modification rat standard rat/mouse bone marrow isolation by supernatant spleen of both animal, mast cells were obtained. Finding the differences of two cells from two species is important in mast cell biology. In the current study, by modification of protocol we achieved high-yield mast cells by using rat splenic cells mitogens and changing concentration with culturing cells in density of 3×10^5 - 1×10^6 cells/ml whereas other data of old yield which published were 1×10^5 , 1×10^6 .

Keywords: Bone Marrow derived mast cells, Cells Culture Supernatant, flowcytometry analysis

ICIA2018\Veterinary Immunology\Poster\8308

8308

Experimental study immunomodulatory effects of *Rosa canina* L. on male rats

Saeede Sadat Mirbageri¹, Hamidreza Hakkakifard², Pouyan houshmandpour³

1. *Young Researchers and Elite Club, Tabriz Branch, Islamic Azad University, Tabriz, Iran*

2. *DVM, Department of Veterinary Clinical Sciences, Tabriz Branch, Islamic Azad University, Tabriz, Iran*

Methods: In this study, 30 male rats were obtained and divided into three groups of 10 rats. First group received 0.9% saline (10 mg/kg), the second group *Rosa canina* L. fruit extract (250 mg/kg) and third group *Rosa canina* L. fruit extract (500 mg/kg) as oral gavages every day for a period of one month. After obtaining blood samples, differential white blood cell (WBC) counts, phagocyte activity (number) of samples were obtained.

Results: Neutrophil and monocyte counts and phagocyte activity significantly increased in comparison with the normal saline group.

Conclusion: The data suggest that the *Rosa canina* L. fruit extract used in traditional medicine might have immunomodulatory effects.

Keywords: *Rosa canina*, Rat, immunomodulator, phagocyte, white blood cells.

ICIA2018\Stem Cells Based Immunotherapy\Poster\8350
8350

Evaluation of Mesenchymal Stem Cells Immunomodulatory Effects on Cytokine Production in Experimental Chronic Colitis

Maryam Heidari¹, Seyed Mahmoud Hashemi^{1,4*}, Sedigheh Pouya¹, Kaveh Baghaei², Mohammad Reza Zali³

1. Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran
2. Basic and Molecular Epidemiology of Gastrointestinal Disorder Research Center, Research Institute for 3. Gastroenterology and Liver Disease, Shahid Beheshti University of Medical Science, Tehran, Iran
4. Gastroenterology and Liver Disease Research Center, Research Institute for Gastroenterology and Liver 5. Disease, Shahid Beheshti University of Medical Science, Tehran, Iran
6. Department of Applied Cell Sciences, School of Advanced Technologies in Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Inflammatory bowel disease (IBD) as a chronic recurrent disorder is characterized by mucosal immune response dysregulation; mesenchymal stem cells (MSCs) are the multipotent cells, which can be effective in immunomodulation via release of anti-inflammatory mediators. In this study, the effects of MSCs were evaluated in Dextran Sodium Sulfate (DSS) induced colitis.

Methods: Chronic colitis was induced in female C57BL/6 mice using 2% DSS in drinking water for 3 cycles, in each cycle, there were 4 days of DSS solution administration followed by 7 days of DSS-free water administration. MSCs, 10⁶ cells per mouse, were injected intraperitoneally (IP) on the last day of DSS in cycles 2 and 3. Colitis Clinical symptoms including body weight changes, bleeding, stool consistency were daily recorded. The colon macroscopic and pathological changes were then analyzed. The IL-17, IL-10 and TGF- β cytokine levels were also evaluated in spleen, mesenteric lymph nodes and serum of mice.

Results: After receiving MSCs in colitis mice, the clinical symptoms and disease activity index (DAI) were improved and the survival rate was increased. In MSCs-treated group, the colon length reduction was mild. The histopathological examination also showed tissue healing and less leukocyte infiltration in comparison to the non-treated group. Additionally, ELISA results indicated, increased level of transforming growth factor-beta (TGF- β), increased level of interleukin (IL)-10, and decreased level of IL-17 following the treatment.

Conclusion: This study demonstrated the ability of MSCs to regulate the inflammatory immune responses in chronic colitis. Production of anti-inflammatory cytokines by MSCs lead to immunosuppression, tissue healing and inflammatory cytokines reduction.

Keywords: Inflammatory Bowel Disease, Dextran Sodium Sulfate, Mesenchymal Stem Cell, Colitis Mouse Model.

ICIA2018\Reproductive Immunology\Poster\8355
8355

Evaluation of TH1 & TH2 cytokines and pregnancy hormones in pregnant women with a history of recurrent abortion and normal pregnancy

Saber Rokhafrooz¹, Mehri Ghafourian^{2,*}, Nasrin Saharkhize³, Sara Iranparast⁴

1. Department of Immunology, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

2. Fertility, Infertility and Perinatology Research Center, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

3. Infertility and In Vitro Fertilization Center, Taleghani Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran

4. Department of Immunology, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

Background: Spontaneous abortion is one of the most common complications of pregnancy. Immunologic factors have been proposed to contribute to the etiology of recurrent abortion. We aimed to measure cytokines in women with a history of recurrent abortion in the first trimester of pregnancy until termination of pregnancy. Moreover, we assessed the relationship between T-helper cytokines and hormones (progesterone and human chorionic gonadotropin [hCG]), involved in normal pregnancy process in the above mentioned groups.

Methods: In this case-control study, normal pregnant women (control group, n=30) and those with history of recurrent abortion (case group, n=43) in the first trimester were enrolled. We assessed TH2 cytokines (IL-4, IL-6 and IL-10) and TH1 cytokines (TNF α and INF γ) with enzyme-linked immunosorbent assay. Karyotyping was performed on the women and their spouses to detect chromosomal abnormalities. We also monitored the case group until the end of their pregnancy.

Results: The mean concentrations of TH2 cytokines differed significantly in women with a history of recurrent abortion whose pregnancy continued successfully compared with those whose pregnancy failed ($P<0.05$). However, no significant difference was observed between the case and control groups in this regard. The concentration of TH1 cytokines including TNF α and INF γ increased in women with recurrent miscarriage whose pregnancy was unsuccessful compared with normal pregnant women. This difference was significant in TNF α levels ($P<0.05$). We found no correlation between hCG levels and TH1/TH2 trends, while a correlation was observed between progesterone levels and TH2 cytokines ratio.

Conclusion: A higher TH2 cytokine level is seen in women with recurrent abortion who have a successful pregnancy, while this trend shifted toward a higher TH1 level in those with unsuccessful pregnancy. Cytokine assessment could be proposed as a marker for predicting success or failure of pregnancy in women with recurrent abortion during the first trimester of pregnancy.

Keywords: Recurrent abortion, TH1&TH2 cytokines, hCG, Progesterone

ICIA2018\Veterinary Immunology\Poster\8371

8371

Investigation of the leukogram and tumor necrosis factor- α in native neonatal calves with diarrhea of Sanandaj area, west of Iran

Mahdieh Raeeszadeh^{1*}, Javad Beheshtipour²

1. Assistant Professor in Pharmacology, Department of Basic Sciences, Sanandaj Branch, Islamic Azad University, Sanandaj, Iran

2. DVM Student, Sanandaj Branch, Islamic Azad University, Sanandaj, Iran

Background: Neonatal calf diarrhea is a sickness affecting the newborn calf under 4 weeks of age. Multiple enteric pathogens (e.g., viruses, bacteria, and protozoa) are involved in the development of this disease. Tumor necrosis factor- α (TNF- α) is secreted chiefly by macrophages. TNF- α possesses a wide range of biological activities, including fever induction; direct activation of neutrophils, T cells, and macrophages; direct tumor cell cytotoxicity; cachexia; and induction of other mediators (e.g., IL-1 and IL-6). Hence, the aim of this study was to determine the leukogram and plasma concentration of TNF- α in native neonatal calves with diarrhea of Sanandaj region.

Methods: This study was conducted during March to July 2017 in Sanandaj area, west of Iran. All native calves under 30 days of age displaying diarrhea were included in the study. Calves were excluded when showing any other diseases. Finally, the diarrhea group consisted of 10 animals. 10 healthy native calves were considered for the control group. Blood samples were collected from the jugular vein into EDTA tubes. The leukogram and neutrophil/lymphocyte ratio (NLR) were measured. The TNF- α plasma concentrations were determined using a commercial enzyme-linked immunosorbent assay (ELISA) kit. Data were analyzed by using an independent t-test.

Results: The neutrophilia was the most important finding in the leukogram of the diarrhea group. The NLR was significantly increased in the diarrhea group compared with control group ($P < 0.05$). TNF- α was detected at low level (23.51 ± 2.35 pg/ml) in control group. TNF- α level was 176.32 ± 13.58 pg/ml in native neonatal calves with diarrhea. The plasma concentration of TNF- α in the diarrhea group was significantly higher than the control group ($P < 0.001$).

Conclusion: According to the obtained results, the level of TNF- α and NLR did not increase before the clinical manifestations of diarrhea in native neonatal calves. Further studies are needed to understand this function.

Keywords: Tumor necrosis factor- α , Neonatal calf diarrhea, Leukogram

ICIA2018\Vaccine\Poster\8372

8372

The Formulation of Hepatitis B Vaccine in MF59 adjuvant with PPD and Its Immunogenicity Comparison with traditional Hepatitis B VaccineReyhane Mirzaee¹, Fahimeh Nematimonsour², Mehdi Mahdavi^{3,*}

1. Department of Biotechnology, Faculty of Advanced Sciences & Technology, Pharmaceutical Sciences Branch, Islamic Azad University, Tehran-Iran (IAUPS)

2. Department of Biotechnology, Faculty of Advanced Sciences & Technology, Pharmaceutical Sciences Branch, Islamic Azad University, Tehran, Iran (IAUPS)

3. Department of Immunology, Pasteur Institute of Iran, Tehran, Iran

Background: Hepatitis B infection is propagated widely in the world population and the proper approach that could be used to control is application of vaccine. The clinical vaccine in all of the world is a vaccine formulated in Alum adjuvant that has not provided suitable immune response in some immunized cases. Herein, we formulated a new HBs vaccine using MF59 adjuvant with PPD and compared cellular and humoral immune responses versus commercial HBs vaccine.

Methods: The HBs Ag was formulated in MF59 adjuvant with PPD and injected s.c. 3 times with two week interval to BALB/C mice. Ten days after last immunization, the levels of cytokines including IL-4, IFN- γ , TNF- α , IL-2 and total antibody were assessed by ELISA method.

Results: There was an increase in IL-4 cytokine of Iranian commercial alum vaccine compares with MF59 amounts of IFN- γ cytokine of Iranian Alum and MF59. The study of TNF- α indicates that Iranian commercial vaccine because of its higher inflammation is more exudated than MF59. There was increase in total antibody of Iranian commercial vaccine as compared to vaccine formulated with MF-59.

Conclusion: There was significant reduction of IL-4 cytokine and also antibodies shows that vaccine formulated with MF59 can induce cellular immunity as compared to humoral immune response. moreover, less adversary effects of MF59 adjuvant as another beneficiary properties could be considered.

Key word: HBs- Ag, vaccine, adjuvant, MF59, alum

ICIA2018\Reproductive Immunology\Poster\8395

8395

Evaluation of vascular endothelial growth factor (VEGF) SNP-634C/G and +936C/T polymorphisms in Iranian women suffering from preeclampsia

Seyed Hojjat Hossaini¹, Mehri Ghafourian^{2,*}, Ata Ghadiri¹, Saber Rokhafrooz¹, Mahin Najafian³, Ahmad Rouhizadeh⁴

1. Department of Immunology, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

2. Fertility, Infertility and Perinatology Research Center, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

3. Department of Obstetrics & Gynecology, Fertility, Infertility and Perinatology Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

4. Cellular and Molecular Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

Background: Preeclampsia is a disorder of pregnancy characterized by high blood pressure and proteinuria after 20 weeks of gestation. This disorder is about 6-8% of all pregnancies. Vascular endothelial growth factor (VEGF), an angiogenic factor, plays a significant role in vascular permeability and vascular proliferation. The aim of this study is to assess the various risk factors related to preeclampsia and the VEGF-634C/G & +936C/T polymorphisms in women with preeclampsia.

Methods: This study is a case-control study that conducted in two groups of pregnant women with preeclampsia (n=135) and healthy pregnant women (n=135) admitted to hospitals of Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran. DNA was extracted from the peripheral blood lymphocytes using phenol-chloroform extraction method. The genotypes of the VEGF-634C/G & +936C/T polymorphisms were detected by polymerase chain reaction restriction fragment length polymorphism (PCR-RFLP). Then, the restriction enzymes (FaqI and NlaIII) were applied to detect +936C/T & -634C/G polymorphisms, respectively. The results were obtained and imaged by 1.5% agarose gel and Gel DOC system. The data were analyzed by SPSS software.

Results: In this study the maternal age, gestational age, maternal hemoglobin and maternal BMI were significantly associated with risk of preeclampsia ($P < 0.05$), while preeclampsia season was not effective in preeclampsia occurrence. There was no significant difference between the case and control groups regarding VEGF-634C/G & +936C/T polymorphisms ($P > 0.05$).

Conclusion: Despite the significant relation in various risk factors with preeclampsia, it seems that VEGF-634C/G & +936C/T polymorphisms are not associated with preeclampsia. However, further studies are necessary to more fully assess the different factors associated with preeclampsia.

Keywords: Preeclampsia, Polymorphism, VEGF, Pregnancy

ICIA2018\Stem Cells Based Immunotherapy\Poster\8429

8429

Overexpression of miR-34a in the Mesenchymal stem cell by Lentiviral vectorsKatayoun Bahman Soufiani¹, Ali Akbar Pourfathollah², Mahin Nikogoftar Zarif³, Ehsan Arefian⁴*1. PhD student, Department of Immunology, Tarbiat Modares University**2. Department of Immunology, Tarbiat Modares University**3. Iranian Blood Transfusion Organization**4. Department of Microbiology, University of Tehran*

Background: Nowadays cell therapy is considered as one of the treatments in many diseases, among which mesenchymal stem cells (MSCs) have a distinctive position. Features such as immunogenicity and targeted migration to inflamed tissues, including cancerous regions, have made these cells the best candidate for therapeutic purposes. MicroRNAs are molecules that influence the expression of the gene can be involved in behavioral and evolutionary characteristics of the cells and pathogenesis of many diseases. Significant reduction of miR-34a in many cancers led to the use of this small non-coding molecule as a therapeutic tool in this study. Therefore, in this study, the miR-34a encoding gene was attempted to be integrated using virus packaging in the MSC cell genome.

Methods: First, the miR-34a (5p) sequence gene was cloned into the GFP+ lentivirus plasmid. The recombinant plasmid was transfected into HEK-293 cell line by the calcium-phosphate method, along with the encoding plasmids of structural proteins and lentiviral envelopes. The viral supernatant was gathered and in the second stage transduction with the condensed virus was performed on MSCs. Total RNA extraction and cDNA development were performed.

Results: The sequence of miR-34a gene into the lentivirus plasmid was confirmed by PCR, enzymatic, colony-PCR and sequencing methods. Real-Time PCR results on cDNA of the transduced MSC cells versus control showed a 34-fold increase in miR-34a expression ($p=0.0000$).

Conclusion: Using the Cloning and Virus Packaging methods makes it possible to enter the gene in the considered MSC genome and achieve a permanent expression in these cells. Increasing the miR-34a expression in MSCs indicates that these techniques are applicable in modifying MSCs.

Keywords: Mesenchymal stem cell, MicroRNA, Virus Packaging

ICIA2018\Veterinary Immunology\Poster\8430
8430

A Passive Surveillance for HP AIVs (H9 & H5s) in Isfahan Birds Garden

Abdolreza Nabinejad¹ (DVM, Ph.D), Nooshin Askarani² (MS)

Isfahan research and education center of agriculture and natural resources Nabinejad@abrii.ac.ir
Isfahan research center of teachers, Software group

Background: Isfahan birds garden is located at the Zayandeh rood river and with about 160 specious and 3000 of different birds annually more than 500000 people around the country and tourists visit the birds. Regarding HP AIs (H5N1 and H5N8) outbreak in the poultry farms (industrial and rural) and immigration of immigrant birds to the central part of the Iran, a passive surveillance were carried out, for increasing immunity of the for the visitors and birds.

Method : A daily shuttle program for sanitization of garden and gates with widely antiseptic material were recommended, and an active quarantine for a sure immunity, meanwhile educational techniques used for training and refreshing the staffs specially the birds nurses which are at the first line of communication to the birds. In the laboratory examination a monthly monitoring for NDV and AIV titer well done by HI test, so 2 ml of wing vein blood in 5-6% of sensitive birds were collected and some swabs of pharynx and stools were prepared in the transporting media for RT-PCR (in suspected birds) molecular detection.

Result: The mean ND titer were 5 with CV of 65% and the mean titer of H9N2 were 6 with a CV equal to 98%, but the highest ND titer (9) were related to chickens and pheasant and fowl, the lowest ND titer (2) were related to water fowls such as Ducks, Pelicans, Flamingos, Gooses. The minimal H9N2 titer were 6 and the maximal value were 10 but no one were positive for In H5N1 and H5N8. The molecular RT-PCR of samples for H5N1 and H5N8 fortunately show negative results...

Conclusion: Regarding to the result a monthly vaccination for ND and AI and continues first level of biosecurity were recommended. Also using the training and intelligent board inside and outside the garden with some regular education course were the assessment of this surveillance.

Key words: AIVs, Birds, Isfahan, Passive, surveillance

ICIA2018\Veterinary Immunology\Poster\8433

8433

Serology of Isfahan Ostrich Chicks for a Complex Downer Syndrome

Abdolreza Nabinejad¹ (DVM,Ph.D), Nooshin Askarani² (MS)

Isfahan Research and Education Center of Agriculture and Natural Resources Nabinejad@abii.ac.ir
Isfahan Research Center of Teachers, Software group

Background: Ostrich rearing is developing in Iran and well increased in Isfahan as the first ranking in the country. It was about 20 years ago that a few Ostrich chicks were imported to Iran from South Africa and at the moment we there are more than 5000 breeders just in Isfahan Province with about 7000 chicks for laying or meat production in future. One of the most complicated disease of the ostrich chicks were downing and constipation fallowed by respiratory problems. The clinical signs goes to some management and nutritional difficulties but the contagious form of disease oriented the infectious habitat of the disease meanwhile the more biosecurity shown the list of occurrence.

Methods: Sampling of bloods done via jugular and wing vein in the infected chicks aged 2 to 6 months from 5 farms in Isfahan city and its around cities, The samples transported to serology laboratory and the sera were prepared for HI test for ND and AI and also ELISA test kit of symbiotic used for chicken Aadenovirus 1 and Borna virus investigation.

Results: Regarding our results Aadenovirus 1 and Borna virus infection were negative comparing to positive and negative controls, but the HI test for ND and AI were positive and valid, in which the CV for ND were 250% with the maximum of 10 ,Minimum were 3 and average of 8 , the CV for AI (H9N2) were 236% with the Maximum titer of 9 ,Minimum titer of 4 and average of 6 , Fortunately no any infection of H5s serotypes were positive According to the clinical and paraclinical results the syndrome were related to per acute ND (enteric form) co working with AI,

Conclusion: Therefore vaccination in the breeders (for maternal antibodies) and checks beside disinfecting and biosecurity of the farms were recommended.

Key words: Downer syndrome, AI, ND, Ostrich Chicks, Isfahan

ICIA2018\Vaccine\Poster\8450

8450

Is there any relationship between birth season with HBs Ab titer and expression level of TLR2, 3, 4 genes?

Kousar Smailnejad ganji^{1,2}, Mehdi Shahbazi², Mousa Mohammadnia Afrouzi²

1. Student research committee, Babol University of medical sciences, Babol, Iran

2. Department of Immunology, School of Medicine, Babol University of Medical Sciences, Babol, Iran

Background: Environmental factors such as birth season can play an important role in the development of the immune system and the immune responses to vaccines. The main objective of this study was to determine the association between the birth season with HBs Ab and the expression level of TLR2, 3, 4 genes in response to the hepatitis B vaccine in 3-5 years old children.

Methods: In this study, 72 children with the age of 3-5 years old were born in the winter and summer of 2011-2013 (36 in each group) were included. Blood samples were collected, then the HBs Ab titer and TLR2, 3, 4 genes expression levels were analyzed by ELISA method and Real Time PCR, respectively.

Results: The results of this study indicated that the average expression of TLR2, 3, 4 genes in the winter-born group was higher than the summer-born group. However, this difference was not statistically significant (TLR2 P-Value = 0.5) (TLR3 P-Value = 0.06) and (TLR4 P-Value = 0.16). The average level of antibody in the summer was greater than the winter. However, this finding was not statistically significant (P-Value = 0.3).

Conclusion: There are no relevance between immune responses to HBV vaccine and birth season. However, many studies showed seasonally variable environmental antigens effect on immune response to vaccination. Further research is required to confirm this relevance.

Keyword: HBs Ab titer, Hepatitis B vaccine, TLR2, 3, 4 and Birth Season

ICIA2018\Veterinary Immunology\Poster\8456

8456

Infectious Stomatitis CBC in a Lizard Caused by Resistant *Pseudomonas* Sp.Abdolreza Nabinejad¹ (DVM, Ph.D), Nooshin Askarani² (MS)*1. Isfahan research and education center of agriculture and natural resources Nabinejad@abii.ac.ir**2. Isfahan research center of teachers, Software group*

Background: Reptiles ecosystemic duty are very important, now a days their exhibitions used for human knowledge mean while rearing crocodiles and lizards is going to be a job for export of meat leather. Mouth rot is probably the worst bacterial infection in reptiles and is the cause of many an animal's loss. Bacteria of the groups *Pseudomonas*, *Aeromonas*, and *Proteus* *Mycobacterium* and *salmonellas* accumulate in the oral mucous membrane and cause infections, swellings, and a cheesy discharge. Most cases of mouth rot occur after an initial injury to the snout and poor physical condition of the lizard. Early diagnosis of infection is vital for recovery. Current reports goes to an infectious stomatitis of lizard with a 19 ages old, 216.5cm length and 116.5Kg BW.

Methods: First of all the animal moved to a hospital and welfare conditions were prepared and the sores of mouth washed using normal saline, the secretion and scratched of mouth mucosa layer with caution sampled and tested for microbiology. The blood sampling of tail vein carried out and CBC and microbial exams.

Results: Regarding to the results the CBC shows some neutrophilia and eosinophilia, the PCV were normal, no bacteria and fungi isolated in blood. The mouth samples showed some G- and some G+ bacteria but no any fungi residues. The differential bacterial cultivation and biochemical test shows *Pseudomonas* Sp. which were resistant to penicillin and the majority of related beta-lactam antibiotics, Enrofloxacin, Aminoglycosides due to its cell wall and efflux pumps but just resistant to Cephalosporines and Fosbac..

Conclusion: The lizard welfare in artificial habitat should be prepared or the infectious agents affect them and zoonotic infections and animal death coming.

Key words: Lizards, Stomatitis, CBC, Resistant, *Pseudomonas*

ICIA2018\Vaccine\Poster\8457

8457

Evaluation of the Level of HBs Antibody after Hepatitis B Vaccine among 3-5 years old children in Babol

Kousar Smailnejad ganji^{1,2}, Mehdi Shahbazi², Mousa Mohammadnia Afrouzi², Mojdeh Ghias Tabari³

1. Student research committee, Babol University of medical sciences, Babol, Iran

2. Department of Immunology, School of Medicine, Babol University of Medical Sciences, Babol, Iran

3. Department of Biochemistry, School of Medicine, Babol University of Medical Sciences, Babol, Iran

Background: Hepatitis B virus (HBV) infection is an important infectious disease worldwide. Vaccination has known to be the most effective methods for the prevention of transmission and prevalence of HBV. The main objective of this study was to examine the effectiveness of HBV vaccine in children aged 3-5 years old.

Methods: The study was conducted in Amirkola Children's hospital during a 6- month period from November 2016 to March 2017. Blood samples were obtained from 120 healthy, 3 to 5-year-old children who had been vaccinated against HBV and HBS antibody concentration was measured by ELISA method. The individuals were divided into two groups according to their anti-HBs titer (responder group who had HBS Ab titer > 10 mIU/ml and non-responder group who had HBS Ab titer ≤ 10 mIU/ ml). Results were reported as number(n) and percentage (%) by using SPSS version 22 software.

Results: 69 of the subjects were boy and 51 of the subjects were girl, respectively. Among the cases, 98(82%) children were responder while, 22(18%) subjects were non-responder.

Conclusion: Our results showed that, the antibody response to HBV vaccine was lower than the similar studies. It is suggested, farther studies should be performed to evaluate HBs antibody levels after vaccination especially in high risk children and booster must be administrated, if require.

Key words: HBs Ab titer, Vaccination, Children and Babol

ICIA2018\Tolerance and Autoimmunity\Poster\8458
8458

The Efficacy of St. John's wort in the Treatment of Experimental Autoimmune Encephalomyelitis in C57BL/6 Mice

Reza Nosratabadi¹, Maryam Rastin², Mohammad Kazemi Arababadi¹, Akbar Anaeigoudari³, Mahmoud Mahmoudi²

1. Department of Immunology, School of Medicine, Rafsanjan University of Medical Sciences, Rafsanjan, Iran

2. Department of Immunology, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

3. Department of Physiology, School of Medicine, Jiroft University of Medical Sciences, Jiroft, Iran

Background: St. John's wort *Hypericum perforatum L (HP)* is an herbal medicine used in traditional medicine due to antidepressant and anti-inflammatory activities.

Methods: Based on these properties, we evaluated the therapeutic efficacy of *Hypericum perforatum extract* (HPE) against experimental autoimmune encephalomyelitis (EAE) an animal model of multiple sclerosis (MS).

Results: The clinical score was reduced from 4.56 to 2.89 after treatment with HPE. This change was associated with a decrease in incidence and infiltration of inflammatory cells into the CNS, as well as a decrease in body weight loss. Additionally, treatment with HPE significantly decreased the level of proinflammatory cytokines (IFN- γ , IL-17A and IL-6) and increased anti-inflammatory cytokines (TGF- β , IL-10 and IL-4), in splenocytes from EAE mice. The flow cytometry analysis revealed HPE-treated mice had markedly higher proportions of CD4⁺ CD25⁺ Foxp3⁺ Tregs in the spleen compared to PBS-treated mice ($10.2 \pm 0.27\%$ versus $7.27 \pm 0.24\%$, $p < 0.01$).

Conclusion: Collectively, the current study demonstrated HPE could potentially reduce the clinical and pathological complication of EAE.

Key words: *Hypericum perforatum L*, Multiple Sclerosis, Experimental Autoimmune Encephalomyelitis (EAE), Myelin Oligodendrocyte Glycoprotein (MOG)

ICIA2018\Veterinary Immunology\Poster\8465

8465

Production of Immunoglobulins Y (IgY) against *Salmonella Typhimurium* in Japanese quail (*Coturnix japonica*)

Elham Sadat Hosseini Nasab¹, Maryam shakoori², Atefeh Esmailnejad³, Bahman Abdi⁴

1. Department of Pathobiology School of Veterinary Medicine Shiraz University Shiraz Iran

2. Department of Pathobiology School of Veterinary Medicine Shiraz University Shiraz Iran

3. Department of Pathobiology School of Veterinary Medicine Shiraz University Shiraz Iran

4. Department of Clinical Science Faculty of Veterinary Medicine Shiraz University Shiraz Iran

Background: Salmonellosis is one of the most common infectious diseases all around the world, with *Salmonella entericaserovars Typhimurium* (*S. Typhimurium*) and *Salmonella entericaserovars Enteritidis* (*S. Enteritidis*) being some of the more common isolates associated with foodborne diseases in humans and animals. Immunoglobulin Y (IgY) is the major serum immunoglobulin in oviparous animals with an overall structure similar to that of mammalian immunoglobulins. As a means of protection to the offsprings, avian transfer their maternal immunoglobulin Y from serum to the egg yolk where it serves as passive immunity of the developing chicks. Chicken egg yolk antibodies have been extensively used for prophylactic, therapeutic detection of microbial contaminants and diagnostic purposes. Oral immunotherapy with specific IgY has been established as an efficient alternative to traditional antibiotic therapy in human and animals. In this study, production and specificity of anti- *Salmonella typhimurium* IgYs in immunized quail eggs were evaluated.

Methods: A total of 30 newly hatched *Salmonella* spp.-free female Japanese quails (*Coturnixcoturnix japonica*) were divided into immunized and control groups. Immunization was carried out with intramuscular injection of formalin and heat inactivated *Salmonella typhimurium* whole bacterial suspensions emulsified with Freund adjuvants. Immunization continued for 3 months to ensure that the antibody had reached a high level. Eggs were collected every week for 100 days. Egg yolk IgY was purified by ammonium sulfate precipitation method. Anti- *S. typhimurium* egg yolk IgY titer and specificity were determined using ELISA and Western blot techniques.

Results: *S. typhimurium* specific IgY antibodies were detected in immunized quail eggs and were significantly higher than the control group ($p=0.001$) which confirmed the immunization procedure. The results also showed higher antibody titer in the group immunized with formalin inactivated immunogen.

Conclusion: The study significantly demonstrated the possibility of generating antibody against *Salmonella typhimurium* in quail egg yolks which can be used for immunotherapeutic and immunodiagnostic purposes.

Abstract Text Maximum 2000 Characters around 300 words. (Times New Roman, font size12)

Keywords: *Salmonella Typhimurium*, Immunoglobulin, IgY, Japanese quail

ICIA2018\Tolerance and Autoimmunity\Poster\8475
8475

Longitudinal interferon beta-1a effects in multiple sclerosis

Parisa Nourbakhsh¹, Ali Ghazavi^{1,2}, Ali Ganji¹, Mohsen Ebrahimi⁴, keyvan Ghasami^{4*}, Ghasem Mosayebi^{1,3}, Sepideh Moradkhani⁵

1. Department of Microbiology and Immunology, School of Medicine, Arak University of Medical Sciences, Arak, Iran

2. Traditional and Complementary Medicine Research Center (TCMRC), Arak University of Medical Sciences, Arak, Iran

3. Molecular and Medicine Research Center, Arak University of Medical Sciences, Arak, Iran

4. Department of Neurology, School of Medicine, Arak University of Medical Sciences, Arak, Iran

5. Department of Immunology, School of Medicine, Kerman University of Medical Sciences, Kerman, Iran

.(Background: Cytokines play an important role in the pathogenesis of multiple sclerosis (MS Interferon beta (IFN- β) has been at the forefront of MS treatment for many years. The aims of this study were .to determine the levels of Th1/Th2/Th9/Th17 serum cytokines, after 1 year of therapy with interferon beta-1a

.Methods: Forty healthy controls and 36 MS patients were enrolled in this longitudinal study Th1/Th2/Th9/Th17 serum cytokine levels were measured using ELISA kit at inclusion and after 12 months of .interferon beta-1a (CinnoVex) treatment

Results: After one year of treatment the recorded significant effects were decreases in IL-17A (917.10 ± 132.05 to 547.83 ± 78.46), IFN- γ (260.80 ± 42.74 to 201.10 ± 51.63) and IL-9 (184.25 ± 12.59 to 142.22 ± 6.73) levels .while no significant change was seen in the IL-5 (165.74 ± 9.65 to 172.96 ± 12.60) level

Conclusion: Based on the findings of this study, a reduction in the Th1, Th9 and Th17 serum cytokine levels were seen after 12 months compared with pre-treatment values, which reflects in vitro and animal studies of IFN- β effector mechanisms.

Keywords: MS, Interferon beta, Cytokine

ICIA2018\Tolerance and Autoimmunity\Poster\8491

8491

Investigation of the Serum Levels of Anti-tetanus Toxin Antibodies in Patients with Hyperthyroidism

Mitra Shokri Majelan¹, Hossain Khorramdelazad¹, Mohammad Taghi Rezayati¹, Azam Bagherizadeh¹, Shekoofeh Moghadam¹, Shima Bazaz¹, Fahimeh Mohammadi¹, Abdollah Jafarzadeh²

1. Department of Immunology, Medical School, Rafsanjan University of Medical Sciences, Rafsanjan

2. Department of Immunology, Medical School, Kerman University of Medical Sciences, Kerman, Iran

Background and Aims: There are receptors for thyroid hormones such as thyroid stimulating hormone (TSH) on lymphocytes, which indicate the ability of hormones to influence the lymphocyte responses. The aim of this study was to evaluate the serum levels of anti-tetanus toxin antibodies (anti-TTA) in patients with hyperthyroidism.

Material and Methods: Totally, 24 patients with hyperthyroidism and 24 age- and gender-matched healthy individuals were enrolled to study. All participants were vaccinated against tetanus according to the national immunization program. The presence of hyperthyroidism confirmed according to the clinical and para-clinical criteria such as TSH levels. The serum samples of participants tested for the levels of anti-TTA by ELISA method. The serum antitoxin concentration 0.1 IU/mL was considered as a protective level of antibody.

Results: The seroprotective rate was 100.0% in healthy group and 94.0% in hyperthyroid patients ($P=0.52$). The mean titer of anti-TTA was 3.28 ± 0.48 IU/ml in healthy group and 4.28 ± 0.50 IU/ml in hyperthyroidism group. There was no significant difference between healthy subjects and hyperthyroid patients regarding the levels of anti-TTA ($P=0.16$). In hyperthyroid men, the titer of anti-TTA was higher than hyperthyroid women (5.32 ± 1.42 IU/ml vs 4.01 ± 0.52 IU/ml) but the difference was not significant ($P=0.20$). Similarly, the titer of anti-TTA in healthy men was higher than healthy women, but the difference was not significant, again (3.74 ± 1.17 IU/ml vs 3.09 ± 0.05 IU/ml, $P<0.35$).

Conclusion: These results showed that hyperthyroidism might has no significant influence on the protective against tetanus. The titer of anti-TTA was not affected by gender of participants.

Keywords: Hyperthyroidism, Protection, Tetanus, Anti-toxin antibody.

ICIA2018\Stem Cells Based Immunotherapy\Poster\8496
8496

Banking of mesenchymal stromal cells: a source for autoimmune-disease studies

Masoumeh Asadi¹, Parvaneh Farzaneh¹, Meysam Ganjibakhsh^{1,3}, Seyed Abolhassan Shahzadeh Fazeli^{1,2}, Neda Sadat Gohari¹, Hedieh Rahmati¹, Mehrnaz Izadpanah^{1,4}, Shiva Mohammadi¹, Maryam Farghadan¹, Sepideh Ashouri Movassagh¹, Zahra Elyasi Gorji¹

1. Human and Animal Cell Bank, Iranian Biological Resource Center (IBRC), ACECR and Tehran, Iran
2. Department of Molecular and Cellular Biology, Faculty of Basic Sciences and Advanced Technologies in Biology, University of Science and Culture, Tehran, Iran
3. Department of Anatomy, Faculty of Medicine, Iran University of Medical Sciences, Tehran, Iran
4. School of advanced technologies in medicine, Tehran University of Medical Sciences, Tehran, Iran

Background: Technological developments in the stem cell field have increased assurance in the possible benefits of these cells for treating diseases. Mesenchymal stromal cells (MSCs) are using in various autoimmune diseases studies such as diabetes and multiple sclerosis. A proper cell banking procedure can provide a suitable resource of cells for research.

Methods: Mesenchymal stromal cells, from different tissues, were isolated by enzymatic digestion. For characterization, specific markers detected by flowcytometry and differentiation ability was evaluated. Multiplex PCR and microbial tests were performed to detect cross-contamination with the cells of other species and microbial contamination, respectively. Short tandem repeat (STR) profiling test was performed to detect cross-contamination with other human cells. Finally, each cell sample was cryopreserved and banked with an identity card.

Results: The cells differentiated into mesodermal lineage including osteogenic and adipogenic lineages and also expressed human MSC markers including CD29, CD90, and CD105 whilst lacking in the expression of CD45, CD34. Quality control tests showed no fungal, bacterial, mycoplasma and viral contamination. All of the cell samples authenticated with unique STR profiling and free of cross-contamination with cells of other species.

Conclusion: Long-term culture of primary cells increases the risk of microbial contamination, senescence, and cross-contamination. Banking of cells can minimize mentioned risks. Therefore, good cell banking procedure may support the quality of autoimmunity investigations.

Keywords: Human mesenchymal stromal cells, Cell banking, Autoimmunity studies

ICIA2018\Tolerance and Autoimmunity\Poster\8499
8499

High Expression of Programmed Death-1 (PD-1) Impairs CD3⁺ CD8⁺ T Cells Activity in Patients with Multiple Sclerosis

Atefeh Najmaldini¹, Mohammad Mehdi Mohammadi², Ladan Langroudi, Hossein ali Ebrahimi

1. Department of Immunology, Kerman University of Medical Science, Kerman, Iran

2. Neurological Neuroscience Research Center, Kerman, Iran

Background: Patients with multiple sclerosis (MS) have a deficiency of peripheral blood CD8⁺ T cells which might reduce control of Epstein–Barr virus (EBV) and lead to MS by allowing EBV infected autoreactive B cells to accumulate in the brain. In patients with MS, T cells progressively declined, consistent with T-cell exhaustion.

Methods: We used flow cytometry to determine the phenotypes and frequency of CD8⁺ cells based on the expression of CD8 and CD3 and PD-1 in 30 healthy subjects and 30 patients with MS who had not received corticosteroids or immunomodulatory therapy in previous 3 months. Additionally, Real-time PCR was used to evaluate EBV load in PBMCs.

Results: Cytotoxic CD8⁺ CD3⁺ T cells are able to remove EBV-infected cells in healthy donors. Our results indicated higher expression of PD-1 on circulating CD3⁺ CD8⁺ T cells in MS patients compared to healthy donors. This phenotype is an indication of exhausted T cells.

Conclusion: Exhausted CTLs in patients suffering from MS are not able to eradicate virus infected B cells. The results suggest that inefficient immune control of EBV in patients with MS during remission may cause exacerbation of the disease. Future studies on the mechanism of T cell exhaustion may aid to better understanding of the disease and design of effective therapies.

Keywords: Multiple sclerosis, Exhausted T Cell, Epstein-Barr Virus, PD-1.

ICIA2018\Reproductive Immunology\Poster\8502

8502

Relative Expression of OX40, OX40L mRNA and OX40L serum levels in women with recurrent spontaneous abortionFateme Rahmani¹, Hossein Hadinedoushan^{1*}, Nasrin Ghasemi²

1. Department of Immunology, Reproductive Immunology Research Center, Shahid Sadoughi University of Medical Science, Yazd, Iran

2. Recurrent Abortion Research Centre, Reproductive Sciences Institute, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

Background: The aim of this study was to determine the role of OX40 and OX40L in Recurrent spontaneous abortion (RSA). We compared the expression of OX40, OX40L genes in peripheral blood mRNA levels and serum levels of OX40L in the women with RSA history and the control group.

Methods: In this case-control study, 40 women with RSA history (case group), and 40 women with no history of abortion (control group) were surveyed. The expression of OX40 mRNA and OX40L mRNA were determined in peripheral blood of the two groups using the quantitative polymerase chain reaction. Also, enzyme-linked immunosorbent assay was used to determine the levels of serum OX40L in the case and the control groups.

Results: There were no significant differences in the maternal age of women in two groups (30.1 ± 4.28 years in the case and 30.03 ± 4.23 years in the control group). There was no difference in the levels of OX40 and OX40L mRNA between the case and the control groups ($p=0.1$ and $p=0.6$ respectively). Also, there was no significant correlation between the expression of OX40 and OX40L mRNA levels with age and the number of abortions. The correlation between OX40 and OX40L mRNA levels was insignificant. Women with RSA history had significantly higher level of serum OX40L than control group ($p=0.03$).

Conclusion: Our findings showed that the expression of OX40 mRNA and OX40L mRNA were similar between women with RSA history and the control group. The elevation of serum OX40L levels may be considered as risk factor for RSA.

Key words: OX40, OX40L, quantitative PCR, RSA

ICIA2018\Stem Cells Based Immunotherapy\Poster\8513
8513

Hanging drop as an optimal method for cancer stem cells enrichment from HT-29 human colorectal cancer cell line in order to use them in cancer immunotherapy

Marzieh Naseri¹, Elmira Gheytauchi², Zahra Madjd^{1,2*}, Mahmood Bozorgmehr^{2*}, Jamshid Hajati³,
Roya Ghods^{1,2}, Jafar Kiani^{1,2}

1. Department of Molecular Medicine, Faculty of Advanced Technology, Iran University of Medical Sciences, Tehran, Iran

2. Oncopathology Research Center, Iran University of Medical Sciences, Tehran, Iran

3. Department of Immunology, School of Medicine, Tehran University of Medical Science, Tehran, Iran

Background: Tumor mass-resident cancer stem cells (CSCs) are suggested to play a key role in recurrence and metastasis of colorectal cancer and its resistance to conventional therapies. On the other hand, immune responses are well known to be the vanguard of natural anti-cancer strategies. However, the inability to target CSCs with current immune approaches may be a significant factor in treatment failures. Dendritic cell (DC) prime with CSC-lysate has demonstrated significant induction of anti-tumor immunity against CSCs both in vivo and in vitro, selectively targeting CSCs. The key step for promoting robust anti-CSC immune responses is their efficient isolation with acceptable purity. Hence, we were about to come up with an optimized and reproducible method for colorectal CSCs enrichment.

Methods: In this study, several colon spheres generation methods including hanging drop, culture in serum free media under non-adherent conditions (poly-HEMA and 2% agarose coated plates) and use of external forces were compared; all the methods were carried out at various cell densities. CSC enrichment was verified using flowcytometry analysis for putative molecular markers of colorectal CSCs. In addition, Real time PCR was used to assess the expression of several well-known stemness genes.

Results: HT29 colon cancer cells grew best using the hanging drop technique. The percentage of CD44⁺ and CD133⁺ cancer cells among spheroids was higher than that of the adherent cells. Also, the expression of stemness genes including OCT4, SOX2 and KLF-4 was markedly increased in the spheroid cells as compared with that of the control adherent cells.

Conclusion: We suggest the hanging drop method as an efficient approach for enrichment of colorectal CSCs to be used in targeted therapeutic modalities. Such suitable isolated CSCs could be utilized for successful generation of lysate-pulsed dendritic cell in future pre-clinical settings.

Keywords: Cancer Stem Cell, Dendritic Cell, Colorectal Cancer, Hanging Drop

ICIA2018\Tolerance and Autoimmunity\Poster\8537
8537

Evaluation of IL-17 Producing Memory Effector CD26 high T Cells in Patients with Psoriasis (Based on PASI Score)

Behnaz Esmacili^{1,2}, Parvin Mansouri³, Alipasha Meysamie⁴, Maryam Izad^{1*}

1. Immunology Department, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

2. Immunology, Asthma and Allergy Research Institute (IAARI), Tehran University of Medical Sciences, Tehran, Iran

3. Skin and Stem Cell Research Center, Tehran University of Medical Sciences, Tehran, Iran

4. Department of Community and Preventive Medicine, Tehran University of Medical Sciences, Tehran, Iran

Background :Psoriasis is a chronic, T cell-mediated inflammatory skin disorder in which Th-17 and its related inflammatory cytokines have a fundamental role in its pathogenesis. Human Th-17 cells express high level of surface CD26. We analyzed the IL-17 producing effector memory T cells (CD26 hi memory T cells) from patients with psoriasis following ex-vivo stimulation with respect to PASI score. Also, to investigate the inflammatory responses that can affect effector T cells differentiation, the level of IL-23, IL-6, TNF α , TGF β and IL-17 cytokines were monitored as well.

Methods: Memory T cells were isolated from 10 patients with psoriasis (6 moderate to severe and 4 patients with mild form) and 10 controls. Ex vivo stimulated IL-17 producing effector (Foxp3+ CD25+ CD26hi) memory T cells were analyzed by flow cytometry. IL-23, IL-6, TNF α , TGF β and IL-17 cytokine levels were also evaluated by ELISA.

Results: Although the IL-17+ effector memory CD26hi T cells reduced in patients compared to controls, it was not significant (P =0.062). However, IL-17+ effector memory CD26hi T cells from patients with mild form showed a significant decrease compared to controls. (p=0.03). We could not find any significant difference regarding IL-23, IL-6, TNF α , TGF β and IL-17 cytokine levels in plasma and cell culture supernatant samples between patients and controls.

Conclusion: Taken together, our results showed that IL-17+ effector memory CD26hi T cells in patients with psoriasis had a decreased trend compared to controls. A significant decrease in IL-17+ effector memory CD26hi T cells only in patients with mild form could be because there are increased number of circulating effector T cells in patients with moderate to severe form.

Key words: Psoriasis, CD26, IL-17

ICIA2018\Tolerance and Autoimmunity\Poster\8558

8558

Diminished serum levels of anti-tetanus toxin antibodies in patients with hypothyroidism

Fahimeh Rostamabadi¹, Hossain Khorramdelazad¹, Mohammad Taghi Rezayati¹, Azam Bagherizadeh¹, Shekoofeh Moghadam¹, Shima Bazaz¹, Fahimeh Mohammadi¹, Abdollah Jafarzadeh²

1. Department of Immunology, Medical School, Rafsanjan University of Medical Sciences, Rafsanjan

2. Department of Immunology, Medical School, Kerman University of Medical Sciences, Kerman, Iran

Background and Aims: Many disorders including decreased weight of lymphoid organs, decreased count of lymphocytes and suppression of the lymphocytes-related immune response were demonstrated in experimental models of hypothyroidism. The aim of this study was to evaluate the serum levels of anti-tetanus toxin antibodies (anti-TTA) in patients with hypothyroidism.

Material and Methods: Totally, 24 patients with hypothyroidism and 24 age- and gender-matched healthy individuals were enrolled to study. All participants were vaccinated against tetanus according to the national immunization program. The presence of hypothyroidism confirmed according to the clinical and para-clinical criteria such as the levels of thyroid hormones. The serum samples of participants tested for the levels of anti-TTA by ELISA method. The serum antitoxin concentration 0.1 IU/mL was considered as a protective level of antibody.

Results: The seroprotective rate against tetanus was significantly lower in hypothyroidism patients than healthy group ($P=0.02$). The mean titer of anti-TTA was in hyperthyroidism group was also lower than healthy group (3.28 ± 0.48 IU/ml vs and 1.67 ± 0.33 IU/ml, $P<0.01$). There was no significant difference between men and women concerning the serum levels of anti-TAA antibodies neither in hyperthyroid patients nor in healthy subjects.

Conclusion: These results showed that hypothyroidism might lead to the lower titer of anti-TTA antibodies and reduce the protection against tetanus. The titer of anti-TTA was not affected by gender of participants.

Keywords: Hypothyroidism, Protection, Tetanus, Anti-toxin antibody.

ICIA2018\Tolerance and Autoimmunity\Poster\8558
8558

Diminished serum levels of anti-tetanus toxin antibodies in patients with hypothyroidism

Fahimeh Rostamabadi¹, Hossain Khorramdelazad¹, Mohammad Taghi Rezayati¹, Azam Bagherizadeh¹, Shekoofeh Moghadam¹, Shima Bazaz¹, Fahimeh Mohammadi¹, Abdollah Jafarzadeh²

1. Department of Immunology, Medical School, Rafsanjan University of Medical Sciences, Rafsanjan
2. Department of Immunology, Medical School, Kerman University of Medical Sciences, Kerman, Iran

Background and Aims: Many disorders including decreased weight of lymphoid organs, decreased count of lymphocytes and suppression of the lymphocytes-related immune response were demonstrated in experimental models of hypothyroidism. The aim of this study was to evaluate the serum levels of anti-tetanus toxin antibodies (anti-TTA) in patients with hypothyroidism.

Material and Methods: Totally, 24 patients with hypothyroidism and 24 age- and gender-matched healthy individuals were enrolled to study. All participants were vaccinated against tetanus according to the national immunization program. The presence of hypothyroidism confirmed according to the clinical and para-clinical criteria such as the levels of thyroid hormones. The serum samples of participants tested for the levels of anti-TTA by ELISA method. The serum antitoxin concentration 0.1 IU/mL was considered as a protective level of antibody.

Results: The seroprotective rate against tetanus was significantly lower in hypothyroidism patients than healthy group ($P=0.02$). The mean titer of anti-TTA was in hyperthyroidism group was also lower than healthy group (3.28 ± 0.48 IU/ml vs and 1.67 ± 0.33 IU/ml, $P<0.01$). There was no significant difference between men and women concerning the serum levels of anti-TAA antibodies neither in hyperthyroid patients nor in healthy subjects.

Conclusion: These results showed that hypothyroidism might lead to the lower titer of anti-TTA antibodies and reduce the protection against tetanus. The titer of anti-TTA was not affected by gender of participants.

Keywords: Hypothyroidism, Protection, Tetanus, Anti-toxin antibody.

ICIA2018\Tolerance and Autoimmunity\Poster\8587

8587

Analysis of Th cell Subsets and Regulatory T Cells in Rheumatoid Arthritis Patients with Different Patterns of Response to Treatment

Seyede Rahele Tabatabaei Jafari¹, Niloofar Alvandpour¹, Ahmad Tahamoli Roodsari², Zahra Basiri², Mehrdad Hajilooi¹, Zohreh Zarei Ghane¹, Ghasem Solgi^{1*}

1. Department of Immunology, School of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran.

2. Division of Rheumatology, Department of Internal Medicine, School of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran.

Background: Rheumatoid arthritis (RA) as a chronic inflammatory autoimmune disease is characterized by infiltration of activated inflammatory cells particularly, CD4⁺ T cells into the rheumatoid joints. This study aimed to evaluate the frequencies of Th1, Th2, Th17 and regulatory T cells (Tregs) in peripheral blood samples of RA patients based on the clinical response to treatment.

Methods: Forty three RA patients under treatment with disease-modifying antirheumatic drugs (DMARDs) and 15 healthy control subjects were recruited in this case-control study. Patients were subdivided to good-responders (n=23) and poor-responders (n=20) according to the therapeutic response. Frequencies of Th cell subsets as well as Tregs were determined in peripheral blood samples and then compared between different groups of the study.

Results: The percentage of Th1 cells in both groups of the patients were found to be significantly higher than controls [good-responders (7.42±4.22%) and poor-responders (7.25±3.75%) vs. 2.93±1.57% in the controls, P<0.001]. However, there were no significant differences in the frequencies of Th2 (1.70±0.73% vs. 1.45±0.73%, P=0.22), Th17 (0.58±0.10% vs. 0.51±0.10%, P=0.39) and Treg cells (0.36±0.19% vs. 0.40±0.19%, P=0.71) in all patients compared to healthy controls.

Conclusion: Our results highlighted an increased frequency of Th1 cells in RA patients regardless of good or poor responses to DMARDs in compare with healthy controls.

Keywords: Rheumatoid arthritis, Th1, Th2, Th17, Regulatory T cells.

ICIA2018\Vaccine\Poster\8595

8595

Designing and engineering of immunogenic peptides of *Acinetobacter baumannii* OmpA

Kobra Mehdinejadani¹, Mojgan Bandehpour², Mohammad mehdi ranjbar³, Ali Hashemi^{4*}, Seyed Amir Jalali¹, Vahid Yardeh⁵, Nariman Mossafa^{1*}

1. Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran
2. Department of Biotechnology, School of Advanced Technologies in Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran
3. Department of Immunology, School of Veterinary Medicine, Tehran University of Medical Sciences, Tehran, Iran
4. Department of Microbiology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran
5. Laleh hospital, Tehran, Iran

Background: *Acinetobacter baumannii* is a Gram-negative bacterium that Recently, it has been being known as a leading nosocomial pathogen. As respects, causing infections by this pathogen is high mortality due to inadequate available treatment. An effective vaccine is an unmet medical need. In this study, considering the fact that outer membrane protein A (OmpA) one of the main vaccine candidates, we present T cell and B cell predicted epitopes by sequence-based epitope prediction tools.

Methods: At first, OmpA was amplified from *A.baumannii* genomic DNA. Then genomic sequence of OmpA was determined and translated to protein sequence. We submitted the full-length OmpA protein in B cell linear epitope predictors (LBtope, SVMTrip, ABCpred, BepiPred, BcePred, BCPred, IEDB). Predicted linear B cell epitopes were resubmitted for T cell epitope predictions using the MHC-II Binding Predictor in the Immune Epitope Database (IEDB). On the other hand, we investigated tertiary and secondary structures of protein that we could select the best epitopes.

Results: Consensus epitopes in different tools with the highest score were selected and engineered. The epitopes contain sequences of F317-332, F122-139, F230-245, F24-50, F197-222.

Conclusion: We reason that mentioned peptides could provide protection against *Acinetobacter baumannii* nosocomial infections thereby functioning as potential vaccine candidates.

Key words: *Acinetobacter baumannii*, OmpA, Immunogenic peptides, Antigenome technology

ICIA2018\Veterinary Immunology\Poster\8602

8602

Avian Influenza Investigation in Some Aquatic Birds

Abdolreza Nabinejad¹ (DVM, Ph.D), Nooshin Askarani² (MS)

1. Isfahan research and education center of agriculture and natural resources nabinejad@yahoo.com
2. Isfahan research center of teachers, Software group

Background: Influenza type A viruses are of the most significance to public health due to their potential to cause an influenza pandemic. Influenza type A viruses are classified into subtypes according to the combinations of different virus surface proteins hemagglutinin (HA) and neuraminidase (NA). So far there are 18 different hemagglutinin subtypes and 11 different neuraminidase subtypes such H9N2, H5N1 and H5N8 or H7N5 as reported in birds of Iran. Aquatic birds are the primary natural reservoir for most subtypes of influenza A viruses. There are a few of aquatic birds included Swan, Pelican, Sea gull, Ducks, Flamingo and Goose in the Isfahan bird garden, current report goes to monitoring of these aquatic birds for AIV by serological and molecular tests.

Methods: Using a sterile siring about 2 ml of blood of wing vein collected and transferred to lab for AIV titration, and coanal cleft and cloacal swabs were prepared for molecular test. monthly monitoring of 5-6% of sensitive birds were examined randomly; The technical method for serology were HI, ELISA and RT-PCR for molecular and confirmation of suspected titers.

Results: All of the birds showed positive titer for H9N2 but the highest titer were 9 for goose and green head duck. The lowest titre was one and related to pelicans, Flamingos. The sera were negative for H5N1 and H5N8 and H7 but the swabs were checked by RT-PCR technique which confirmed the negative titers

Conclusion: This monitoring monthly is carried out and sanitization of garden and gates and with widely antiseptic material carried out daily.

ICIA2018\Veterinary Immunology\Poster\8603
8603

Investigation about Prevalence of Anthrax in the Western Cities of Kermanshah Province, During 2010- 2015

Mehrdad Pooyanmehr¹, Erfan Bahrami², Mohammad Reza Khedri², Amir Amiri Paryan^{*3}

1. Department of Basic Sciences and Pathobiology, Microbiology section, Faculty of Veterinary Medicine, Razi University, Kermanshah, Iran

2. Veterinary Student, Clinical Sciences, Faculty of Veterinary Medicine, Razi University, Kermanshah, Iran

3. Veterinary Student, Pathobiology & Basic Sciences, Faculty of Veterinary Medicine, Razi University, Kermanshah, Iran

Background: Zoonotic diseases constitute a public health problem throughout the world. Anthrax is a potentially fatal zoonotic disease and is an important zoonotic disease which is primarily a job-related infection caused by *Bacillus Anthracis*. The diagnosis of cutaneous anthrax (CA) may be very difficult, particularly in atypical presentations and non-endemic regions. Aim this study, anthrax cases seen between 2010 –2015 (six years) in the western area of Kermanshah province.

Methods: Medical records of infected in patients with CA (Clinical, Para clinical information and pathological diagnosis of disease CA such as microbiologic procedures) who had been operated in referral Health Centers Western cities areas of Kermanshah province (Ravansar, Javanrood, Pavehe and Salasebabajany) were collected in the period six years (2010 – 2015). One patient with CA was included in this study.

Results: Results indicated that All patient's referent one patient were who diagnosed CA in 4 cities medical center registries, (Ravansar 1, Javanrood 0, Pavehe 0 and Salasebabajany 0 patients) at six years' period respectively, per 100,000 (0.001%) person. One patient with a diagnosis of CA was followed up. One patient had a history of animal contact. The clinical presentation of CA was typical in one patient in 2015 (1 case) in Ravansar. Patient was initially misdiagnosed with insect bites or angioedema. Cultures from the lesions were positive for *Bacillus Anthracis* in one case. Gram stain from the lesions revealed Gram-positive rods in one case. Patient were diagnosed by clinical presentation and a history of contact with sick animals or contaminated animal products.

Conclusion: CA is a very contagious and important infectious disease worldwide. Early and accurate diagnosis dramatically affects the prognosis of the disease. The diagnosis of CA may be difficult, especially in atypical presentations and non-endemic areas. Thus, CA should be kept in mind, especially in these situations.

Keywords: *Bacillus Anthracis*, Zoonotic diseases, Western Cities Area, Kermanshah Province

ICIA2018\Tolerance and Autoimmunity\Poster\8604

8604

Investigation of the Frequency of Natural Killer Cells Expressing CD16 bright in Peripheral Blood of Multiple Sclerosis and Neuromyelitis Optica Syndrome Disease Patients by Flow Cytometry

Leila Khani¹, Mir Hadi Jazayeri², Mahmood Bozorgmehr³, Seyed Masood Nabavi⁴, Leila Janani⁵

1. MS student, Department of Immunology, School of Medicine, Iran University of Medical Science, Tehran, Iran

2. Assistant Professor, (a) Department of Immunology, School of Medicine, (b) Immunology Research Center Iran University of Medical Science, Tehran, Iran

3. Oncopathology Research Center, Iran University of Medical Science, Tehran, Iran

4. Associate Professor of Neurology, (MS Fellow, Clinical Epidemiology Expertise Degree) Neurology Group, Department of Regenerative Biomedicine, Cell Science Research Center, Neuroscience and Cognition Center, Royan Institute for Stem Cell Biology and Technology, ACECR, Tehran, Iran

5. PhD Department of Biostatistics, School of Public Health, Iran University of Medical Science, Tehran, Iran

Background: Multiple Sclerosis (MS) and Neuromyelitis Optica Syndrome Disease (NMOSD) are inflammatory diseases of central nervous system. Despite the similar manifestations of these two diseases, immunopathogenesis and the treatment of these two conditions are totally different. Natural killer cells (NK) are immune cells which play an important role in defending and regulating immune responses. There are two distinct subsets of NK cell based on CD16 and CD56 expression in peripheral blood. CD16^{bright} CD56^{dim} has high cytotoxic activity (90%). NK cells expressing CD16^{dim} CD56^{bright} are usually located in lymph organs (10%), with low cytotoxic activity, in spite of high cytokine production, whereas there is conflicting evidence of NK cells roles in the pathophysiology of MS and NMO. The aim of this study was to compare the frequency of NK cells in peripheral blood of MS and NMO patients.

Methods: 15 Naïve MS, 15 relapsing-remitting (RR) MS, 15 NMOSD and 15 healthy subjects were included in our study. The percent of NK cells were evaluated by flow cytometry.

Results: The CD16^{bright} median of RR MS, Naïve MS, NMOSD and healthy groups was 7.72, 4.92, 2.13 and 8.10, respectively. There was a significant difference between RR MS and Naïve MS ($p=0.029$), RR MS and controls ($p=0.002$), although no significant differences were found between RR MS and NMOSD ($p=0.00$), Naïve MS and NMOSD ($p=0.056$), controls and NMOSD ($p=0.868$), Naïve MS and controls ($p=0.187$) groups.

Conclusion: The present study demonstrated there is a lower frequency of NK cells expressing CD16^{bright} in peripheral blood of Naïve MS and NMO patient comparison to healthy and RR MS subjects. Actually, it seems immunomodulatory drugs prescribed to RR MS patients lead to increases CD16^{bright} frequency. Therefore, more studies are needed to understand the role of different NK cell subsets in pathogenesis and their uses in therapeutic processes.

Keywords: Multiple Sclerosis, Neuromyelitis Optica Syndrome Disease, Natural Killer Cell Subsets, Flow Cytometry

ICIA2018\Reproductive Immunology\Poster\8634
8634

Investigation the relationship between C677T polymorphism in MTHFR gene and plasma homocysteine concentration in recurrent fetal miscarriage

Omid Kooshkaki¹, Gholamreza Ananisarab², Yalda Zarafshan³

1. Student Research Committee, Birjand University of Medical Sciences, Birjand, Iran

2. Department of immunology-School of medicine-Birjand University of medical sciences
Birjand University

Background: Recurrent miscarriage is a painful experience for both the physician and the parent. It occurs in 2 to 5 percent of pregnant women. The causes of recurrent abortion are genetic, anatomical, hormonal, and immunological. Recurrent miscarriage etiology is not known in 40 to 50% of cases. Single nucleotide polymorphisms in the coding genes of the enzymes, which regulate metabolic pathways such as methylene tetrahydrofolate reductase (MTHFR), are considered as one of the important contributors to the development of thrombophilia. Thrombosis seems to be a disturbance in circulation between mother and fetus in the pairs of capillaries and ultimately leads to abortion. The aim of this study is to determine the association between C677T polymorphism in MTHFR gene and plasma homocysteine concentration in recurrent fetal miscarriages.

Methods: Overall, 100 women were included in the research, the case group comprised of 50 women who had a history of spontaneously recurrent miscarriage with unspecified cause, whereas the controls consisted of 50 women with a history of at least two successful fertility. Methods used in the study included PCR-RFLP with limited effective Hinf-I enzyme in order to investigate polymorphism and ELISA analysis to investigate plasma homocysteine concentration.

Results: Based on our results, in the control group, 15 (30%) had heterozygote and 4 (8%) had mutated homozygotes, i.e., a total of 19 subjects (38%) were abnormal. Also, ELISA results showed that, the mean homocysteine level in the case group was higher than that in the control group ($P=0.002$).

Conclusion: Studies in different populations have shown different results for the association between MTHFR polymorphisms and recurrent miscarriages. This relationship observed in our study.

Keywords: Polymorphism, MTHFR gene, plasma homocysteine, frequent abortion

ICIA2018\Veterinary Immunology\Poster\8655
8655

Identification of Avian Influenza Virus in Broiler Chickens in Gilan Province

Omid ramezani¹, Jalal shayegh², Hossein hosseini³, Marzie sadeghi^{4*}

1. DVM, Department of Veterinary Clinical Sciences, Shabestar Branch, Islamic Azad University, Shabestar, Iran

2. Department of Clinical sciences, Faculty of the Veterinary Medicine, , Shabestar Branch, Islamic Azad University, Shabestar, Iran

3. Department of Clinical sciences, Faculty of the Veterinary Medicine, Karaj Branch, Islamic Azad University, Karaj, Iran

4. student, Department of Clinical Sciences, Faculty of the Veterinary Medicine, University of Shahrekord, Shahrekord, Iran

Background: Avian Influenza is a contagious and acute respiratory disease of great importance to public health, which causes many economic losses. Several methods have been proposed for identifying known viruses such as ELISA and HI test but in terms of the speed of RT-PCR detection, it is probably one of the most important methods. The purpose of this study is to identify and detect viruses in Gilan province.

Methods: For this purpose, 66 samples of swabs from different cities of Gilan province were collected and analyzed using RT-PCR method.

Results: The results showed that 22.73% of the samples were positive, and biosecurity measures should be taken to prevent the spread of the disease.

Keywords: broiler chicken, RT-PCR, influenza, Gilan

ICIA2018\Tolerance and Autoimmunity\Poster Discussion\8665
8665

Expression Analysis of PD-1 and Tim-3 Immune Checkpoint Receptors in Vitiligo Patients

Ali Rahimi¹, Hadi Hossein Nattaj¹, Zohreh Hajheidari², Abolghasem Ajami^{1*}, Hossein Asgarian Omran^{1, 3*}

1. Department of Immunology, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

2. Department of Dermatology, Faculty of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

3. Immunogenetics Research Center, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

Background: Vitiligo is a skin-related chronic autoimmune disease that degrades melanocytic cells, causing colorless and asymptomatic stains. In recent years, the contribution of immune checkpoint receptors has been addressed in the pathogenesis of multiple autoimmune diseases. In the present study, the expression profile of T-cell immunoglobulin and mucin-domain containing-3 (Tim-3) and programmed cell death-1 (PD-1) checkpoint molecules was investigated on CD8⁺ T cells of patients with vitiligo as the main pathogenic effector cells in this autoimmune complication.

Methods: A total of 30 vitiligo patients and 30 sex- and age-matched healthy controls were included in this study. To determine the frequency of Tim-3⁺/PD-1⁺/CD8⁺ T cells, PBMCs were stimulated with PHA for 72 h and a three-color flow cytometry method was applied. To measure the cytokines production, PBMCs were stimulated with PMA/ionomycin for 18 h and concentrations of IL-4, IFN- γ and TNF- α were measured in culture supernatants by ELISA. CD8⁺ T cells were then positively isolated from all participants by magnetic beads separation method and the mRNA expression of PD-1 and Tim-3 molecules was determined by TaqMan based Real-Time PCR.

Results: Vitiligo patients have significantly showed more expression of Tim-3 and PD-1 on the surface of their CD8⁺ T cells than that of normal controls. While, the production levels of TNF- α and IFN- γ were found higher by vitiligo patients than those of controls, IL-4 production was lower in patients with vitiligo. Expression analysis of Tim-3 and PD-1 mRNA confirmed the results obtained from flow cytometry and showed more expression of Tim-3 and PD-1 in CD8⁺ T cells of vitiligo patients compared to normal individuals.

Conclusion: Our results indicate that Tim-3 and PD-1 are involved in immune dysregulation mechanisms of CD8⁺ T cells in vitiligo. Further studies are needed for better understanding of Tim-3 and PD-1 roles in immunopathogenesis of vitiligo and may introduce useful biomarkers for disease progression and/or immunotherapy.

Keywords: Vitiligo, CD8⁺ T cells, Tim-3, PD-1, Flow cytometry, Real-Time PCR

ICIA2018\Research and Development\Poster\8670
8670

Establishment and introducing of cell line collections to support the immunological investigations in Iran

Parvaneh Farzaneh¹, Seyed Abolhassan Shahzadeh Fazeli^{1, 2}

1. Human and Animal Cell Bank, Iranian Biological Resource Center (IBRC), ACECR, Tehran, Iran

2. Department of Molecular and Cellular Biology, Faculty of Basic Sciences and Advanced Technologies in Biology, University of Science and Culture, Tehran, Iran

Background: Continuous and primary cell cultures are excellent models to study the different biological and pathological aspects of *in vitro* studies. Establishment, collection and distribution of several collections of authenticated contamination free cell samples for various immunological studies can boost research.

Methods: To support all topics of immunological investigations we divided the cell collections in two categories. The first one consists of continuous standard cell lines with specific identity which are distributing worldwide by cell banks and the second kinds of required collections were prepared from Iranian native population or laboratory animals. Collections of lymphoblastoid cell lines established by EBV transformation of patients with type one immunodeficiency, also collections of different autoimmune diseases and several types of cancers established by this method. Primary cultures of solid tumor, stromal stem cells of different tissues were two other collections with complete cell characterization and identity.

Results: Human and animal cell bank is distributing different cell collections such as cells related to autoimmunity, immunodeficiency, cancer and laboratory animals, which support immunological investigations.

Conclusion: Introducing of standard cell line collections may improve and facilitate immunological investigations in Iran.

Keywords: Cell line, Collection, Immunology, Investigation

ICIA2018\Tolerance and Autoimmunity\Poster\8677
8677

Evaluation of ATP and HIF-1-alpha in Brain of EAE Mouse Model of Multiple Sclerosis Disease

Seyedeh Mahboobeh Mosavi Mehr, Seyed Alireza Mesbah Namin

1. Department of Clinical Biochemistry, Faculty of Medical Sciences, TarbiatModares University, Tehran, Iran

Background: Multiple sclerosis (MS) is a chronic autoimmune and inflammatory neurological disease of the central nervous system. Dysfunction of mitochondria of the nervous system to produce enough energy also affect the complete myelination, followed by inflammation in neurons occurred in patients with MS. The aim of this study is evaluation of ATP and hypoxia-inducible factor 1-alpha (HIF1-a) which is considered as the master transcriptional regulator of cellular and developmental response to hypoxia, in brain tissue of mouse model of MS disease.

Methods: The C57BL/6 mice were purchased from the Pasture Institute and Hooke kit used for inducing experimental autoimmune encephalomyelitis (EAE), the mouse model of MS disease. High Score of this model (3 to 4) used in this study. The brain tissues of mice were examined for two sets of experiments, one for estimation of HIF1-a using special Elisa kit from Zell Bio GmbH Company and other for measurement of ATP using especial assay kit from Sigma-Aldrich company, after deproteinizing of the samples in 3 group of sham, negative control and EAE mice.

Results: Amount of ATP, in contrast to HIF1-a, was significantly decreased in EAE model in comparison with sham and negative control and significant reducing of HIF-1 a, indicating of existence of hypoxia in brain of EAE mice, as expected.

Conclusion: Significant attenuation of ATP and existence of hypoxia in brain of EAE mice indicated that the function of mitochondria of the nervous system of EAE impaired to produce of energy affecting of neurogenerative process in this model of MS disease.

Keywords: Multiple sclerosis, EAE, ATP, Hypoxia

ICIA2018\Stem Cells Based Immunotherapy\Poster\8687

8687

Optimizing the differentiation of human umbilical cord Wharton's jelly mesenchymal stem cells into perimordial germ-like cells

Samira Ghasemi^{1,2}, Mohammad Shafiee^{1,3}, Mohsen Saeedi^{1,4}, Naser Mobarra^{1,5}

1. Stem Cell Resaerch center, Golestan University of Medical Sciences, Gorgan. Iran.

2. Department of Advanced Medical Technologies, Golestan University of Medical Sciences, Gorgan. Iran.

3. Department of Medical Genetics, Golestan University of Medical Sciences, Gorgan. Iran.

4. Department of Medical Imonology, Golestan University of Medical Sciences, Gorgan. Iran.

5. Department of Clinical biochemistry, Golestan University of Medical Sciences, Gorgan. Iran.

Background: Infertility affects 15-25% of couples of reproductive age. An increasing prevalence of infertility is one of the many problems in countries around the world. Assisted reproductive technologies are not successful in some cases. In recent years, the quantity and quality of germs has been reduced. A novel approach to treating infertility is the use of stem cells. In this study, optimizing the differentiation of human umbilical cord Wharton's jelly mesenchymal stem cells into perimordial germ-like cells has been performed.

Methods: Human warton's Jelly Mesenchymal Stem Cells (hWJ-MSCs) were prepared. Cell surface markers and their differentiation potential were evaluated. Then the induction of the differentiation of these cells into germ like cells was performed using BMP4 and RA. Expression changes of Oct-4, c-kit, Stella, and Vasa genes were evaluated by Real Time PCR. All data were significant.

Results: The morphological changes of hWJ-MSCs, as well as the expression of Oct-4, c-kit, Stella, and Vasa mRNA and protein markers indicate the successful differentiation of these cells into primordial germ like-cells. Also all data was significant. (P value <0.05).

Conclusion: Our results showed that the differentiation of human umbilical cord Wharton's jelly mesenchymal stem cells into perimordial germ-like cells well done with a slight variation in existing differentiation protocols. It can be used to treatment of infertility problems caused by impairment in the structure or function of germ cells.

Key words: differentiation, PG- like Cell, hWJ-MSC

ICIA2018\Reproductive Immunology\Poster\8691
 8691

The effect of active form of 1,25VitD3 on Treg /Th17 axis (percentages, gene expressions and cytokines level) in patients with Unexplained Recurrent Spontaneous Abortion (URSA)

Elham Abdollahi^{1,2}, Houshang Rafat panah^{3,4}, Seyed Abdolrahim Rezaee⁴, Maryam Rastin⁵

1. Immunology Research Center, Department of Medical Immunology, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

2. Student Research Committee, Mashhad University of Medical Sciences, Mashhad, Iran.

3. Research Center for HIV/AIDS, HTLV1 and Viral Hepatitis, Iranian Academic for Education, Culture and Research (ACECR), Mashhad Branch, Mashhad, Iran.

4. Inflammation/Inflammatory Diseases Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

5. Immunology Research Center, Bu-Ali institute , Mashhad University of Medical Sciences, Mashhad, Iran

Background: Unexplain Recurrent Spontaneous abortion (URSA), defined as three or more consecutive pregnancy losses before the 20th week of gestation, occurs in 1%–5% of women of reproductive age. CD4⁺CD25⁺ CD127⁻ FoxP3⁺ Treg cells constitute a minority of the CD4⁺ T-cell population in peripheral blood. T helper 17 (TH17, CD3⁺ CD8⁻ IL-17⁺) cells involve in pathogenesis of chronic inflammatory diseases as well as URSA. Treg/Th17 ratio increases in patients with URSA compared to healthy pregnant women. Vitamin D3 has been shown to inhibit Th17 cell responses and to induce differentiation of Tregs and/or expansion of fork head box protein 3 (FOXP3) regulatory T cells. Considering the imbalance of Th17/Treg in RPL patients, and the effects of Vitamin D3 on these two cell subsets. The aim of this study was to evaluate the effect of vitamin D on the alternation of Treg/Th17 ratio in patients with URSA.

Methods: 20 patients with URSA were sampled for 10 ml whole blood to isolate peripheral blood mononuclear cells (PBMCs) using ficoll-hypaque density gradient centrifugation. Isolated cells were cultured in the presence of 50 nM 1,25VitD3. T reg and Th17 cells were analyzed by flowcytometry after and before treatment with 1,25VitD3.

Results: TGF-β gene expression and IL-10 gene expression also increased in patients with URPL (P value<0.05) after treatment with vitD (P value<0.05). Significant increase in IL-10 level also observed in patients. The results indicated that vitamin D increase Treg/Th17 ratio. There was a significant difference between the percentage of CD4⁺CD25^{bright} CD127⁻ T cells before and after treatment with vitamin D (cells (0.59% vs1.24% P<. 05).

Conclusion: This study showed that the role of vitamin D3 seems to be to provide an anti-inflammatory condition which is in favor of pregnancy maintenance. We can conclude this metabolite can exert as a *supplementary therapeutic* in patients with URSA.

Key words: URSA, T regulatory cells, Th17 cells, 1,25VitD3

ICIA2018\Tolerance and Autoimmunity\Poster\8724
8724

Expression of miR-326 Relapsing-remitting Multiple Sclerosis Responders and Nonresponders to Interferon Beta Therapy

Mahtab Fattahi¹, Nahid Eskandari^{*2}, Fattah Sotoudehnejad¹, Vahid Shaygannejad³

1. Department of biology, science and research branch, Islamic Azad University

2. Department of Immunology, School of Isfahan Medical, Isfahan University of Medical Sciences

3. Department of Neurology, Isfahan Neurosciences Research Center, Isfahan University of Medical Sciences

Background: Multiple sclerosis (MS) is a chronic, inflammatory and demyelinating disease that affects the central nervous system. Th17 cells are the main mediators of inflammatory process in autoimmunity disease including MS. miRNAs are likely involved in most biological processes, and play important roles in the immune system function. Dysregulated expression of miRNAs is associated with pathological conditions, including autoimmune diseases. Interferon beta (IFN- β) represents one of the most commonly administered drugs for the treatment of the RR-MS patients. The purpose of this study was to evaluate miR-326 expression in responders and nonresponders to interferon beta treatment.

Methods: A total of 70 patients including responders and nonresponders to INFB (35) were enrolled. We analyzed the expression level of miR-326 using peripheral blood from RR-MS patients at 12 months after starting with IFN- β therapy. Real-time qPCR was performed to analyze miR-326 expression.

Results: These results indicated miR-326 expression particularly reduced in responders compared to nonresponders, although this was not statistically significant.

Conclusions: Overall, miR-326 expression levels may act as a marker of the biological effects of IFN- β therapy.

Keywords: Interferon-beta, Multiple sclerosis, miR-326

ICIA2018\Reproductive Immunology\Poster\9737
9737

Antisperm antibodies in the serum of unmarried woman (virgin): a case study

Ali Al-fahham¹

Department of Basic Science Faculty of Nursing University of Kufa

Background: The study of antisperm antibodies (ASA) etiology is very important in the diagnosis and treatment of infertility. Studies of presence of antisperm antibodies in the bodies of unmarried women (virgins) are very rare, so that this article aims at studying, in more details, the possible causes and interpretations behind the development of antisperm antibodies in virgins.

Methods: This descriptive case study included 35-years old single women with positive ASA. Description and clinical history of the patients were assessed by special questionnaire provided for this purpose. The concentration of serum ASA was estimated by using Enzyme-Linked Immunosorbent Assay (ELISA). Complete blood count was done by using HumaCount System. Total serum IgG and IgM was calculated by Radial immunodiffusion assay. The isolated bacteria were identified according to microscopically, biochemical tests and VITEK 2 system. All laboratory investigations and diagnostic procedures were done in the hospital between 1st to 15th December 2017.

Results: The mean serum antisperm antibody concentrations is (62.1 IU/ml), it is considered as positive titer. Complete blood count is normal, except for slight increase in WBC count and percentage of basophils, monocytes and lymphocytes. This study recorded high concentrations of serum total IgG and IgM levels (1870 IU/ml and 255 IU/ml respectively). The UTI was confirmed by counting total bacterial concentration (167220 CFU/ml) in the urine, and diagnosis of suspected causes showed the following species: *Escherichia coli*, *Klebsiella pneumonia*, *Staphylococcus haemolyticus* and *Proteus mirabilis*.

Conclusion: There is two suggested mechanisms to explain ASA in virgins: (1) antigen cross-reactivity between sperm and bacterial antigens to which antibodies can react; (2) induction of the immune system by antigens of sperm ingested into the gastrointestinal tract with contaminated food and drink.

Keywords: Antisperm, Antibodies, virgins, Infertility, Case-Study

ICIA2018\Reproductive Immunology\Poster\9740

9740

Evaluation of the expression of virus related Toll like receptors in placenta from hydatidiform mole patients

Mahboobeh Zolala¹, Mohhamad Mahdi Mohhamadi¹, Shahriyar Dabiri², Maryam Iranpoor², Manzoomeh Shamsi Meymandi³, Fatemeh Raeisi Nejad⁴

1. Department of Immunology School of Medical University of Kerman

2. Department of Pathology School of Medical University of Kerman

3. Department of Physiology School of Medical University of Kerman

4. Department of Immunology School of Medical University of Jiroft

Background: Mole hydatidiform is an abnormal pregnancy which is characterized by over proliferation of trophoblasts and vesicular edema in placental villi with or without embryo. The basic etiology of gestational trophoblastic neoplasia is unknown. During trophoblastic disease a considerable over proliferation in placenta trophoblasts in seen which in some cases cause malignancy. In this study, we examined viral TLRs (TLR3,7,8,9) in HM placenta samples using IHC technique to possibly find a correlation between viral infection and HM. in this study, The expression levels of viral TLRs were evaluated in normal placenta and compared with HM ones.

Method: In a case/control study, 39 samples of HM and 43 samples of control obtained from archive of pathology laboratory of target hospital. Certain viral TLRs were monitored using IHC assays. The obtained data were statistically analyzed using Mann Whitney U method.

Results: The differences in TLR3 and TLR9 expression levels between HM and control samples were statistically significant (P value = 0.0001 and 0.03, respectively). In case of TLR7 and TLR8 no significant difference were observed.

Conclusion: Results of this study elucidated the increased expression of TLR3 and TLR9 markers in HM samples comparing control ones. We concluded that HM may be related with dsRNA and CpG DNA virus infections. As TLR9 response is observed during both viral and bacterial infections, it seems that a bacterial infection may be also involved in generation of HM. The expression levels of TLR7 and TLR8 markers in HM samples were same as control ones. So, it could be concluded that ssRNA virus are not involved in HM generation.

Key words: mole hydatidiform, Toll like receptor, immunohistochemistry, placenta

ICIA2018\Reproductive Immunology\Poster\9756
9756

Pro-inflammatory Th17 type and regulatory T cell cytokines in women with unexplained recurrent spontaneous abortion

Narges Roomandeh^{1, 2}, AboTaleb Saremi², Javad Arasteh³, Fatemeh Pak⁴, Majid Mirmohammadkhani⁵, Parviz Kokhaei^{4, 6}, Ahad Zare^{2, 7}

1. Student Research Committee and Department of Immunology, Faculty of Medicine, Semnan University of Medical Sciences, Semnan, Iran
2. Sarem fertility and infertility research center (SAFIR) and Sarem Cell Research Center (SCRC), Sarem Women's Hospital, Tehran, Iran
3. Department of Biology, Faculty of Basic Sciences, Islamic Azad University Central Tehran Branch, Tehran, Iran
4. Cancer Research Center and Department of Immunology, Faculty of Medicine, Semnan University of Medical Sciences, Semnan, Iran
5. Research Center for Social Determinants of Health, Semnan University of Medical Sciences, Semnan, Iran
6. Immune and Gene Therapy Lab, Cancer Centre Karolinska, Karolinska University Hospital, Stockholm, Sweden
7. Immunology, Asthma and Allergy Research Institute, Tehran University of Medical Sciences, Tehran, Iran

Background: Elevated evidences showed that unexplained recurrent spontaneous abortion (URSA) is associated with inflammatory responses and breakage of immunological autotolerance. Th17 cells have an important role in autoimmune disease and loss of tolerance to the fetus. Regulatory T cells play an important role in the maintenance of tolerance during pregnancy. Therefore the balance between Th17 and Treg cells may elucidate the pathophysiology of URSA. In this study we aimed to investigate the serum concentration of regulatory and inflammatory cytokines in both normal and URSA females.

Methods: Forty-six women with URSA and 28 non-pregnant control women with at least one successful pregnancy were included in this study. Serum was obtained from both groups and stored at -70°C. The serum concentration of IL-17, IL-21, IL-22, IL-10 and TGF- β were quantitatively determined by ELISA.

Results: The levels of IL-17, IL-21 and IL-22 in sera were significantly higher in URSA women compared with these in normal controls ($P < 0.001$, $P = 0.01$ and $P < 0.001$ respectively). Also, TGF- β serum concentration as a Treg cytokine, was significantly lower in URSA patients than that in normal women ($p = 0.02$). However, the difference in serum IL-10 level between URSA women and control group was not statistically significant ($P = 0.49$).

Conclusion: Our results suggest that according to higher serum levels of IL-17, IL-21 and IL-22 and lower level of TGF- β in cases compared with healthy controls, enhancement in Th17 associated cytokines levels and reduction in Treg type cytokine may be one of the factors that indicate increased Th17 cell function and reduced Treg activity in URSA.

Key words: IL-21, IL-22, Lymphocyte therapy, URSA

ICIA2018\Tolerance and Autoimmunity\Poster\9773
9773

The Relationship between Serum Level of IL-33 and IL-35 with Amount of Thyroid Auto-antibodies in Pregnant Women with Multiple Sclerosis

Mahdavi R¹, Jamali M², Rostami Rad M³

1. Department of Immunology, Birjand University of Medical Sciences, Birjand, Iran

2. Department of Immunology, Medical School, Kerman University of Medical Sciences, Kerman, Iran

3. Department of Microbiology, Medical School, Tehran University of Medical Sciences, Tehran, Iran

Background: Multiple sclerosis (MS) is an autoimmune-mediated demyelinating disease of the central nervous system (CNS). Interleukine-35 (IL-35) has been described as an immune suppressive cytokine, also IL-33 is a cytokine with both pro- and anti-inflammatory effects. These two cytokines have been implicated in the immune pathogenesis of MS. Thyroid auto-antibody hormones also change during pregnancy and postpartum but these changes may be affected by MS disease. The aim of this study was to determine the association between serum levels of IL-33 and IL-35 with amount of thyroid hormones during early pregnancy and at six months postpartum period in pregnant MS women compared to healthy pregnant women.

Methods: In case-control study, we analyzed 55 pregnant MS women and 55 healthy pregnant women as a control group from January 2016 to December 2017 in the Afzalipoor Hospital, Kerman (a city located in the southeast of Iran). The serum levels of IL-33 and IL-35 were measured by enzymes linked immunosorbent assay (ELISA) and level of thyroid auto-antibody hormones (anti-TG and anti-TPO) were assessed by electrochemiluminescence (ECL) method.

Results: The serum levels of IL-33 and IL-35 in pregnant MS women during the first trimester were significantly higher compared to control group ($P<0.001$) and pregnant women with MS presented with significantly lower anti-TG and anti-TPO in the beginning of pregnancy ($P<0.01$)

Conclusions: We conclude that elevated levels of IL-33 and IL-35 at the beginning of pregnancy may be associated with decreased thyroid auto-antibodies in the first trimester and association between these cytokines with the levels of thyroid hormones at six months postpartum period were reversed.

Keywords: Interleukine-33, Interleukine-35, Multiple sclerosis, Thyroid hormone, Pregnancy

ICIA2018\Research and Development\Poster\9786

9786

Compression of different methods for the rapid establishment of B lymphoblastoid cell lines

Masoumeh Asadi¹, Parvaneh Farzaneh¹, Mehrnaz Izadpanah^{1,4}, Samaneh Mahmood Aghdam¹, Shiva Mohammadi¹, Zahra Elyasi Gorji¹, Seyed Abolhassan Shahzadeh Fazeli^{1,2}, Mansoureh Farhangniya¹

1. Human and Animal Cell Bank, Iranian Biological Resource Center (IBRC), ACECR and Tehran, Iran

2. Department of Molecular and Cellular Biology, Faculty of Basic Sciences and Advanced Technologies in Biology, University of Science and Culture, Tehran, Iran

3. Department of Anatomy, Faculty of Medicine, Iran University of Medical Sciences, Tehran, Iran

4. School of advanced technologies in medicine, Tehran University of Medical Sciences, Tehran, Iran

Background: Recent studies have been reported that Epstein-Barr virus (EBV) immortalization is a Gold Standard method to keep high-molecular-weight DNA in long-term. Therefore, banking of EBV-derived cells has been providing an appropriate reference material for immunologic diseases studying. Since a variety of methods exist for the immortalization of B-lymphocytes by EBV, in this study four different methods were compared to the success rate of immortalization.

Method: Blood samples were collected under sterile conditions and thirty samples were considered for each method. Group A: Peripheral Blood Mononuclear Cells (PBMCs) were isolated and then infected with EBV. Group B: PBMC pellets were frozen using 90% FBS, 10% DMSO for two months. Then, the cells were thawed and infected. Group C: whole blood was frozen and then, the frozen blood was thawed and infected. Group D: whole blood was washed and directly infected.

Result: Lymphoblastoid cell lines (LCL) were simply generated by fourth methods. All the established the cell lines showed typical rosette morphology while single cells were also seen in the all groups. The success rate of the generated LCL in each group was 86%, 93%, 60%, and 70%, respectively.

Conclusion: Our results showed a success rate of the group B was highest. Furthermore, by this method can save time and cost but this method is not applicable for the low volume of blood. Also, it is suggested using D method for blood volume less than 1 ml.

Keywords: Lymphoblastoid cell lines, EBV, Success rate

ICIA2018\Veterinary Immunology\Poster\9808

9808

Evaluating the effects of low molecular weight sodium alginate as an immunostimulator, also as a novel prebiotic combined with Bactocell® (a probiotic) on humoral and mucosal immune responses of Asian seabass (*Latescalcalifer*) juveniles

GhasemAshouri^{1, 3*}, NasrollahMahboobiSoofiani¹, Seyed Hossein Hoseinifar², Seyed Amir Hossein Jalali^{1, 3}, Amir Mani¹, Hien Van Doan⁴

1. Department of Fisheries, Faculty of Natural Resources, Isfahan University of Technology, Isfahan 84156–83111, Iran

2. Department of Fisheries, Faculty of Fisheries and Environmental Sciences, Gorgan University of Agricultural Sciences and Natural Resources, Gorgan, Iran.

3. Department of Research Institute for Biotechnology and Bioengineering, Isfahan University of Technology, Isfahan 84156–83111, Iran

4. Department of Animal and Aquatic Sciences, Faculty of Agriculture, Chiang Mai University, Chiang Mai 50200 Thailand

Background: Using immunostimulants as bio-friendly and environmentally safe agents is an alternative approach to combat diseases in aquaculture. Therefore, the present study was performed to evaluate the potential effect of individual or combined administration of dietary low molecular weight sodium alginate (LMWSA) and Bactocell® on haematological, humoral and skin mucosal immune responses of *L. calcarifer* juveniles.

Methods: Fish (12.0 ± 0.2 g) were fed experimental diets as follows: Control (diet 1, basal diet), 5 g kg⁻¹ LMWSA (diet 2), 10 g kg⁻¹ LMWSA (diet 3), 0.9×10^7 CFU g⁻¹ Bactocell® (diet 4), 5 g kg⁻¹ LMWSA + 0.9×10^7 CFU g⁻¹ Bactocell® (Diet 5), and 10 g kg⁻¹ LMWSA + 0.9×10^7 CFU g⁻¹ Bactocell® (Diet 6). At the end of the trial, blood samples from the caudal vein and skin mucus were collected for evaluation of immunological parameters.

Results: Results indicated a significant ($P < 0.05$) increase in innate immune parameters including serum lysozyme, bactericidal, hemolytic and respiratory burst activities as well as mucosal immune responses including lysozyme and bactericidal activities when diet was supplemented with immunostimulants. Moreover, the combined effects of LMWSA with Bactocell® resulted in more pronounced immunological responses compared to the control and singular administration. Red and white blood cell counts significantly increased with either singular or combined administration of LMWSA and Bactocell® compared with the control group ($P < 0.05$).

Conclusion: These results indicated that combined administration of LMWSA and Bactocell® can be considered as a beneficial feed additive and immunostimulant in *L. calcarifer* juveniles.

Keywords: Asian sea bass, Bactocell®, Sodium alginate, Mucosal immunity, Innate immunity, Haematological parameters

ICIA2018\Vaccine\Poster\9816

9816

Immunogenicity of Oma87 in murine model against *Acinetobacter baumannii* infectionsRaziyeh Abdolhamidi¹, Iraj Rasooli²

1. M.Sc. Student, Department of Biology, Faculty of Basic Sciences, Shahed University, Tehran, Iran

2. Professor, Department of Biology, Faculty of Basic Sciences, Shahed University, Tehran, Iran

Background: *Acinetobacter baumannii* is an emerging gram negative coccobacillus causing nosocomial infections such as pneumonia, meningitis and bacteremia in intensive care units. *A.baumannii* infections are becoming a serious health problem with the rapid advent of drugresistant strains, resulting in urgent need to develop strategies based on non-antibiotic. The purpose of this study is to identify if an outer membrane protein with molecular weight of about 87 kDa (oma87) has the potentials to be an efficient vaccine candidate to combat *A.baumannii* infection.

Methods: Oma87 which has a molecule length of 841 amino acids is conserved in more than 95% of *A.baumannii* strains. Gene encoding an outer membrane protein (ABAYE1583) was identified and was then cloned and expressed in *E.coli*. A recombinant protein with molecular weight of 87 kDa was detected and analyzed with SDS-PAGE. The purified protein was used as an antigen in mice groups.

Results: The antigen elicited antibodies against *A.baumannii*.

Conclusion: In conclusion, Oma87 is a highly safe and protected protein that can serve as a valuable antigen for developing an effective vaccine or antibody preparation to control *A. baumannii* infections.

Keywords: *A.baumannii*, Immunogenicity, Vaccine, OMA87

ICIA2018\Stem Cells Based Immunotherapy\Poster\9835

9835

Wharton's Jelly-Derived Mesenchymal Stem Cell's Conditioned Media Increase Neutrophils Viability and Function

Mohammad Mahmoudi¹, Mahsa Taghavi², Seyed Mahmoud Hashemi^{1,3}, Nima Rezaei^{2,4,5}

1. Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Department of Immunology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

3. Department of Applied Cell Sciences, School of Advanced Technologies in Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

4. Research Center for Immunodeficiencies, Children's Medical Center, Tehran University of Medical Sciences, Dr. Gharib St, Keshavarz Blvd, Tehran, Iran

5. Network of Immunity in Infection, Malignancy and Autoimmunity (NIIMA), Universal Scientific Education and Research Network (USERN), Sheffield, UK

Background: Mesenchymal stem cells (MSCs) have immunomodulatory properties on a variety of leukocyte subsets through direct interaction or releasing soluble factors. They have mostly been defined as immunosuppressive, but recent studies have shown that they have supportive effects on the functions of some immune cells such as Neutrophils, which play an important role in the acute inflammatory response and in the clearance of bacterial infections.

Methods: We investigated the effects of Wharton's jelly-Derived Mesenchymal stem cell's conditioned media on the viability and function of neutrophils. Neutrophils were derived from healthy donors and cultured with or without WJ-MSC's conditioned media to evaluate neutrophil's viability, apoptosis and respiratory burst. The treatment of neutrophils with WJ-MSC' conditioned media resulted in increased viability, decreased apoptosis and enhanced the respiratory burst.

Conclusions: Thus, we conclude that the use of WJ-MSC's conditioned media may be a promising therapy for increasing immunity by promoting the anti-microbial activity of Neutrophils.

Keywords: Mesenchymal Stem Cell, Neutrophil, Conditioned Media, Viability

ICIA2018\Reproductive Immunology\Poster\9863
9863

Association between rs1049174 NKG2D gene polymorphism and unexplained recurrent spontaneous abortion (RSA) in Iranian women

1. Department of Immunology, Faculty of Medicine, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.
2. department of immunology, shahid sadoughi university of medical science , international campus, yazd, iran
3. Department of Genetics and Molecular Biology, Faculty of Medicine, Shahid Sadoughi University of Medical Sciences, Yazd, Iran
4. Reproductive Immunology Research Center, Shahid Sadoughi University of Medical Sciences, Yazd, Iran
5. Recurrent Abortion Research Center, Research and Clinical Center for Infertility, Shahid Sadoughi University of Medical Sciences, Yazd, Iran
6. Research Center for Food Hygiene and Safety, Shahid Sadoughi University of Medical Sciences, Yazd, Iran
7. Department of Genetics and Molecular Biology, School of Medicine, Isfahan University of Medical Science, Isfahan, Iran
8. Zoonotic Diseases Research Center, School of public health, Sahid Sadoughi University of Medical Sciences, Yazd, Iran

Background: According to previous investigations, maternal immune responses may affect the fetus, and cause recurrent pregnancy loss or infertility. Natural killer (NK) cells are the main immune cell population of decidua. They function as an important part of the uterine immune system and play a significant role in outcome of pregnancy. NK cells express inhibitory and activating receptors that regulate their function. NKG2D is one of the best known activating receptor of NK cells which recognizes its ligand on abnormal or stressed cells and activates NK cells to kill them. Objective: We have studied the polymorphism of NKG2D gene to determine alleles and genotypes distribution in Iranian women and whether it could be associated with risk of RSA.

Methods: The genetic polymorphism of NKG2D gene was studied by PCR-RFLP in 140 women with RSA and 175 control women with at least one successful pregnancy and without any known pregnancy loss. Statistical analysis was done after study assessment and data extraction.

Results: A significant difference was observed between patients and controls in regard with GG-genotype frequency of the NKG2D gene. The distribution of genotypes frequencies was also proved to be significantly different between the cases and controls ($p < 0.05$).

Conclusion: This study demonstrated a significant association between NKG2D gene polymorphism (rs1049174 G/C) and risk of RSA in Iranian women.

Keywords: RsnKg2dpolymorphismrpl

ICIA2018\Reproductive Immunology\Poster\9864

9864

Survey of CTLA-4 gene polymorphism in women with recurrent pregnancy loss

Vaziri Nezamdoust F, Hadinedoushan H, Ghasemi N.

1. Department of Immunology, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

2. Recurrent Abortion Research Center, Yazd Reproductive Sciences Institute, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

Background: Recurrent pregnancy loss (RPL) is one of the most important pregnancy problems. Epidemiologic studies have revealed that 1-2% of women experience recurrent pregnancy loss. An aberrant regulation of immunological, metabolic, vascular and endocrine processes leads to RPL. Cytotoxic T lymphocyte-associated antigen 4 (CTLA-4) expresses on active T cells and causes of inhabitation and toleration of active T cells. CTLA-4 gene expression in fibroblast cells of the placenta throughout pregnancy and during pregnancy is increased and causes to tolerance during pregnancy.

Methods: This study was carried out on two groups of women who were referred to the Yazd Reproductive Sciences Institute. 150 women with a history of at least 2 RPL as the case group and 150 women who didn't have a history of abortion with at least one healthy child as the control group. Five milliliter peripheral blood was obtained from all samples and then DNA was extracted. A SNP rs 3087243 of CTLA-4 gene was analyzed using the RFLP-PCR method.

Results: There was no difference between the two groups regarding age of women (26.7 ± 3.5 vs. 28.6 ± 3.6 yr). In the case group, the genotype frequencies of rs 3087243 polymorphisms were AG (46%), AA (24%), and GG (30%), and in the control group, they were AG (59.7%), AA (23.3%), and GG (17.3%). There was a significant difference between the genotypes of AA, AG, and GG in two groups ($p=0.022$).

Conclusion: Our result indicates that the rs 3087243 AG genotype may contribute to RPL development in Iranian women.

Keywords: Recurrent pregnancy loss, Single nucleotide polymorphisms, CTLA-4 gene.

ICIA2018\Stem Cells Based Immunotherapy\Poster\9868
9868

The level of CD133 expression in colorectal cancer cell lines

Zahra Asadzadeh^{1,2}, Elham Safarzadeh^{1,2}, Behzad Mansoori¹, Behzad Baradaran^{1,2*}

1. Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

2. Faculty of Medicine, Department of Immunology, Tabriz University of Medical Sciences, Tabriz, Iran

Introduction: Colorectal cancer is one of the most common malignant tumors worldwide. Colorectal cancer cells have previously been demonstrated to contain a subpopulation of CD133+ tumor cells that have the ability to initiate tumor growth and are thus referred to colorectal cancer stem cells. In this study, we have proposed to evaluate the expression of CD133 in colorectal cancer cell lines.

Methods: Colorectal cancer cell lines HT29, HCT116 and SW480 were cultured in RPMI-1640 medium. Total RNA was isolated from cultured colorectal cancer cells via Trizol reagent (Ambion, TX, USA) according to the manufacturer protocol. Then cDNA was synthesis by the cDNA synthesis kit (Biofact, Korea) and CD133 expression levels were measured using Quantitative reverse transcription PCR (RT-qPCR).

Results: we measured CD133 expression levels in three colorectal cell lines. RT-PCR analysis showed the levels of CD133 expression in HT-29, HCT-116, and SW-480 cell lines which were 98.30 ± 0.20 , 69.62 ± 1.51 , and 0.23 ± 0.06 respectively.

Conclusion: In conclusion, the CD133 was found to be expressed in high levels in HT-29 cell line of colorectal cancer. The results showed the HT-29 cell line might be a good option for researchers to analysis on cancer stem cell biology and therapy in colorectal cancer.

Keywords: CD133 expression, colorectal cancer, Quantitative reverse transcription PCR

ICIA2018\Research and Development\Poster\9874
9874

Association of major depressive disorder with Lipid peroxidation (LPO)

Abdolghader Taene¹, Ghazaleh Khalili Tanha², Gholamreza Anani Sarab³

1. Student Research Committee, Birjand University of Medical Sciences, Birjand, Iran

2. MSc of cellular and molecular biology, Cellular and Molecular Research Center, Birjand university of medical science, Birjand, Iran

3. Cellular and Molecular Research Center, Birjand University of Medical Sciences, Birjand, Iran

Background: Major Depression Disorder (MDD) is one of the most common psychiatric disorders and has been ranked by some estimates as the fourth leading cause of disease burden worldwide. Increasing numbers of studies indicate that oxidative stress play an essential role in pathophysiology and some symptom of MDD. One of the most important damaging effects of free radicals is the onset of lipid peroxidation which leads to destruction of cell membranes and LPO is an indicator for lipid peroxidation.

Methods: Blood samples from forty MDD patients, which 20 patients with treatment and 20 without treatment in first episode, and 20 healthy volunteers as control were collected. Patients diagnosed with MDD according to the Diagnostic and Statistical Manual of Mental Disorders- V (DSM-V). Samples were collected from Department of psychiatry Emam Reza Hospital, in Birjand. The level of Serum LPO was measured by lipid peroxidation kit (Navandsalamat, Iran) according to the manufacturer's instructions. The differences in mean LPO between groups were analyzed.

Results: The result revealed a notable rising in the average of serum MDA in without treatment patient as compared to control group ($P=0.001$), this level showed no difference with treatment group ($P=0.15$). Also the results showed an increase of BDI (Beck Depression Inventory) score (36.42 ± 10.90) in without treatment patients in compare to treatment group (24.47 ± 15.64) and control group (9.56 ± 8.87) ($P<0.005$).

Conclusion: In the present study, we demonstrated oxidative stress in patients with MDD, which are reflected by increased levels of MDA as a marker of oxidative stress.

Keywords: Major depressive disorder, Lipid peroxidation (LPO), oxidative stress

ICIA2018\Veterinary Immunology\Poster\9881
9881

Evaluation of an Indirect ELISA Using a Tachyzoite Surface Antigen SAG1 for Diagnosis of *Toxoplasma gondii* Infection in Cats

Farzaneh Hosseini¹, Morteza Hosseininejad¹

1. Professor, Department of Clinical Sciences, Faculty of Veterinary Medicine, Shahrekord University, Shahrekord, Iran

2. DVM Student of Veterinary Medicine, Shahrekord University, Shahrekord, Iran

Corresponding author's e-mail: hosseini-m@vet.sku.ac.ir

Background: *Toxoplasma gondii* infection is very common in cats throughout the world. Most cats are sub-clinically infected and potentially fatal clinical disease occurs in some of them. The aim of this study is to develop an indirect enzyme linked immunosorbent assay (ELISA) test using an affinity purified tachyzoites surface antigen (SAG1) to detect *T.gondii* infection in cats.

Methods: Six sero-negative kittens were used in this study; four kittens received 10⁴ *T.gondii* tachyzoites of NED strain (type III) and the remaining two were used as uninfected controls. Serum samples were collected within 41 days and were evaluated for anti- *T. gondii* antibodies using indirect fluorescent antibody test (IFAT) and ELISA method.

Results: IgG antibodies were detectable at least from eight days after tachyzoites inoculation and an increasing pattern in both serum ELISA indices (SIns) and IFAT titers were detected. SIns were significantly different in sera of cats presenting different IFAT titers. In order to evaluate the performance of ELISA to detect anti-*T. gondii* antibodies of naturally infected cats, serum samples were also collected from household and stray cats and evaluated in the same way. IFAT was regarded as the standard test and sensitivity and specificity of the ELISA to detect the infection in naturally infected kittens were analyzed using two-graph receiver operating characteristic (TG-ROC) analysis. An area under curve (AUC) of 0.996 revealed the test as a highly accurate test with relative sensitivity and specificity of 100 and 96% for a cut-off value of 0.10 for SIn.

Conclusion: This study develops an ELISA test for the serological diagnosis of *T. gondii* infection in cats and can be used for serological evaluation of the infection with relatively high sensitivity and specificity.

Key words: *Toxoplasma gondii*, surface antigen 1 (SAG1), P30, enzyme linked immunosorbent assay (ELISA), cats.

ICIA2018\Vaccine\Poster\9901
9901

Immunogenic role of ABAYE2132 recombinant protein against *A.baumannii*

Zeinab Mahmoodi¹, Iraj Rasooli²

1. M.Sc. Student, Department of Biology, Faculty of Basic Sciences, Shahed University, Tehran, Iran

2. Corresponding author, Department of Biology, Shahed University, Tehran, Iran

Background: *Acinetobacter baumannii*, an aerobic Gram-negative, non-flagellated opportunistic coccobacillus is one of the most common and highly antibiotic resistant pathogens. It is associated with high morbidity and mortality rates. Adherence of microorganisms to host cells is an initial and important step in the pathogenesis of *Acinetobacter baumannii*. This has led to investigations development of vaccine to prevent or treat *A.baumannii* infections.

Methods: In the present study, a fimbrial protein known as ABAYE2132 which was predicted as vaccine candidate by in silico analysis and was found to be conserved among the *A.baumannii* strains, was cloned and expressed in *E.coli* BL21(2566). The protein was purified by Ni-NTA chromatography. Immunization with ABAYE2132 generated high antibody titers at >64000 serum dilutions. The mice were immunized with recombinant protein and were challenged with *A. baumannii*. The number of bacterial cells in the immunized mice was determined 24h after the challenge.

Results and conclusion: These results indicate that ABAYE2132 vaccination could be a vaccine development strategy to prevent *A.baumannii* infection.

Keywords: *A.baumannii*, Pili, Adhesion, Recombinant protein, Vaccine

ICIA2018\Veterinary Immunology\Poster\9916
9916

Molecular Detection of *Wolbachia pipientis* in *Dirofilaria* infected dogs using HRM Real-Time PCR

FatemeManshori¹, FatemeJalousian^{*1}, Seyed Hossein Hosseini¹, Minoo Soltani²

1. Department of parasitology, Faculty of veterinary medicine, University of Tehran

2. Department of mycology, Faculty of veterinary medicine, University of Tehran

Background: *Dirofilariasis*, one of the most important vector-borne zoonotic infections. This worm causes a severe and fatal disease called heartworm disease in dog; it also creates pulmonary nodules in humans. *Dirofilaria immitis*, harbour intracellular bacteria of the genus *Wolbachia pipientis* (Rickettsiales). These bacteria play an important role in the pathogenesis and immune response to filarial infection.

Objective: The main objective of the present study is to investigate the occult infection of *dirofilariasis* in dogs based on the detection of *Wolbachia*.

Methods: In the present study 138 blood samples from stray dogs from Gilan, Mazandaran and Ghazvin provinces was collected. A sample which had microfilar in peripheral blood was confirmed as *Dirofilaria immitis* positive sample based on morphological and molecular character of microfilar. In this sample, FTSZ gene of *Wolbachia pipientis* was amplified by specific primers and PCR product was sequenced. This sample was introduced as Gold standard and HRM-Real TIME PCR was optimized based on the melting temperature of the Gold standard sample. Other samples was assayed by HRM Real- TIME PCR.

Results: Melting temperature was determined 58.5°C for FTSZ gene of *Wolbachia*. The results showed that in Gilan province 77.3% of samples were positive, in Mazandaran 57.6% and in Ghazvin 28.3% of samples were positive. However using parasitology method (modified knott) it was shown 45.5% infection in Gilan, 15.2% infection in Mazandaran and 2.5% in Ghazvin.

Conclusion: The high percentage of infection based on *wolbachia* detection showed the cases of occult infection of *dirofilariasis* in dogs. This result suggest that alternative diagnostic tests should be applied to determine the occult infections. In addition to, the presence or absence of these bacteria in association with microfilaria is of importance in the management and control of *dirofilariasis* as a zoonotic disease.

Keywords: *Dirofilaria immitis*, *Wolbachia pipientis*, HRM-Real Time PCR, Iran

ICIA2018\Veterinary Immunology\Poster\9935
9935

Serological Evaluation of Experimental *Toxoplasma gondii* Infection in Cats by Using Immunoblotting Based on an Affinity Purified Surface Antigen

Mohammadreza Alijani^{*1}, Morteza Hosseini², Farzaneh Hosseini²

1. DVM Student of Veterinary Medicine, Shahrekord University, Shahrekord, Iran

2. Professor, Department of Clinical Sciences, Faculty of Veterinary Medicine, Shahrekord University, Shahrekord, Iran

Corresponding author's e-mail: mohammadrezaalijani1992 @gmail.com

Background: *Toxoplasma gondii* is an apicomplexan parasite that infects human and almost all warm-blooded animals. The life cycle of the parasite includes an asexual reproduction in intermediate hosts (Mammals and birds) and a sexual reproduction in definitive hosts. Cats are both the intermediate and the definitive host for *T. gondii*. The aim of this study was to investigate anti-*T. gondii* antibodies in cats by using an immunoblot method based on a major surface antigen; SAG1.

Methods: Six sero-negative kittens were infected intraperitoneal by Rh strain of *T. gondii*. Serum samples were collected and evaluated for the presence of IgG antibodies by using two techniques; immunoblotting and indirect fluorescent antibody test (IFAT). SAG1 based immunoblotting was able to detect anti-*T. gondii* antibodies at least eight days after infection.

Results: The results obtained from immunoblotting shows that it is comparable with the standard IFAT test so that infection is detectable in early stages of the disease when serum samples are evaluated with this immunoblotting method.

Conclusion: According results of this study shows that Immunoblotting using affinity purified *T. gondii* SAG1 antigen is a sensitive diagnostic test and is capable to diagnose early *Toxoplasma gondii* infection.

Keywords: Cats; Immunoblot; P30; SAG1; *Toxoplasma gondii*.

ICIA2018\Reproductive Immunology\Poster\9940
9940

***Aloe vera*-induced modulation of nitric oxide signaling pathway and androgenesis in the testis rat leydig cells**

Samira Asgharzade ^{1*}, Mahmoud Rafieian kopaei ², Amin Mirzaeian ¹

1. Cellular and Molecular Research Center, Basic Health Sciences Institute, Shahrekord University of Medical Sciences, Shahrekord, Iran

2. Medical Plant Research Center, Basic Health Sciences Institute, Shahrekord University of Medical Sciences, Shahrekord, Iran

Background: Nitric oxide (NO), a product of inducible nitric oxide synthase (iNOS), contributes in germ cell apoptosis. This study was aimed to evaluate the effects of *Aloe vera* gel (AVG) on male Wistar rat reproductive organ, serum NO level, and expression of iNOS gene in leydig cells.

Methods: Adult male Wistar rats (n=36) were used for experiments in three groups. The experimental groups were orally administered with the AVG extract solution once-daily as follow: 150 mg.kg-1; group A, 300 mg.kg-1; group B, and only normal saline; group C (control group). They were mated with untreated females and the reproductive and chemical parameters were assessed for each group, including semen quality, serum testosterone, sperm fertility, gonad and body weight, serum NO concentration (by the griess method), and iNOS gene expression (using RT-PCR).

Results: The testes weight, serum testosterone, as well as sperm count and fertility of the AVG treated groups were significantly reduced when compared to the control ($P<0.001$). Concentration of serum NO was significantly increased ($37.1\pm4.63\ \mu\text{M}$) in the administrated group with higher AVG concentration, compared to the control group ($P<0.001$; $10.19\pm0.87\ \mu\text{M}$); however, iNOS mRNA expression was increased in the treated animals ($P<0.001$).

Conclusion: iNOS may play a functional role in spermatogenesis via apoptosis, reducing sperm count, but further studies are needed to illustrate the mechanisms by which AVG exerts its negative effects on spermatogenesis and sperm quality.

Keywords: Nitric oxide, *Aloe vera*, rat, leydig cells

ICIA2018\Vaccine\Poster\10024
10024

Design and in silico analysis of a new antigen, based on heat stable toxin, to be incorporated in candidate vaccines against enterotoxigenic *Escherichia coli* (ETEC)

Abbas Hajizade^{1*}, Ali Hatef Salmanian², Jafar Amani³, Firouz Ebrahimi¹, Ayoob Arpanaei⁴, Yousof Tarverdizade¹

1. Biology Research Centre, Faculty of Basic Sciences, Imam Hossein University, Tehran, Iran

2. Plant biotechnology department, National Institute for Genetic Engineering and Biotechnology (NIGEB), Tehran, Iran

3. Applied Microbiology Research Centre, Baqiyatallah University of Medical Sciences, Tehran, Iran

4. Department of Industrial and Environmental Biotechnology, National Institute of Genetic Engineering and Biotechnology, Tehran, Iran

Background: As one of the main bacterial leading causes of morbidity and mortality in children under 5 years of age, especially in developing countries, enterotoxigenic *Escherichia coli* (ETEC) are of the main causative agents of infectious diarrhea that play a significant role in public health. Indeed, the bacteria are responsible for the majority of traveler's diarrhea. So, it is necessary to develop an efficient vaccine against the pathogen. Pathogenic strains of ETEC produce two different enterotoxins: heat-labile and/or heat-stable toxins along with many colonization factors. All pathogenic strains of ETEC express at least one of these toxins. There are many proposed candidate vaccines against ETEC based on LT toxin. However, few candidate vaccines are introduced against STa. Progresses in bioinformatics have led to the development of many in silico tools which can help us to design and evaluate the immunogenicity of the vaccine candidates.

Methods: Here, in this study we designed a new antigen based on a mutated form of STa toxin. The physico-chemical parameters and the secondary and tertiary structures of the antigen were calculated. Prediction of the antigenic B-cell and T-cell epitopes were performed by appropriate softwares. The protein was then reverse-translated to DNA and the resulted DNA was codon optimized according to the codon preference of *E. coli* B121(DE3) host cells.

Results: The results showed that the structure of the designed antigen has a proper 3D structure in which the main epitopes are exposed and the domains are separated. There are potent B- and T-cell epitopes in the protein. Physicochemical properties of the protein as well as the structure of mRNA showed that the designed gene can be translated in *E. coli* host cells.

Conclusion: In conclusion, the designed antigen structurally and immunologically has almost all factors to be incorporated in candidate vaccines.

Keywords: Heat stable toxin (STa), enterotoxigenic *Escherichia coli* (ETEC), Candidate vaccine, Immunoinformatics

ICIA2018\Reproductive Immunology\Poster\10027

10027

Relation of Gene's expression Profile of IL-17 and IL-23 in Cumulus cells and their Follicular Fluid Concentrations in Infertile Patients at the Risk of Ovarian Hyperstimulation Syndrome (OHSS)

Leila Motamedzadeh ^{1, 3}, Hossein Hadinedoushan ², Ehsan Farashahi Yazd ³, Farzaneh Fesahat ³, Mohammad Mahdi Mohammadi ^{1, 4}

1. Department of Immunology & Allergy, Kerman University of Medical Sciences, Kerman, Iran

2. Department of Immunology, Reproductive Immunology Research Center, Shahid Sadoughi University of Medical Science, Yazd, Iran

3. Research and Clinical Center for Infertility, Shahid Sadoughi University of Medical Science, Yazd, Iran

4. Physiology Research Center, Kerman University of Medical Sciences, Kerman, Iran

Background: Ovarian hyperstimulation syndrome (OHSS) is one of major complications during assisted reproductive technology treatment. This study aims to determine the concentrations of interleukin IL-17 and IL-23 in follicular fluid (FF) as well as their mRNA levels in cumulus cells of infertile women at the risk of OHSS.

Methods: The cumulus cells of oocytes from 80 infertile women were divided into two groups of case (n=40) with an average age 29.6 ± 1.4 years and control (n=40) with an average age 29.2 ± 1.1 according to the patients' infertility etiologies. The control group was the infertile patients with male factor infertility etiologies while; infertile women at the risk of OHSS were defined as case group in this study. IL-17 and IL-23 concentrations in FF were determined. mRNA expression levels of IL-17 and IL-23 of samples were measured using real time polymerase chain reaction method in all subjects.

Results: The concentrations obtained from FF demonstrated the higher levels of IL-17 in case group ($P=0.04$), while, it showed no remarkable difference in IL-23 ($P=0.3$). The mRNA levels of IL-17 and IL-23 showed no significant differences between two groups. There was a positive significant correlation between FF concentrations of IL-23 and the oocytes maturation rates in case group ($P=0.01$). The numbers of follicles, MII oocytes, immature oocytes, fertilized oocytes and formed embryos were significantly higher in case group.

Conclusion: Our finding showed the mRNA expression of IL-17 and IL-23 were similar in two groups, and IL-17 may be considered as risk factor for OHSS.

Key Words: OHSS, IL-23, IL-17

ICIA2018\Reproductive Immunology\Poster\10033
10033

Evaluation of Regulatory B cell in Blood of Women with Recurrent Spontaneous Abortion (RSA)

Farzad Nazari¹, Davar Amani¹, Esmaeil Mortaz¹

Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: The main function of regulatory B cells (Breg) is producing of IL-10 and this cytokine control immune balance during early human pregnancy, so Breg cells emerge as important players in pregnancy. The function of Breg in immune-mediated reproductive disorders (e.g. Recurrent Spontaneous Abortion (RSA)) is unknown. The aim of this study was to evaluate the Breg cells in RSA patients.

Methods: Fresh blood of 10 RSA patients and 12 Control were harvested and then the surface expression of CD19⁺, CD24^{hi}, and CD38^{hi} on B cells were measured by flowcytometry (FACS calibur), and about 10,000 events were measured in all cases.

Results: Significant upregulation of B cells with CD19⁺ CD24^{hi} CD38^{hi} phenotype (Breg cells) was seen in normal pregnant (mean: 7.83%) whereas downregulation of markers was seen in these cells in women suffered from RSA (mean: 0.9%) at the first trimester. Statistical analysis is going.

Conclusion: Our data show that decrease in Breg cells in early gestation might have important role in recurrent spontaneous abortion via regulation of immune response.

Keywords: Regulatory B cell, IL-10, Abortion, Reproductive immunology, RSA

ICIA2018\Reproductive Immunology\Poster\10034
10034

Evaluation of regulatory B cell in blood in women with recurrent spontaneous abortion (RSA) and Normal Pregnant Individual

Farzad Nazari¹, Davar Amani¹, Esmacil Mortaz¹

Department of Immunology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Regulatory B lymphocytes are important players of immune response by producing of IL-10 during human normal pregnancy, whereas the function of this cell during Recurrent Spontaneous Abortion (RSA) is unknown. The goal of this investigation was to determine Breg cells percent in RSA patients in comparison of normal pregnant individuals.

Methods: Fresh blood of 10 RSA patients and 10 Control (non-pregnant women) was harvested and then the surface expression of CD19⁺, CD24^{hi} and CD38^{hi} on B cells were measured by flow cytometry (FACS calibur), and about 10,000 events were measured in all cases.

Results: The percent of Breg cells (4.18%) in non-pregnant women was significantly higher than RSA patients (0.9%). $P < 0.05$

Conclusion: The result of this study show that Breg cells might have important role in RSA.

Keywords: Regulatory B cell, IL-10, Abortion, Reproductive immunology, RSA

ICIA2018\Veterinary Immunology\Poster\10045

10045

Evaluation of the production level of the polyclonal antibody against *Listeria monocytogenes* in animal models

Mansoor Khakpoor¹, Yasin Bagheri^{2*}, Alireza Barati

1. Department of Pathobiology, Tabriz Branch, Islamic Azad University, Tabriz, Iran.

2. Young Researchers And Elite club, Tabriz Branch Islamic Azad university, Tabriz, Iran

Background: *Listeria monocytogenes* is a short basillary Gram-positive, spore-free and intracellular parasitism which transmitted through contaminated vegetables, milk, cheese and meat to human or animal. Immunoglobulin Y (IgY) is a great and cheap source for polyclonal antibodies that obtained from egg yolk of immunized laying hens. These antibodies can be an appropriate replacement for other antibodies that produce by laboratory animals. Recent years, IgY is applied to preventing, diagnosis and treatment of disease in human medicine researches.

Methods: In current study, 12 laying hens were divided into 2 groups: Control group (2 hens) and Treatments (10 hens). Treatments immunized with *Listeria monocytogenes* antigens in 3 steps.

Results: Finally, by measuring total protein and yolk globulins of collected eggs were proved that there was no significant difference in levels of globulins obtained in 9 and 10 weeks after first injection while they showed a significant difference compared to the control group ($p < 0.05$).

Conclusion: The amount of globulin and total protein in the eleventh week had significant reduction and showed no significant difference with controls. Accordingly, it could be concluded that hens' immunization increased yolk globulin levels in the ninth and tenth weeks after injection.

Key words: polyclonal antibodies, *Listeria monocytogenes*, egg yolk

ICIA2018\Vaccine\Poster\11063

11063

Induction of Immune Responses by DNA Vaccines Formulated With Dendrimer and Poly (Methyl Methacrylate) (PMMA) Nano-Adjuvants in BALB/c Mice Infected With Leishmania major

Fatemeh Tabatabaie¹, Nasim Samarghmarghandi², Somayeh Zarrati³, Fatemeh Maleki⁴, Mehdi ShafieeArdestani⁵, Taher Elmi¹, Sayed Hussain Mosawi⁶

1. Department of Parasitology and mycology, School of Medicine, Iran University of Medical sciences, Tehran, Iran

2. Department of Immunology, Faculty of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

3. Department of Microbiology, Science & research Branch, Islamic Azad University, Tehran, Iran

4. Department of Parasitology, Faculty of Para Medical Sciences, Iran University of Medical Science, Tehran, Iran

5. Department of Radiopharmacy, Faculty of Pharmacy, Tehran University of Medical sciences, Tehran, Iran

6. Department of Biology and Microbiology, Faculty of Medical Technology, Khatam AL Nabieen University, Kabul, Afghanistan

Background: Leishmaniasis is a parasitic disease induced by a protozoan from the genus Leishmania. No effective vaccine has yet been developed against the disease.

Methods: After the plasmid construction and preparation of adjuvants, three intramuscular injections of the nano-vaccin (100 µg) and the recombinant TSA protein (20 µg) were subcutaneously performed. Eventually, the challenged animals were infected with the parasites (1×10^6 promastigotes). After the last injections of the nano-vaccines, the responses of their antibody subclasses and cytokines were assessed via ELISA method before and after the challenge.

Results: This study revealed that the new nano-vaccines were strong and effective in inducing specific antibody and cellular responses and reducing the parasite burden in the spleen compared to the control groups of Leishmania major-infected BALB/c mice.

Conclusion: Based on the results, we can suggest that the formulated vaccines are suitable candidates for further studies in the field of leishmaniasis control.

Keywords: PMMA, Dendrimer, Leishmaniasis, TSA

ICIA2018\Veterinary Immunology\Poster\11088

11088

Immunogenicity of the attenuated line of *Leishmania infantum* in dog

Zeynab Fallah¹, Hamid Daneshvar², Hossin Kamiabi³

1. Department of Parasitology, Medical School of Kerman University, Kerman

2. Department of Immunology, Medical School of Kerman University, Kerman

3. Department of Parasitology, Medical School of Kerman University, Kerman

Background: *Leishmania infantum* (*L. infantum*), a causative agent of visceral leishmaniasis (VL), is endemic in Iran and several Middle East (1). Control interventions involving immunization of dogs to prevent infection could be beneficial as dogs are both the natural host and the main reservoir of *L. infantum* (2). Control of VL in dogs can lead to a reduced prevalence of disease in associated human populations (3).

Methods: Promastigotes of *L. infantum* JPCM5 (MCAN/ES/98/LIM-877) were cultivated in HOMEM with 10% FCS in the presence of gentamicin. The attenuated line of parasite was used at a concentration of 10^9 cells/mL in PBS. Serum was prepared from the blood samples taken derived from the peripheral veins of the dogs. Serum *Leishmania*-specific IgG was measured using ELISA.

Results: All dogs presented increased levels of IgG antibodies over the 12-month period of observation.

Conclusion: The attenuated line of *L. infantum* was immunogen and induced anti leishmanial-IgG.

Keywords: Attenuated line of *Leishmania infantum*, dog, immunogenicity

ICIA2018\Immunology of chemical victims\11171

11171

Evaluation of serum levels of Pro-inflammatory cytokines)TNF- α , IL1- β (in Sulfur Mustard(SM) exposed patients with long term pulmonary complications

Ashraf vafa , Tooba Ghazanfari1, Naghizadeh MM

1. Immunoregulation Research Center, Shahed University, Tehran, Iran

Background: Exposure to mustard gas leads to acute and chronic toxic effects on the pulmonary system. The results of various studies have been suggested the role of Pro-inflammatory cytokines including IL-1 β and TNF- α in the pathogenesis of Pulmonary inflammatory diseases. IL- β and TNF- α are recognized as a key mediator of systemic inflammatory reactions and numerous changes including altered cell signaling, migration, changing on cytokine production and fever.

Aim: The aim of this study is the evaluation of serum levels of IL-1 β and TNF α in SM-exposed patients with long term pulmonary complications.

Materials and Methods: In this study, 65 SM-exposed patients as exposed group and 69 unexposed people as control group were studied. The both exposed and control groups were homogeneous in terms of gender, age and body mass index. Pulmonary function test was performed by spirometry. The serum levels of IL-1 β , TNF- α were measured by ELISA method.

Results: The serum levels of TNF- α was high in SM-exposed group compared to control group. But There were no significant differences between the serum levels of IL-1 β in SM exposed group and the control group.

Conclusion: Due to increased serum levels of TNF- α in exposed group compared to the control group, It may play an important role in pulmonary inflammatory complications. It is a proof of TNF α effects on pulmonary system as a mediator of inflammatory reactions. Although there were no significant differences in the serum levels of IL-1 β in SM-exposed group compared to the control group, It doesn't rollout the role of IL-1 β in pathogenesis of long term pulmonary complications, because it may have local changes in pulmonary system and long term complications. Therefore, locally assessment of IL-1 β would be valuable.

Keywords: Mustard Gas; TNF α ; IL-1 β ; pulmonary complications ;

ICIA2018\Immunohematology \12215
12215

Long-term effects of sulfur mustard on viability of peripheral blood mononuclear cells

Ensieh Sadat Mirsharif ¹, Tooba Ghazanfari ^{1*}

1. Immunoregulation Research center, Shahed University, Tehran, Iran

Background: Sulfur mustard (SM) or (bis-[2-chloroethyl] sulfide) is a chemical warfare agent that attacks mainly skin, eye and lungs. Although Medical effects of exposure to SM in humans include ocular, dermal, respiratory, reproductive and developmental toxicity, gastrointestinal injuries, hematological, and cancer, but the effects of SM-exposure on body are not restricted to these known injuries. The aim of this study is examine the long-term effects of SM on viability of peripheral blood mononuclear cells in SM victims of the Iraq-Iran war (1980–1988).

Materials and methods: During a case -control study, 110 SM-exposed individuals and 109 unexposed as control group were recruited. PBMC were purified from whole blood samples of participants using Ficol-Paque gradient. Cells were suspended in culture medium RPMI 1640, 10% FBS and cultured in the absence and presence of PHA in 96-well plate (1×10^6 cells/ml) at 37 °C and 5% CO₂ for 48 h. Cell viability was assessed with MTT assay. Stimulation index was calculated and compared between groups.

Results: The Stimulation index (Percentage of surviving cell) of PBMC in SM-exposed group and control groups were 111.83 ± 20.71 and 114.62 ± 22.68 , respectively. There were no significant differences in exposed group compared with controls. There was no significant difference in the viability of PBMC in SM-exposed group compared to their matched control subjects.

Conclusion: The results of this study indicate that long-term complications of mustard gas may not be associated with viability of the blood cells.

Key words: Sulfur mustard, long-term effect, peripheral blood cells, viability

ICIA2018\Innate immunity and inflammation\12216
12216

Serum level of IL-17A and long term pulmonary complications (27 years after sulfur mustard exposure)

Ali Mohammad Mohseni Majd^{1*}, Tooba Ghazanfari¹, Mohammad Mehdi Naghizadeh¹

1. Immunoregulation research center of shahed University, Tehran, Iran

Background: Interleukin 17A (IL-17A) is the one of key cytokines produced by human activated Th17 and is the most important cytokine in family of Th17 cytokines about lung immunity. This cytokine as the pro-inflammatory cytokine was described recently and particularly as an important parameter for chronic inflammatory diseases. Growing up of IL-17A level of serum has been associated with chronic inflammatory of lung complications and it can be effective on sulfur mustard (SM)-lung as distinct disease.

Methods: We collected serum samples from 112 healthy individuals and 109 SM-injured patients with prolonged pulmonary complications and were classified based on Auscultation, Pulmonary assessment, pulmonary symptoms and respiratory consequence severity. Serum level of IL-17A was investigated by ELISA method.

Results: The serum level of IL-17A in SM-exposed group and non-exposed group showed no significant difference. Our results also show there are no significant auscultation comparison between SM-exposed group and control group. Furthermore, the results have not showed any significant comparison in cough and hemoptysis groups between SM-exposed group and non-exposed group.

Conclusion: Generally, IL-17A may not be a certain agent in pathogenesis of lung complications but could intensify the effects of disease and make it worse; also the mutation on STAT3 gene can cause to decrease of IL-17A level. This toxic gas could affect on genes expression of different pathways in long term. STAT3 gene deficiency can cause to decrease of TH-17 cells number by non-differentiation in origin, subsequently the level of IL-17A as a main cytokine of TH-17 will be in low level.

Keywords: Interleukin 17A, Th17, lung, inflammation, chronic disease.

ICIA2018\Immunoparasitology \12217

12217

Effects of hydroalcoholic extract of *Glycyrrhiza glabra* and glycyrrhizic acid on nitric oxide production in peritoneal macrophages of *Leishmania major* infected BALB/c mice

Shima Sheikhi¹, Ali Khamesipour², Tayebah Radjabian³, Tooba Ghazanfari¹

1. Immunoregulation Research Center, Shahed University, Tehran, Iran

2. Center for Research and Training in Skin Diseases and Leprosy, Tehran University of Medical Sciences, Tehran, Iran

3. Department of Biology, Faculty of Basic Sciences, Shahed University, Tehran, Iran

Background: Cutaneous leishmaniasis is a neglected disease caused by different species of *Leishmania*. The macrophage lineage is the main cells for the growth of *Leishmania* and nitric oxide(NO) production by them is one of the ways to destroy microorganisms. There are some natural drugs, including *Glycyrrhiza glabra* extract and its components, which showed to act as immunomodulator and enhance macrophage activation. The aim of this study was to explore the effect of hydroalcoholic extract of *G. glabra* and its main component glycyrrhizic acid on macrophage NO production in *Leishmania major* infected BALB/c mice.

Methods: 35 *Leishmania major* infected BALB/c mice were randomly divided into 5 groups; the groups of mice were treated intraperitoneally for one month as following; 200 or 600 mg/kg of *G. glabra* root extract, its component glycyrrhizic acid (200mg/kg), *Glucantime* (160 mg/kg) or phosphate buffered saline as a negative control. The mice were scarified and the peritoneal macrophages were removed and the nitric oxide level was measured using Griess test.

Results: The mean production of nitric oxide in macrophages of groups received 200 or 600 mg/kg of the extracts, glycyrrhizic acid, *Glucantime* and negative control was 11.7, 9.7, 5.1, 6.9 and 3.6nmol/ml, respectively. The highest amount of nitric oxide was produced in the groups that received the extract, and it was statistically significant in comparison with the negative control, or glycyrrhizic acid or *Glucantime* treated groups.

Conclusion: The hydroalcoholic root extract of *G. glabra* induced macrophage activation, which resulted in an increase in nitric oxide production in macrophages of *Leishmania major* infected BALB/c mice.

Keyword: Nitric Oxide, Macrophage activation, *Leishmania major*, *Glycyrrhiza glabra*

ICIA2018\Immunology of chemical victims\12218
12218

Evaluation of IL-2 and IL-10 level in peripheral blood mononuclear cells of sulfur mustard exposed Victims

Nafiseh Zand, Tooba Ghazanfari¹

1. Immunoregulation Research Center, Shahed university, Tehran, Iran

Background: Sulfur mustard (SM) is a chemical agent which has short and long term toxicity against many organs. The respiratory tract is one of the main targets, and is the most disabling long term complication of SM. IL-10 is an anti-inflammatory cytokine that is important in the regulation of immune response and increased in chronic inflammation and IL-2 has a critical role in regulating both cellular and humoral chronic inflammatory responses. These cytokines play a main role in the pathogenesis of chronic airway inflammation in asthma and chronic obstructive pulmonary disease. The aim of this study is to evaluate the level of IL-2 and IL-10 in PBMC of sulfur mustard exposed Victims.

Methods: 100 SM exposed patients and 109 age and sex matched control subject were included in case-control study. Their PBMC were isolated from whole blood by using ficol-paque gradient. The cells were allowed to proliferate and secrete cytokines through cultured in 37°C for 24h with 5 % CO₂ with and without PHA as a mitogen stimulator. The levels of IL-2 and IL-10 of supernatant of culture were measured by ELISA methods.

Results: The baseline of IL-2 and IL-10 in exposed group were significantly higher than control group ($P < 0.05$). But PHA-induced IL-2 and IL-10 level were not significantly different between control and exposed group ($P > 0.05$).

Conclusion: Based on the results of present study, Mustard gas affects the production of these cytokines in the exposed Victims even after years of exposure. So, this dis-regulation of immune system may has important role on the respiratory and immune systems at SM exposed people in long term.

Keywords: Sulfur mustard, IL-2, IL-10, chronic inflammation

ICIA2018\ Psychoneuroimmunology and immunoendocrinology \12219
12219

The effect of instability stress and exogenous oxytocin on the number of the natural killer cells in peripheral blood and spleen in male rats

Zahra Omid¹, Mohammad Reza VaezMahdavi², Mohammad Vojgani³, Marzieh Eghtedar doost¹, Nayere Askari⁴, Tooba Ghazafari¹

1. Immunoregulation Research Center, Shahed University, Tehran. Iran

2. Department of physiology, Faculty of Medical Sciences, Shahed University, Tehran. Iran

3. Department of Immunology, Faculty of Medical Sciences, Tehran University, Tehran. Iran

4. Department of physiology, Faculty of Basic Sciences, Shahid Bahonar University, Kerman. Iran

Background: Natural killer cells are innate immune lymphocytes that they play a major role in defense against tumors and viral infections. Many factors such as social stress act on the immune system. This type of stress is an important factor in the etiology of psychopathological disorders, including depression and anxiety. Oxytocin is a 9-amino acid hormones synthesize in the central nervous system and some of the peripheral tissue and plays a main role in response to stress, and its injection reduces depression and stress-induced anxiety. The aim of this study was the evaluation of instability stress effect and oxytocin on the number of NK cells in rats exposed to this stress.

Methods: WISTAR rats were subjected to instability stress for 21 days, hence their cage-mate were changed every three days. From the 11th day, a group of rats received 20 microliter and the other group received 40 microliter 1 mg/ml oxytocin and the control group received normal saline by intranasal rout. At the end of study, the animals were anesthesia and then were killed. Blood sample and spleen tissue were obtained and the number of NK cells was counted by flow cytometry by CD3-CD161 + markers.

Results: The number of NK cells in the peripheral blood and spleen was 2.67% and 4.12% (received low dose) and 3.80% and 4.69% (received high dose) and the normal saline group were 5.20% and 4.48% respectively. Statistically, there was no significant difference in the NK cells count in between groups that receiving oxytocin with different doses and with control group.

Conclusion: Instability stress had not effect on the number of NK cells in the spleen and blood, and also the doses of 20 and 40 µL of 1mg/ml oxytocin did not a significant effect on the count of NK cells.

Keywords: Natural killer cell, Instability stress, Oxytocin

ICIA2018\Mucosal and Gastrointestinal Immunology\Oral\12224
 12224

Study the correlation of serum IL-6 with stages of tumor in patient suffered from colorectal cancer

Hossein Ali Khazaei¹, Alireza Khazaei², Naeges Sarlogzaei³, Ehsan Khazaei⁴, Amin Khazaei⁵, Bahman Khazaei⁶, Amin Roustami⁶, Fariba Hejazynia⁷, Sina Nabi Zadeh⁶

1. Clinical Immunology Research Center, Department of Immunology and Hematology, Zahedan University of Medical Sciences, Zahedan-Iran
2. Associated prof in Surgery at Faculty of dentistry. Zahedan University of Medical Sciences. Zahedan-Iran
3. Associated prof in Social medicine Faculty of Medicine, Zahedan University of Medical Sciences. Zahedan, Iran
4. Zahedan University of Medical Sciences, Zahedan-Iran
5. Dentist
6. MD
7. Ali Ebneh Abi Taleb Hospital of Zahedan University of Medical Sciences, Zahedan-Iran

Background: Colorectal cancer is the second leading cancer in the five most common cancer in men and women. Cytokine including IL-6 is involved in cancer and various autoimmune diseases. Increased levels of IL-6 in serum and tumor tissue of patients with colorectal cancer, ovarian, breast, and prostate cancers have been reported. The aim of this study was to evaluate serum levels of IL-6 with stages of tumor in patients suffered from colorectal cancer.

Materials and Methods: This cross-sectional study was carried out on 40 adult patients with colorectal cancer referred to a hospital clinic in Zahedan Ali Ebnh Abi Talib. Serum levels of interleukin-6 in these patients was assessed using ELISA method. Results obtained were analyzed by using ANOVA and Tukey tests.

Results: The mean serum levels of IL-6 was obtained as 0.86 ± 2.012 pg/ml that was varied from at least 0.86 to up to 4.1 pg/ml. Based on tumor size in patients with colorectal cancer, the results showed in patients with tumor size of less than 20 mm, the concentrating of IL-6 was 1.36 ± 0.42 pg/ml and in patients with more than tumor size of 20 mm was obtained as 2.2 ± 0.86 pg/ml. There was significant difference between the two averages as confirmed by t-test. In other words, the levels of IL-6 in patients with large tumor size is significantly higher. Significant differences was observed based on the levels of IL-6 and pathologic condition of tumor. Comparison of the two stages according to Tukey test showed that the serum levels of IL-6 level was similar in stage I and II and had lowest amount of IL-6. Stage IV had significantly higher levels of IL-6 and the levels of IL-6 in stages III had significantly higher than in stage I and II and lower than in stage IV. In other words, by increasing the level of staging, the amount of interleukin-6 was significantly increased (P value = 0.000).

Conclusion: The results of this study showed that between the amount of serum levels of interleukin-6 and tumor stage and tumor size were significant correlation. Our findings emphasize the importance and immunological role of inflammatory cytokines, particularly interleukin-6 in this disease.

Keywords: IL-6, stages of tumor and colorectal cancer

ICIA2018\Immunology of Chemical Victims and Environmental Pollution\Poster Discussion\7354
7354

Evaluation of some of the factors associated with immune system following effect of magnetic field in rats

Amin Lafzi¹, Mansour Khakpur²

1. Dvm, Veterinary Medicine Clinician, Tabriz, Iran

2. Department Of Pathobiology, Tabriz Branch, Islamic Azad University, Tabriz, Iran

Abstract:

This study was designed to investigate the effects of electromagnetic fields (EMF) uniformly on the immune system of 12 Norwegian male Wistar rats were used as experimental model. These rats were divided into three control, treatment one and treatment two groups. The first treatment group was subjected to electromagnetic field for four hours and fifteen days. The second group received this treatment for eight hours and fifteen days. At the end of the fifteenth day, blood samples from rats were taken and were prepared to be analyzed. Then, blood cells were counted and after conducting biochemical tests, the data were analyzed using SPSS computer application, one-way ANOVA and Tukey range tests. The results revealed that the level of WBC increased in the first experimental group compared to control group; yet, it decreased in the second experimental group. The level of total protein relatively decreased compared to control group; however, level of albumin decreased significantly and at the same time Gamma globulin increased. Eventually, the increase in the number of lymphocytes and decrease in the number of neutrophils and monocytes are the most significant effects of electromagnetic field on immune system.

Keywords: Immune system; Magnetic field; Rats. ICIA2018\Immunology of Chemical Victims and

Environmental Pollution\Poster Discution\7621

7621

Effect of extremely low frequency electromagnetic fields (ELF-EMFs) on serum levels of IL-6, IL-21 and expression of BCL-6, AID genes in spleens of rats

Mohammad Gholamian Hamadan^{1,2}, Mahdi Alahgholi Hajibehzad¹, Hanie Mahaki^{1,3}, Zohreh Zareighane¹, Elaheh Talebi Ghane⁴, Alireza Zamani^{1,3}

1. Department of Immunology, School of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

2. Student Research Committee, Hamadan University of Medical Sciences, Hamadan, Iran

3. Molecular Immunology Research Group, Research Center for Molecular Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

4. Department of Biostatistics, School of Paramedical Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: The spleen is the important organ in humoral immune system against blood antigens. Some environmental factors such as extremely low frequency electromagnetic fields (ELF-EMFs) may cause harmful effects on it. The purpose of this study was assessing the role of long-term exposure of ELF-EMFs in different intensities on expression of B-cell lymphoma 6 protein (BCL-6) gene, and Activation-induced cytidine deaminase (AID) gene.

Methods: Eighty adult male rats were exposed to ELF-EMFs in intensities of 1,100,500 and 2000 μ T at frequency of 50 Hz for 2 hours daily for up to 60 days.

Results: The results showed that AID gene expression was declined in low intensity of 1 μ T compared with control group, while no changes were observed after ELF-EMFs exposure on expression levels in other intensity and other gene. Serum level of IL-6 increased in 500 μ T group in compared with other groups. Also, serum level of IL-21 have no significant change in all groups.

Conclusion: ELF-EMFs exposure does not alter differentiation of T_{FH} (Tcell Follicular Helper), the most important factor for initiation specific humoral immune system but can affect on Centroblasts.

Keywords: BCL-6, AID, IL-6, IL-21, T_{FH}, ELF-EMFs.

ICIA2018\Immunology of Chemical Victims and Environmental Pollution\Poster Discussion\7725
7725

Maternal exposure to A.lwoffii F78 prevents allergic asthma in offspring

Najme frozan nia ¹

1. Department of basic science School of Medicine University of mashhad Islamic azad university

Background: Asthma is a chronic inflammatory disease. epigenetic factors affecting the asthma phenotype have been lately investigated. Cowshed, farm animals and their products which are found in farming environment must have an important role in preventing asthma.

A.lwoffii f78 is a gram-negative cowshed bacterium which causes allergoprotective effect in prenatal model of transmaternal experimental asthma protection.

Methods: For this review we searched for asthma in titles and abstracts and for A.lwoffii F78 in whole papers at PubMed and Elsevier and Google scholar. We found 25 articles 12 of which were irrelevant to our subject or had a low impact.

Results: Maternal exposure to A.lwoffii F78 prevents allergic asthma in the mice with homozygous asthmatic mother. The exposure first causes a transient increase in proinflammatory cytokines and toll like receptor messenger RNA but later it shows lower amounts in cord. Its effect is dependent on maternal toll like receptor signaling.

Dendritic cells stimulated by A.lwoffii f78 cause to induce T_H1 polarising gene in naïve T cells. In the mice treated by A.lwoffii F78 T_H1 immune response shifting and decrease of T_H2 antibodies titration rather than T regulatory cells response were observed.

Exposure to A.lwoffii causes immune suppression and regulatory mechanisms by IL8, TNF, IL12 cytokines.

Conclusion: The articles proved that the strong influence of prenatal exposure to A.lwoffii F78 will prevent allergic asthma as it was claimed in hygiene hypothesis.

Keywords: Allergy protection, A.lwoffii F78, Hygiene hypothesis, T_H1 response Shifting, epigenetic

Allergy and immunotherapy\Poster Discussion\7962
7962

Evaluation of lymphocyte migration to inflamed skin using CD3, CCR4, and CCR10 markers following the Paederus induced Linear dermatitis in rat

Fereshteh Hosseini Valiki¹, Abolghasem Ajami², Mohammad Mahdi Mohammadi³

1. Department of Immunology, Faculty of Medicine, Kerman University of Medical Sciences, Kerman, Iran

2. Department of Immunology, Faculty of Medicine, Mazandaran University of Medical Sciences, Mazandaran, Iran

3. Department of Immunology, Faculty of Medicine, Kerman University of Medical Sciences, Kerman, Iran

Background: Paederus dermatitis is a blistering disorder that is caused by a small insect of the genus Paederus especially P.fuscipes. The disease spread around the world including Iran. It is more common in humid and hot areas, especially in the provinces of Gilan, Mazandaran and Khuzestan. This study aimed to investigate the reaction of the Adaptive immune system regarding the recruitment of CD3, CCR4 and CCR10 markers which are specifically expressed on the surface of T lymphocytes.

Methods: In this experimental study, 24 female rats were divided into two groups as the test and the negative control. In the test group, Paederus dermatitis was induced by making insects in contact with shaved rat skin. Biopsies were obtained 24, 72 and 120 hours after induction. In negative control group physiological saline was applied. Specimens were evaluated by Immunohistochemical staining method. Antibodies against CD3, CCR4 and CCR10 were used. distribution and staining intensity of CD3, CCR4 and CCR10 markers were estimated by the HSCORE index and findings were analyzed using the Kruskal-Wallis and Wilcoxon statistical tests.

Results: Based on the results of immunohistochemistry, expression of CD3, CCR4 and CCR10 in test group at 24, 72, 120 hours compared to the control group showed significant increase ($p = 0.0006$, $p = 0.001$ and $p < 0.0001$ Respectively). The peak of expression of all markers was in 72 hours after exposure. Hematoxylin-eosin staining also confirmed the fact that the majority of lymphocyte infiltration was occurred at 72 hours post exposure.

Conclusion: The expression of CD3, CCR4 and CCR10 on cells present in Paederus dermatitis lesions could indicate that T lymphocytes are recruited to the site of inflammation by chemokine-chemokine receptor interactions and hence provides evidence for the response by the adaptive immune system following a paederus dermatitis.

Keywords: Paederus dermatitis, Type IV hypersensitivity, Immunohistochemistry, CD3, CCR4, CCR10

Index

A		
A. A. Amirzargar	993	Abdolkarim Pajoumand 813
A. Aghaie	643	Abdolkarim Sheikhi 998
Abass Rezaei	615, 617	Abdollah Ardebili 782
Abazar Roustazadeh	162	Abdollah Jafarzadeh 524
Abbas Ahmadi	911	Abdollah Jafarzadeh 104, 377, 437, 644, 678, 913, 940, 942, 1016, 1022, 1023
Abbas Ahmadi Vasmehjani	640	Abdollah Jafarzadeh 998
Abbas Alamian	277	Abdollah Jafarzadeha 367, 761
Abbas Azadmehr	159, 483, 485, 487, 557, 616, 680, 682, 799, 860, 861	Abdollah Salimi 667
Abbas Bahador	370	Abdolrahim Nikzamir 186, 464
Abbas Fayezi	581	Abdolrahim Rezaei 381
Abbas Gerami	559	Abdol Rasoul Talei 623, 650
Abbas Ghaderi	28, 29, 78, 173, 297, 301, 623, 624, 633, 646, 650, 653, 654, 690	Abdol-Rasoul Talei 301
Abbas Hajifathali	537, 538	Abdolreza Esmaeilzadeh 65, 988
Abbas Hajizade	1054	AbdolReza Fazel 303, 668
Abbasian A	30, 613	Abdolreza Nabinejad 1009
Abbasi Ardeshtir	311, 915, 935	Abdolreza Nabinejad 1008, 1011, 1026
Abbas Jamshidian	555	Abdolreza Sotoodeh Jahromi 756, 892
Abbas Lotfi	990	Abdossaleh Zar 756
Abbas Mirshafiey	376, 402, 432, 433, 445, 720, 869	Abolfazl Yari 568, 569, 572
Abbas Mirshafieyd	367, 761	Abolghasem Ajami 531, 652, 824, 900, 1031, 1071
Abbas Rezaei	35, 183, 217, 232, 255, 259, 480, 497, 498, 589, 677, 957, 960	Abolghasem Allahyari 665
Abbas Sahebghadam Lotfi	553	AboTaleb Saremi 1039
Abbas Shirdel	665	Aboulfazl Rahmani 799
Abdolghader Taene	214, 814, 815, 885, 1048	Abtahi Froushani 337
Abdolkarim Hosseini	320	Abtahi Froushani Seyyed Meysam 915, 935
Abdolhussein Shakurnia	621, 626, 767, 770, 775	A.Dalimi 585
		A.Dehghan 823
		Adeleh Adelipour 259
		Adeleh Mahdizadeh 140

Adel Fekri	829, 975	Ahmad Zavaran Hosseini	240, 670
Adel Mohammadzadeh	24, 251	Ahmed Al-Humadi	717
Adibe Rahmani	556	Ahmed Sedegh	241
Afsaneh Aghaie	90	Ajami A	102
Afsaneh Jahangirian	244	ajediKhanian, M.	729
Afsaneh Jamali	103	Akbar Anaeigoudari	1013
Afsaneh Sharifian	805	Akbar Dorgalaleh	88, 745
Afshin Amari	835, 965	Akbar Ghorbani	60
Afshin Derakhshani	598, 599, 601	Akbar Safaei	28
AfshinDerakhshani	603	Akram Abouie Mehrizi	265, 546, 971
AfshinMoniri	331	AkramAbouie Mehrizi	969
Afshin Moradi	136, 412, 448, 465	Akram A. Mehrizi	266, 982
Afshin Namdar	262	Akram Davoodi	605
Afshin Nikkhoo	195, 930, 931, 932, 934	Akram Firouzi Amandi	167
Afshin Ostovar	23	Akram Jamali	38
Afshin Tahmasebi	377	Alaleh RezaLotfi	36
Aghdas Mirjalili	147	Ali Aghebati Maleki	53
Agheel Tabarmolahasan	833	Ali Ahmad Bayat	58, 174, 323
Aghil Bahramian	782	Aliakbar Amirzargar	260, 261, 967, 968
Ahad Mokhtarzadeh	469, 734	Ali Akbar Amirzargar	129, 262, 659
Ahad Zare	24, 25, 1039	Ali Akbar Delbandi	227, 356, 393, 507, 508, 509,
A. Heravian	802		739
ahereh Taheri	420	Aliakbar Haddad Mashadrizeh	318, 458, 460
Ahmad Asoodeh	145	Aliakbar Haddad-Mashadrizeh	714
Ahmad Bagheri	556	Aliakbar Movassaghpour	537
Ahmad Gharehbaghian	743	Ali Akbar Nasiri	550
Ahmad Hosseini	32	Ali Akbar Pourfathollah	90, 747, 1007
Ahmad Karkhah	447, 707, 994	Ali Akbar Pourfatollah	746
Ahmad Majd	25, 281, 284	Ali Akbar Rahim Rahimi	456
Ahmad Oryan	420	Ali Akbar Saboor Yaraghi	452
Ahmad Rahimi	38	Ali Akbar Velayati	398, 399
Ahmadreza Jamshidi	132	Ali Al-fahham	1037
Ahmad Reza Jamshidi	96, 128, 129	Ali Amiri	394
Ahmad-Reza Mahmoudi	41	Ali Anisian	249
Ahmad Rouhizadeh	1006	Ali Anissian	175, 296
Ahmad Tahamoli Roodsari	1024	Ali Arabkhazali	914
Ahmad Tajemiri	744	Ali Ariaifar	33, 295, 653



14th International Congress of Immunology and Allergys (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

Ali asadirad	66	Ali Nohekhan	487
Ali Asadirad	187, 517	Alipasha Meysamie	1021
Ali Asadi Rad	448	Ali Pourmohammad	396
Ali Asghar Tehrani	198	Ali Rahimi	531, 1031
Ali Dabbagh	171	Ali Ramezani	116
Ali Eshaghi	460	Ali Rastegari	195, 931, 932, 934
Ali farahi	815	Ali RezaFarahani	277
Ali Farahi	814	Alireza Abdollahi	298
Ali Farazmand	433	Alireza Ahmadi	996
Ali Fattahi Bafghi	147	Alireza Andalib	183, 615, 617, 677
Ali Fouladvand	570	AlirezaAndalib	305
Ali Ganji	113, 1015	alireza asef	914
Ali Ghajari	574	Alireza Azargoon	394
Ali Ghazavi	113, 1015	AlirezaBarati	1058
Ali Gorgin Karaji	584, 586, 587, 590, 591, 605	Alireza bari	665
Ali Grgin karaji	22	Alireza Biglari	53, 65
Ali Hashemi	271, 782, 783, 1025	Alireza Esmaeili	404
Ali Hatef Salmanian	264, 979, 981, 1054	AlirezaEsmaeili	127
Ali H. Salmanian	982	Alireza Ghanavatinejad	223, 224
Ali jalili	285, 923	Alireza Ghavami	292
Ali Jalili	468, 634, 635, 645, 924	Alireza Izadi	373
Alijan Tabarraei	162	Alireza Khabbazi	834
Ali Khamesipour	547, 1064	Alireza Khazaei	1067
Ali Khodadadi	155, 203, 619	Alireza Khoshdel	990
Ali Mahdavinezhad	349	Ali Reza Khosravi	110
Ali Masjedi	195, 930, 931, 932, 934	Alireza Mardomi	294
Ali Mohamadi	684	Alireza Mehdizadeh	731
Ali Mohammad Behroozikhah	277	Alireza Mirzaei	651
Ali Mohammadi	31, 43, 291, 479, 692, 694	Alireza Najafi	803, 816
Ali Mohammad Mohseni Majd1	1063	Alireza Pasdar	140
Alimohammad Varzi	87	Alireza Rafiee	395
Ali Mosayebian	19	Alireza Rafiei	328, 403, 848, 858
Ali mostafaie	359	Alireza Razavi	296, 303, 668, 847
Ali Mostafaie	444, 556, 744, 904	Ali Reza Razavi	452
Ali Mostafaie3, Kamran Mansouri	869	Alireza Refeie	655
Ali Movahedi	140	Alireza Rezaei Manesh	592
Ali Niazi	230	Alireza Rezaieymanesh	129, 130, 131, 410, 831, 832

Alireza Rezvani	32	Amir Ali Hamidieh	67, 238
Alireza salami	824	Amir Amiri Paryan	780, 843, 875, 1027
Alireza Shekohifar	693	Amir asghari	939
AlirezaShoae-Hassani	67	Amir Avan	140
Alireza Tahamtan	116	Amir Baghbanzadeh	469
Alireza Vakilian	940, 942	Amir Ghaemi	308
Ali Reza Zali	247	Amir-hassan Zarnani	94
Alireza Zamani	231, 306, 362, 493, 526, 527, 609, 751, 766, 857, 867, 970, 1069	Amir Hassan Zarnani	178, 217, 223, 224, 296, 475, 476, 497, 501
AlirezaZamani	977	Amir-Hassan Zarnani	41, 175, 498, 502
Ali R Khoshdel	762	Amirhossein Alimohammadian	327, 696
Ali Saeedi Boroujeni	203	Amir Hossein Faghfour	731
Ali Salmani	974	Amir Hossein Jahangiri	334
Ali Soleimanzadeh	950, 951, 952, 955, 956	Amir Mani	1042
Ali Vahid Dastjerdi	976	Amir Mizbani	146
Ali Zekri	671	Amir Moghadam Ahmadi	990
Allahyari Abualqasem	786	Amir Mohammad Alborzi	385
Alvanegh Mahmood Tavallaei	60	Amir Mohammad Malvandi	95
Amani Saeede	243, 1000	Amir Rahimi	713
Amein Hosseini	651	AmirReza AghebatKhayr	123
Amin Farhang	984	AmirReza Dehghanian	654
Amin Ghanivash	912	Amir Reza Heidari	252
Amin Isanejad	759	Amir Sadeghi	922
Amin kamrani	302, 948	Amir Salek Farrokhi	226, 405, 501, 502, 510, 660
Amin Khazaei	1067	Amir Shakouri	179, 883
Amin Lafzi	749, 972, 1068	Amirtaher Eftekharrosadat	660
Amin Mirzaeian	656, 1053	Amir Tukmechi	771
Amin Oveisi	307	A. Mohammadi 1	105
Amin Reza Nikpoor	789	Amouzegar A	102
Amin Roustami	1067	Amrolah Mostafazade	352, 517
Amin Safa	913	Amrollah Mostafazadeh	862
Amin Seyfaddiny	123	AmrollahMostafazadeh	241, 616
Amin Shafizad	611	Anahita Abbasi	540
Amin soltani	856	Anahita Najafalizadeh	232, 615, 617
Amin Soltani	610, 612	And Abdol mohamad Rostami	70
Amirabbas Salmani	18	Anders Malmström	138
AmirabbasSalmani	570	Andisheh Niakan	29

Andreas Kurz	117	Asadollah Baesi	626, 767, 770, 775
Anne Puel	76	Asadollah Mohamadi	665
Anoshirvan Kazemnejad	62	Asadollah Mohammadi	382, 389, 793, 812
Antimo D'Aniello	215	Asal Golchin	180
Antoine Haddad	800	Asgarian-Omran H	102
Antonie Zwiers	138	Asghar Abdoli	387, 807, 811
AnwarFathollahi	242	Asghar Afshari	148, 275
A. Radmanesh	802	Asghar Aghamohammadi	75, 341, 343, 344, 345
Aram Mohammadi	206, 208, 491	AsgharAghaMohammadi	74
Aram Mohammadi	204, 484	Asghari M	137
Arash Arashkia	68, 477, 547	A.Shahmohammadi	823
Arash Kalantari	593, 595, 804	A. Sharifzadeha	781
Arash Khojasteh	207	Ashkan Rasouli	806
Arash Pourgholaminejad	149	Ashraf vafa	1061
Arash Samoudi	572	Ashraf Vafa	754
Arbabi Mohsen	777	Asl iranifam Neda	1000
Arda Kiani	329	Asma khanzade	939
ArdaKiani	332	Asrin Safari	119
Ardalan Asadollah pour	33	Asyeh Javanian	581
Ardeshir abbasi	357, 888	Atabati Hadi	786
Ardeshir Hesampour	514	Ata Ghadiri	155, 621, 835, 838, 1006
Aref Hosseinian Amiri	403	Atefe Alirezaee	670
Arezki Chekaoui	169	Atefe Ghamar Talepoor	165
Arezo Dabiri	978	Atefeh Abedini	329, 332
Arezo Firoozeh	50	Atefeh Esmailnejad	560, 1014
Arezo Gohari Shabgah	390	Atefeh Najmaldini	1018
Arezo Gowhari Shabgah	896	Atefeh Sadeghi Shermeh	393
Arezo Naderi	168	Atefeh Yaftiyan	334
Arezou Abbasi	824	Atefe Mehrmanesh	590
ArezouDabiri	728	Atena Mozafari	979, 981
Arghavan Asghari	830	Atieh Ehsani	205
A.R.Khoshdel	985	Atiye Mellati	825
A.R. Khosravia	781	Atoosa Gitiara	448
Arron Mungela Foma	262	Atousa Homafar	631
Arsalan Jalili A	449	Atri Ghods	650
Aryaeian Naheed	409, 884	Ava Nasrolahi	355
Arya Sohrabi	198	Ayoob Arpanaei	1054

Ayssouda Hossein zade	368	Bahareh Rostami	933
Azad bahrapour	937	Bahare Niknam	246, 520, 521
Azad Bahrapour	539, 881	Bahare Shokoohian	234, 317
Azam Bagherizadeh	940, 942, 1016, 1022, 1023	Bahman Abdi	1014
Azam bolhassani	933	Bahman akbari	457
Azam Bolhassani	769, 989	Bahman Khalesi	277
Azam heidari	571	Bahman Khazaei	1067
Azam Jamshidian	523	Bahman Mansouri Miotlagh	870
AzamMoradi Zarmehri	980	Bahram Kazemi	237
Azam Rahimi	534	Bahram Nikkhoo	634, 788
Azam Rahimpour	535	Bahram Yavari	578
Azam Roohi	608	Bakhshi Joibari Fahimeh	150
Azar Baradarn	143	Bakhtiari Zahra	1000
A. Zavarani Hosseini	481	Baratali Mashkani	383, 792
Azin Aghamajidi	616	Bassam Badra	800
Azita Talebzadeh Bonakdar	262	Batol Zamani	408
Azizi F	102	Batoul Pourgheysari	620
B		BayazidGhaderi	634
Babak Abdolkarrimi	87	B. Baradaran	105
Babak Afkari	826	BehnamKalali	241
Babak Beikzadeh	276	Behnam Sanei	143
Babak Ghalehbaghi	191	Behnaz Esmaili	1021
Babak Khodadadi	593, 595, 804	Behnaz khani	960
Babak Negahdari	58	Behnoosh Tayebi	25
Babak Vazirianzade	155	Behnoush Baradaran	817
Bahador Bagheri	160, 563	Behrooz Ilkhanizadeh	24
Baharak Qa'edi	485, 487	Behrooz Vaziri	233, 425
Bahare Fazeli	812	Behrouz Gharesi fard	503, 959
Bahareh Abd Nikfarjam	431	Behrouz Gharesi Fard	220, 504, 961
Bahareh Fazeli	384	Behrouz Jalali Ghassam	460, 714
Bahareh Golkaran	549	Behrouz Nikbin	96
Bahareh Hasani Sabzevar	318, 714	Behzad baradaran	734
Bahareh Kardideh	406, 407	Behzad Baradaran	31, 37, 43, 130, 131, 410, 469,
Bahareh Keshavarzi	41		479, 674, 679, 681, 682, 684, 691, 692, 694, 831,
Bahareh Mohebi	187		832, 1047
Bahareh Pakpour	941	BehzadBaradaran	310
Bahareh Razeghi Haghighi	289	Behzad Baradaran1	689

Behzad Einollahi	533	Doroudchi M	79
BehzadEinollahi	974	Doroudchi, M	729
Behzad Ghareyazie	233	Dr. Abdolrahim Rezaee	795
Behzad Hatami	448	Dr. Ali Reza Moradi	386
Behzad mansoori	31	Dr.Ali Shirifi Zarchi	54
Behzad Mansoori 43, 291, 479, 682, 684, 692, 694, 1047		Dr. Houshang Rafatpanah	790
Behzad Nazel Khosroshahi	996	Dr Kambiz Pirnia	490
Beikzadeh L	808	Dr. Kambiz Pirnia	386
Bijan Bambai	230	Dr.Kaveh Kavousi	54
Bijan Khademi 78, 297, 646		Dr.Maryam Darayee	708
Bijan Pirnia 386, 490		Dr. Mehran Bakhshesh	810
Bijan Sedighi Moghaddam 622		Dr. Mohammad Mehdi Ranjbar	810
Bogoljub Ciric 70		Dr. Rasool Roshan	386
B. Sedighimoghadam 643		Dr. Reza Boostani	790
C		Dr. Reza mansouri	397
ChafiaTouil-Boukoffa 169		D. Shanehbandi	105
Charbonnier LM 26		E	
chardouli R 102		Ebadi Maryam	777
Chatila TA 26		Ebrahim Rahimi	640
Christopher A McCulloch 209		Ebrahim Eftekhari	895
Cynthia Aslan 697, 698		Ebrahim Mazloomi	24
Cyrus Jalili 407		Ebrahim Mirzadeghan	41
D		Ebrahim Sakhinia	826
Danial Ghasemi 997		Ebrahim Taghizadeh	168
Dariush Shanehbandi 291, 692, 694		Ebrahim Zabihi	616, 625
Davar Amani 49, 66, 182, 192, 202, 352, 353, 517, 661, 750, 753, 756, 922, 1056, 1057		EbtekarMasume	753
		E. Davarpanah	270
Davood Bashash 676		Eelco D Keuning	250
Davood Jafari 262, 422		Effat Farzanehnia	229
Davood Javanmard 813		Ehsan Arefian	706, 1007
Davood Rostamzadeh 32		Ehsan Farashahi Yazd	946, 1055
Davoud Ghaffarian 381, 786		Ehsan Khazaei	1067
Delavari Mahdi 777		Ehsan Nazemalhosseini Mojarad	472
Delirez Nowruz 311		Ehsan Nazemol hosseini Mojarad	189
Diba karimi 983		Ehsan Nazemolhosseini Mojarad	471
Dimitrios Balomenos 415		Ehsanollah Ghaznavi Rad	774
Djanette Hakem 169		Ehsan Sohrabi	738

Ehsan Soltaninejad	260, 967	Elham Moases Ghafari	430
Ehsan Zare Mehrjardi	740	Elham Mohammadi	544
Ehssan Shariat bahadori	144	Elham Mohit	325, 459
Eisa Salehi	99	ElhamNadimi	675
Elahe Aleebrahim Dehkordy	610, 612	Elham Rahimi	838
Elahe Alikhani	238	Elham Rajaei	835
Elahe Amani	119	Elham Razani	676
Elahe Amirinezhad Fard	444, 904	Elham Roshandel	537, 538
Elahe Davarpanah	425	Elham Sadat Hosseini Nasab	560, 1014
Elaheh Amani	596	Elham Safarzadeh	43
Elaheh azizi	471	Elham Safarzadeh1	1047
Elaheh Safari	235	Elham Shokati	438
Elaheh Talebi Ghane	751, 1069	Elham Zangane	240
Elahe Izadi	827	Elmira Gheytanchi	1020
Elahe Mansoori	895	Elmira shiuokhi	351, 738
Elahe Safari	438, 839	Elnaz Asghari	736
Elham Abbasi	871	Elnaz Farahani	249
Elham Abdollahi	1035	Emad kordbacheh	984
Elham Aghamohamadi	465, 922	Emami AH	30, 613
Elham Aghamohammadi	186, 464	Ensieh Lotfali	574
Elham Ahmadi	865	Ensieh Sadat Mirsharif	99, 1062
Elham Alipour fayez	393	Erfan babaei	285, 923
ElhamAmirchaghmaghi	741	Erfan Bahrami	780, 843, 875, 1027
Elham Atabati	830	Erfan Sadeghi	84, 840
Elham Bagbani	43	Erna Domsgen	62
Elham baghbani	734	Esfahanian F	102
Elham Baghbani	469, 691, 692, 694	Eskandarian Maryam	221, 222
Elham Bagherinia	586, 587, 590	Eskandar Kamali-Sarvestani	70
Elham Boustanipour	671	Eskandaryan M	721
Elham Gholami	268, 425	Esmaeil Allahmoradi	101, 293
Elham Harati	847	EsmaeilAsli	277
Elham Jafari	494	Esmaeil Ebrahimie	230
ElhamKashafi Jahromi	733	Esmaeil Jarrahi	461
Elham Khalaf Adeli	747	Esmaeil mortaz	914
Elham Mahabadi Ashtiyani	527	Esmaeil Mortaz	121, 122, 299, 329, 332, 336, 346, 347,
ElhamMahabadi-Ashtiyani	977		393, 398, 399, 516, 1056, 1057
Elham Mahlooji	20	EsmaeilMortaz	331

Esmail Saneei Moghaddam	745	Farhad Salari	22, 584, 590, 591, 605, 697
Esmail Mortaz	66	FarhadSalari	586, 587
Ezzati Mehmandust Olya Mohammad	777	Farhad Seif	191, 592, 640, 836
F		Farhad Shahsavari	394
Fabiano Oliveira	268	Fariba Asgharpour	872
Faeze Fakour.k	795	Fariba ghorbanzadeh	789
Faezeh Abbasi Rad	342, 392, 960	Fariba Hejazynia	1067
Faezeh Asghari	689	Fariba Keramat	107, 766, 778
Faezeh Maghsood	614	Fariba Mirjalili	740
Faeze Khodadadi	663, 664	Fariba Ramezanali	741
Fahime Aeine negin	151	Fariba Rashid	473
Fahimeh Arabpour	611	Fariba sohrabi	744
Fahimeh Dadkhah	732	Fariborz Bahrami	395
Fahimeh Khadem	392, 960	Fariborz Mokarian	232, 615, 617
Fahimeh Lavi	256, 519	Fariborz Zandieh	284
Fahimeh Mohammadi	1016, 1022, 1023	Farideh Talebi	139, 255, 257, 413, 414
Fahimeh Nematimonsour	1005	Faride Jowkar	300, 647
Fahimeh Ranjbar Kermani	798	Faride Sadeghkhani	706
Fahimeh Rostamabadi	1022, 1023	Farid Semsarha	588
Fanuel Songwe	282	Farid Soleimani Mohammadi	788
Farah Bokharai Salim	803, 816	Farimah Masoumi	257
Farahmand M	808	Farin Malekifard	859, 950, 951, 952, 955, 956
FarahnazBodaghkhan	292	FarnazFahimi	180
Farahnaz Ghahremanfard	636, 642	Farnaz Hajizadeh	211, 944
farahnaz motamedi	991	Farnaz Malekifard	423, 424
Farahzad Jabbari Azad	18, 570	Farnaz Zahedifard	145
Faramarz Sour	864	Farrin Babaei Balderlou	366
Fardin Ahmadkhani	372	Farrokh Ghahremaninejad	281
Fardin Fathi	667	Farshad Foroughi	260, 261, 967
Farhad Abolnezhadian	283	Farshid Fathi	84, 417, 840
Farhad Babaie	130, 131, 410, 831, 832	Farshid Mossallaie	997
Farhad Gharibdoost	82, 128	Farshid noorbakhsh	260
Farhad Jadidi Niaragh	195, 930, 931, 934	Farshid Noorbakhsh	98, 139, 255, 257, 343, 413,
Farhad Jadidi-Niaragh	167, 932		414
FarhadJadidi-Niaragh	44	Farshid Saadat	57
Farhad Rezaei	376	Farshid Yeganeh	208, 242, 249, 421
Farhad Riazi Rad	395	Farzad Nazari	1056, 1057

Farzad Yazdani	298	Fatemeh Heidarnejad	449
Farzaneh Chahardahcherik	385	Fatemeh Hemati	580
Farzaneh Dasturian	730	Fatemeh Hesam Pour	504
Farzaneh Farshdari	325, 459	Fatemeh Hosseini	685
Farzaneh Fesahat	946, 1055	Fatemeh Izadi Jahromi	654
Farzaneh Ghaderi Shabankareh	173	Fatemeh Kazemi Lomedasht	480, 566, 703
Farzaneh Ghorbani	212	Fatemeh khadir	420
Farzaneh Hosseini	1049, 1052	Fatemeh Khaki Khatibi	894
Farzaneh Peighambarzadeh	232	Fatemeh Maleki	1059
Farzaneh Rostami	458	Fatemeh Mami	249
Farzaneh Saffari	678	Fatemeh Mardani	135, 254, 256, 404
Farzaneh Tafvizi	926	Fatemeh Mirzaeyan	656
Farzaneh Tofighi Zavareh	432	Fatemeh Mohammadizadeh	943
Farzane Ordoni aval	680, 682	Fatemeh Motevalli	989
Farzin Javid	168	Fatemeh Nasri	504
Fateme akbari mehni	597	Fatemeh Noparast	559
Fateme Babaha	75	Fatemeh Pak	636, 642, 1039
Fatemeh Adjaminejad	116	Fatemeh Pourgholi	442
Fatemeh Ahmadi	756	Fatemeh Raeesi Nejad	494
Fatemeh Ahmadi Motamayel	349	Fatemeh Raeisi Nejad	1038
Fatemeh Atyabi	195, 931, 932, 934	Fatemeh Ranjbar	577
Fatemeh Ayoobi	104, 990	FatemehRanjbar	576
Fatemeh Boroun	573	Fatemeh Razavi	329
Fatemeh B. Rassouli	663, 664, 876, 877, 880, 889	Fatemeh Rezaei	225, 505, 964
Fatemeh Dadgar Pakdel	648	Fatemeh Rezayat	589, 957
Fatemeh Doroudian	422	Fatemeh Sadeghifar	416
Fatemeh Ezzatifar	328	Fatemeh Saffari	678
Fatemeh Faraji	178	Fatemeh Samiezadeh Mashadi	318
Fatemeh Ghaemimanesh	58, 174, 323	Fatemeh Sarlati	208
Fatemeh Goudarzi	777	Fatemeh Sefid	711, 712
Fatemeh Hadi	316, 704, 787	FatemehSefid	710
FatemehHadi	52	Fatemeh Shafiee	620
Fatemeh Hajari Taheri	477	Fatemeh Shahhosseini	178
FatemehHajari Taheri	68	Fatemeh shojaei jeshvaghani	264
Fatemeh Hajighasemi	710, 711, 712	Fatemeh Tabatabaie	1059
Fatemeh Hajmanochehri	799	FatemehTazari	567
Fatemeh Hasheminia	458	Fatemeh Vafashoar	827

Fatemeh Yari	234, 317	Forough Nejatollahi	63, 675
Fatemeh Yarian	454, 455	Forough Golsaz Shirazi	453, 608
FatemehYarian	187	Forough Golsaz-Shirazi	40, 44, 177
FatemehYazdian	61	Forouzan Yousefi	519
FatemeJalousian	1051	Forouz Keshani	485
FatemeManshori	1051	F. Rahmani	993
Fateme Mousavi	281	Frank A. Redegeld	299
Fateme pak	465	Frank Momburg	176
Fateme Rahmani	1019	Friborz Hovanloo	756
fateme ranjbar	606	F. Vafashoar	133
Fateme ranjbar	597	G	
Fateme Sadat Dashti	873	Garshasb Rigi	702, 725
Fateme Sadri Ardalani	314	Georges Snounou	982
Fateme Shaabanpour Aghamaleki	687, 688	Gert Folkerts	122
FatemeShaabanpour Aghamaleki	55, 657	Geshvad Nasiri Pirbazari	980
FatemeZare	324	Ghada Kchour	776, 800
Fatma Zohra Mazari	169	GhasemAshouri	1042
Fattah Sotoodehnejad Nematalahi	454	Ghasemi N	1046
Fattah Sotoudehnejad	1036	Ghasem Janabaei	293
Fazeli MS	30, 613	Ghasem Mosayebi	113, 1015
Fazel Shokri	40, 44, 116, 177, 224, 269, 314, 452, 453, 476, 608	Ghasem Solgi	36, 107, 349, 493, 778
Fereshteh Abdolmaleki	229	Ghasem Solgi1	1024
Fereshteh Alsahebhosoul	81, 184, 525, 732	Ghazaleh Bordbari	244
Fereshteh Alsahebhosoul	978	Ghazaleh Khalili Tanha	214, 1048
Fereshteh Bahmani	408	Ghazal Fatemi-Nasab	116
Fereshteh Hosseini Valiki	1071	Ghazanfari T	851, 852, 878, 879
Fereshteh Mehdipour	301, 623, 646, 650	Gholamalikardar	235
Fereshteh Pourabdolhossein	447	Gholamali Kardar	281
Fereshteh Rasa	895	Gholam Ali Kardar	20, 25, 282
Fereshte Salami	345	Gholamhossein Hasanshahi	326, 768
Ferial Taleghani	486	Gholamhossein Hassanshahi	495
FGhaffarifar	585	Gholam Hossien HasanShahi	123
F. Gorzin	993	Gholamreza Ahmadian	725
Firoozeh Khamoushi	591, 605	Gholamreza Ananisarab	1029
Firouz Ebrahimi	278, 1054	Gholamreza Anani Sarab	89, 91, 158, 693, 885, 887,
Firoz Ebrahimi	555	Gholamreza AnaniSarab	1048
			214

Gholamreza Asadikaram	373	Hadis Moradjodaki	110
Gholamreza Azizi	215, 341, 345	Hadi Zeidabadi	896
Gholamreza Habibi	148, 275	Hafezeh Davari	140
Gholamreza Hatam	426	Haideh Namdari	434, 863
Gholamreza Javadi	323, 473	HajarAbbasi	72
Gholamreza Nikbakht Brojeni	279	Hajar Safari	625
Gholamreza Nikbakht Brujeni	273	Hajar Shali1	360
Gholamreza Pourmand	259	Hajar Yaghoobi	550
Gilda Parsamanesh	352, 517	Hakim Azizi	550
Giti Farsi	762, 990	Haleh Sarrafnia	805
Golbahar Hosseini	983	Halimeh Hassanzadeh	416
Golnaz Babamiri	596	Hamed Bagheri	929
Golnaz Rashidi	155, 619	Hamed Bashiri	992
Golshan Sheikhsaria	947	Hamed Farhadi Kohan	766
Gordon A. Ferns	140	Hamed Fouladseresht	726, 727
H		Hamed mohammadi	131
Habbib Dastmalchi Saeed	907	Hamed Mohammadi	37, 130, 410, 831, 832
Habibollah Mirzaei	116	Hamid Abtahi	113
Habib Onsoni	674, 679	Hamid Ahanchian	18
Habib Soheyli	774	Hamid Aria	108, 371, 772, 773
Hadi Atabati	381	Hamid Asadzadeh Afgdaei	136
Hadi Esmaili Govarchin Galee	855	Hamid Asadzadeh Aghdaee	353
Hadi Ghazanfari	370	Hamid Asadzadeh Aghdaei	69, 187, 188, 190,
Hadi Hassannia	685		412, 448, 470, 473, 805
HadiHassannia	44	Hamid Asadzadeh Aghdayi	352, 517
Hadi Hossein Nataj	403	Hamid Badali	372
Hadi Hossein-Nataj	19, 46, 184	Hamid Daneshvar	151, 1060
Hadi Hossein Nattaj	309, 531, 662, 1031	Hamide danesh	597
Hadi Mirzaeei	550	Hamideh Rahimi	144, 900
Hadi Mohagheghian Yaghoubi	356	Hamide Kouhpeikar	877, 880
HadiMoheb Alian	580	Hamid Eshaghi	74
Hadinedoushan H	1046	Hamid Farahmand	561
Hadi Poormoghim	827	Hamid Farhangi	980
Hadiseh Farahani	422	Hamid M. Niknam	422, 547, 548
Hadiseh Rezazadeh	46	Hamid Mohamad Jafari	328
Hadis khamevar	375	Hamidreza Abdolsamadi	349
Hadi smaili Govarchin Ghalee	910	Hamid Reza A. Naghash	736

Hamid Reza Bidkhori	390, 416	Hedieh Rahmati	722, 1017
Hamidreza Hakkakifard	1001	Hediyeh Ghalikhani	491
Hamidreza Jamaati	329	Hediyeh Gholampour Sardashti	91
Hamidreza mohajerani	925	Hediyeh Rafighdoust	454, 455
Hamid Reza Mohammadimotlagh	406	Helia Azimi	801
Hamid-Reza Mohammadi-Motlagh	869	Helya Sadat Mortazavi	813
Hamid reza Mortazavi	22	Hematabadi M	102
Hamidreza Nouri	467	Hemati N	541
Hamid Reza Nouri	447, 625, 707, 709, 872, 994	Hengameh Akhavan	204
Hamid Reza Rashidinejad	913	Henrike Veninga	138
Hamid Reza Sadeghipour	847	HesamBabaei Khameneh	616
Hamidreza Safari	158	Hesam Karimi	545
Hamid Reza Taheri Nakhost	386	Hien Van Doan	1042
Hamid R. Golkar	408	Hiva Mohammadizad	114, 115
Hamid Salimi	385	Hiva Sharebiani	50
Hamid Shamsi	658	H. Jafarinejad	802
Hamzeh Rezaei	349	H. Jafarinezhad	643
Haniehheydarlou	219	H. Noormehr	481
Hanieh Nikkhah	866, 907, 908	Hobbenaghi Rahim	243
Hanie jafardokht bonjar	597	Hoda Kavosi	128
Hanie Mahaki	306, 362, 751, 1069	Hoda Rahmani	49, 271
Haniye Ghasiyari	182	Hodjatallah Rabbani	58, 174
Hasan Borji	580	Hodjattallah Rabbani	323
Hasan Namdar Ahmadabad	506	Homa Davoodi	627, 683
Hashemi, A	918	Homa Mohseni Kouchesfahani	534
Hashemi Seyyed Mahmoud	915	Homa Sadri	575
Hashem Marawne	738	Hoorieh Soleimanjahi	103, 545, 817
Hassan Abolhassani	345	Hosain AqaHosaini	59
Hassan Mahmoudi	766	Hosein Ghambarian	66
Hassan Malekinejad	859	Hosein Ghanbarian	338, 450
Hassan Nikoueinejad	533, 974	Hosein Honari	929
Hassan Rokni-Zadeh	53	Hosein Khan Ahmad	677
Hatam GholamReza	150	Hosein Zolfagharian	279
Hatami H	878, 879	Hosna salmani	595
Hedayatollah shirzad	612	Hossain Khoramdelazad	104
Hedayatollah Shirzad	185, 463, 856	Hossain Khorramdelazad	644, 940, 942, 1016, 1022,
HedayatollahShirzad	620		1023

Hossein Aazami	191	Hossein Vazini	996
Hossein Akbari	533, 974	Hossein Yousofi Darani	143
Hossein ali Ebrahimi	1018	Hossien Hafezi	417, 840
Hossein Ali Ghaffaripour	914	Hossien Mortazavi	322
Hossein Ali Khazaei	1067	Hossien Motedayyen	417, 840
Hossein Asgarian Omran	293, 309, 403, 531, 659, 662, 900, 1031	Hossien Rezvan	419
Hossein Asgarian-Omran	19, 46	Hossin Kamiabi	1060
Hossein Batooli	20	Houda Belguendouz	169
Hossein Dalili	62	Houri Edalat	60
Hossein Fallahi	52, 316, 704, 787	Housein Malekafzali Ardakani	322
Hossein Ghaedamini	925	Housein Malekafzali Ardekani	146
Hossein Ghanbarian	253, 629, 987	Houshang Rafat panah	1035
Hossein Golmoghdam	388	Houshang Rafatpanah	382, 383, 384, 389, 390, 664, 665, 792, 793, 812, 876, 877, 880, 889, 906
Hossein Golmoghaddam	901	HoushangRafatpanah	663
Hossein Goudarzi	782, 783	Houshang Rafat Panah	381
Hossein Hadinedoushan	873, 946, 1019, 1055	H. Poormoghim	133
HosseinHadinedoushan	324	Hussam Al-Humadi	125, 717
Hossein Hafezi	84	Hyeryon Lee	145
Hossein Hodjati	108, 371, 772, 773	I	
Hossein hosseini	1030	Ian M Adcock	121
Hosseini F	562	Ian M. Adcock	122, 331, 398, 399
Hossein Javdani	814	Ibrahim Toktaz	702
Hossein Javdani Javdani	815	Ilnaz Rahimmanesh	183, 677
Hossein Khanahmad	183, 480	Iman Farahani	113
Hossein Khan-Ahmad	35	Iman Naamipouran	965
Hossein Khorramdelazad	495	Ingrid Muller	322
Hossein Mahmoudvand	427, 428	Iradj Maleki	652
Hossein Modirrousta	148, 275	Iraj Rasooli	1043, 1050
Hossein Montazerghaem	730	Iraj Salehi	362, 493, 526
Hossein Mortazavi	146	IrajSalehi	977
Hossein Motedayyen	84, 154, 202, 217, 497, 498, 720	Izad M	102
Hossein Nikkhah	355	J	
Hosseinpur Z	851, 852, 878, 879	Jafar Ai	534
Hossein Rahavi	368	Jafar Amani	264, 982, 1054
Hossein Rahimi	665	Jafar kiani	335
		Jafar Kiani	1020



14th International Congress of Immunology and Allergy (ICIA2018)

26-28 April 2018, Tehran, I.R.Iran

Jafar Mahmoudian	175	JavadMohammadi	174
Jafar Majidi	109, 131, 831	Javad Shokri Shirvani	467, 921
Jafar Majidi. PhD	109	Javid Morad Abbassi	285
Jafar Salimian	354	JavidSadeghi	180
Jahangiri A.H	721	Jean Villard	536
Jahangiri Amir Hossein	221, 222	Jesus G Valenzuela	268
Jalal Abdolalizadeh	179, 883	J. G. Valenzuela	270
Jalal Khoshnoodi	40, 44, 177, 269, 453, 608	Johan Garssen	122, 331, 398, 399
Jalal shayegh	1030	Joke M. M. den Haan	138
JalehBarar	180	Joo Hwan No	145
Jaleh Zolghadr	503	K	
Jalil Adabi	119, 596	Kalantari-Khandani B	678
Jalil Kuhpayehzadeh	827	Kalantar Kurosh	150
Jalil Mehrzad	95	Kambiz Bagheri	92
Jalil Tavakkol Afshar	789	KambizYousofzadeh Eshkevari	330
Jalil Tavakol Afshari	18, 896	Kamran Mansouri	744
JalilTavakol Afshari	570	Kamran Mousavi Hosseini	798
Jamali M	737, 1040	Kamran Rostami	182, 465
Jamal Mohammadi Ayenehdeh	246, 520, 521	Karima Lahmar-Belguendouz	169
Jamileh Norouzi	354	Karimi Ali	1000
Jamshid Hadjati	48, 313	Karimi MH	541
JamshidHadjati	673	Karin Mink	241
Jamshid Hajati	1020	Kasra Esmaeilnia	148
Jamshidian Tehrani Narges	541	Katayoon Bidad	96
Jamshidi Fatemeh	409, 884	Katayuon Bahman Soufiani	1007
Jamshid Yazdani	796, 797	Kaveh Baghaei	69, 136, 187, 352, 412, 448, 466, 517, 622, 1002
Jasper J. Koning	138		
Javad Absari	533	Kaveh Baghai	66
JavadAbsari	974	Kaveh Khazaeel	962, 963
Javad Akhtari	848	KavehTari	645
Javad Alizadeh	771	Kayhan Azadmanesh	176
Javad Arasteh	1039	Kazem Ahmadi	111, 236, 439, 882
Javad Beheshtipour	558, 1004	Kazem AhmadiI	378
Javad Dadgar Pakdel	648	Kazem Mashayekhi	353
Javad Fathi	984	Kazem Nemati	727
Javad Hosseini	912	Kazem Parivar	370
Javad Masoumi	940	KeivanGohari Moghadam	99

Keivan Shams	996	Leila Janani	1028
Kenji Chamoto, Tasuku Honjo	42	Leila Karimi	291
Kenneth Lundstrom	547	Leila Khani	1028
keramati, M	918	Leila Mahdavinejad	362
Khadijeh Koushki	353	Leila Mobasheri	89
Khaje-Bishak Y	137	Leila Moradi	74
Khatereh Kafshdouzan	563	Leila Motamedzadeh	946, 1055
Khatereh Noormohammadi	366	Leila Nejatbakhsh Samimi	286
Khodakaram Jahanbin	965	Leila Nematollahi	325
Khodamorad Jamshidi	651	Leila Parvaneh L	74
Khoshfetrat Ali Baradar	243	Leila Shakerian	62, 413
Khosro Sardanian	306	Leili Aghebati Maleki	948
Khosrow Ebrahim	756	Leili Aghebati-Maleki	167, 947, 973
Kiarash Ghazvini	50, 819	Lennart Hammarstrom	62
Kimia Taghavi	332	Lisa G M van Baarsen	250
Kimia Zandeh	174	Lotfollah Behroo	594
Kobra Entezami	438	M	
Kobra Mehdinejadiani	271, 320, 1025	M.2, Karimi	79
Kokhaei P	30, 613	Maede Hamidi	621
Kosar Abouhamzeh	215	Maedeh Moradi	699
Kosar Salehitorabi	601	M.A.Fahmidekar	985
Kousar Smailnejad ganji	1010, 1012	Mahboobeh Nazari	175
Kousar Smailneja ganji	552, 899	Mahboobeh Razmkhah	297
Koushansineh sepehr	303, 668	Mahboobeh Zolala	1038
Kurosh kalantar	388	Mahboobe Razmkhah	633
Kurosh Kalantar	118, 157, 426, 997	Mahboubeh Nikookalam	874
L		Mahbubeh Zolala	494
Ladan Langroudi	34, 1018	Mahdaneh Roshani	782, 783
Laleh Babaeekhou	265, 546	Mahdavi R	737, 1040
Laleh Mavaddatiyan	909	Mahdi Abdollahi	158
Laleh Sharifi	376, 445	Mahdi Alahgholi Hajibehzad	362, 493, 526, 527,
Latif Gachkar	813		970, 977, 1069
Leila Bidar L	959	Mahdi Alahgholi-Hajibehzad	857
Leila Eini	518	Mahdi AlahgholiHajibehzad	751
Leila Farhadi	911, 912	Mahdi Asmar	785
Leila Hadadi	659	Mahdi Barati	160
Leila Hatami Giklou Jajan	233	Mahdi Behdani	178, 480

Mahdi Dehghanmanshadi	324	Mahmood Bozorgmehr	223, 224, 237, 518, 1020, 1028
Mahdieh Amiri	567	Mahmood Jeddi Tehrani	269, 314, 476, 608
Mahdieh Azizi	421	Mahmood Jeddi-Tehrani	40, 44, 177
Mahdieh khazae	135, 254	Mahmood Saffari	202, 720
Mahdieh Khazae	404	Mahmood Sheikfatollahi	990
Mahdieh Molanouri Shamsi	103	Mahmoud Hassani	68, 477
Mahdieh Naghavi Alhosseini	627, 683	Mahmoudi Mahdi	409, 884
Mahdieh Raeeszadeh	558, 1004	Mahmoud Jeddi Tehrani	453
Mahdi Fasihi-ramandi	236	Mahmoud Khaksary Mahabady	962, 963
Mahdi Fasihi Ramandi	553	Mahmoud Khayatani	730
Mahdi Fasihi-Ramandi	111, 378	Mahmoud Mahmoudi	254
Mahdi Ghasemi artiyani	764	Mahmoud Mahmoudi	127, 135, 252, 256, 404, 519,
Mahdi Ghasemiartiyani	763		1013
Mahdi Ghatrehsamani	612, 856	Mahmoud Mokbel	776
Mahdi Goudarzvand	215	Mahmoud Rafieian kopaei	610, 1053
Mahdi Habibi-Anbouhi	178	Mahmoud Vahidi	237
Mahdi Kholoujini	996	Mahnaz Ghaebi	499
Mahdi Mahmoudi	82, 96, 128, 129, 132, 402	Mahnaz Sourani	759
Mahdi Pouramir	625	Mahnoosh Foroughi	171
Mahdi Rohani	614	Mahsa Alem	161, 435, 905
Mahdi Samani	489	Mahsa Eshkevar Vakili	997
Mahdi Shabani	314	Mahsa Hajivalili	442
Mahdi Shekarabi	227	Mahsa Jafari	397
Mahdi Taghaddosi	828, 829, 995	Mahsa Javadian	697, 698, 699, 700
Mahdi Taghadosi	406, 407, 821, 822, 975	Mahsa Mahdian	349
Mahdi Tarhini	776, 800	Mahsa Mohammadzadeh	58
Mahdi Vojdani	129, 132	Mahsa najafzadeh	604
Mahdiye Saadi	447, 994	Mahsa Najafzadeh	601, 602
Mahiar Jeloudari Mamaghani	276	Mahsa Rahgoshay	768
Mahin Najafian	1006	Mahsa Rahimzadeh	611, 730, 895
Mahin Nikogoftar	238, 900	Mahsa Rezaeepoor	81, 525
Mahin Nikogoftar Zarif	1007	Mahsa Saeedi Niasar	805, 927
Mahla jafari	597	Mahsa Safari	868
Mahlajafari	606	Mahsa Sobhani	231
Mahla Jafari	576	Mahsa Taebe	367, 761
MahlaJafari	577	Mahsa Taghavi	1044
Mahmood Bakhtiyari	171	Mahsa Tanha	94

Mahshid Mehdizadeh	538	Manizheh Habibpour	88
Mahshoid Mohammadipour	234, 317	Mansoor Khakpoor	1058
Mahsima Shabani	343	Mansoureh Farhangnia	735
Mahtab Fattahi	1036	Mansoureh Farhangniya	1041
Mahtab Gholamzadeh	806	Mansour Khakpur	749, 972, 1068
Mahtab Maghsudlu	798	Mansour Salehi	732
Mahyar Malekzadeh	653, 654	Manzoomeh Shamsi Meymandi	1038
Mahyar Nourian	470	Maral Nazarifar2	588
Mahyat Jafari	264, 979, 981	Marco Maccarana	138
MaisamGhaedi	295	Marjan Mansourian	81, 525
Majgan Bagherzadeh	552, 899	Marjan Nassiri Asl	431
Majid Ahmadi	218, 219, 499, 500, 949, 958	Marjan Zamani	790
Majid Changi-Ashtiani	53	Markus Gerhard	241
Majid Faridgozar	533, 974	Martin Skerhut	241
Majid Ghayour-Mobarhan	140	Martje Erkelens	250
Majid Khoshmirsafa	13, 191, 286, 393, 592, 640,	Maryam Abedi	300
	836	Maryam Abolghazi	422
Majid Marjani	122	Maryam Abolhasani	175
MajidMarjani	331	Maryam Ahmadzadeh	325, 459
Majid Mirmohammadkhani	1039	Maryam Ajami	746
Majid Mirsadrai	580	Maryam Akhtari	132
Majid Momeni Moghaddam	416	Maryam Akrami Somea Bozorg	42
Majid Mossahebi Mohammadi	303, 668	Maryam Alinejad Dizaj	346, 347
Majid Naderi	88	Maryam Ameri Torzani	265
Majid Sirati Sabet	186, 464	Maryam Asadi	873
Majid Tebianian	124	Maryam Azarafza	652
Majid Validi	707	Maryam Azimi Mohamadabadi	529
Malek Hossein Asadi	628	Maryam.Azizpour.m	551
Malihe Aghasizade	140	Maryam Bahrami	847
Maliheh Moradzadeh	162, 289, 350, 638, 639, 733	Maryam Bayat	159, 436
Malika Terrahi	169	Maryam Behfar	67
Mana Momeni	581	Maryam Darayee	713
Mandana Sattari	202, 203, 204, 205, 206, 207, 208, 484,	Maryam Davarpanah	18
	486, 488, 489, 491	Maryam Diansaei	168
Manijeh Khanmohammadi	41	Maryam Ekhtelat	838
Mani Ramzi	536	Maryam Erfanmanesh	988
Manizhe Faghih	192, 922	Maryam Fakhimi	33, 633, 647

Maryam Fakhmi	300	Maryam Moslehi Baharanchi	427, 428
Maryam Farghadan	1017	Maryam Naseroleslami	313
Maryam Ghahremani Piraghaj	338, 450	Maryam Nemati	104, 437, 913
Maryam Ghane	455	Maryam Nezafat Firizi	506
Maryam Golar	269	Maryam Nikoonezhad	538
Maryam Habibzadeh	467	Maryam Nourizadeh	62, 187, 622
Maryam Heidari	69, 136, 412, 420, 466, 1002	Maryam Rastin	127, 135, 252, 254, 256, 404, 1013,
Maryam Hemmatzade	831		1035
Maryam Hemmat zadeh	37	Maryam Roozbehkia	402
Maryam Hemmatzadeh	131	Maryam Rostamtabar	709
Maryam Hosseini	844	Maryam Sahebari	127
Maryam Iranpoor	1038	Maryam sahlolbei	335
Maryam Izad	434, 529, 863, 1021	Maryam Salimi	449
Maryam Kafizadeh	768	Maryam Shabani	170, 446
Maryam Karami	626	Maryam Shahhoseini	741
Maryam Kargar	452	Maryam shakoori	1014
Maryam karimi	788	Maryam Sharifshoushtari	20, 25
Maryam keykhaee	803, 816	Maryam Sharif Shoushtari	284
Maryam Khabareh	363	Maryam Sohrabi	857
Maryam Kheirandish	244	Maryam Tajabadi Ebrahimi	941
Maryam khoshkhooi	570	Maryam Tavakoli	223, 224
Maryam Khoshkhooi	18	Maryam Vaezjalali	801
Maryam Khosravi	385	Maryam Valizadeh	746
Maryam Mahdiyar	640	MaryamYousefi	175
Maryam Mahjoub	592	Maryam Zare	961
Maryam Mehri	487	Maryan Skandaryan	334
Maryam Mhammad Ganji	305	Marzie Askari	493
Maryam Moazamigoodarzi	113	Marzieh Ebrahimi	36
Maryam Mobini	403, 608	Marzieh Eghtedar doost	1066
Maryam Moghaddam Matin	416	Marzieh Heidarzadeh Arani	20
Maryam Mohammad-Ganji	35	Marzieh Mansouri	536
Maryam Mohammadian	847	Marzieh Mohammadian Manesh	564
Maryam mollaee alamooti	991	Marzieh Naseri	1020
Maryam Moogooei	368	Marzieh Nejati Fard	485
Maryam Moossavi	598, 599	Marzieh Norouzian	611
Maryam Moradi	581, 992	Marzieh Rezaei	624
Maryam Moradi Chaleshtori	695	Marzie Motaghy	123, 326

Marzie Norouzian	646	Matin Aghalar	38
Marzie sadeghi	1030	Mazarzadeh	823
Marzie Zilae	283	Mazdak Ganjalikhani hakemi	525
Marziyeh Mohammadi Kordkhai	437	Mazdak Ganjalikhani-hakemi	81, 183, 480
Marziyeh Mohammadi-Kordkhai	104	Mazdak Ganjali khani-Hakemi	35
Marziyeh mohammadi kordkheyli	524	Mazdak Ganjalikhani Hakemi	232, 255, 305, 342, 344, 392, 978
Masha Hoseini	961		
Masomeh Tehrani	626, 770	Mazdak Ganjalikhani-Hakemi	75
Masomeh Asadifard	89	MazdakGhanjalikhan Hakemi	76
Masomeh Baghaei	195, 931, 932, 934	M.Badri	585
Masome Keykha	452	Mehdi Aleahmad	224
Masoud Etemadifar	81, 525, 732	Mehdi Aleahmad	223
Masoud Ghorbani	233	Mehdi Alikhani	461
Masoud haghshenas	452	Mehdi Barati	636, 642
Masoud Moghimi	515	Mehdi Behdani	477
Masoud Movahedi	284	Mehdi Dadashpour	167
Masoud Movassaghi	121	Mehdi Ebrahimiyan	374
Masoud Nafari	461	Mehdi Fouladi	180
Masoud Reza Seyfi Abad Shapouri	274	Mehdi Ghahary	111
Masoud Shahdroodian	980	Mehdi Golchin	544
Masoud Shirmohammadi	699, 700	Mehdi Kadivar	614
Masoud Soleimani	514	Mehdi Kalani	108, 371, 772, 773
MasoudSoleimani	34	Mehdi KazemporDizaji	331
Masouma Mowahedi	85	Mehdi Kazempour Dizaji	332
Masoume Asadi	722, 735, 1017, 1041	Mehdi Mahdavi	549, 637, 874, 999, 1005
Masoume Douraghi	269, 376	Mehdi Mahmoodi	762, 990
Masoume Ebtekar	65	Mehdi Malek Jamshidi	432, 433
Masoume Ebtekara	545	Mehdi Mirsaeidi	121
Masoume Mazandarani	627	Mehdi Montazer	28
Masoume Rajabibazl	535	Mehdi Raie	84, 417
Masoume Tavakoli Yarak	651	Mehdi Ramazanpour	329
Massood Hosseinzadeh	630	Mehdi ShafieeArdestani	1059
Massoud AH	26	Mehdi shahbazi	921
Massoud Saidijam	766	Mehdi Shahbazi	114, 115, 552, 705, 791, 899, 1010, 1012
Masume Ebtekar	661, 750		
Masumeh Ahanjan	779	Mehdi Shamsara	238
Masumeh Avari	794	Mehdi Shekarabi	191, 393, 592

Mehdi Shkarabi	836	Meraj Tabatabaei	41
Mehdi soltanzadeh Yamchi	921	Meysam Abtahi Foroushani	658, 779
Mehdi Talebi	537	Meysam Amidfar	728
Mehdi Tourani	467	Meysam Ganjibakhsh	722, 1017
Mehdi Yousefi	149, 211, 212, 218, 219, 401, 432, 433, 442, 499, 500, 528, 825, 844, 944, 947, 948, 949, 958, 973	M. Faranoosh	643
		M.H	79
		M. Hedayat	993
Mehdi Yusefi	109, 194	Mihan Pourabdollah Toutkaboni	332
Mehdi Yusefi. PhD	109	Milad Ameri	666, 920
Mehran Akbari	318	Milad Asadi	43, 291, 697
Mehran Falahpour	887	Milad Gholami	83
Mehran Gholamin	245	Milad Hemati	580
Mehrangiz Dezfoulia	927	Milad Moloudizargari	299
Mehran Habibi-Rezaei	891	MiladMoloudizargari	331
Mehran Moharreri	383, 792	Milad sabzevary-ghahfarokhi	185
Mehrdad Gholami	376	Mina Alimohammadi	655
Mehrdad Hajilooi	107, 349, 362, 778, 1024	Mina Biria	205
Mehrdad Hashemi	58, 174	Mina Ghorbanpour	260
Mehrdad Iranshahi	663, 664, 876, 877, 880, 889	Mina Mirzaee	999
Mehrdad Payande	744	Mina Motallebnejad	872
Mehrdad Payandeh	359	Mina Rahmanpour	107
Mehrdad Pooyanmehr	780, 843, 875, 1027	Mina Rezaei	51
mehrdad ravanshad	991	Minoo Adib	669
Mehri Barabadi	261, 967	Minoo Mohamadi	941
MehriBarabadi	260	Minoo Shaddel	658
Mehri Ghafourian	767, 775, 835, 965, 1003, 1006	Minoo Soltani	1051
Mehri Hajiaghay	431	Mir Davood Omrani	80
Mehri Shahir	336, 516	Mir Hadi Jazayeri	1028
Mehrnaz Izadpanah	735, 1017, 1041	Mir Saeed Yekaninejad	260, 262, 282, 967, 968
Mehrnoosh Doroudchi	108, 165, 371, 388, 441, 726, 727, 772, 773, 901, 961	Mitra Sabetghadam Moghadam	354
		Mitra Salehi	189
Mehrnoosh Maalhash	892	Mitra Shokri Majelan	1016
Mehrnoosh pashaei	636, 642	M. Javadian	105
Mehrnoosh Ramezani	41	M. J. Mahmoudi	993
Mehrnoosh Zagheh	970	M.Khojasteh	823
Mehrzaad Ghari	323	M.Mahmoodi	823, 985
Meinoosh Shabani	813	M. Nadji	133

Moazzeni Seyyed Mohammad	221, 222	Mohammad Ghasemi	155, 283
Moazzeni S. M	721	Mohammad Gholamian Hamadan	751, 1069
Moein Farshchian	416	Mohammad Hadi Sadeghian	663, 664, 877, 880,
Mohaddase Jarrahi	757, 758		980
MohaddesehGhezelcheh Khamsiyan	307	Mohammad Haghighi	493
Mohadesehsadat Hosseini	551	Mohammad Hasan Bemanian	559
Mohadeseh Toghyani Khorasgani	723	Mohammad Hasan Motedayen	279
Mohamad Ali asadi	388	Mohammad Hossein Baradaran	360
Mohamad Ali Assarehzadegan	827	Mohammad Hossein Dabbaghmanesh	654
Mohamad Ali Ibrahim zade	858	Mohammad Hossein Ghahremani	175
Mohamad Ali Ibrahim Zade	848	Mohammad Hossein Ghorbani	190
Mohamad Amin Pourhoseingholi	465	Mohammad Hossein karimi	959
Mohamad Reza Fazlollahi	25	Mohammad Hossein Karimi	540
Mohamad reza Haghshenas	300, 647	Mohammad Hossein Kazemi	269, 537
Mohamadreza Khosravi Farsani	759	Mohammad Hossein Modaresi	68
Mohammad Abbasi	609	Mohammad Hossein Nicknam	96, 129
Mohammad Abdi	912	Mohammad Hossein Niknam	968
Mohammad Ali Haghighati	568, 569, 572	Mohammad Hossein Nikoo	727
Mohammad Ali-Hasanzadeh	959	MohammadHossein Sanei	183
Mohammad Ali Judaki	269, 453	Mohammad Hossein Somi	660
Mohammad Ali Mansournia	798	Mohammadhossein Yazdi	637
Mohammad Ali Mazloomi	268	Mohammad Hossein Yazdi	874
Mohammad Alimoradi	744	Mohammadi MM	678
Mohammad Ali Sahraian	530	Mohammadi Rahim	243
Mohammad Ali Shokrgozar	178	Mohammadi, S.A	918
Mohammad Amin Behmanesh	962	Mohammad Jafari	356, 739, 803, 816, 996
Mohammad amin Boshagh	924	Mohammad Jahantigh	555
Mohammad Amin Boshagh	468	Mohammad Javad Abasszadeh Afshar	494
Mohammad Amin Pourhosseingholi	182	Mohammad Javad Ashraf	297
Mohammad Amir Alizade Tabrizi	483	Mohammad Javad Fattahi	654
Mohammad Ayubi Rad	762	Mohammad JavadGharavi	575
Mohammad Azizi	277	Mohammad Javad Motamedi	979, 981
Mohammad DavoodzadehGholami	235	Mohammad Javad Nasiri	396
Mohammad Ebrahim Ghaffari	766	Mohammad Javad Zibaeenezhad	165, 727
Mohammad Fayyad Kazan	776, 800	Mohammad Kamali	326
Mohammad Fereidouni	582, 583, 598, 599, 601, 602,	Mohammad Kamalinejad	847
	603, 604, 830	Mohammad Kazemi	732

Mohammad Kazemi Arababadi	147	Mohammad Reza Gharaati	461
Mohammad KazemiArababadi	1013	Mohammad Reza Hafezi Ahmadi	893
Mohammad Keramatipour	648	Mohammad Reza Haghshenas	297
Mohammad Khojasteh	865	Mohammadreza Hajizadeh	990
Mohammad Khosravi	573, 954, 962, 963	Mohammad Reza Hedayati Moghaddam	390
Mohammad Mahdi Eftekharian	213, 231, 362	Mohammadreza Jalali	449
Mohammad Mahdi Hayatbakhsh	896	Mohammad Reza Keramati	980
Mohammad Mahdi Mohammadi	262, 494, 656, 946,	Mohammad Reza khedri	875
	1055, 1071	Mohammad Reza Khedri	780, 843, 1027
Mohammad Mahdy Ranjbar	279	Mohammad Reza Mirzaei	628
Mohammad Mahmoudi	1044	Mohammad Reza Mirzaeic	367, 761
Mohammad Majidi	449	Mohammad Reza Mofateh	601
Mohammad Masoud Shoushtarian	833	Mohammad RezaMofatteh	603
Mohammad Mehdi ADIBZADEH SERESHGI		Mohammad Reza Mokhber Dezfouli	276
	378	Mohammad Reza Mozdianfard	20
Mohammad Mehdi Akbarin	665	Mohammad Reza Nejadmoghaddam	175
Mohammad Mehdi Amiri	40, 44, 177, 314, 608	MohammadrezaSharifi	305
Mohammad Mehdi Koshayar	665	Mohammad Reza Shokri	223
Mohammad Mehdi Mohammadi	913, 1018	Mohammad-Reza Shokri	224
Mohammad Mehdi Naghizadeh	1063	Mohammadreza Talebi Ardakani	484
Mohammad mehdi ranjbar	1025	MohammadrezaTohidkia	180
Mohammad Mehdi Soltan Dallal	376, 445	Mohammad Reza Tohidkia	456
Mohammad mirzakhani	921	Mohammad Reza VaezMahdavi	1066
Mohammad Moafi	419	Mohammad Reza Yazdani	441
Mohammad Mofateh	582	Mohammad reza Zali	69, 136, 187
Mohammad Nemati	573	Mohammad Reza Zali	72, 182, 186, 190, 353, 412,
Mohammad Nouri	456		448, 464, 465, 466, 470, 805, 922, 926, 927, 1002
Mohammad Rafiee	609, 901	Mohammad R Hajizadeh	762
Mohammadraman Molodi	911	Mohammad Rostami Nejad	182, 186, 192, 353,
Mohammad Rashno	155, 965		464, 465, 926
Mohammad Rastegar Moghadam	318	Mohammad Rostami-Nejad	922
Mohammad Reza Abbaszadegan	245	Mohammadsadegh Naghizadeh	604
Mohammadreza Alijani	152, 1052	Mohammad Sadegh Naghizadeh	582, 599, 887
Mohammad Reza Atashzar	640	Mohammad sadegh Soltani Zangbar	503
Mohammad Reza Bolouri	393	Mohammad Salehi	374
Mohammad-Reza Bolouri	41	Mohammad Shafiee	1034
Mohammad Reza Fazlollahi	282	Mohammad Shahbazzadeh	868

Mohammad Shahrooei	53, 581	Mohsen Rastegari Pouyani	869
Mohammad Shojaei	892	Mohsen Rokni	452, 980
Mohammad Taghikhani	660	Mohsen Roshani	49
Mohammad Taghi Rezayati	940, 1016, 1022, 1023	Mohsen Saeedi	1034
Mohammad-Taghi Rezayati	104, 644, 942	Mohsen Salesi	760
Mohammad Taher Afshounpour	103	Mohsen Shahmoradi	620
Mohammad Taheri	80, 213, 559	Mohsen Tadayyon	390
Mohammad Taher Tahoori	405	Mohsen Tehrani	46, 293, 652, 662, 824
Mohammad Tajfard	140	Mohtaram Nasrollahi	785
Mohammad Varahram	122, 398, 399	Mojdeh Ghias Tabari	552, 899, 1012
Mohammad Varhram	331	Mojde Kazemi	913
Mohammad Vodjgani	82, 128, 648	Mojgan Astaneh	309
Mohammad Vojgani	1066	Mojgan Bagherzade	791
Mohammad Yousef Alikhani	609, 766	Mojgan Bagherzadeh	114, 115, 921
Mohammad Zamanian	762	Mojgan bandehpour	455
mohammad zare mehrjerdi	564	Mojgan Bandehpour	237, 271, 695, 1025
Mohammad Ziaee	87	Mojgan Ghaedi	368
Mohammad Mahdi Mohhamadi	1038	Mojgan Mohammadi	18, 570, 896
Mohammad Hosein Karimi	542	Mojgan Mortazavi	259
Mohammad Sadegh Falahatkar	799	Mojgan Naghitorabi	838
Mohsen Abolhassani	68, 233, 477	Mojgan Sanjari	494
Mohsen Akhiani	824	Mojgun Mokhtari	184
Mohsen Arabpour	623	Mojtaba Bagheri	886
Mohsen Badalzadeh	25	Mojtaba Farahbakhsh	806
Mohsen Ebrahimi	1015	Mojtaba Mirzaei	278, 555
Mohsen Farhang	638, 639	Mojtaba Nofeli	929
Mohsenfarhang zargar	289	Mojtaba Salamatian	854, 898
Mohsen Hamidpour	743	Mojtaba Shahnazi	860, 861
Mohseni Mahboobeh	222	Mojtaba Shohan	463
Mohsen Jafarian dehkordi	976	Momeni Mohammad	966
Mohsen Khalili	902	Mona Abbasi kalabe	460
Mohsen Masjedi	156, 184, 720	Mona Abbasi kalabeh	458
Mohsen Molaei	605	Mona Amani	839
Mohsen Nafar	968	Mona Hemmati Chegeni	87
Mohsen Najimi	555	Mona Moshiri	376, 445
Mohsen Naseri	158, 449	Mona Sadeghalvad	406
		Mona Salahshoor dahr	454

Mona Salahshoor Dahr	455	Mostafa Jarahian	176
Mona Salimi	145	Mostafa Manian	140
Monireh Golpour	655	Mostafa Moin	20, 25, 284
Monireh Jahantigh	71	Motahareh Bahadori	40
Monireh Kazemimanesh	176	Motahareh Ehsani	552, 705, 791, 899
Monireh Mohsenzadegan	191	Motahhreh Ghasemi	700
Morad Rahimi	556	Mousa Mohammadnia afrouzi	791
Mortaza Ghasemi	526	Mousa Mohammadnia Afrouzi	921, 1010
Mortaz , Esmaeil	1000	Mousa Mohammadnia-Afrouzi	114, 115, 552, 899
Mortaz Esmaeil	243	MousaMohammadnia Afrouzi	1012
Morteza Abkar	199, 200, 553	Mousa Mohammadnia afrouzi	705
Morteza Bagheri	731	Mousa Tavassoli	423, 424
Morteza Djamali	20	Mozafar Mohammadi	63
Morteza Fallahpour	286	Mozhdeh Namvarpour	124
Morteza Hosseini	562, 1049	M.R.Hajizade	985
MortezaHosseini	1052	M.Rohban	985
Morteza Hosseinzadeh	967, 968	M.R.Shafipoor	985
MortezaHosseinzadeh	260	Mrs.Zahra Sasanipour1	708
Morteza Jafarinia	902	M.S.Fazaeli	985
Morteza Mahmoudi	175	M. Soltani	105
Morteza Oladnabi	680, 682	M. Soltania	781
Morteza Sagharjoghi Farahani	595, 804	M.Vakili	823
MortezaSagharjoghi Farahani	593	N	
Morteza Salimian	743	Nadereh Naderi	611, 730, 895
Morteza Samadi	85, 127, 301	Nadereh Tae	427, 428
Morteza Shamsizadeh	745	Nader Tajik	178, 217, 246, 259, 497, 498, 507, 508, 509, 520, 521
Mosayeb Rostamian	422, 548		
Moslem Ahmadi	314, 959	Nadia Manavi	714
Moslem Kulivandi	986	Naeges Sarlogzaei	1067
Mostafa Aalipour	568, 569, 572	Nafise Esmael	677
Mostafa Ejtehadifar	113	Nafiseh Esmaeil	232, 342, 392, 615, 617, 856, 960
Mostafa Haji-Fatahaliha	844	Nafiseh Tabasi	135, 254, 256, 404
Mostafa Haji molahoseini	171	NafisehTabasi	127
Mostafa Haji Mollahoseini	803, 816	Nafiseh Zand	1065
Mostafa Haji Molla Hoseini	21, 242, 421	Naghi Jabarivasal	306
Mostafa Haji MollaHoseini	249	Naghizadeh MM	754, 851, 878, 1061
Mostafa Hosseini	453	Naghmeh Sattarahmady	199, 200

Naghmeh Shirafkan	684	Nasiri Z	852
NaghmehShirafkan	310	Nasrin Ghasemi	1019
Naghmeh Zamani	926	Nasrin Hosseinahli	694
Nahid aboutaleb	864	Nasrin Iranshahi	822, 828, 829, 975, 995
Nahideh Afzale Ahangaran	71, 903, 907	Nasrin Iranshai	821
Nahid Ekandari	417	Nasrin Masoudzadeh	146
Nahid Eskandari	154, 156, 340, 525, 677, 732, 978, 1036	Nasrin Moazzen	18
NahidEskandari	140, 641	Nasrin Nouroollahzadeh	779
NahidGhaedAmini	140	Nasrin Saharkhize	1003
NahidJivad, HoseinTeymoori	330	Nasrin Zare	641
Nahid Rezaei	255	Nasrollah Erfani	297, 300, 647
Naiyer Hosseinahli	692, 694	NasrollahMahboobiSoofiani	1042
Najaf Allahyari Fard	230	Nasser Gholijani	118
Najibeh Shekari	689, 697, 698, 699, 700	Nassim Kheshtchin	48, 673
Najme frozan nia	1070	Nassim rahmani	230
Najmefrozannia	752	Nastaran Mohseni	566, 703
NajmeShahinfard	610	Nastaran Sadat Savar	547
Nakisa Zarrabi Ahrabi	448	Navid Dashti	128
Namdar Ahmadabad Hasan	150	Navid D. D jadid	982
Narges Amirjamshidi	616	Navid D. Djadid	266
Narges Arkian Boroujeni	981	Navid Dinparast Djadid	971
NargesArkian Boroujeni	979	NavidDinparast Djadid	969
Narges Roomandeh	1039	Navideh Haghnava	689
Narges Rostami	195, 930, 931, 932, 934	Navideh Mirakhani	951
Narges Valizadeh	384, 390, 819	Navid Mijani	849
Nariman Mosaffa	41, 49, 188, 271, 720	Navid Shomali	698
Nariman Mossafa	1025	NavidShomali	310, 684
Narjes Hasan Fakhrabadi	278, 555	Nayere Askari	1066
Narjes Soleimanifar	96	Nazanin Amiri	190
Naseh Sigari	923	Nazanin Atieh Kazemi sefat	319
Naser Mobarra	1034	Nazanin Fathi	155, 619
NaserSamadi	180	Nazanin Gholampour Farogi	460
NaserShagerdi Esmali	631	Nazanin Gholampour Faroji	318, 458, 714
Nasimmohammad kheshtchin	313	NazaninJafarpour	674
Nasim Rahmani Kukia	888	Nazanin mojtavavi	48
Nasim Samarghandi	669	Nazanin Mojtavavi	296, 827
Nasim Samarghmarghandi	1059	Neda Ahmadi afshar	423, 424

Neda Dezfuli	399	Nima Rezaei	75, 343, 344, 402, 1044
Neda Heidari	72, 414	Nima Rostami Koshki	374
Neda Mahavar	582, 603, 604, 830	Nima Taefehshokr	168, 307
Neda Mohsenpour	53	Niusha Amiri	489
Neda Mokhberian	629	N. Mojtavavi	133
NedaMousaviNiri	313	Nodehi M	102
Neda Sadat Gohari	1017	Nooshin Ahmadzadeh	132
Neda Shajari	681	Nooshin Askarani	1008, 1009, 1011, 1026
Negara Azarpira	630	Norouz Delirezsh	161, 435
Negar Behnamfar	165	Norouz DelireZh	849
Negar seyed	268	NosratollahZarghami	313
Negar Seyed	425	Nour Amirmozafari	785
Negar Vanaki	132	Nourizadeh M	30, 613
Negar Varae	515	Nouroz Delerizh	686
Negin Maleki	327, 696	Noushin Lotfi	417, 840
negin mohammadpoor	910	NoushinLotfi	84
Negin Mohammadpoor	870	Nouzar Nakhaee	373
Negin Nokhandani	683	Nowruz Delirezsh	24, 197, 478, 557, 859, 956
Nesa Rashidi	356	N. Rezaei	993
Neyshabouri	862	N. Seyed	270
Nezafat Ferizi Maryam	150	N. Shekari	105
N. Ghanizadeh	643	O	
Nika Khoshnevis	588	ohan Garssen	121
Nikoo H	79, 729	Omid Emadian	652
Nikoo Hossein-khan nazer	207	Omid Kooshkaki	1029
Nikoo Hossein Khan Nazer	489	Omid ramezani	1030
Nilofar Keyhanmehr	417	Omidreza Salehi	760
Niloofer Alvandpour	1024	Ostadrahimi A	137
Niloofer Basrehizad	588	owsar Smaeilnejad	791
Niloofer Hassannejad	370	P	
Niloofer Sadat Ebrahimi	472	Padide Ebadi	542
Niloofer Taghipour	188	Padideh Ebadi	540
Niloufar Keyhanmehr	154	Pak F	30, 613
Niloufar Rezvani	546	Pantea Mohammadi	556
Niloufar Sadat Miri	483, 485, 487, 557, 799, 860,	Parastoo Shakerin	622
	861	Pardis Bayat	427, 428
NimaParvaneh	74	Pardis Sadeghipour	322

Paria Jorbozohdar	286	Pooyanmehr Mehrdad	966
Parinaz mahdavi	468	Pordel, S	79, 729
Parisa Amir Kalvanagh	163, 545	Pouria Karimi	568, 569, 572
Parisa Esmailie	992	Pouriya Dadebighlu	168
Parisa Khodabandeh Shahraki	630	Poursani Shiva	409, 884
Parisa Nourbakhsh	592, 1015	Pourya Tahmasebi	618, 903
Parisa Vahidi	95	Pouya Lahoori	567
Parisa Zafari	821, 822, 828, 829, 975, 995	Pouyan houshmandpour	1001
Parissa Farnia	331	Pro.AR.Khosravi	375
Parizad Bavandpour-Baghshahi	115	P. Safari	133
Parsova Tavasolian	343	R	
Parvaneh Afsharian	741	Radjabian T	851, 852, 878, 879
Parvaneh Farzaneh	722, 735, 1017, 1032, 1041	Rafal Al-Saigh	717
Parvaneh Jafari	551	Rafal J. Al-Saigh	125
Parvin Ayremlou	731	Rafatpanah Houshang	786
Parvin Mansouri	1021	Raheleh Shokohi Shoormasti	25
Parvin Nikaein	653	Rahele Miri	390
Parviz Farzamifar	700	Rahim Hobbenaghi	197, 478, 859
Parviz Kokhaei	160, 465, 545, 563, 636, 642, 1039	Rahman Emamzadeh	715
Parviz Zoulghadri	799	Rahmani Kukia Nasim	915, 935
Pascale Kropf	322	RahmaniKukia Nasim	311
Payahoo L	137	Rahman Shokri	415
Payam Saadat	682	Ramazan Ali Sharifian	659
Payam Tabarsi	121, 122, 331, 393, 398, 399	Ramazan Rezaei	82
Payman Tavakoli	658	Ramezan-Ali Sharifian	44
Pedram Ahmadpoor	260, 262, 967, 968	Ramin Farajpour	809
Pedram Azimzadeh	353	Ramin Ghahremanzadeh	175, 475
Pedram Nazari	385	Ramin Lotfi	584, 591
Pejman Allahvaisy	596	RaminPourakbari	973
Pejman Allahveisey	119	Ramin Shekariz	46, 293, 309
Peter Krustrup	756	RaminShekariz	662
Peyman Bakhshaei	269	Ramin Yaghobi	540
Phipatanakul W	26	Ramona Nitui	241
P Jorbozedar	133	Ramyar Azizi	958
P. Kokhaei	643	Rana Ezzeddini	405, 660
Poorya Foroutan	924	RaphaellaSemper	241
Poorya foroutan pajoohian	468	Rashin Mohseni	67

Rasool Baharloo	286	Reza Gheitasi	893
Rasool Rahimi Junqani	152	Reza Hadavi	58, 174, 323
Rasoul Baharlou	640	Reza Hajiaghadee	159, 860, 861, 862
Rasoul Godini	52, 316, 704, 787	Reza Hajiasghar Sharbaf	662
Rasoul Jafari	143	Reza Hosseini Ghatar	608
Rasoul Nasiri kalmarzi	285, 923	Reza Hosseini-Ghatar	40, 44
RasoulSalehi	140	Reza Jafari	259, 806
Rasoul Zarrin	731	Reza Malk poor	494
Rassoul Dinarvand	175	Reza Mansouri	20, 61, 124, 368, 397
Rayhaneh Ahangar Parvin	437	Reza Marvinam	324
Rayhaneh Ahangar-Parvin	104	RezaMirfakhraie	83
Razie Alibeigi	312	Reza Nadafi	138, 250
Razieh Amini	867	Reza Nedaeinia	140
Razieh Dawodi	379	Reza Norian	557
Razieh Dilmaghanian	99	Reza Nosratabadi	252, 1013
Razieh Farazmand	458	Reza Pasandideh	274
Razieh Nejadi	486	Reza Ramazani Khoorshiddoost	588
Razieh Rezaeianpour	591	Reza Rezaie	983
Razie Mahmoodian	743	Reza Shahbazi	493
Raziyeh Abdolhamidi	1043	Reza Valadan	652
Raziyeh Karami	612	Reza yazdani	74
RaziyehKarami	610	Reza Yazdani	75, 344, 345
Reihaneh Pirjani	83	Rezvan Mohammadi Nezhad	564, 953
Reina E Mebius	250	Rezvan MohammadiNezhad	561
Reina E. Mebius	138	Rezvan Pilpa	50
Reyhaneh Babaei Khorzoughi	170, 446	R.Mansori	823
Reyhaneh Ghorbanpour	273	Robabeh Taheripanah	83
Reyhaneh Hoseinpoor	535	Robabe Latifi	860, 861
Reyhane Mirzaee	1005	Roghayeh babaei	816
Reza Aflatoonian	259	Roghayeh Pirzadeh	679
RezaAflatoonian	741	Rogier M. Reijmers	138, 250
Reza Ahangari Cohan	933	Rokhsareh Sadeghi	209, 486
Reza Alizadeh Navaei	824, 900	Rooholla Abdollahpour	757, 758
Reza Boostani	381, 382, 384, 389, 793, 812, 906	Roonak azizbeigi	937
Reza Falak	8, 13, 41, 235, 259, 286, 836	Roonak Azizbeigi	539, 881
Reza Farid Hosseini	18, 381, 383, 570, 792, 906	Rostam Golmohammadi	970
Reza Gharakhanlou	103	Rostami Rad M	737, 1040

Rouzbeh Sanaei	343	Saeed Malek Hosseini	388
Roya Bishe Kolaei	757, 758	Saeed Mirzaee	942
Roya Dolatkhah	355	Saeed Namaki	72, 136, 412
Roya Ghods	41, 175, 518, 1020	Saeed Nazeri	971
Roya khanahmadi	939	Saeed noorolyai	469, 734
Roya Khosravi Eghbal	269	Saeed Oraee Yazdani	247
Roya Najafi Vosough	970	Saeedreza Vazirian	292
RoyaShamsi	658	Saeed Talebi	319
Roya sherkat	340	Saeed Zeinali Heris	883
Roya Sherkat	76, 342, 392, 419	Saeid Abediankenari	13, 19, 294, 796, 797
Roya Taleban	419	Saeideh Erfanian	892
Roya Yaraee	449	Saeideh Milani	58, 174, 323
Rozhin Jamali	865	Saeideh Tabesh	282
Rozita Abolhasan	194	Saeid Pishgoee	896
Rozita Doosti	530	Saeid Taghiloo	101, 293, 824, 900
R. Safaralizadeh	270	Safar farajnia	457
R. Yarmohamadi	802	Safa Tahmasebi	988
S		Safdar Mehdizadeh Shermeh	833
saam torkan	976	Safoora Nikzad	231
Saber Rokhafrooz	1003, 1006	Safoora pordel	287, 288
Saboor-Yaraghi AA	102	Sahand Fattahi	307
Sadegh Farhadi	665	Sahar Bayat	83
Sadegh Fattahi	616	Sahar etemadi	910
Sadegh Feizollahzadeh	183	Sahar Etemadi	870
Sadegh Feyzollahzadeh	589, 957	Sahar Ghaffari	563
SadraAnsaripour	610	Sahar Karimzadeh	726
Saeed Alamian	199, 200, 553	Sahar Khorasani	135, 254, 256, 404
Saeed Amel Jamehdar	819	Sahar Mashayekhi	151
Saeed Azandeh	515	Sahar Mirzaee	874
Saeed Babaie	113	Sahar Mohammadi	774
Saeede Ghasemi	834	Sahar Mosalla	860, 861
Saeedeh Erfanin	289	Sahar Raoofi Mohseni	453
Saeedeh Mehraban Moghadam	384	Sahar Salimi	98
Saeedeh Torabi Goudarzi	109	Sahar Shariati	301, 650
Saeede Sadat Mirbageri	1001	Sahar Tahaghoghi Hajghorbani	848, 858
Saeede Shabanitabar	533	Sahel Heydari	356, 508
Saeed Farhadzadeh	168	Sahel Khazae	727

Saied Abediankenari	659	Samira Torki	856
Saied Bokaie	445	Samira Zand	779
Saiedeh Erfanian	162, 350, 638, 639, 733	Sanam Dolati	194, 211, 212, 401, 528, 825, 944
Sajad Dehnavi	18, 570	Sanaz Ahmadi Gezeldasht	390
Sajad Rashidi	426	Sanaz Baghban Rahimi	308
SajediKhanian	79	SanazDehghani	259
Sajjad Ahmadi Renani	439, 882	Sanaz Hesami	530
Sajjad Esmaeili	887	Sanaz Jourghasemi	107, 778
Sakineh Pirahmadi	969	SanazKeshavarz shahbaz	967
SakoMirzaie	634	SanazKeshavarzshahbaz	260
Saleh Ramazani Khorshiddoost	588	Sanaz Keshavarz Shahbaz	261
Saloomeh Fouladi	156	Sanaz Rasolzadeh	844
Samaeh Khorami	438	Sanaz Rasolzadeh Nori	149
Samaneh Abdolmohammadi Vahid	949	Sanaz Sheikhzadeh	197, 478
Samaneh Arab	48	Sanaz Zarasvand Nia	274
Samaneh Beigi	187	Sara Abdolmaleki	269
Samaneh Delavari	203, 621, 835	Sara Aghakhani	730
Samaneh Gholami	833	Sara Ghafarpour	647
Samaneh H. Shabani	266, 982	Sara Ghaffarpour	99
Samaneh khorami	839	Sara Golbahari	779, 853, 897
Samaneh Mahmood Aghdam	1041	Sara Hajimiri	50
Samaneh Mahmoud Aghdam	735	S.A Rahim Rezaee	50
Samaneh Maleknia	54	Sara Iranparast	767, 770, 775, 1003
Samaneh Salehi pour	767, 775	Sara Mohamadi Saani	883
Samaneh Tokhanbigli	352, 448	Sara Mohamadnezhad	785
Samane Mohamadzade	677	Sara Parsa	667
Saman Hosseinkhani	95	Sara Parsapour	97
Saman Nikeghbalian	624	Sara Rasouli	794
Saman Seyedabadi	887	Sara Shirahmadi	911
SamanSoleimanpour	50	Sara Soudi	8, 103, 240, 253, 338, 450, 513, 514, 670, 695, 817, 987
Samira Asgharzade	1053	Sara Yaghotian	581
Samira Derakhshan	298	Sarira Shahnavaz	208, 484, 491
Samira Ghasemi	1034	Sarira Shah Navaz	204
Samira Ghorbani	139, 255, 257, 413, 414	S.A.R Rezaee	819
Samira Mirbaha	715	Sayed Ali Hosseini Siyar	857
Samiramis Qavam	893	Sayed Hussain Mosawi	1059
Samira Mohammadi Yeganeh	320, 695		

Sayed Mohammad-Hassan Emami Najaffi Dehkordi		Sayed Alireza Esmaeili	254, 256
	184	Sayed-Alireza Esmaeili	135
Sayyed Hamid Zarkesh Esfahani	59, 715	Sayed Alireza Mesbah Namin	530, 1033
Sayyed Mohamad Hossein Ghaderian	83	Sayed Alireza Razavi	282
Sayyed Vahab Azizi	104	Sayed Amir Hossein Jalali	561, 564, 953, 1042
S. Bayanolhagh	802	Sayed Amir Jalali	312
Sedigheh Amini Kafi Abad	798	Sayed Amir Jalali	45, 271, 1025
Sedigheh Arbabian	471	Sayed Davood Hosseini	551
Sedigheh Nabian	559	Sayedeh Forough Sadeghian-Kiadehi	101
Sedigheh Pouya	69, 136, 412, 466, 1002	Sayedeh Mahboobeh Mosavi Mehr	1033
Sedigheh Rahrotaban	298	Sayedeh Mehrnaz Mostatabzadeh	850
Sedigheh Younessi Amirabad	92	Sayedeh Naemeh javid	627
Sedigheh Zakeri	265, 266, 546, 969, 971, 982	Sayedeh Naemeh Javid	683
Sepideh Ashouri Movassagh	1017	Sayed Ehsan Asadi	38
Sepideh Haghdooost	421	Sayedeh Tahereh Mohaddes	665
Sepideh Khodaverdi	227, 508, 739	Sayedeh Zohreh Jalali	57
Sepideh Maralbashi	697, 698	Seyede Rahele Tabatabaei Jafari	1024
Sepideh Mohseni	235	Seyede Zeinab Hosseini	788
Sepideh Moradkhani	592, 1015	Sayed Fazlollah Mousavi	233
Sepideh Sadat Hosseini	416	Sayed Hamid Aghajanzadeh	833
Sepideh Shahbazi	989	Sayed Hamidreza Monavari	813
Sepideh Soukhtezari	368	Sayed Hamidreza Mortazavi	586, 590, 591, 605
Sepide Khodaverdi	507, 509	Sayed Hamidreza Mortazavi	584, 587
Sepide Salehipour Bavarsad	274	Sayed H. Ghaffari	671
Setareh Haghighat	549, 999	Sayed Hojjat Hossaini	1006
Setareh Moazen	675	Sayed Hossein Hoseinifar	1042
setare Mamishi	804	Sayed Hossein Hosseini	1051
Sayed Abdolrahim Rezaee	383, 384, 792, 1035	Sayed Javad Hasheminia	728
Sayed Abdolrahim Rezaee	665	Sayed Javad Mowla	628
Sayed Abdolreza Mortazavi-Tabatabaei	67	Sayed. Latif Mousavi Gargari	379
seyed Abolhassan Shahzadeh Fazeli	735	Sayed Mahmoud Hashemi	34, 66, 69, 72, 99, 136, 187,
Sayed Abolhassan Shahzadeh Fazeli	722, 1017, 1032,		242, 246, 253, 303, 335, 336, 338, 412, 448, 450,
	1041		466, 514, 516, 520, 629, 668, 695, 987, 1002, 1044
Sayed Ahmad Hosseini	283	Sayed Mahmoud Hashemi,	66
Sayed Ali ghaffarikhalaftmohammadi	998	Sayed. Mahmoud Hashemi	513
Sayed Ali Javad Mousavi	286	Sayed Mahmoud Masiha Hashemi	207
Sayed Ali Mirhosseini	383, 792	Sayed Mahmud Hejazi	763

Seyed Masih E'temad aubi	929	S.Faghihzadeh	99
Seyed Masood Nabavi	1028	Shaban Alizadeh	88
Seyed Mehdi Ebrahimi	592	Shabnam Abtahi	654
Seyed Mehdi Sadat	769	Shabnam Ghazanfari-Sarabi	891
Seyed Mehran Modaresadegi	76	Shabnam Kazemian	805
Seyed Mehrdad Jalali	90	Shabnam Shahrokh	190, 470
Seyed Mohammad HadiBagheri Rostami	595	Shaden Kamhawi	268
Seyed mohammad hossein Musavi shenas	302	Shadi Abbasnia	50, 819
Seyed Mohammad Mehdi Peyghambari	900	Shadi Eghbalfard	500
seyed Mohammad Moazzeni	510	Shadi Kesmati	416
Seyed Mohammad Moazzeni	8, 225, 226, 501, 502, 505, 964	Shadi Sadri	189
Seyed Mojtaba Amiri	828, 829, 995	Shadi Tabibian	745
Seyed Morteza Seifati	740	Shadi Zamanian	382
Seyed Nezamodin Hosseini	233	Shafiei Reza	150
Seyed Nooreddin Faraji	63	Shaghayegh Askarian amiri	680, 682
Seyed Reza Mirhafez	140	Shaghayegh Derakhshani	117
Seyed Reza Mohebbi	805, 927	Shaghayegh Haghjoo Javanmard	641
Seyed Rohollah Arab	383, 792	Shaghayegh Pezeshki Naraghi	488
Seyed saadi hosseini	351	Shaghayegh Tajik	25
SeyedSaadihosseini	738	Shaghik Barani	78
Seyed Shahabeddin Mortazavi Jahromi	432, 433	Shahabeddin Mortazavi Jahromi	869
Seyed Shahabeddin Mortazavi Jahromib	367, 761	Shahab Shahgaldi	226, 510
Seyed Shapoor Reza Shojaei	736	Shahdad Khosropanah	441, 726, 901
Seyed Vahid Ganjiani	560	Shahin Bonakdar	449
Seyyed Abdolrahim Rezaee	390	Shahram Farhad	409, 884
Seyyed Amin Ayatollahi Mousavi	373	Shahram Nazarian	379, 984, 986
Seyyed Amir Yasin Ahmadi	394	Shahram Parvane	556
Seyyede Masoume Athari	578, 579	Shahram Samie	631
Seyyed Meysam	337	Shahram Shahabi	24
Seyyed Meysam Abtahi	618, 903	Shahrbanoo Rostami	656
Seyyed Meysam Abtahi Foroushani	366	Shahriar Alipour	826
Seyyed Meysam Abtahi Froushani	71, 430, 512, 632, 686, 846, 849, 850, 853, 854, 855, 866, 870, 888, 897, 898, 905, 907, 908, 910	Shahriar Zeighami	33
		ShahriarZeighami	295
		Shahriyar Dabiri	1038
		Shahrokh Navidpour	556
Seyyed Mohammad Moazzeni	334	Shahrooz Rasoul	243, 1000
Seyyed Shamsadin Athari	578, 579	Shahryar Hashemzadeh	43

Shahrzad Nojehdehi	514	Shohreh Nikoo	41, 94, 223, 224
Shaker Ardakani M	30, 613	Shokofeh Rahimpourkoldeh	978
Shakiba Y	30, 613	Shokoofeh Almasi	810
Shamila D Alipoor	121	Shokooh Salimi Moghadam	954, 962, 963
Shamila D. Alipoor	122, 398, 399	Shokrolah Farrokhi	392
ShamimSarhaddi	180	Shokrollah Farrokhi	23, 287, 288
Sharareh Bordbari	244	Sholeh Etehad asnaf	443
Sharifeh Khosravi	81	Siamak Asri Rezaei	198
Shayan Mostafaei	96, 132, 368	Siavash Abedi	19
Shayesteh Mehdinejadani	271, 320	Siavash Mashhour	512, 846
Shayesteh Z	808	Sibylle Schneider-Schaulies	117
Sheikh Sajjadih Mohammadreza	141	SimaBalouchi Anaraki	646
Shekoofeh Moghadam	940, 942, 1016, 1022, 1023	Sima Ghaffari	699
Shideh Khangholi	637	Sima Habibi	673
Shideh Mehrmofakham	484, 489, 491	Sima Habibzadeh	146, 268, 322, 425, 842
Shideh Mehr Mofakham	204	Sima Kazemi	766
Shila Jalalpour	377	Sima Rafati	145, 146, 268, 322, 420, 425, 842
Shima Bahador	715	Sima Shahmohammadi Farid	296, 355
Shima Bastaki	195, 930, 931, 932, 934	Simin Ahmadvand	28
Shima Bazaz	1016, 1022, 1023	Sina Abdolrahim Poor Heravi	341
Shima Farshbaf	408	Sina Alizadeh	769
Shima Khajouee Ravari	628	Sina Nabi Zadeh	1067
Shima Nahidi	425	Sina Taefeshokr	168, 307
Shima Shapoori	81, 525	Sina Vakili	902
Shima Sheikhi	1064	S.Meysam Abtahi frushani	909
shirin Assar	829, 995	Sogand Nayeri	567
Shirin Farivar	51, 55, 657, 687, 688	Sogol Mazhari	448
Shirin Farjadian	536, 630	Soheila Ajdary	395
Shiva Borzouei	526, 527, 977	Soheila Arefi	356
Shiva Farajpour	809	Soheila Golchin	45
Shiva Irani	769, 933	Soheila Rahmani	386
Shiva Khezri	909	Soheila Rohani	712
Shiva Mohammadi	1017, 1041	SoheilaRohani	710
Shiva Poursani	132	Soheyla Rohani	711
Sho'aub Dehghani	596	Soleimani S	678
Shohreh Fakhari	468, 634, 635, 924, 992	Solmaz khansalar	388
Shohreh Mohammadi	417, 840	Solmaz Khansalar	997

Solmaz Shirjang	31	Tahereh Jalilvand	506
Somaieh Kazemnejad	41, 223, 224	Tahereh Kalantari	70
Somaieh Nasereslami	387, 807, 811	Tahereh Mohammadi	227, 507, 509, 739
Somaye Aletaha	402	Tahereh Naji	94
Somayeh Abolhassani	644	Tahereh Soltantoyeh	40, 608
Somayeh Ahmadi afshar	161	Tahereh Taheri	268, 425
Somayeh Ahmadiafshar	435, 905	Tahereh Zeinali	692, 694
Somayeh Behjat Khatouni	979, 981	Taher Elmi	1059
Somayeh Ebrahimi Barough	534	Taher Nejdassattari	25
Somayeh Fallahnejad	682	Tajaddin akbarzadeh khiavi	718
Somayeh Ghotloo	177, 202, 720	Takayoki Yoshimoto	65
Somayeh Hamd Ghadareh	667	Talat MokhtariAzad	116
Somayeh Pourtalebi	644	Tanja Konijn	138
SomayehRezaeifard	295	Tannaz Bahadori	453
Somayeh Sadeghi	49	Tannaz Bahadori	608
Somayeh Shamlou	62	Taravat Bamdad	387, 807, 811
Somayeh Zarrati	1059	Taravat Talebi	53
Somaye Mami	364	Tavakoli Hajar	409, 884
Somaye pooranfar	760	Taybeh Tavakoli	644
Somayyeh Ghadimi	236	Tayebeh Mehdipour	456
Soodabeh Aghapour	744	Tayebeh Nowras	158
SoolmazHabibi	741	Tayebeh Radjabian	1064
Soraya Bilvaieh	22	Tayebeh Sohrabi	452, 980
Sorayya Salimi	130, 410, 832	ThureAdler	241
Soroor Erfanimanesh	782	Tina Shahani	53
Sousan Ghasemi	988	T. Kazemi	105
S. Rafati	270	T. Mousavi	133
S. Rostami	643	Tohid Kazemi	689, 697, 698, 699, 700
S. Sadati	993	Tooba Ghazafari	1066
S. Soudi	481	Tooba Ghazanfari	49, 98, 99, 363, 364, 754, 886, 1062,
Su Ling Li	307		1063, 1064, 1065
Sussan K. Ardestani	99	Tooba Ghazanfari1	1061
Suzan Amini	581	Toomaj Sabooteh	394
T		Toran Sepehrzad	559
Tabatabaei Yazdi SM	30, 613	T. Taheri	270
Taghi Golmohammadi	132	V	
Taghi Naserpour Farivar	431	Vahedeh Hosseini	271, 320

Vahid Baghdadi	360	Yahya Asemani	159
Vahid Bagheri	245, 495	Yalda Zarafshan	1029
Vahid Baniasadi	176	Yasaman Alimohammadi	925
Vahid Chaleshi	186, 190, 464, 465, 470, 473	Yasaman salimi	936
Vahid Jajarmi	629	Yasaman Taslimi	146, 268, 322, 425
VahidKhazeShahgoli	43	Yasamin Pahlavan	834
Vahid Kia	320	Yaser Azizi	864
Vahid Mashayekhi	146, 322	Yaser jafarikhataylou	952, 955
Vahid Mirzaee	377	Yaser Moghaddam	305
Vahid Mohammadi	854, 898	Yaser Panahi	215
Vahid R Haghaniat	762	Yaser Piralii	152
Vahid Salimi	116, 269	YasharAzari Key	307
Vahid Shayeghanejad	978	Yasin Bagheri	1058
Vahid Shaygannejad	523, 1036	Yasmin Vahidi	33, 690
VahidShaygannejad	728	Yasmin Zare	630
Vahid Yardel	1025	Yazdani Neyshabouri Sh	808
VahidYounesi	149	Yekaninejad S	102
Vajiheh Ostadi	76, 340	Yousef Shahali	281
Valiollah Mehrzad	641	Younes Pilehvar-Soltanahmad	167
Vazeh H	808	Yousef Daneshmand pour	401
Vaziri Nezamdoust F	1046	Yousef Mortazavi	266, 982
Vedad Alkhamis	666, 920	Yousef Tarverdizadeh	986
Versa Omraninava	797	Yousef yousefzadeh	939
Versa OmraniNava	796	Yousof Tarverdizade	1054
V. Eskandari1	993	Yunes Panahi	302
Vida Hashemi	195, 405, 930, 931, 932, 934	Z	
Vida Homayouni	35, 305, 480	Zahra Adnani Sadati	340
Vida Sheikh	526, 527, 977	Zahra Aghazadeh	433
W		Zahra akbari jonoush	61
Wytske J. Fokkens	23	Zahra Alizadeh Tabari	485, 487
X		Zahra Alizade Tabari	483
Xanthi N. Stachte	138	Zahra Amirghofran	157, 159, 292, 434, 436, 863, 997
Y		Zahra Arabi	437
Yadollah Farshi	745	Zahra Asadi	886, 925
YadollahOmidi	180	Zahra Asadzadeh	1047
Yaftiyan A	721	Zahra Asghari	583
Yaftiyan Atefeh	221, 222		

Zahra Badiei	980	Zahra Radaee	867
Zahra Basiri	1024	Zahra Rashid	475, 476
Zahra Elyasi Gorji	722, 1017, 1041	Zahra RezaeiYazdi	127
Zahra Faghih	11, 28, 29, 33, 295, 300, 690	Zahra Sadat Delbari	876, 889
Zahra farzaneh	388	Zahra Sadr	671
Zahra Farzaneh	118, 157	Zahra Samimi	406, 407
Zahra Fereydouni	444, 904	zahra Shaker	753
Zahra Gholami Mahmoodian	857	Zahra Shaker	661, 750
Zahra Hamidi Esfahani	344	Zahra Sharifzadeh	68, 477
Zahra Hashemi	395	Zahra Soleimani	906
Zahra Hooshyar	533	Zahra Taghipour	104
Zahra Hoseini Khah	858	Zahra Taghpour	437
Zahra Hosseinikhah	96	Zahra Telikani	526
Zahra Kanannejad	220	Zahra Zeinaly	337, 632, 686, 855
Zahra karami	571	Zakaria Vahabzadeh	911, 912
Zahra Kashani Khatib	88	Zakiyeh Azadi	25
Zahra Khademi	523	Zangeneh Mohamad Mahdi	966
Zahra khosrowpour	513	Zarrin Minucheher	230
Zahra Koolivand	359, 744	Zatollah Asemi	408
Zahra Madjd	518, 1020	Zeinab Barartabar	186
Zahra Matluobi	163	Zeinab Kavyani	635
Zahra Meshkat	140	Zeinab Mahmoodi	1050
Zahra Mirsanei	48, 673	Zeinab Rostamy	158
Zahra Moeinibakhsh	588	Zeinab Salehian	874
Zahra Mohtasham Amiri	806	Zeynab Baghernejadan	21
Zahra Mojtahedi	32, 297, 624	Zeynab Fallah	1060
Zahra Momeni	537	Zhima Akhavansales	739
Zahra Moradi	227, 507, 509, 739	ziba rahimi	357
Zahra Mozaheb	665	zivar zangeneh	916
Zahra nekooee zadeh	542	Zohair Mohammad Hasan	661, 750, 753
Zahra Nori Sabzikar	198	Zohre Eftekhari	276
Zahra Noroozbabaei	376	Zohreh Babaloo	32, 130, 167, 410, 826, 832
Zahra Norouzbabaie	445	Zohreh Bolandi	253, 987
Zahra Omid	1066	Zohreh Hajheidari	531, 1031
Zahra Pourmoghadam	218	Zohreh Hekmati	442
Zahra pourpak	20, 25	Zohreh Hesari	160
Zahra Pourpak	62, 281, 284	ZohrehHesari	563



Zohreh Koohini	19, 403	Zohre Poursina	382
Zohreh Mortezaei	490	Zohre Zaerieghane	362, 970
Zohreh Nasiri	464	Z. Rahmati	993
zohreh rahimi	357	Z. Rahnemoun	993
Zohreh Sharifi	234, 798	ZS.Hasan	585
Zohreh Vahidi	135, 254	Zuhair M. Hassan Saraf	65
ZohrehVahidi	127	Zuhair Mohammad Hassan	303, 668
Zohreh Zareighane	493, 751, 1069	Zuhair MohammadHassan	34
Zohreh Zarei Ghane	349, 1024	zuhair saraf hasan	936
Zohre Mohammadi	171	Zuhair Saraf Hasan	163
Zohre Najafi	274		



Actolukast
Montelukast 5 mg Chewable tablet, 10mg tablet

- Once daily dosing
- Rapid onset of action compared to corticosteroids
- Safe for chronic use
- Safe in pregnancy (Category B)
- An alternative to steroids in children unable to use inhaled corticosteroids or suffering from poor growth

ACTOVERCO

Together for a healthy future

No. 58, 8th St., Kooye Nasr (Gisha St.), Tehran, IR Iran, Postal Code: 1446863914, Tel: +98 (21) 4163 70 00
ACT.MKT.INT.07.1396.0231

www.actoverco.com

☒ **BRING OUT THE BEST OF YOU**



fastover

Tablets 30, 60, 120, 180 mg FEXOFENADINE

- Once daily administration, 24 hour relief
- Lowest sedative effect
- Multi-symptom relief
- Safe in children over 6 months

ACTOVERCO

Together for a healthy future

No. 58, 8th St., Kooye Nasr (Gisha St.), Tehran, IR Iran, Postal Code: 1446863914, Tel: +98 (21) 416 370 00
ACT.MKT.INT.07.1396.0230

www.actoverco.com

INCOMPARABLE IMPRESSION, more than your trust



Suprimun
Mycophenolate mofetil Tablet 500mg

First generic of mycophenolate mofetil with the most
extended clinical experience launched in 2003



EMA APPROVED

European Medicines Agency



Together for a healthy future.



www.actoverco.com

Let's control allergy all year round



LETIZEN®
Cetirizine Tablet 10mg

- Once daily
- Fast onset of action
- Suitable for all types of seasonal and perennial allergies
- Drug of choice for chronic urticaria
- A long-lasting effect

ACTOVERCO
Pharmaceuticals

KRKA

No. 58, 8th St., Kooye Nasr (Gisha St.), Tehran, IR Iran, Postal Code: 1446863914, Tel: +98 (21) 4163 70 00
ACT.MKT.INT.10.1396.0304
www.actoverco.com

CinnoRA®
Adalimumab

Back to the Beauty



AIVOPOX®
Paclitaxel

Alvocade
Bortezomib

kidiPeg®
Pegliposome

Lenoma®
Lenalidomide

Cabotax® 60
Cabazitaxel

PROGIO
Biotin 100 mg

Dalfyrol®
Fampridine 10 mg

Stivant™
Bevacizumab

Bendanib®
Bendamustine HCl

AryoTrust™
Trastuzumab

CinnaPoietin®
Erythropoietin beta

AlvoXal®
Oxaliplatin

Clasto®
Celecoxib

PegaGen®
Pegfilgrastim

AlvoPem®
Pemetrexed

ReciGen®
Interferon B-1a

nano-D
Vitamin D3

cinnafact®
Buserelin

MYELOID

Suprotac 1
Tacrolimus

CinnoMer® 40
Glarimer Acetate

Kidi Mab®
Adalimumab

CinnoRA®
Adalimumab

AlvoTere®
Docetaxel

AryoSeven®
Factor VII

CinnaTropin®
Somatropin

Leukaza®
Azacitidine

Cinnalf®
Folic Acid

CinnoTec®
Dimethyl Fumarate

Zadiva®
Dimethyl Fumarate

Zytux™
Rituximab

Alvinotec®
Irinotecan Hydrochloride

Zobis®
Zoledronic acid

Leudribine®
Chlorthalidone

CinnoVex®
INTERFERON BETA - 1b

Altebrel™
Etanercept

CinnoPar®
Teniparide

Xetabin®
Capecitabine

IdanThra®
Idarubicin (as hydrochloride)

IMMUNOREGULATION

Volume 2. Supplement 1. April 2018

ISSN: 2588-5200



ACTOVERCO
Together for a healthy future

شرکت سیناژن
CinnaGen

